ERRATA

Erratum to: Zero Preservation for a Family of Multivalued Functionals, and Applications to the Theory of Fixed Points and Coincidences

T. N. Fomenko^{a,*} and Yu. N. Zakharyan^{b,**}

Presented by Academician of the RAS S.V. Matveev May 23, 2020

Received May 23, 2020; revised May 23, 2020; accepted June 2, 2020

DOI: 10.1134/S1064562421010087

The sentence in p. 273 (left column, line 21 from below) should read as follows:

A multivalued functional $\Phi: Y \rightrightarrows \mathbb{R}_+$ is called (α, β) -search on Y if, for any point $x \in Y$ and any $c \in \Phi(x)$ and $r > 0$ such that $B(x, r) \subseteq Y$ and $c \leq (\alpha - \beta)r$, there exists a point $x' \in Y$ and a value $c' \in \Phi(x')$ for which $d(x, x') \leq \frac{1}{\alpha}c$ and $c' \leq \frac{\beta}{\alpha}c$.

The second condition in Theorem 3 should read as follows:

(ii) For any $x \in \bar{U}$ and any $y \in T_t(x)$ and $r > 0$ such that $B(x; r) \subseteq \bar{U}$ and $d(y, Q) \leq (\alpha - \beta)r$, there exists a point $x' \in \bar{U}$ and a value $y' \in T_t(x')$ for which $\rho(x, x') \leq \frac{1}{\alpha}d(y, Q)$ and $d(y', Q) \leq \frac{\beta}{\alpha}d(y, Q)$.

All the results of the paper are correct. We sincerely apologize to the editorial board and the readers for the admitted inaccuracies.

The original article can be found online at <https://doi.org/10.1134/S1064562420040225>

^a Faculty of Computational Mathematics and Cybernetics,
Lomonosov Moscow State University, Moscow, 119991 Russia

^b Faculty of Mechanics and Mathematics,
Lomonosov Moscow State University, Moscow, 119991 Russia

*e-mail: tn-fomenko@yandex.ru

**e-mail: yuri.zakharyan@gmail.com