
ERRATA

Erratum: Investigation of Nanocomposites Based on Hydrated Calcium Phosphates and Cellulose *Acetobacter xylinum*, *Fiz. Khim. Stekla*, 2008, vol. 32, no. 2, pp. 248–258 [Glass Phys. Chem. (Engl. transl.), 2008, vol. 32, no. 2, pp. 192–200]

**A. K. Khripunov^a, Yu. G. Baklagina^a, V. A. Sinyaev^b, E. S. Shustikova^b, B. A. Paramanov^c,
D. P. Romanov^d, R. Yu. Smyslov^a, and A. A. Tkachenko^e**

^a *Institute of Macromolecular Compounds, Russian Academy of Sciences, Bol'shoi pr. 31, St. Petersburg, 199004 Russia*

^b *Bekturov Institute of Chemical Sciences, Academy of Sciences of Kazakhstan,
ul. Ualikhanova 106, Almaty, 480100 Kazakhstan*

^c *St. Petersburg Medical Academy of Postgraduate Studies, ul. Kirochnaya 41, St. Petersburg, 191015 Russia*

^d *Grebenshchikov Institute of Silicate Chemistry, Russian Academy of Sciences,
nab. Makarova 2, St. Petersburg, 199034 Russia*

^e *St. Petersburg State University, Universitetskaya nab. 7/9, St. Petersburg, 199034 Russia*

Received June 29, 2007

DOI: 10.1134/S1087659609020175

Figure 1 contains a graphic misprint. The correct version of the figure is given below.




Fig. 1. Schematic diagram illustrating the biosynthesis of the cellulose *Acetobacter xylinum* gel film. (a) Brown's scheme [30] of the liner sequence (LS) of terminal complex (TC) triplets (TCTs) for *Acetobacter xylinum* and the hierarchy of the structure of microfibrillar ribbons (MFRs) composed of 10–100 nanocrystalline subunits (NCSs) with a cross section 13×7 nm in size according to Fink et al. [31]. (b) Electron microscope image of microfibrillar ribbons (K₁–K₂) formed by an *Acetobacter xylinum* bacterium during the synthesis for 3 h. (c) Model representation of cellulose macrochain packing. The (-110) and (110) faces forming nanocrystalline subunits and the primary OH groups are shown.