Erratum

A surprising method for polarising antiprotons

Th. Walcher^{1,2,a}, H. Arenhövel¹, K. Aulenbacher¹, R. Barday¹, and A. Jankowiak¹

¹ Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany

² Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati (Rome), Italy

Original article: Eur. Phys. J. A **34**, 447 (2007) DOI: 10.1140/epja/i2007-10462-x

Received: 12 December 2008

Published online: 5 January 2009 – © Società Italiana di Fisica / Springer-Verlag 2009

Triggered by a criticism of A.I. Milstein, S.G. Salnikov and V.M. Strakhovenko [1] we have searched for the reason of the dramatic difference of our triple-spin transfer cross-section in the scattering of spin-1/2 hadrons from leptons [2]. We found a numerical problem in our calculation of the spin-dependent cross-sections [3,4]. After the correction of this problem the cross-sections are smaller by about 14 orders of magnitude. The corrected figs. 1–4 of ref. [2] are repeated in this erratum (in the following page).

Though some features as the ratio $\langle S_2^- \rangle / \langle S_2 \rangle = 0.49$ are more favorable now, the huge reduction of the cross-section kills the proposal. In order to arrive at similar polarisation times and figures of merits as in ref. [2] an increase of the positron density of about 13 orders of magnitude had to be achieved. As a realistic increase four orders appear to be possible today [5].

However, there still remains a difference between our and the calculation of ref. [1]. Whereas we get $\sigma_{-\lambda,\lambda,+}^h \approx 30$ mbarn for $\lambda = -$ and 0.5 mbarn for $\lambda = +$ at $T_h = 1.7$ keV A.I. Milstein, S.G. Salnikov and V.M. Strakhovenko get $\sigma_{total} = \sigma_{-,+,+}^h + \sigma_{+,-,+}^h = 0.75$ mbarn. This difference may be due to their averaging over all relative initial directions of the hadron momentum with respect to the lepton spin.

References

1. A.I. Milstein, S.G. Salnikov, V.M. Strakhovenko, Nucl. Instrum. Methods B **266**, 3453 (2008) [arXiv:0802.3766 [hep-ph]].
2. Th. Walcher, H. Arenhövel, K. Aulenbacher, R. Barday, A. Jankowiak, Eur. Phys. J. A **34**, 447 (2007) [arXiv:0706.3765 [physics.acc-ph]].
3. H. Arenhövel, Eur. Phys. J. A **34**, 303 (2007) [arXiv:0706.3576 [nucl-th]].
4. H. Arenhövel, Erratum to Eur. Phys. J. A **34**, 303 (2007) (this issue).
5. K. Aulenbacher, H. Arenhoevel, R. Barday, A. Jankowiak, T. Walcher, AIP Conf. Proc. **980**, 170 (2008).

^a e-mail: walcher@kph.uni-mainz.de

Fig. 1. The integrated spin-flip cross-sections $\langle \sigma_{-\lambda, \lambda, +}^h \rangle$ ($\lambda = \pm$) for antiproton and proton electron scattering as a function of the proton lab kinetic energy T_h for $b = 10^{10}$ fm in PW and DW. The cross-section for proton and antiproton are identical in PW. It is easy to calculate the lepton kinetic energy in the proton rest system using the nonrelativistic relation $T_l = T_h(m_l/m_h)$.

Fig. 3. The integrated spin-flip cross-sections $\langle \sigma_{-\lambda, \lambda, +}^h \rangle$ ($\lambda = \pm$) for proton electron scattering as a function of the proton lab kinetic energy T_h for $b = 10^{10}$ fm in PW and DW.

Fig. 2. The integrated spin-flip cross-sections $\langle \sigma_{-\lambda, \lambda, +}^h \rangle$ ($\lambda = \pm$) for antiproton electron scattering as a function of the proton lab kinetic energy T_h for $b = 10^{10}$ fm in PW and DW.

Fig. 4. Ratio of $\langle S_2^- \rangle / \langle S_2 \rangle$ for proton electron scattering as a function of the proton lab kinetic energy T_h for the DW approximation.