Signature of a strong coupling with the continuum in $^{11}\text{Be} + ^{120}\text{Sn}$ scattering at the Coulomb barrier

L. Acosta^{1,a}, M.A.G. Álvarez^{2,5}, M.V. Andrés², M.J.G. Borge³, M. Cortés³, J.M. Espino², D. Galaviz³,
J. Gómez-Camacho^{2,5}, A. Maira³, I. Martel¹, A.M. Moro², I. Mukha², F. Pérez-Bernal¹, E. Reillo³, D. Rodríguez¹,
K. Rusek⁴, A.M. Sánchez-Benítez¹, and O. Tengblad³

¹ Departamento de Física Aplicada, Universidad de Huelva, E-21071 Huelva, Spain

² Departamento de Física Atómica Molecular y Nuclear, Universidad de Sevilla, E-41080 Sevilla, Spain

³ Instituto de Estructura de la Materia, CSIC, Madrid, E-28006 Madrid, Spain

⁴ The Andrzej Sołtan Institute for Nuclear Studies, 00-681 Warsaw, Poland

⁵ Centro Nacional de Aceleradores, Universidad de Sevilla-CSIC-Junta de Andalucía, E-41092 Sevilla, Spain

Original article: Eur. Phys. J. A **42**, 461 (2009) DOI: 10.1140/epja/i2009-10822-6

Received: 17 June 2009

Published online: 4 July 2009 – © Società Italiana di Fisica / Springer-Verlag 2009

Due to an error during writing the manuscript, the information given for both the beam and the energy used for normalizing the data are not correct.

The phrase “... of a beam of α particles at the energy of 12 MeV ...” in the last paragraph of sect. 2 should be replaced by “... of a beam of ^{12}C at the energy of 27 MeV ...”.

This change is not substantial for the study described and does not affect to the final conclusions presented in this work.

^a e-mail: luis.acosta@dfa.uhu.es