STAR inner tracking upgrade—a performance study

Jan Kapitán^a

for the STAR Collaboration

Nuclear Physics Institute ASCR, Rez/Prague, Czech Republic

Received: 27 September 2008 / Revised: 5 March 2009 / Published online: 20 March 2009
© Springer-Verlag / Società Italiana di Fisica 2009

Abstract Anisotropic flow measurements have demonstrated development of partonic collectivity in 200 GeV Au + Au collisions at RHIC. To understand the partonic EOS, thermalization must be addressed. Collective motion of heavy-flavor (c, b) quarks can be used to indicate the degree of thermalization of the light-flavor quarks (u, d, s). Measurement of heavy-flavor quark collectivity requires direct reconstruction of heavy-flavor hadrons in the low p_T region. Measurement of open charm spectra to high p_T can be used to investigate heavy-quark energy loss and medium properties. The Heavy Flavor Tracker (HFT), a proposed upgrade to the STAR experiment at midrapidity, will measure v_2 of open-charm hadrons to very low p_T by reconstructing their displaced decay vertices. The innermost part of the HFT is the PIXEL detector (made of two low mass monolithic active pixel sensor layers), which delivers a high precision position measurement close to the collision vertex. The Intermediate Silicon Tracker (IST), a 1-layer strip detector, is essential to improve hit identification in the PIXEL detector when running at full RHIC-II luminosity. Using a full GEANT simulation, open charm measurement capabilities of STAR with the HFT will be shown. Its performance in a broad p_T range will be demonstrated on v_2 ($p_T > 0.5$ GeV/c) and R_{CP} ($p_T < 10$ GeV/c) measurements of D^0 meson. Results of reconstruction of Λ_C baryon in heavy-ion collisions are presented.

1 Introduction

Produced by the initial hard scattering, heavy quarks (c, b) are an ideal tool to probe early stages of heavy ion collision [1]. They derive their mass from the Higgs field, and therefore are not modified by the surrounding QCD medium (they stay heavy even in the case of chiral symmetry restoration).

Previous studies have identified the development of partonic collectivity in heavy-ion collisions at RHIC [1], but they have not yet demonstrated thermalization of the created matter. The study of heavy quark collectivity may allow us to address this issue. Measurement of elliptic flow (v_2) of open charm hadrons to low transverse momentum (p_T) is of particular interest.

Suppression of high p_T hadron production at RHIC [2–4] is commonly thought to arise from partonic energy loss in dense matter due to induced gluon radiation [5, 6]. Radiative energy loss of heavy quarks is expected to be suppressed (dead cone effect) [7].

Most heavy-flavor measurements at RHIC use semileptonic decay modes of heavy-flavor hadrons, but lack of precise kinematical information about the parent hadron makes it difficult to study dynamics at low p_T [8]. Measurements of nuclear modification factor (R_{AA}) of heavy-flavor decay electrons at high p_T [9, 10] indicate a significant energy loss of heavy quarks. However, knowledge of the relative contributions of charm and bottom decays to electron spectra is crucial to interpret these results. Electron spectra from open charm hadron decays are also sensitive to relative Λ_C/D meson yield [11], due to different branching ratios of their inclusive electron decay channels.

In central Au + Au collisions at RHIC, a baryon/meson enhancement has been observed in the intermediate p_T region ($2 < p_T < 6$ GeV/c) [12–14]. These results are usually explained by a hadronization mechanism involving collective multi-parton coalescence rather than independent vacuum fragmentation. The success of the coalescence approach implies deconfinement and possibly thermalization of the light quarks prior to hadronization.

Since Λ_C is the lightest charmed baryon, and its mass is not far from that of the D^0 meson, a similar pattern of baryon/meson enhancement is expected in charm sector [15]. Λ_C/D^0 enhancement is also believed to be a signature of a strongly coupled quark-gluon plasma [16]. Therefore it would be very interesting to measure R_{CP} of Λ_C baryon and compare it to R_{CP} of D^0 mesons.

^a e-mail: kapitan@rf.rhic.bnl.gov

Direct reconstruction of open charm hadrons is necessary for these measurements. Given the large combinatorial backgrounds in heavy-ion collisions, topological reconstruction is needed to achieve reasonable signal significance. The Heavy Flavor Tracker (HFT), a proposed upgrade [17] to the STAR experiment, will enable measurement of open charm hadrons by reconstructing their displaced decay vertices.

2 Heavy flavor tracker design

The HFT detector consists of two subsystems: The PIXEL detector (2 layers) and Intermediate Silicon Tracker (IST, 1 layer). The midrapidity tracking system of STAR further includes the existing Silicon Strip Detector (SSD) and the large Time Projection Chamber (TPC). Compared to the previous version, the current design of the HFT has been optimised for lower mass and better hit resolving, however it has not been fully simulated yet. In the following, simulation results are presented for the previous design. The parameters of these are displayed in Table 1.

The PIXEL detector is made of low-mass monolithic active pixel sensors (MAPS, see Fig. 1) and enables high precision measurement close to the primary collision vertex, featuring 18 μm pixel pitch and a thickness of only 0.28% X_0 per layer. It is fabricated by CMOS technology.

The detector configuration is shown in Fig. 2. Inner layer sensors facing the beam pipe will improve STAR's di-lepton capability by rejecting e^+e^- pairs from photon conversions (beam pipe thickness being only $\approx 0.15\%$ X_0).

As there is a very large number of pixels, the read-out of the PIXEL detector is $\approx 200 \mu\text{s}$. Given the high luminosity projected for the future RHIC-II upgrade ($50 \cdot 10^{26} \text{ cm}^{-2} \text{ s}^{-1}$), the PIXEL detector will integrate over ≈ 10 minimum bias collisions. The IST detector is essential to improve hit identification at PIXEL2 layer in this pile-up environment. The IST detector consists of 1 layer of fast single-sided strip sensors.

Table 1 Hit position resolutions of SSD + HFT layers for the two design versions. IST2 (simulated design) has two layers (A, B) with crossed strips

layer	current design		simulated design	
	r (cm)	hit resol. ($r - \phi \times z$) ($\mu\text{m} \times \mu\text{m}$)	r (cm)	hit resol. ($r - \phi \times z$) ($\mu\text{m} \times \mu\text{m}$)
SSD	23	30×699	23	30×699
IST2-B	—	—	17	17×12000
IST2-A	—	—	17	12000×17
IST1	14	115×2900	12	17×6000
PIXEL2	8	5×5	7	9×9
PIXEL1	2.5	5×5	2.5	9×9

Fig. 1 MAPS principle of operation: electrons created in the epitaxial layer thermally diffuse towards low potential n-well region. A small contribution to the total signal also exists from electrons created in the p++ substrate

Fig. 2 One half of the PIXEL detector with its support structure. It will be mounted on one side, where also all the cables and cooling air comes from

3 Simulation results

HIJING central Au + Au events at $\sqrt{s_{NN}} = 200 \text{ GeV}$ with added D^0 and Λ_C were filtered through GEANT and STAR detector response simulators. To assess the impact of pile-up, pseudo-random hits were added to PIXEL1 and PIXEL2 layers, corresponding to a minimum bias (MB) collision rate of 80 kHz and a primary vertex diamond size $\sigma_{PV,Z} = 15 \text{ cm}$. This is upper estimate of pile-up, in fact current RHIC-II projections show smaller luminosity (corresponding to MB collision rate $\approx 50 \text{ kHz}$) and bigger $\sigma_{PV,Z}$.

The decay modes used for reconstruction were $D^0 \rightarrow K^- + \pi^+$ (B.R. 3.8%) and $\Lambda_C \rightarrow K^- + \pi^+ + p$ (B.R. 5.0%). A possibility of using a Λ_C decay through a resonant intermediate $\Lambda(1520)$ state has been investigated, and a significant improvement of S/B ratio could be expected. However, a full simulation has not been performed, and results shown hereafter are for the non-resonant decay.

As already mentioned in Sect. 2, the track impact parameter (pointing) resolution is mainly delivered by the PIXEL detector, which is consistent with results from full simulation shown in Fig. 3. For particle identification (PID) of daughter particles, $K - \pi$ separation for $p_T < 1.6$ GeV/c and $(K + \pi) - p$ separation for $p_T < 3.0$ GeV/c is expected using the MRPC TOF detector [18], with an efficiency $\approx 90\%$.

Reconstruction efficiencies for D^0 and Λ_C are shown in Fig. 4. The daughter tracks are required to be well reconstructed in the TPC and have correctly associated hits in both PIXEL layers. Efficiency for Λ_C is smaller due to the fact, that there are three daughter tracks with, on average, lower p_T than in the case of D^0 .

For Λ_C reconstruction, primary track combinatorial background is huge due to its very short decay length ($c\tau = 59.9 \mu\text{m}$) and three-body decay. To reduce this background, PID information of daughter tracks is required for

Fig. 3 Track impact parameter resolution. Symbols: full detector simulation, lines: calculation for PIXEL detector only [17]

Fig. 4 Efficiency of reconstructing daughter tracks of D^0 and Λ_C in $|\eta| < 1$. Note that no cuts to isolate signal from combinatorial background have been applied here

$p_T < 5$ GeV/c, limiting the acceptance. It is possible to reconstruct Λ_C at higher p_T , but this has not been studied in detail yet.

Given the overall uncertainty of $c\bar{c}$ cross section at RHIC, D^0 $dN/dy = 0.002$ per binary collision has been used for signal estimates, which is half of the value measured by the STAR Collaboration [19]. Λ_C/D^0 ratio 0.2 was assumed for the case of no enhancement [15]. The shape of p_T spectra for D^0 and Λ_C in central Au + Au collisions was assumed to be a power-law, with $\langle p_T \rangle = 1.0$ GeV/c and $n = 11$ (with C being a normalisation constant):

$$\frac{1}{p_T} \frac{dN}{dp_T} = C \cdot \left(1 + \frac{p_T}{p_0}\right)^{-n} \quad (1)$$

$$p_0 = \langle p_T \rangle \cdot \frac{n-3}{2} \quad (2)$$

To reduce the large combinatorial background, topological cuts have been applied: only tracks with Distance of Closest Approach to event Primary Vertex $DCA_{PV} > DCA_{PV, cut}$ have been used, where the cut is in the range 40–80 μm depending on transverse momentum of reconstructed D^0 (Λ_C). The DCA of daughter tracks to the decay vertex was required to be less than 2σ and D^0 (Λ_C) momentum was required to point back to the event primary vertex. Finally, a 2 (3) particle invariant mass cut has been applied.

Signal significance of $D^0 + \bar{D}^0$ from 100×10^6 central Au + Au collisions is shown in Fig. 5. To estimate background for D^0 (Λ_C) at different centralities, $(N_{part})^2$ ($(N_{part})^3$) scaling was used. For the signal, the same R_{CP} was assumed as measured by STAR for charged hadrons [3]. This gives a factor 4 higher yield at high p_T in peripheral (60–80%) collisions than given by N_{bin} scaling. Ratio of N_{bin} between central (0–10%) and peripheral (60–80%) collision is ≈ 44 .

Fig. 5 D^0 signal significance in central Au + Au events. The effect of pile-up is visible in low and medium p_T region

For measurements of $D^0 v_2$ and R_{CP} , statistics of 500×10^6 minimum bias Au + Au events was assumed. This is the amount of data the STAR detector can take in about 1 month of running (with DAQ rate 500 Hz and 40% beam up time).

Estimated statistical errors on v_2 (for $D^0 + \bar{D}^0$, $|\eta| < 1$) are shown in Fig. 6. For $p_T > 1.0$ GeV/c, the HFT will be able to distinguish between extreme elliptic flow scenarios in the coalescence picture in the first year of operation.

500×10^6 minimum bias events include 62.5×10^6 central and 125×10^6 peripheral events for R_{CP} measurement. D^0 signal significance (100×10^6 central events) in $9 < p_T < 10$ GeV/c bin is about 15σ (consistent with Fig. 5). The background is negligible here, so signal significance in 100×10^6 peripheral events will be $\sqrt{(44/4)} \approx 3.3$ times smaller. In a similar way, the calculation was done for lower p_T (taking into account the background here), and the esti-

mated statistical errors on $D^0 R_{CP}$ measurement are shown in Fig. 7. At highest p_T , the relative error is 20%, which is a good precision for this region.

For measurements of Λ_C , statistics of 2×10^9 minimum bias and 250×10^6 central triggered events was assumed, giving 500×10^6 central and 500×10^6 peripheral events used for R_{CP} . Λ_C reconstruction cuts were optimized in p_T range 2–5 GeV/c, estimated mass peak for the middle p_T bin is shown in Fig. 8. Estimated signal significance for this p_T bin in 500×10^6 central Au + Au collisions is 8σ .

To estimate errors of Λ_C/D^0 measurement, two scenarios have been used: 1. no enhancement, ratio equal to 0.2 and flat in p_T , 2. the same enhancement as Λ/K_S^0 [14]. D^0 has much larger yield and $c\tau$ than Λ_C , statistical errors coming from its measurement are therefore negligible. Λ_C/D^0 ratio in peripheral collisions is assumed flat (in rough agreement

Fig. 6 Two scenarios for D^0 elliptic flow (charm quark flow zero or equal to the flow of light quarks) and estimated statistical errors

Fig. 8 Expected invariant mass distribution of Λ_C , $3 < p_T < 4$ GeV/c, for 500×10^6 central events

Fig. 7 Estimated statistical errors for R_{CP} measurement of D^0 meson, in 500×10^6 minimum bias Au + Au events

Fig. 9 Expected errors on Λ_C/D^0 measurement, assuming 500×10^6 central and 500×10^6 peripheral collisions. Except for the lowest p_T bin, the errors are dominated by measurement of Λ_C in peripheral collisions

with Λ/K_S^0 [14]), with statistical errors given by rescaling signal and background from central to peripheral collisions, as described above.

Statistical errors on $R_{CP}(\Lambda_C)/R_{CP}(D^0)$ are shown in Fig. 9, where the two extreme cases can be well distinguished. Thus, baryon/meson ratio in charm sector will be measured with good precision, for the first time in heavy ion collisions.

4 Conclusions

The HFT detector will measure open charm hadrons over a broad p_T range, enabling precision study of charm collectivity, energy loss and baryon/meson ratios. These are important ingredients for a systematic study of the dense medium created in heavy-ion collisions at RHIC.

This will be achieved by using low mass MAPS sensors (PIXEL) together with a fast strip detector (IST), delivering high efficiency and ultimate pointing resolution at low p_T , even in the high luminosity environment of RHIC-II. Improved reconstruction efficiency is expected due to improved detector design and to possibly lower pile-up hit densities in RHIC-II luminosity, to be fully simulated in the future.

Acknowledgements This work was supported in part by the IRP AV0Z10480505, by GACR grant 202/07/0079 and by grant LC07048 of the Ministry of Education of the Czech Republic.

References

1. J. Adams et al., Nucl. Phys. A **757**, 102 (2005)
2. C. Adler et al., Phys. Rev. Lett. **89**, 202301 (2002)
3. J. Adams et al., Phys. Rev. Lett. **91**, 172302 (2003)
4. J. Adams et al., Phys. Rev. Lett. **91**, 072304 (2003)
5. R. Baier et al., Annu. Rev. Nucl. Part. Sci. **50**, 37 (2002)
6. M. Gyulassy et al., [arXiv:nucl-th/0302077](https://arxiv.org/abs/nucl-th/0302077)
7. Y.L. Dokshitzer, D.E. Kharzeev, Phys. Lett. B **519**, 199 (2001)
8. S. Batsouli, S. Kelly, M. Gyulassy, J.L. Nagle, Phys. Lett. B **557**, 26 (2003)
9. B.I. Abelev et al., Phys. Rev. Lett. **98**, 192301 (2007)
10. A. Adare et al., Phys. Rev. Lett. **98**, 172301 (2007)
11. P.R. Sorensen, X. Dong, Phys. Rev. C **74**, 024902 (2006)
12. B.I. Abelev et al., Phys. Rev. Lett. **97**, 152301 (2006)
13. B.I. Abelev et al., Phys. Lett. B **655**, 104 (2007)
14. J. Adams et al., [arXiv:nucl-ex/0601042](https://arxiv.org/abs/nucl-ex/0601042)
15. A. Andronic et al., Phys. Lett. B **571**, 36 (2003)
16. S.H. Lee et al., [arXiv:0709.3637v3](https://arxiv.org/abs/0709.3637v3) [nucl-th]
17. E. Anderssen et al., http://rnc.lbl.gov/hft/docs/hft_final_submission_version.pdf
18. O. Barannikova et al., Nucl. Instrum. Methods A **558**, 419–429 (2006)
19. B.I. Abelev et al., [arXiv:0805.0364v2](https://arxiv.org/abs/0805.0364v2) [nucl-ex]