Determinations of $|V_{cb}|$ and $|V_{ub}|$ from baryonic Λ_b decays

Y. K. Hsiao^{1,2}, C. Q. Geng^{1,2,3,a}

¹ School of Physics and Information Engineering, Shanxi Normal University, Linfen 041004, China

² Department of Physics, National Tsing Hua University, Hsinchu 300, Taiwan

³ Synergetic Innovation Center for Quantum Effects and Applications (SICQEA), Hunan Normal University, Changsha 410081, China

Received: 1 August 2017 / Accepted: 1 October 2017 / Published online: 26 October 2017

© The Author(s) 2017. This article is an open access publication

Abstract We present the first attempt to extract $|V_{cb}|$ from the $\Lambda_b \rightarrow \Lambda_c^+ \ell \bar{\nu}_\ell$ decay without relying on $|V_{ub}|$ inputs from the B meson decays. Meanwhile, the hadronic $\Lambda_b \rightarrow \Lambda_c M_{(c)}$ decays with $M = (\pi^-, K^-)$ and $M_c = (D^-, D_s^-)$ measured with high precisions are involved in the extraction. Explicitly, we find that $|V_{cb}| = (44.6 \pm 3.2) \times 10^{-3}$, agreeing with the value of $(42.11 \pm 0.74) \times 10^{-3}$ from the inclusive $B \rightarrow X_c \ell \bar{\nu}_\ell$ decays. Furthermore, based on the most recent ratio of $|V_{ub}|/|V_{cb}|$ from the exclusive modes, we obtain $|V_{ub}| = (4.3 \pm 0.4) \times 10^{-3}$, which is close to the value of $(4.49 \pm 0.24) \times 10^{-3}$ from the inclusive $B \rightarrow X_u \ell \bar{\nu}_\ell$ decays. We conclude that our determinations of $|V_{cb}|$ and $|V_{ub}|$ favor the corresponding inclusive extractions in the B decays.

1 Introduction

In the standard model (SM), the unitary 3×3 Cabibbo–Kobayashi–Maskawa (CKM) matrix elements present the coupling strengths of quark decays, with the unique physical weak phase for CP violation. Being unpredictable by the theory, the matrix elements as the free parameters need the extractions from the experimental data. Nonetheless, there exists a long-standing discrepancy between the determinations of $|V_{cb}|$ based on the exclusive $B \rightarrow D^{(*)} \ell \bar{\nu}_\ell$ and inclusive $B \rightarrow X_c \ell \bar{\nu}_\ell$ decays, given by [1–3]

$$\begin{aligned} |V_{cb}| &= (39.18 \pm 0.99) \times 10^{-3}, & (B \rightarrow D \ell \bar{\nu}_\ell) \\ |V_{cb}| &= (38.71 \pm 0.75) \times 10^{-3}, & (B \rightarrow D^* \ell \bar{\nu}_\ell) \\ |V_{cb}| &= (42.11 \pm 0.74) \times 10^{-3}. & (B \rightarrow X_c \ell \bar{\nu}_\ell) \end{aligned} \quad (1)$$

From the data in Eq. (1), we see that the deviations between the central values of the inclusive and exclusive decays are around $(2-3)\sigma$. For the resolution, the analysis in Ref. [4] suggests that the $B \rightarrow D^*$ transition form factors developed

by Caprini, Lellouch and Neubert (CLN) [5] may underestimate the uncertainty that is associated with the extraction of $|V_{cb}|$. Moreover, it has been recently pointed out that the theoretical parameterizations of the $B \rightarrow D^{(*)}$ transitions given by Boyd, Grinstein and Lebed (BGL) [6] are more flexible in reconciling the difference [7, 8]. Similar to the data for $|V_{cb}|$ in Eq. (1), there also exists a tension as regards the determination of $|V_{ub}|$ between the exclusive and inclusive B decays, which has drawn a lot of theoretical attentions to search for the solutions in the SM and beyond [9–14].

On the other hand, the baryonic Λ_b decays could provide some different theoretical inputs for the CKM matrix elements, which are able to ease the tensions between the exclusive and inclusive determinations. Indeed, to have an accurate determination of $|V_{ub}|/|V_{cb}|$ the LHCb Collaboration has carefully analyzed the ratio of [15],

$$\mathcal{R}_{ub} \equiv \frac{\mathcal{B}(\Lambda_b \rightarrow p \mu \bar{\nu})_{q^2 > 15 \text{ GeV}^2}}{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ \mu \bar{\nu})_{q^2 > 7 \text{ GeV}^2}} = \frac{|V_{ub}|^2/|V_{cb}|^2}{R_{FF}}, \quad (2)$$

where $\mathcal{B}$ denotes the branching fraction and q is the certain range of the integrated energies for the data collection. In Eq. (2), $\mathcal{R}_{ub}$ by relating $\mathcal{B}(\Lambda_b \rightarrow p \mu \bar{\nu})$ to $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \mu \bar{\nu})$ reduces the experimental uncertainties, while R_{FF} is the ratio of the $\Lambda_b \rightarrow \Lambda_c$ and $\Lambda_b \rightarrow p$ transition form factors, calculated by the lattice QCD (LQCD) model [16] with a smaller theoretical uncertainty.

In this work, we would like to first explore the possibility to determine $|V_{cb}|$ from the baryonic decays. In particular, we use the observed branching ratios of $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$ and $\Lambda_b \rightarrow \Lambda_c M_{(c)}$ with $\ell = e^-$ or μ^- , $M = (\pi^-, K^-)$ and $M_c = (D^-, D_s^-)$, which have never been used in the previous studies. The full energy-range measurement of the semileptonic $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$ decay is given by [17]

$$\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell) = \left(6.2_{-1.3}^{+1.4}\right) \times 10^{-2}, \quad (3)$$

^a e-mail: geng@phys.nthu.edu.tw

Fig. 1 Feynman diagrams depicted for **a** $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$ and **b** $\Lambda_b \rightarrow \Lambda_c M_{(c)}$

and the decay branching ratios of $\Lambda_b \rightarrow \Lambda_c^+ M_{(c)}$ are observed as [17]

$$\begin{aligned} \mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ \pi^-) &= (4.9 \pm 0.4) \times 10^{-3}, \\ \mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ K^-) &= (3.59 \pm 0.30) \times 10^{-4}, \\ \mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ D^-) &= (4.6 \pm 0.6) \times 10^{-4}, \\ \mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ D_s^-) &= (1.10 \pm 0.10) \times 10^{-2}. \end{aligned} \quad (4)$$

The above modes in Eq. (4) can be regarded to proceed through the $\Lambda_b \rightarrow \Lambda_c$ transition together with the recoiled mesons, such that the theoretical estimations give

$$\begin{aligned} \frac{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ \pi^-)}{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ K^-)} &\simeq R(M) \left(\frac{V_{ud}}{V_{us}} \right)^2 \left(\frac{f_\pi}{f_K} \right)^2 = 13.2, \\ \frac{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ D_s^-)}{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ D^-)} &\simeq R(M_c) \left(\frac{V_{cs}}{V_{cd}} \right)^2 \left(\frac{f_{D_s}}{f_D} \right)^2 = 25.1, \end{aligned} \quad (5)$$

where $f_{M_{(c)}}$ are the meson decay constants and $R(M_{(c)})$ are the rates to account for the mass differences from the phase spaces. Note that the ratios in Eq. (5) remarkably agree with $(13.6 \pm 1.6, 24.0 \pm 3.8)$ from the data in Eq. (4), respectively. This implies that the theoretical calculations of $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c M_{(c)})$ can be reliable to be involved in the fitting of $|V_{cb}|$. Moreover, it is proposed in the soft-collinear effective theory (SCE) [18] that the non-leptonic $\Lambda_b \rightarrow \Lambda_c \pi^-$ decay can be related to the semileptonic $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$ decays if the factorization works. This leads to the predictions of $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell) \approx 6 \times 10^{-2}$ and $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \pi^-) = 4.6 \times 10^{-3}$, which agree well with the data in Eqs. (3) and (4), respectively. Particularly, the data in Eq. (4) have the significances of $(8-12)\sigma$, which apparently benefit the precise determination of $|V_{cb}|$. As a result, the extraction of $|V_{cb}|$ from the data in Eqs. (3) and (4) can be an independent one besides those from the $B \rightarrow D^{(*)} \ell \bar{\nu}_\ell$ and $B \rightarrow X_c \ell \bar{\nu}_\ell$ decays. With the newly extracted $|V_{cb}|$ value, we will then be able to determine $|V_{ub}|$.

2 Formalism

As seen in Fig. 1, in terms of the effective Hamiltonian at quark level for the semileptonic $b \rightarrow c \ell \bar{\nu}_\ell$ and non-leptonic $b \rightarrow c \bar{\alpha} \beta$ ($\bar{\alpha} = \bar{u}(\bar{c})$ and $\beta = q = d, s$) transitions by the W -boson external emissions, the amplitudes of the $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$ and $\Lambda_b \rightarrow \Lambda_c M_{(c)}$ decays are found to be [16, 19]

$$\begin{aligned} \mathcal{A}(\Lambda_b \rightarrow \Lambda_c^+ \ell \bar{\nu}_\ell) &= \frac{G_F}{\sqrt{2}} V_{cb} \langle \Lambda_c^+ | \bar{c} \gamma_\mu (1 - \gamma_5) b | \Lambda_b \rangle \bar{\ell} \gamma^\mu (1 - \gamma_5) \nu_\ell, \\ \mathcal{A}(\Lambda_b \rightarrow \Lambda_c^+ M_{(c)}) &= \frac{G_F}{\sqrt{2}} V_{cb} V_{\alpha\beta}^* a_1^{M_{(c)}} i f_{M_{(c)}} q^\mu \langle \Lambda_c^+ | \bar{c} \gamma_\mu (1 - \gamma_5) b | \Lambda_b \rangle, \end{aligned} \quad (6)$$

where G_F is the Fermi constant, $V_{\alpha\beta} = V_{u(c)q}$ ($q = d, s$) for $M_{(c)} = \pi^- (D^-)$, $K^- (D_s^-)$, and the matrix elements of $\langle M_{(c)} | \bar{\beta} \gamma^\mu (1 - \gamma_5) \alpha | 0 \rangle = i f_{M_{(c)}} q^\mu$ have been used for the meson productions. The parameters $a_1^{M_{(c)}} = c_1^{\text{eff}} + c_2^{\text{eff}} / N_c^{\text{eff}}$ are derived by the generalized factorization approach with the effective Wilson coefficients $(c_1^{\text{eff}}, c_2^{\text{eff}}) = (1.168, -0.365)$ and color number N_c^{eff} [20].

In the helicity-based definition, the matrix elements of the $\Lambda_b \rightarrow \Lambda_c$ transition are given by [21]

$$\begin{aligned} \langle \Lambda_c | \bar{c} \gamma_\mu b | \Lambda_b \rangle &= \bar{u}_{\Lambda_c}(p', s') \left[f_0(q^2) (m_{\Lambda_b} - m_{\Lambda_c}) \frac{q^\mu}{q^2} \right. \\ &\quad \left. + f_+(q^2) \frac{m_{\Lambda_b} + m_{\Lambda_c}}{s_+} \right. \\ &\quad \left. \times \left(p^\mu + p'^\mu - \left(m_{\Lambda_b}^2 - m_{\Lambda_c}^2 \right) \frac{q^\mu}{q^2} \right) + f_\perp(q^2) \right. \\ &\quad \left. \times \left(\gamma^\mu - \frac{2m_{\Lambda_c}}{s_+} p^\mu - \frac{2m_{\Lambda_b}}{s_+} p'^\mu \right) \right] u_{\Lambda_b}(p, s), \\ \langle \Lambda_c | \bar{c} \gamma_\mu \gamma_5 b | \Lambda_b \rangle &= -\bar{u}_{\Lambda_c}(p', s') \gamma_5 \left[g_0(q^2) (m_{\Lambda_b} + m_{\Lambda_c}) \frac{q^\mu}{q^2} \right. \\ &\quad \left. + g_+(q^2) \frac{m_{\Lambda_b} - m_{\Lambda_c}}{s_-} \right] \end{aligned}$$

Table 1 Inputs of the experimental data

Branching ratios	Experimental data [17]
$10^2 \mathcal{B}(\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell)$	$6.2^{+1.4}_{-1.3}$
$10^3 \mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ \pi^-)$	4.9 ± 0.4
$10^4 \mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ K^-)$	3.6 ± 0.3
$10^4 \mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ D^-)$	4.6 ± 0.6
$10^2 \mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ D_s^-)$	1.1 ± 0.1

$$\begin{aligned} & \times \left(p^\mu + p'^\mu - \left(m_{\Lambda_b}^2 - m_{\Lambda_c}^2 \right) \frac{q^\mu}{q^2} \right) + g_\perp(q^2) \\ & \times \left(\gamma^\mu + \frac{2m_{\Lambda_c}}{s_-} p^\mu - \frac{2m_{\Lambda_b}}{s_-} p'^\mu \right) \Big] u_{\Lambda_b}(p, s), \end{aligned} \quad (7)$$

where $q = p - p'$, $s_\pm = (m_{\Lambda_b} \pm m_{\Lambda_c})^2 - q^2$, and $(f_0, f_+, f_\perp)$ and $(g_0, g_+, g_\perp)$ are form factors. The momentum dependences of $f = f_j$ and g_j ($j = 0, +, \perp$) are written as [16]

$$f(t) = \frac{1}{1 - t/(m_{\text{pole}}^f)^2} \sum_{n=0}^{n_{\text{max}}} a_n^f \left[\frac{\sqrt{t_+ - t_0} - \sqrt{t_+ - t_0}}{\sqrt{t_+ - t} + \sqrt{t_+ - t_0}} \right]^n, \quad (8)$$

where $(n_{\text{max}}, t_+, t_0) = (1, (m_{\text{pole}}^f)^2, (m_{\Lambda_b} - m_{\Lambda_c})^2)$ with m_{pole}^f representing the corresponding pole masses. In terms of the equations in Ref. [17], one is able to integrate over the variables of the phase spaces in the two-body and three-body decays for the decay widths.

The inclusions of $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c M_{(c)})$ can refrain the theoretical uncertainties for the fit of $|V_{cb}|$ as the $\Lambda_b \rightarrow \Lambda_c$ transition form factors in Eq. (7) are associated with both decays of $\Lambda_b \rightarrow \Lambda_c M_{(c)}$ and $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$. For the demonstration, we define

$$\begin{aligned} \mathcal{F}(\Lambda_b \rightarrow \Lambda_c) &= \frac{1}{|V_{cb}|^2} \\ &\times \int_{q^2} \frac{\hat{\tau}_{\Lambda_b}}{(2\pi)^3 32 m_{\Lambda_b}^3} \frac{d\Gamma(\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell)}{dq^2} dq^2, \end{aligned} \quad (9)$$

where $\hat{\tau}_{\Lambda_b} \equiv \tau_{\Lambda_b}/(6.582 \times 10^{-25})$, which leads to the total branching fraction of $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell) = |V_{cb}|^2 \mathcal{F}(\Lambda_b \rightarrow \Lambda_c)$ with $q^2 > (m_\ell + m_{\bar{\nu}})^2$.

3 Numerical results and discussions

For the numerical analysis, we perform the minimum χ^2 fit with $|V_{cb}|$ being a free parameter to be determined. The theoretical inputs for the CKM matrix elements, decay constants and $a_1^{M_{(c)}}$ are given by [17, 20]

$$\begin{aligned} (|V_{cd}|, |V_{cs}|) &= (0.220 \pm 0.005, 0.995 \pm 0.016), \\ (|V_{ud}|, |V_{us}|) &= (0.97417 \pm 0.00021, 0.2248 \pm 0.0006), \\ (f_\pi, f_K) &= (130.2 \pm 1.7, 155.6 \pm 0.4) \text{ MeV}, \\ (f_D, f_{D_s}) &= (203.7 \pm 4.7, 257.8 \pm 4.1) \text{ MeV}, \\ a_1^{M_{(c)}} &= 1.05 \pm 0.12. \end{aligned} \quad (10)$$

The experimental values in Eqs. (3) and (4) are accounted to be five data points, summarized in Table 1. Note that the $\Lambda_b \rightarrow \Lambda_c$ form factors in Eq. (8) are adopted from Ref. [16]. For the minimum χ^2 fit, we have

$$\chi^2 = \sum_{i=1}^6 \left(\frac{\mathcal{B}_{\text{th}}^i - \mathcal{B}_{\text{ex}}^i}{\sigma_{\text{ex}}^i} \right)^2 + \sum_j \left(\frac{\mathcal{F}_{\text{fit}}^j - \mathcal{F}_{\text{th}}^j}{\sigma_{\mathcal{F}_{\text{th}}}^j} \right)^2, \quad (11)$$

where $\mathcal{B}_{\text{th}}^i$ and $\mathcal{B}_{\text{ex}}^i$ stand for the branching ratios from the amplitudes in Eq. (6) and experimental data inputs in Table 1, σ_{ex}^i illustrate the experimental errors, $i = 1, 2, \dots, 5$ correspond to the five observed decay modes involved in the fit, and $\mathcal{F}_{\text{th}}^j$ and $\sigma_{\mathcal{F}_{\text{th}}}^j$ denote the central (best fit) value and error for the j th theoretical input in Eq. (10) along with the $\Lambda_b \rightarrow \Lambda_{(c)}$ transition form factors in Eq. (8), respectively. Subsequently, we obtain

$$\begin{aligned} |V_{cb}| &= (44.6 \pm 3.2) \times 10^{-3}, \\ (a_1^M, a_1^{M_c}) &= (1.19 \pm 0.08, 0.87 \pm 0.06), \\ \mathcal{F}(\Lambda_b \rightarrow \Lambda_c) &= 31.18 \pm 0.64, \\ \chi^2/\text{d.o.f} &= 7.3/4 \simeq 1.8, \end{aligned} \quad (12)$$

where d.o.f. represents the degrees of freedom. In Eq. (12), our fit with $\chi^2/\text{d.o.f} \sim 1.8$ indicates a good one, and the value of $|V_{cb}|$ clearly agrees with the inclusive result in Eq. (1) from $B \rightarrow X_c \ell \bar{\nu}_\ell$. Note that this is the first attempt to extract $|V_{cb}|$ from the $\Lambda_b \rightarrow \Lambda_c^+ \ell \bar{\nu}_\ell$ decay without relying on $|V_{ub}|$ inputs from the B meson decays. Clearly, it leaves the room for the more future accurate measurements of $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$ to improve the fit. On the other hand, like the $|V_{ub}|$ extraction from $\mathcal{R}_{ub}$ in Eq. (2) using the input of $|V_{cb}|$ from the B decays, one can inversely use the value of $|V_{ub}|$ from the $B \rightarrow \pi \ell \bar{\nu}$ decay to extract that $|V_{cb}| = (44 \pm 3) \times 10^{-3}$ [22], where the central value and error are in agreement with our results in Eq. (12).

For the initial input of $a_1^{M_{(c)}}$ in Eq. (10), the central value with $N_c^{\text{eff}} = 3$ corresponds to the factorizable contributions, and the error with N_c^{eff} ranging from 2 to ∞ is to accommodate the non-factorizable effects. In the two-body B meson decays, this empirical estimation has been demonstrated to agree with the sub-leading calculations like the QCD factorization [23–25], where the similar a_1 is modified by the detailed calculation for the non-factorizable effects but with the number to fall into the allowed range with N_c^{eff} from 2 to ∞ in the generalized version of the factorization [20]. While the theory depends on the data to

Table 2 The fitting results for the different scenarios in comparison with the experimental data

	$\chi^2/\text{d.o.f}$	$ V_{cb} \times 10^3$	$ V_{ub} \times 10^3$	$\mathcal{F}(\Lambda_b \rightarrow \Lambda_c)$
S1	1.8	44.6 ± 3.2	4.3 ± 0.4	31.18 ± 0.64
S2	2.3	45.1 ± 4.5	4.3 ± 0.5	31.23 ± 0.64
$B \rightarrow D\ell\bar{\nu}_\ell$ [1]		39.18 ± 0.99		
$B \rightarrow D^*\ell\bar{\nu}_\ell$ [1]		38.71 ± 0.75		
$B \rightarrow X_c\ell\bar{\nu}_\ell$ [2]		42.11 ± 0.74		
$B \rightarrow \pi\ell\bar{\nu}_\ell$ [17]			3.72 ± 0.19	
$B \rightarrow X_u\ell\bar{\nu}_\ell$ [17]			4.49 ± 0.24	
LQCD [16]				31.19 ± 1.33

test its validity, in the Λ_b decays the generalized factorization has been used to explain the data well [19,26–29]. However, it is clear that the sub-leading calculations for the non-factorizable effects have not been well developed in $\Lambda_b \rightarrow \Lambda_c M_{(c)}$ yet. Nonetheless, in the soft-collinear effective theory [18], the tree-level diagram as in Fig. 1b is estimated to dominate over the non-factorizable ones, where the predictions of $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \pi^-) = 4.6 \times 10^{-3}$ and $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \ell^- \bar{\nu}_\ell) \approx 6 \times 10^{-2}$ are consistent with the values of 4.66×10^{-3} and 6.43×10^{-2} from our fit, respectively. The determinations of $a_1^{M_{(c)}}$ in Eq. (12) that give around $\mathcal{O}(10\%)$ deviations from the initial inputs in Eq. (10) implies that the decay is insensitive to the non-factorizable effects. Note that, with $a_1 = 1$ as the input, together with the form factors computed by the light-front quark model, one obtains that $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \pi^-, \Lambda_c K^-) = (4.22, 3.41) \times 10^{-3}$ in the heavy quark limit [30], which are close to our results of $(4.66, 3.63) \times 10^{-3}$, respectively. We remark that based on the QCD factorization and the same $\Lambda_b \rightarrow \Lambda_c$ transition form factors as our inputs [16], the next-next-leading-order calculations result in [31] $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c \pi^-) = 2.85 \pm 0.54 \times 10^{-3}$ and $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c K^-) = 2.21 \pm 0.40 \times 10^{-4}$, which are all lower than the data in Eq. (4).

By bringing the newly extracted $|V_{cb}|$ into $|V_{ub}|/|V_{cb}| = 0.095 \pm 0.005$ as the most recent ratio from the exclusive B and Λ_b decays [17], we get

$$|V_{ub}| = (4.3 \pm 0.4) \times 10^{-3}, \quad (13)$$

which is consistent with the inclusive result of $(4.49 \pm 0.24) \times 10^{-3}$ from $B \rightarrow X_u \ell \bar{\nu}_\ell$ [17] but different from the exclusive one of $(3.72 \pm 0.19) \times 10^{-3}$ from $B \rightarrow \pi \ell \bar{\nu}_\ell$ [17]. To compare our fitting results with different data inputs, we set two scenarios:

$$\begin{aligned} (S1) \quad & \mathcal{B}(\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell) + \mathcal{B}(\Lambda_b \rightarrow \Lambda_c M_{(c)}), \\ (S2) \quad & \mathcal{B}(\Lambda_b \rightarrow \Lambda_c M_{(c)}), \end{aligned} \quad (14)$$

where S1 corresponds to the fitting shown in Eqs. (12) and (13), which gives the lowest uncertainty for $|V_{cb}|$ along with the best value of $\chi^2/\text{d.o.f}$. In Table 2, we summarize our results as well as the data from the B decays. In Eq. (12)

and Table 2, the result of $\mathcal{F}(\Lambda_b \rightarrow \Lambda_c) = 31.18 \pm 0.64$ compared to the value of 31.19 ± 1.33 as the initial theoretical input in the LQCD models presents that the fit with $\mathcal{B}(\Lambda_b \rightarrow \Lambda_c M_{(c)})$ is indeed able to reduce the theoretical error from 1.33 to 0.64 for the extraction of $|V_{cb}|$. Besides, the nearly identical central values of $\mathcal{F}(\Lambda_b \rightarrow \Lambda_c)$ hints that the form factors claimed to be less uncertain at $q^2 > 7 \text{ GeV}^2$ [16] are also suitable for $\Lambda_b \rightarrow \Lambda_c M_{(c)}$ at a small q^2 . Note that $a_1^{M_{(c)}}$ slightly deviated from the initial value by the fitting.

Finally, we remark that if we take the $\Lambda_b \rightarrow \Lambda_c$ transition form factors in the forms of $f(q^2) = f(0)/[1 - a(q^2/m_{\Lambda_b}) + b(q^2/m_{\Lambda_b})^2]$, adopted from Refs. [32], we obtain a lower value of $|V_{cb}| = (35.7 \pm 4.0) \times 10^{-3}$ with $\chi^2/\text{d.o.f} = 0.8$ by keeping the five data points in Table 1 in the fitting. In this case, less flexible inputs for the form factors with only central values for $(f(0), a, b)$ are used, leading to the result similar to the extraction from the exclusive $B \rightarrow D^{(*)} \ell \bar{\nu}_\ell$ decays with the CLN parameterization for the $B \rightarrow D^{(*)}$ transitions [5].

4 Conclusions

In sum, since the extractions of $|V_{cb}|$ showed the $(2-3)\sigma$ deviations between the exclusive $B \rightarrow D^{(*)} \ell \bar{\nu}_\ell$ and inclusive $B \rightarrow X_c \ell \bar{\nu}_\ell$ decays, we have performed an independent determination from the exclusive $\Lambda_b \rightarrow \Lambda_c \ell \bar{\nu}_\ell$ and $\Lambda_b \rightarrow \Lambda_c M_{(c)}$ decays. We have obtained $|V_{cb}| = (44.6 \pm 3.2) \times 10^{-3}$ to agree with the extraction in $B \rightarrow X_c \ell \bar{\nu}_\ell$. With the improved ratio of $|V_{ub}|/|V_{cb}|$ from the LHCb and PDG, we have derived that $|V_{ub}| = (4.3 \pm 0.4) \times 10^{-3}$, which is close to the result from the inclusive decays of $B \rightarrow X_u \ell \bar{\nu}_\ell$. Consequently, we have demonstrated that our extractions of $|V_{cb}|$ and $|V_{ub}|$ support those from the inclusive B decays. Clearly, the reliabilities for the determinations of $|V_{cb,ub}|$ from the exclusive B decays should be reexamined.

Acknowledgements We would like to thank Dr. Juergen Koerner for bringing Ref. [18] to our attention and for useful discussions. This work was supported in part by National Center for Theoretical Sci-

ences, MoST (MoST-104-2112-M-007-003-MY3), and National Science Foundation of China (11675030).

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

Funded by SCOAP³.

References

1. Y. Amhis et al., Averages of b -hadron, c -hadron, and τ -lepton properties as of summer 2016. [arXiv:1612.07233](https://arxiv.org/abs/1612.07233) [hep-ex]
2. P. Gambino, K.J. Healey, S. Turczyk, Phys. Lett. B **763**, 60 (2016)
3. A. Alberti, P. Gambino, K.J. Healey, S. Nandi, Phys. Rev. Lett. **114**, 061802 (2015)
4. F.U. Bernlochner, Z. Ligeti, M. Papucci, D.J. Robinson, Phys. Rev. D **95**, 115008 (2017)
5. I. Caprini, L. Lellouch, M. Neubert, Nucl. Phys. B **530**, 153 (1998)
6. C.G. Boyd, B. Grinstein, R.F. Lebed, Phys. Rev. D **56**, 6895 (1997)
7. D. Bigi, P. Gambino, S. Schacht, Phys. Lett. B **769**, 441 (2017)
8. B. Grinstein, A. Kobach, Phys. Lett. B **771**, 359 (2017)
9. A. Crivellin, Phys. Rev. D **81**, 031301 (2010)
10. A.J. Buras, K. Gemmler, G. Isidori, Nucl. Phys. B **843**, 107 (2011)
11. X.W. Kang, B. Kubis, C. Hanhart, U.G. Meißner, Phys. Rev. D **89**, 053015 (2014)
12. T. Feldmann, B. Müller, D. van Dyk, Phys. Rev. D **92**, 034013 (2015)
13. A. Crivellin, S. Pokorski, Phys. Rev. Lett. **114**, 011802 (2015)
14. Y.K. Hsiao, C.Q. Geng, Phys. Lett. B **755**, 418 (2016)
15. R. Aaij et al. [LHCb Collaboration], Nat. Phys. **11**, 743 (2015)
16. W. Detmold, C. Lehner, S. Meinel, Phys. Rev. D **92**, 034503 (2015)
17. C. Patrignani et al. [Particle Data Group], Chin. Phys. C **40**, 100001 (2016)
18. A.K. Leibovich, Z. Ligeti, I.W. Stewart, M.B. Wise, Phys. Lett. B **586**, 337 (2004)
19. Y.K. Hsiao, C.Q. Geng, Phys. Rev. D **91**, 116007 (2015)
20. A. Ali, G. Kramer, C.D. Lu, Phys. Rev. D **58**, 094009 (1998)
21. T. Feldmann, M.W.Y. Yip, Phys. Rev. D **85**, 014035 (2012). Erratum: [Phys. Rev. D **86**, 079901 (2012)]
22. M. Wingate, PoS CKM **2016**, 043 (2017). [arXiv:1704.00673](https://arxiv.org/abs/1704.00673) [hep-lat]
23. M. Beneke, G. Buchalla, M. Neubert, C.T. Sachrajda, Phys. Rev. Lett. **83**, 1914 (1999)
24. M. Beneke, G. Buchalla, M. Neubert, C.T. Sachrajda, Nucl. Phys. B **591**, 313 (2000)
25. M. Beneke, G. Buchalla, M. Neubert, C.T. Sachrajda, Nucl. Phys. B **606**, 245 (2001)
26. Y.K. Hsiao, P.Y. Lin, C.C. Lih, C.Q. Geng, Phys. Rev. D **92**, 114013 (2015)
27. Y.K. Hsiao, P.Y. Lin, L.W. Luo, C.Q. Geng, Phys. Lett. B **751**, 127 (2015)
28. C.Q. Geng, Y.K. Hsiao, Y.H. Lin, Y. Yu, Eur. Phys. J. C **76**, 399 (2016)
29. Y.K. Hsiao, Y. Yao, C.Q. Geng, Phys. Rev. D **95**, 093001 (2017)
30. H.W. Ke, X.Q. Li, Z.T. Wei, Phys. Rev. D **77**, 014020 (2008)
31. T. Huber, S. Krnkl, X.Q. Li, JHEP **1609**, 112 (2016)
32. T. Gutsche, M.A. Ivanov, J.G. Körner, V.E. Lyubovitskij, P. Santorelli, N. Habel, Phys. Rev. D **91**, 074001 (2015). Erratum: [Phys. Rev. D **91**, 119907 (2015)]