

Two Ramond–Ramond corrections to type II supergravity via field-theory amplitude

Hamid R. Bakhtiarizadeh^a 

Department of Physics, Sirjan University of Technology, Sirjan, Iran

Received: 18 August 2017 / Accepted: 16 November 2017
© The Author(s) 2017. This article is an open access publication

Abstract Motivated by the standard form of the string-theory amplitude, we calculate the field-theory amplitude to complete the higher-derivative terms in type II supergravity theories in their conventional form. We derive explicitly the $O(\alpha'^3)$ interactions for the RR (Ramond–Ramond) fields with graviton, B-field and dilaton in the low-energy effective action of type II superstrings. We check our results by comparison with previous work that has been done by the other methods, and we find exact agreement.

1 Introduction

Higher-derivative corrections to string theories and M-theory are importantly studied in various ways: string amplitude [1–7], non-linear sigma model [8,9], superfield and noether's method [10–17], duality completion [18–21], and so on. Each of these approaches has been employed in different formalisms such as the RNS (Ramond-Neveu-Schwarz) [22–24], GS (Green-Schwarz) [17,20,26,27] and pure-spinor [28] formalisms to determine the higher-order terms.

For many purposes, it is enough to use only the lowest-order terms in the theory, but there are some situations for which one must go beyond the lowest-order supergravity actions and higher-order corrections play an important role. For example, the origin of induced Einstein–Hilbert terms are traced to R^4 couplings in ten dimensions [29,30]. Furthermore, in black-hole and black-brane physics, considering higher-derivative terms leads to modifications of the thermodynamics [31–33]. Another important set of such applications can be found in the context of the gauge/gravity duality [34–36]. There are a vast number of applications of higher-derivative terms, but the mentioned examples are sufficient to illustrate the importance of having a good understanding of higher-derivative terms [22].

It is well known that the low-energy effective action of superstring theory is given by supergravity describing only the interactions of massless modes in the string-theory spectrum. This can be shown explicitly by calculating the field-theory amplitudes of massless states [25]. One must go beyond this low-energy limit to capture truly stringy behavior. Significant information as regards string and M-theory can be extracted from the corresponding low-energy effective actions, in particular once one considers corrections that go beyond the leading order. Subleading terms in type II effective actions start at order α'^3 or eight-derivative level. In order to determine the structure of the higher-derivative terms, we will calculate the scattering amplitude of the massless states.

As mentioned above, the scattering amplitudes of massless states in superstring theory include corrections to their corresponding low-energy effective actions. These terms contain α' corrections to the supergravity which arise due to the length of the fundamental string ℓ_s and string coupling constant g_s which correspond to string quantum corrections in spacetime. In type IIA superstring theory, one of those corrections first obtained at the tree level from the four-graviton scattering amplitude as well as from the σ -model beta function approach can be written

$$\mathcal{L}_{R^4} = -\frac{e^{-2\phi}}{2\kappa_{10}^2} \frac{\ell_s^6}{2^8 \cdot 4!} \zeta(3) \left(t_8 t_8 R^4 + \frac{1}{4 \cdot 2!} \epsilon_{10} \epsilon_{10} R^4 \right). \quad (1.1)$$

Here $2\kappa_{10}^2 = (2\pi)^7 \ell_s^8 g_s^2$, t_8 is a product of four Kronecker deltas, and $t_8 t_8 R^4$ denotes an abbreviation of a product of two t_8 tensors and four Riemann curvature tensors.

The contents of this paper is as follows: In Sect. 2, we briefly review type II supergravity theories and fix our conventions and notations. In Sect. 3, we establish the formalism and explain the procedure needed to find the necessary Feynman rules for the processes we want to compute. By employing these rules we then calculate the tree-level four-point amplitude for two RR–two NSNS scattering to find higher-derivative corrections to type II supergravities in their

^a e-mail: bakhtiarizadeh@sirjantech.ac.ir

conventional form up to the R^4 terms. We restrict our attention to different external NSNS states, namely the metric, antisymmetric tensor and dilaton. Finally, we compare our results with previous work and find exact agreement.

2 Brief review of type II supergravity

We begin by reviewing the low-energy effective actions of both type II superstring theories [37]. These are supergravity theories that describe interactions of the massless fields in the string-theory spectrum. The action in the Einstein frame for the type IIA supergravity is given by

$$S_{\text{IIA}} = \frac{1}{2\kappa^2} \int d^{10}x \sqrt{-G} \left(R - \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \frac{1}{2} e^{-\Phi} |H|^2 - \frac{1}{2} \sum_{n=2,4} e^{\frac{5-n}{2}\Phi} |\tilde{F}^{(n)}|^2 \right) - \frac{1}{4\kappa^2} \int B \wedge dC^{(3)} \wedge dC^{(3)}, \quad (2.1)$$

where R is the scalar curvature, Φ is the dilaton field and H is the B-field strength $H = dB$. The RR field strengths are defined in terms of RR potentials as $\tilde{F}^{(2)} = dC^{(1)}$ and $\tilde{F}^{(4)} = dC^{(3)} - H \wedge C^{(1)}$. The above action is the reduction of 11-dimensional supergravity on manifold $R^{1,9} \times S^1$. By a Weyl rescaling of the metric, this action can be transformed to the Einstein frame in which the Einstein term has the conventional form¹. It also contains three distinct types of terms. The first three terms in the first line involve NSNS fields, which are common to both type II superstring theories. The last term contains RR fields and as a whole are named kinetic terms. The second line is called the Chern–Simons term.

In the type IIB supergravity, the presence of the self-dual five-form introduces a significant complication for writing down a classical action for type IIB supergravity. In other words, it is hard to formulate the action in a manifestly covariant form. There are several different ways of dealing with this problem. One of them is to find an action which reproduces the super-symmetric equations of motion when the self-duality condition is imposed by hand. The type IIB supergravity action in the Einstein frame is given as

$$S_{\text{IIB}} = \frac{1}{2\kappa^2} \int d^{10}x \sqrt{-G} \left(R - \frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \frac{1}{2} e^{-\Phi} |H|^2 - \frac{1}{2\alpha} \sum_{n=1,3,5} e^{\frac{5-n}{2}\Phi} |\tilde{F}^{(n)}|^2 \right) - \frac{1}{4\kappa^2} \int H \wedge dC^{(2)} \wedge C^{(4)}, \quad (2.2)$$

where $\alpha = 1$ for $n = 1, 3$ and $\alpha = 2$ for $n = 5$. The RR field strengths here are defined as $\tilde{F}^{(1)} = dC^{(0)}$, $\tilde{F}^{(3)} = dC^{(2)} - HC^{(0)}$ and

$$\tilde{F}^{(5)} = dC^{(4)} - \frac{1}{2} C^{(2)} \wedge H + \frac{1}{2} B \wedge dC^{(2)}, \quad (2.3)$$

respectively. The self-duality condition, which must be imposed in the equations of motion by hand, is given by

$$\tilde{F}^{(5)} = \star \tilde{F}^{(5)}. \quad (2.4)$$

This condition has to be imposed as an extra constraint, manually. Without that, one cannot find any consistency between the field-theory and string-theory S-matrix elements.

3 Field-theory amplitude

Having had the supergravity actions, one can easily read different vertices and propagators and accordingly calculate the Feynman amplitude of two RR–two NSNS massless states. To that end, suppose that the massless fields are small perturbations around the flat background, *i.e.*,

$$\begin{aligned} g_{\mu\nu} &= \eta_{\mu\nu} + 2\kappa h_{\mu\nu}; \\ B^{(2)} &= 2\kappa b^{(2)}; \\ \Phi &= \phi_0 + \sqrt{2}\kappa\phi; \\ C^{(n)} &= 2\kappa c^{(n)}, \end{aligned} \quad (3.1)$$

where κ is the gravitational coupling constant. Substituting these perturbation expansions into the supergravity actions (2.1) and (2.2), and expanding them up to cubic and quartic powers in κ , one obtains three-point and four-point interactions, respectively. The amplitude of two RR–two NSNS fields typically has the following form:

$$\begin{aligned} \mathcal{A} &= \mathcal{A}_s + \mathcal{A}_u + \mathcal{A}_t + \mathcal{A}_c \\ &= \frac{\mathcal{A}_s}{s} + \frac{\mathcal{A}_u}{u} + \frac{\mathcal{A}_t}{t} + \mathcal{A}_c \\ &= \frac{1}{sut} (tu\mathcal{A}_s + st\mathcal{A}_u + su\mathcal{A}_t + sut\mathcal{A}_c), \end{aligned} \quad (3.2)$$

where \mathcal{A}_s , \mathcal{A}_u and \mathcal{A}_t are the amplitudes in s , u and t channels, respectively, and \mathcal{A}_c is the contribution of contact terms which must be added in order for the total scattering amplitude to be gauge invariant. The Mandelstam variables are defined by $s = -4\alpha' k_1 \cdot k_2$, $u = -4\alpha' k_1 \cdot k_3$, $t = -4\alpha' k_2 \cdot k_3$, and they satisfy the identity $s + t + u = 0$.

What motivates us to write the field-theory amplitude in the above form originates from the general form of string-theory amplitude. The structure of the string-theory amplitude consists of the well-known Gamma functions in terms of Mandelstam variables multiplied by a kinematic factor \mathcal{K} [1, 38]. To find the couplings which are reproduced by the amplitude on the string-theory side, it is convenient to expand the Gamma functions at low energy, *i.e.*,

¹ The Einstein frame is related to a string frame by $g_{\mu\nu}^E = e^{-\Phi/2} g_{\mu\nu}$.

$$\frac{\Gamma(-e^{-\phi_0/2}s/8)\Gamma(-e^{-\phi_0/2}t/8)\Gamma(-e^{-\phi_0/2}u/8)}{\Gamma(1+e^{-\phi_0/2}s/8)\Gamma(1+e^{-\phi_0/2}t/8)\Gamma(1+e^{-\phi_0/2}u/8)} \sim -\frac{2^9 e^{3\phi_0/2}}{stu} - 2\zeta(3) + \dots \tag{3.3}$$

where the expansion parameter is α' , and ϕ_0 is the constant dilaton background. The right-hand side in the above expression gives the low-energy limit of the amplitude. The first term on that side just corresponds to amplitudes of exchanging two RR–two NSNS massless fields in the s, t and u channels as well as contact terms in type II supergravity [25]. The second term contains a Riemann zeta function $\zeta(3)$, which is irrational and contributes as a stringy correction to the supergravity. In this way, it is basically possible to derive higher-derivative corrections from string scattering amplitudes.

By comparing the amplitude (3.2) with the leading term of string-theory amplitude, one finds the following relation between the field-theory amplitude and the string-theory kinematic factor:

$$\mathcal{K} = -2^{-9} e^{-3\phi_0/2} (tu\mathcal{A}_s + st\mathcal{A}_u + su\mathcal{A}_t + sut\mathcal{A}_c). \tag{3.4}$$

Multiplying this factor by $-2\zeta(3)$ and transforming it to the spacetime, one then finds the couplings of two RR–two NSNS fields at order α'^3 .

4 $(\partial F^{(n)})^2 R^2$ couplings

After explaining our strategy in the previous section, we are now in a position to find various couplings. For the first one, we are going to find the couplings containing two RR n -form field strengths with $n = 1, 2, 3, 4, 5$ and two Riemann curvatures. To this end, we consider the elastic scattering process of two RR fields into two gravitons. When the two RR forms have the same rank, the supergravity actions (2.1) and (2.2) dictate that the massless poles in the s and u channels and the contact terms are given by the following expressions:

$$\begin{aligned} A_s &= \left(\tilde{V}_{F_1^{(n)} F_2^{(n)} h}\right)^{\mu\nu} \left(\tilde{G}_h\right)_{\mu\nu,\lambda\rho} \left(\tilde{V}_{hh_3h_4}\right)^{\lambda\rho}, \\ A_u &= \left(\tilde{V}_{F_1^{(n)} h_3 C^{(n-1)}}\right)^{\mu_1 \dots \mu_{n-1}} \left(\tilde{G}_{C^{(n-1)}}\right)_{\mu_1 \dots \mu_{n-1}}^{\nu_1 \dots \nu_{n-1}} \\ &\quad \times \left(\tilde{V}_{C^{(n-1)} F_2^{(n)} h_4}\right)_{\nu_1 \dots \nu_{n-1}}, \\ A_c &= \tilde{V}_{F_1^{(n)} F_2^{(n)} h_3 h_4}. \end{aligned} \tag{4.1}$$

The t channel amplitude A_t can now be obtained from A_u by permuting the particles lines 3 and 4.

For the process we want to calculate, we need the two RR–one graviton, three gravitons, one RR–one graviton–one RR and two RR–two gravitons vertex functions. These vertices can be obtained from the supergravity actions (2.1) and (2.2) by expanding to cubic and quartic powers in κ , respectively. Fortunately, these vertex functions as well as

the graviton and RR propagators (in Feynman-like gauge) have been derived previously in the literature [23–25, 39, 40] up to an overall factor:

$$\left(\tilde{V}_{F_1^{(n)} F_2^{(n)} h}\right)^{\mu\nu} = \frac{i\kappa}{n!} \left(2n F_1^{(\mu}{}_{\mu_1 \dots \mu_{n-1}} F_2^{\nu)\mu_1 \dots \mu_{n-1}} - \eta^{\mu\nu} F_{1\mu_1 \dots \mu_n} F_2^{\mu_1 \dots \mu_n}\right), \tag{4.2}$$

$$\left(\tilde{G}_h\right)_{\mu\nu,\lambda\rho} = -\frac{i}{2k^2} \left(\eta_{\mu\lambda}\eta_{\nu\rho} + \eta_{\mu\rho}\eta_{\nu\lambda} - \frac{1}{4}\eta_{\mu\nu}\eta_{\lambda\rho}\right), \tag{4.3}$$

$$\begin{aligned} \left(\tilde{V}_{h_1 h_2 h}\right)^{\lambda\rho} &= -2i\kappa \left[\left(\frac{3}{2}k_1 \cdot k_2 \eta^{\lambda\rho} + k_1^{(\lambda} k_2^{\rho)} - k^\lambda k^\rho\right) \text{Tr}(h_1 \cdot h_2) \right. \\ &\quad - k_{1 \cdot} h_2 \cdot h_1 \cdot k_2 \eta^{\lambda\rho} + 2k_2^{(\lambda} \eta_2^{\rho)} \cdot h_1 \cdot k_2 \\ &\quad + 2k_1^{(\lambda} \eta_1^{\rho)} \cdot h_2 \cdot k_1 + 2k_1 \cdot h_2^{(\lambda} \eta_1^{\rho)} \cdot k_2 \\ &\quad \left. - k_1 \cdot k_2 (h_1^\lambda \cdot h_2^\rho + h_2^\lambda \cdot h_1^\rho) \right. \\ &\quad \left. - k_1 \cdot h_2 \cdot k_1 h_1^{\lambda\rho} - k_2 \cdot h_1 \cdot k_2 h_2^{\lambda\rho} \right], \end{aligned} \tag{4.4}$$

$$\left(\tilde{V}_{F_1^{(n)} h_2 C^{(n-1)}}\right)^{\mu_1 \dots \mu_{n-1}} = \frac{2i\kappa}{(n-1)!} F_{1\mu} [{}^{\mu_1 \dots \mu_{n-1}} h_2^{\mu}{}_{\nu}] k_\nu, \tag{4.5}$$

$$\left(\tilde{G}_{C^{(n)}}\right)_{\mu_1 \dots \mu_n}^{\nu_1 \dots \nu_n} = -\frac{in!}{k^2} \eta_{[\mu_1}{}^{\nu_1} \eta_{\mu_2}{}^{\nu_2} \dots \eta_{\mu_n]}{}^{\nu_n}, \tag{4.6}$$

$$\begin{aligned} \tilde{V}_{F_1^{(n)} F_2^{(n)} h_3 h_4} &= \frac{4i\kappa^2}{n!} \left[F_{1\mu_1 \dots \mu_n} F_2^{\mu_1 \dots \mu_n} \left(-\frac{1}{4} h_{3\lambda\rho} h_4^{\lambda\rho} \right. \right. \\ &\quad \left. \left. + \frac{1}{8} h_3^\lambda h_4^\rho h_{\lambda\rho} \right) \right. \\ &\quad - \frac{1}{2}(n+1) F_{1\mu_2 \dots \mu_n} F_{2\nu}{}^{\mu_2 \dots \mu_n} h_3^\lambda h_4^{\mu\nu} \\ &\quad + \frac{1}{2} n(n-1) F_{1\mu\lambda\mu_3 \dots \mu_n} F_{2\nu\rho}{}^{\mu_3 \dots \mu_n} h_3^{\mu\nu} h_4^{\lambda\rho} \\ &\quad \left. + F_1^{\mu\mu_2\mu_3 \dots \mu_n} F_2^{\nu}{}_{\mu_2\mu_3 \dots \mu_n} h_3^\lambda h_4^{\nu\lambda} \right] \\ &\quad + (3 \leftrightarrow 4). \end{aligned} \tag{4.7}$$

Our notation is such that $\text{Tr}(h_1 \cdot h_2) = h_1^{\mu\nu} h_{2\nu\mu}$, $k_1 \cdot h_2^\lambda = k_{1\rho} h_2^{\rho\lambda}$ and $h_1^\rho \cdot k_2 = h_1^{\rho\lambda} k_{2\lambda}$. The bracket (parenthesis) notation over indices means antisymmetrization (symmetrization) with a factor 1/2. k_ν in (4.5) denotes the momentum of internal leg *i.e.*, $k_\nu = k_{1\nu} + k_{2\nu}$.

The next step is to write the amplitude in terms of independent variables. This imposes all symmetries including mono-term symmetries (antisymmetry property of RR potentials and symmetry property of graviton polarizations) as well as multi-term symmetries (the Bianchi identities governing on the RR field strengths and Riemann tensors). In doing so, we have first written the RR field strengths and Riemann tensors in terms of RR potentials and graviton polarizations, respectively. Then we have manipulated with some of the terms. For instance, momentum conservation $\sum_{i=1}^4 k_i = 0$ as well as the mass-shell and on-shell relations for momenta, $k_i^2 = 0$ and $k_i \cdot \epsilon_i = 0$, are imposed. We have also rewritten some

terms using the relation $s + t + u = 0$. Furthermore, we have applied the tracelessness property of external graviton polarizations.

After doing the above steps to canonicalize the amplitude, the final result is simplified in terms of some Mandelstam variables, momenta, two RR potentials and two graviton polarizations. When $n = 1$, the result is simplified as

$$\begin{aligned} & \frac{1}{2} \left(s^2 u^2 h_3^{\alpha\beta} h_{4\alpha\beta} + 2su^3 h_3^{\alpha\beta} h_{4\alpha\beta} + u^4 h_3^{\alpha\beta} h_{4\alpha\beta} \right. \\ & + 16su^2 h_{3\beta}^\gamma h_{4\alpha\gamma} k_1^\alpha k_2^\beta + 16u^3 h_{3\beta}^\gamma h_{4\alpha\gamma} k_1^\alpha k_2^\beta \\ & - 16s^2 u h_{3\alpha}^\gamma h_{4\beta\gamma} k_1^\alpha k_2^\beta - 32su^2 h_{3\alpha}^\gamma h_{4\beta\gamma} k_1^\alpha k_2^\beta \\ & - 16u^3 h_{3\alpha}^\gamma h_{4\beta\gamma} k_1^\alpha k_2^\beta + 64u^2 h_{3\gamma\delta} h_{4\alpha\beta} k_1^\alpha k_2^\beta k_2^\delta \\ & - 128su h_{3\alpha\gamma} h_{4\beta\delta} k_1^\alpha k_2^\beta k_2^\delta - 128u^2 h_{3\alpha\gamma} h_{4\beta\delta} k_1^\alpha k_2^\beta k_2^\delta \\ & + 64s^2 h_{3\alpha\beta} h_{4\gamma\delta} k_1^\alpha k_2^\beta k_2^\delta + 128su h_{3\alpha\beta} h_{4\gamma\delta} k_1^\alpha k_2^\beta k_2^\delta \\ & \left. + 64u^2 h_{3\alpha\beta} h_{4\gamma\delta} k_1^\alpha k_2^\beta k_2^\delta \right), \end{aligned} \tag{4.8}$$

where the RR polarizations have been written in terms of momenta. To rewrite the amplitude (4.8) in terms of RR field strengths and Riemann tensors, we consider all possible contractions between two RR 1-form field strengths with a partial derivative acting on each one and two Riemann tensors. Then we make an ansatz of all these contractions by multiplying them by unknown constant coefficients:

$$\begin{aligned} & C_1 F^{a,b} F^{c,d} R_{ac}{}^{ef} R_{bdef} + C_2 F^{a,b} F^{c,d} R_{ac}{}^{ef} R_{bdef} \\ & + C_3 F^{a,b} F^{c,d} R_a{}^e{}_c{}^f R_{bdef} + C_4 F^{a,b} F^{c,d} R_a{}^e{}_c{}^f R_{bfde} \\ & + C_5 F_a{}^c F^{a,b} R_b{}^{def} R_{cdef} + C_6 F^{a,b} F^{c,d} R_a{}^e{}_b{}^f R_{cedf} \\ & + C_7 F_a{}^c F^{a,b} R_b{}^{def} R_{cedf} + C_8 F_{a,b} F^{a,b} R_{cedf} R^{cedf} \\ & + C_9 F_{a,b} F^{a,b} R_{cedf} R^{cedf}. \end{aligned} \tag{4.9}$$

By comparing Eq. (4.8) with the result obtained from writing (4.9) in terms of independent variables with the same steps as mentioned above, one obtains some algebraic equations among unknown constant coefficients:

$$\begin{aligned} & \{2C_5 + C_6 + C_7 + 8C_8 + 4C_9 = 0, \\ & 2C_5 + 2C_6 + C_7 = 0, \\ & -512 + 2C_1 + C_2 + C_3 + 2C_6 = 0, \\ & 2C_1 + C_2 - C_4 + 2C_5 + 3C_6 + C_7 = 0, \\ & 2C_1 + C_2 - C_4 + C_6 = 0, \\ & 2C_1 + C_2 - C_4 + 2C_5 + C_6 + C_7 + 16C_8 + 8C_9 = 0, \\ & 512 + 4C_1 + 2C_2 - C_3 - 3C_4 + C_6 = 0, \\ & 2C_1 + C_2 - C_4 + 8C_8 + 4C_9 = 0, \\ & -1024 + 2C_1 + C_2 + 2C_3 + C_4 + 2C_5 + 5C_6 \\ & + C_7 = 0, -512 + C_3 + C_4 + C_6 = 0, \\ & -1024 + 4C_1 + 2C_2 + 2C_3 - 2C_5 + C_6 - C_7 \\ & + 8C_8 + 4C_9 = 0, \\ & 6C_1 + 3C_2 - 3C_4 - 2C_5 + C_6 - C_7 = 0, \end{aligned}$$

$$\begin{aligned} & -512 + C_3 + C_4 + 2C_5 + 3C_6 + C_7 = 0, \\ & -4096 + 16C_1 + 8C_2 + 8C_3 - 8C_5 + 8C_6 - 4C_7 = 0, \\ & 512 + 2C_1 + C_2 - C_3 - 2C_4 = 0, \\ & 2048 + 2C_1 + C_2 - 4C_3 - 5C_4 + 2C_5 - C_6 + C_7 = 0 \}. \end{aligned} \tag{4.10}$$

After solving the above 18 equations, simultaneously, one finds the unknown constant coefficient C_3 to be 512. The other coefficients are free parameters which can be set to zero.

Having available the unknown coefficients and substituting them into Eq. (4.9), we can obtain the couplings between two RR 1-form field strengths and two Riemann tensors². The effective action, which reproduces the corresponding amplitude, takes the following form in the string frame:

$$\begin{aligned} S_{(\partial F^{(1)})^2 R^2} &= \frac{\gamma}{3 \cdot 2^4 \kappa^2} \\ &\times \int d^{10}x e^{2\phi_0} \sqrt{-G} F^{a,b} F^{c,d} R_a{}^e{}_c{}^f R_{bdef}, \end{aligned} \tag{4.11}$$

where $\gamma = \alpha'^3 \zeta(3)/2^5$, and comma refers to a partial derivative with respect to the index afterwards. Note that, to acquire the standard sphere-level dilaton factor $e^{-2\phi_0}$ in the string frame, it is convenient to normalize the RR potentials and graviton polarizations with a factor e^{ϕ_0} . The normalization of RR fields in the above action is consistent with the supergravities (2.1) and (2.2). Moreover, the couplings of two RR 2-form field strengths and two Riemann curvatures appear in the following effective action:

$$\begin{aligned} S_{(\partial F^{(2)})^2 R^2} &= \frac{\gamma}{3 \cdot 2^5 \kappa^2} \\ &\times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(2F_a{}^{d,e} F^{ab,c} R_b{}^f{}_d{}^g R_{cfeg} \right. \\ &+ F^{ab,c} F_c{}^{d,e} R_{ad}{}^{fg} R_{befg} \\ &+ 2F^{ab,c} F^{de,f} R_{acf}{}^g R_{bdeg} \\ &\left. + 2F^{ab,c} F^{de,f} R_{afd}{}^g R_{bgce} \right). \end{aligned} \tag{4.12}$$

Hence, there is no unique way of writing down the expressions (4.11) and (4.12) because of different presentations of tensor polynomials using the symmetries of the individual tensors, but we believe the above forms are the most economical one. The number of terms has been reduced as much as possible using the algorithm introduced in [24]. We also find the couplings between two RR 3-form field strengths and two Riemann curvatures:

² The calculations in this study have been carried out by the Mathematica package xAct [41].

$$\begin{aligned}
 S_{(\partial F^{(3)})^2 R^2} &= \frac{\gamma}{3^2 \cdot 2^7 \kappa^2} \\
 &\times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(8F_{abc}{}^{,e} F^{abc,d} R_d{}^{fgh} R_{efgh} \right. \\
 &- 3F_{abd,c} F^{abc,d} R_{efgh} R^{efgh} \\
 &+ 12F_{ab}{}^{,e,f} F^{abc,d} R_{cf}{}^{gh} R_{dehg} \\
 &- 12F_{ab}{}^{,e,f} F^{abc,d} R_{ce}{}^{gh} R_{dfgh} \\
 &+ 24F_{ab}{}^{,e,f} F^{abc,d} R_{cd}{}^{gh} R_{efgh} \\
 &- 12F_{ab}{}^{,e,f} F^{abc,d} R_{cd}{}^{gh} R_{efgh} \\
 &+ 24F_{ad}{}^e{}_{,b} F^{abc,d} R_c{}^{fgh} R_{efgh} \\
 &- 24F_a{}^{ef,g} F^{abc,d} R_{bde}{}^h R_{cfhg} \\
 &+ 48F_a{}^{ef,g} F^{abc,d} R_{bce}{}^h R_{dfgh} \\
 &+ 24F_a{}^{ef,g} F^{abc,d} R_{bce}{}^h R_{dfgh} \\
 &- 24F_a{}^{ef,g} F^{abc,d} R_{bcd}{}^h R_{efgh} \\
 &- 12F^{abc,d} F_d{}^{ef}{}_{,a} R_{be}{}^{gh} R_{cfhg} \\
 &\left. + 12F^{abc,d} F_d{}^{ef}{}_{,a} R_{bc}{}^{gh} R_{efgh} \right). \tag{4.13}
 \end{aligned}$$

Furthermore, our calculations show that the couplings which include two RR 4-form field strengths and two Riemann curvatures have the following form:

$$\begin{aligned}
 S_{(\partial F^{(4)})^2 R^2} &= -\frac{\gamma}{3^2 \cdot 2^6 \kappa^2} \\
 &\times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(3F_{abef}{}^{,i} F_{cdgi,h} R^{abcd} R^{efgh} \right. \\
 &- 9F_{abe}{}^i{}_{,f} F_{cdgi,h} R^{abcd} R^{efgh} \\
 &- 9F_{abe}{}^i{}_{,g} F_{cdhi,f} R^{abcd} R^{efgh} \\
 &+ 6F_{be}{}^{gh,i} F_{dfgi,h} R_a{}^e{}_{,c}{}^f R^{abcd} \\
 &+ 3F_{ce}{}^{gh,i} F_{dfgi,h} R_{ab}{}^{ef} R^{abcd} \\
 &+ 6F_{be}{}^{hi}{}_{,c} F_{dfhi,g} R_a{}^{efg} R^{abcd} \\
 &- 6F_{cf}{}^{hi}{}_{,b} F_{dghi,e} R_a{}^{efg} R^{abcd} \\
 &- 2F_b{}^{ghi}{}_{,e} F_{dghi,f} R_a{}^e{}_{,c}{}^f R^{abcd} \\
 &+ 3F_{ce}{}^{gh,i} F_{dghi,f} R_{ab}{}^{ef} R^{abcd} \\
 &- 24F_{bce}{}^{hi,i} F_{fghi,d} R_a{}^{efg} R^{abcd} \\
 &- 6F_{bc}{}^{hi}{}_{,e} F_{fghi,d} R_a{}^{efg} R^{abcd} \\
 &\left. + 9F_{be}{}^{hi}{}_{,c} F_{fghi,d} R_a{}^{efg} R^{abcd} \right). \tag{4.14}
 \end{aligned}$$

We are now going to find the couplings containing two RR 5-form field strengths and two Riemann tensors. The amplitude in this case is somewhat different, because of the presence of the 5-form field strength $F^{(5)}$, which is self-dual. This condition should be imposed manually as an extra condition in physical quantities like the equations of motion as well as S-matrix.

We expect that imposing the self-duality condition $F^{(5)} \rightarrow (F^{(5)} + \star F^{(5)})/2$, which is equivalent to (2.4), in the couplings with structure $(\partial F^{(5)})^2 R^2$ not only gives the correct overall factor but also results into another coupling with

structure $\epsilon_{10}(\partial F^{(5)})^2 R^2$. This coupling is not calculated here, because the number of indices is too large, which makes it difficult to calculate all possible contractions for this structure. However, the result for the couplings between two RR 5-form field strengths and two Riemann curvatures is

$$\begin{aligned}
 S_{(\partial F^{(5)})^2 R^2} &= \frac{\gamma}{3^2 \cdot 2^8 \kappa^2} \\
 &\times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(18F_{abe}{}^{ij}{}_{,g} F_{cdfij,h} R^{abcd} R^{efgh} \right. \\
 &- 3F_{abef}{}^{i,j} F_{cdghi,j} R^{abcd} R^{efgh} \\
 &- 12F_{be}{}^{ghi,j} F_{dfghj,i} R_a{}^e{}_{,c}{}^f R^{abcd} \\
 &- 6F_{ce}{}^{ghi,j} F_{dfghj,i} R_{ab}{}^{ef} R^{abcd} \\
 &- 8F_{be}{}^{hij}{}_{,c} F_{dfhij,g} R_a{}^{efg} R^{abcd} \\
 &+ 8F_{cf}{}^{hij}{}_{,b} F_{dghij,e} R_a{}^{efg} R^{abcd} \\
 &+ 2F_b{}^{ghij}{}_{,e} F_{dghij,f} R_a{}^e{}_{,c}{}^f R^{abcd} \\
 &+ 4F_{ce}{}^{ghi,j} F_{dghij,f} R_{ab}{}^{ef} R^{abcd} \\
 &- 48F_{bce}{}^{hi,j} F_{fghij,d} R_a{}^{efg} R^{abcd} \\
 &+ 8F_{bc}{}^{hij}{}_{,e} F_{fghij,d} R_a{}^{efg} R^{abcd} \\
 &\left. - 12F_{be}{}^{hij}{}_{,c} F_{fghij,d} R_a{}^{efg} R^{abcd} \right), \tag{4.15}
 \end{aligned}$$

where the product of two Levi-Civita tensors implicitly has been replaced by the generalized Kronecker delta according to: $\epsilon^{m_1 \dots m_d} \epsilon_{n_1 \dots n_d} = -\delta_{[n_1}{}^{m_1} \dots \delta_{n_d]}{}^{m_d}$.

In the next sections we follow a similar approach to obtaining the other couplings. The details of the calculations are omitted for the sake of brevity.

5 $(\partial F^{(n)})^2(\partial\phi)^2$ couplings

In this section, the couplings with structure $(\partial F^{(n)})^2(\partial\phi)^2$ will be calculated. Similar to the previous section, there are also five types of couplings here, *i.e.*, with $n = 1, 2, 3, 4, 5$. When we have two RR field strengths, in which the RR forms have the same rank, and two dilatons, the actions (2.1) and (2.2) indicate that the amplitudes in the s and u channels and the contact terms become

$$\begin{aligned}
 A_s &= \left(\tilde{V}_{F_1^{(n)} F_2^{(n)} h} \right)^{\mu\nu} \left(\tilde{G}_h \right)_{\mu\nu,\lambda\rho} \left(\tilde{V}_{h\phi_3\phi_4} \right)^{\lambda\rho}, \\
 A_u &= \left(\tilde{V}_{F_1^{(n)} \phi_3 C^{(n-1)}} \right)^{\mu_1 \dots \mu_{n-1}} \\
 &\quad \times \left(\tilde{G}_{C^{(n-1)}} \right)_{\mu_1 \dots \mu_{n-1}}{}^{\nu_1 \dots \nu_{n-1}} \\
 &\quad \times \left(\tilde{V}_{C^{(n-1)} F_2^{(n)} \phi_4} \right)_{\nu_1 \dots \nu_{n-1}}, \\
 A_c &= \tilde{V}_{F_1^{(n)} F_2^{(n)} \phi_3 \phi_4}, \tag{5.1}
 \end{aligned}$$

respectively. Here also the amplitude in the t channel is the same as A_u in which the particle labels of the RR fields are

interchanged, *i.e.*, $A_t = A_u (3 \leftrightarrow 4)$. The required three- and four-point interactions are given by

$$\left(\tilde{V}_{\phi_1\phi_2h}\right)^{\mu\nu} = -2i\kappa \left(k_1^{(\mu}k_2^{\nu)} - \frac{1}{2}k_1.k_2\eta^{\mu\nu}\right), \tag{5.2}$$

$$\left(\tilde{V}_{F_1^{(n)}\phi_2C^{(n-1)}}\right)_{v_1\dots v_{n-1}} = i\kappa \frac{1}{\sqrt{2}(n-1)!} (5-n) F_{1\mu v_1\dots v_{n-1}} k^\mu, \tag{5.3}$$

$$\tilde{V}_{F_1^{(n)}F_2^{(n)}\phi_3\phi_4} = -2i\kappa^2 \frac{1}{n!} \left(\frac{5-n}{2}\right)^2 F_{1\mu_1\dots\mu_n} F_2^{\mu_1\dots\mu_n}. \tag{5.4}$$

The graviton and RR propagators are given by Eqs. (4.3) and (4.6), respectively. Substituting these expressions into the amplitude (5.1), one finds the couplings consisting of two RR 1-form field strengths and two dilatons as follows:

$$S_{(\partial F^{(1)})^2(\partial\phi)^2} = \frac{\gamma}{3.2^5\kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(2F^{a,b}F^{c,d}\phi_{a,b}\phi_{c,d} - F_{a,b}F^{a,b}\phi_{c,d}\phi^{c,d}\right). \tag{5.5}$$

One can also write down the couplings between two RR 2-form field strengths and two dilatons as

$$S_{(\partial F^{(2)})^2(\partial\phi)^2} = \frac{\gamma}{3.2^8\kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(16F_a^{d,e}F^{ab,c}\phi_{b,d}\phi_{c,e} + 18F_a^d{}_{,c}F^{ab,c}\phi_b{}^e\phi_{d,e} - 9F_{ac,b}F^{ab,c}\phi_{d,e}\phi^{d,e}\right). \tag{5.6}$$

In the same way, the couplings of two RR 3-form field strengths and two dilatons can be written as

$$S_{(\partial F^{(3)})^2(\partial\phi)^2} = -\frac{\gamma}{3^2.2^6\kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(F_{def,c}F^{def}{}_{,b}\phi_a{}^c\phi^{a,b} - 3F_a{}^{e,f}{}_{,b}F_{cef,d}\phi^{a,b}\phi^{c,d} - 3F_a{}^{ef}{}_{,c}F_{def,b}\phi^{a,b}\phi^{c,d}\right). \tag{5.7}$$

Furthermore, the couplings of two RR 4-form field strengths and two dilatons can be introduced as

$$S_{(\partial F^{(4)})^2(\partial\phi)^2} = -\frac{\gamma}{3^2.2^9\kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(F_{cdeg,f}F^{cdef,g}\phi_{a,b}\phi^{a,b} - 2F_b{}^{def,g}F_{cdef,g}\phi_a{}^c\phi^{a,b} - 16F_a{}^{efg}{}_{,c}F_{befg,d}\phi^{a,b}\phi^{c,d}\right). \tag{5.8}$$

Finally, for $n = 5$ case only the contribution of the *s* channel exists and is given by the first line in (5.1). The amplitudes in the *t* and *u* channels and the contact terms here are zero. Once more, using the self-duality condition, one can easily

find the following coupling for two RR 5-form field strengths and two dilatons with correct overall factor:

$$S_{(\partial F^{(5)})^2(\partial\phi)^2} = \frac{\gamma}{3^2.2^7\kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} F_a{}^{efgh}{}_{,c}F_{befgh,d}\phi^{a,b}\phi^{c,d}. \tag{5.9}$$

Note that imposing the self-duality condition also leads to a new coupling of the form $\epsilon_{10}(\partial F^{(5)})^2(\partial\phi)^2$, but we ignore it for the reasons already mentioned. We use the algorithm introduced in [24] to reduce the tensor polynomials and rewrite the couplings of this section in the minimal-term form.

6 $(\partial F^{(n)})^2(\partial\phi)R$ couplings

Let us now consider the couplings with structure $(\partial F^{(n)})^2(\partial\phi)R$. At the first glance, it seems that there are five types of couplings in this section, *i.e.*, with $n = 1, 2, 3, 4, 5$, but the contribution of $\partial F^{(1)}\partial F^{(1)}\partial\phi R$ and $\partial F^{(5)}\partial F^{(5)}\partial\phi R$ vanishes as can be seen from the action (2.2). When the two RR forms have the same rank, from the type II supergravity actions, it is concluded that the amplitudes in the *s* and *u* channels and the contact terms are given by the following expressions:

$$\begin{aligned} A_s &= \tilde{V}_{F_1^{(n)}F_2^{(n)}\phi} \tilde{G}_\phi \tilde{V}_{\phi\phi_3h_4}, \\ A_u &= \left(\tilde{V}_{F_1^{(n)}\phi_3C^{(n-1)}}\right)^{\mu_1\dots\mu_{n-1}} \left(\tilde{G}_{C^{(n-1)}}\right)_{\mu_1\dots\mu_{n-1}}{}^{v_1\dots v_{n-1}} \\ &\quad \times \left(\tilde{V}_{C^{(n-1)}F_2^{(n)}h_4}\right)_{v_1\dots v_{n-1}}, \\ A_c &= \tilde{V}_{F_1^{(n)}F_2^{(n)}\phi_3h_4}, \end{aligned} \tag{6.1}$$

respectively. The amplitude in the *t* channel is the same as A_u in which the particle labels of the RR fields are interchanged, *i.e.*, $A_t = A_u (3 \leftrightarrow 4)$. Here, the dilaton propagator and vertex operators which are needed in the calculation of amplitude, are

$$\tilde{V}_{F_1^{(n)}F_2^{(n)}\phi} = -i\kappa \frac{5-n}{\sqrt{2}n!} F_{1\mu_1\dots\mu_n} F_2^{\mu_1\dots\mu_n}, \tag{6.2}$$

$$\tilde{G}_\phi = -\frac{i}{k^2}, \tag{6.3}$$

$$\tilde{V}_{h_1\phi_2\phi} = -2i\kappa k_2.h_1.k, \tag{6.4}$$

$$\left(\tilde{V}_{F_1^{(n)}\phi_2C^{(n-1)}}\right)^{\mu_1\dots\mu_{n-1}} = -i\kappa \frac{5-n}{\sqrt{2}(n-1)!} F_1^{\lambda\mu_1\dots\mu_{n-1}} k_\lambda, \tag{6.5}$$

$$\begin{aligned} \tilde{V}_{F_1^{(n)}F_2^{(n)}\phi_3h_4} &= \frac{i\kappa^2(5-n)}{\sqrt{2}n!} \left(F_{1\mu_1\dots\mu_n} F_2^{\mu_1\dots\mu_n} h_{4\mu}{}^\mu \right. \\ &\quad \left. - 2n F_{1\mu\mu_2\dots\mu_n} F_{2\nu}{}^{\mu_2\dots\mu_n} h_4^{\mu\nu}\right). \end{aligned} \tag{6.6}$$

The RR propagator is given by Eq. (4.6) as well. After doing some algebra, we have

$$S_{(\partial F^{(2)})^2(\partial\phi)R} = \frac{\gamma}{2^{\frac{11}{2}}\kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(2F_a^{d,e} F^{ab,c} \phi_b^{,f} R_{cdef} - 2F^{ab,c} F_c^{d,e} \phi_e^{,f} R_{dabf} - F_a^d{}_{,c} F^{ab,c} \phi_e^{,f} R_{bedf} \right). \tag{6.7}$$

Doing the same steps as above, one finds the couplings with structure $(\partial F^{(2)})^2(\partial\phi)R$ in the type IIA theory. In the string frame, they are

$$S_{(\partial F^{(3)})^2(\partial\phi)R} = -\frac{\gamma}{3 \cdot 2^{\frac{9}{2}}\kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(2F_a^{cd,e} F_c^{f,g} \phi^{a,b} R_{bf dg} - F_a^{cd}{}_{,b} F_c^{ef,g} \phi^{a,b} R_{dgef} - 2F_a^{cd,e} F_c^{fg}{}_{,b} \phi^{a,b} R_{dfeg} + 2F_c^{fg}{}_{,b} F^{cde}{}_{,a} \phi^{a,b} \right). \tag{6.8}$$

Finally, we have found the following result for the couplings containing two RR 4-form field strengths, one dilaton and one Riemann curvature in the string frame:

$$S_{(\partial F^{(4)})^2(\partial\phi)R} = \frac{\gamma}{32 \cdot 2^{\frac{13}{2}}\kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(2F_b^{fgh}{}_{,d} F_c{}_{fgh,e} \phi^{a,b} R_a{}^{cde} - 3F_{bc}{}^{fg,h} F_{defg,h} \phi^{a,b} R_a{}^{cde} + F_c{}^{efg,h} F_{defg,h} \phi^{a,b} R_a{}^c{}_{b,d} \right). \tag{6.9}$$

Here also the algorithm introduced in [24] has been used to rewrite the couplings of this section in their minimal-term form.

7 $(\partial F^{(n)})^2(\partial H)^2$ couplings

Now, we consider the couplings with structure $(\partial F^{(n)})^2(\partial H)^2$. There are also five types of couplings in this section, *i.e.*, with $n = 1, 2, 3, 4, 5$. When $n = 1, 2, 3$, it can be seen from the actions (2.1) and (2.2) that the s and u channel amplitudes as well as the contact terms are given by

$$\begin{aligned} A_s &= \left(\tilde{V}_{F_1^{(n)} F_2^{(n)} h} \right)^{\mu\nu} \left(\tilde{G}_h \right)_{\mu\nu,\lambda\rho} \left(\tilde{V}_{hb_3b_4} \right)^{\lambda\rho} \\ &\quad + \tilde{V}_{F_1^{(n)} F_2^{(n)} \phi} \tilde{G}_\phi \tilde{V}_{\phi b_3b_4}, \\ A_u &= \left(\tilde{V}_{F_1^{(n)} b_3 C^{(n+1)}} \right)^{\mu_1 \dots \mu_{n+1}} \left(\tilde{G}_{C^{(n+1)}} \right)_{\mu_1 \dots \mu_{n+1}}{}^{\nu_1 \dots \nu_{n+1}} \\ &\quad \times \left(\tilde{V}_{C^{(n+1)} F_2^{(n)} b_4} \right)_{\nu_1 \dots \nu_{n+1}}, \\ A_c &= \tilde{V}_{F_1^{(n)} F_2^{(n)} b_3 b_4}, \end{aligned} \tag{7.1}$$

respectively. The t channel amplitude A_t can now be obtained from A_u by permuting the external B-fields lines 3 and 4. The vertices, which have not been introduced in the previous sections, are

$$\left(\tilde{V}_{b_1 b_2 h} \right)^{\mu\nu} = -2i\kappa \left[\frac{1}{2} (k_1 \cdot k_2 \eta^{\mu\nu} - k_1^\mu k_2^\nu - k_1^\nu k_2^\mu) \text{Tr}(b_1 \cdot b_2) - k_1 \cdot b_2 \cdot b_1 \cdot k_2 \eta^{\mu\nu} + 2k_1^{(\mu} b_2^{\nu)} \cdot b_1 \cdot k_2 + 2k_2^{(\mu} b_1^{\nu)} \cdot b_2 \cdot k_1 + 2k_1 \cdot b_2^{(\mu} b_1^{\nu)} \cdot k_2 - k_1 \cdot k_2 (b_1^\mu \cdot b_2^\nu + b_2^\mu \cdot b_1^\nu) \right], \tag{7.2}$$

$$\tilde{V}_{b_1 b_2 \phi} = -\sqrt{2}i\kappa [2k_1 \cdot b_2 \cdot b_1 \cdot k_2 - k_1 \cdot k_2 \text{Tr}(b_1 \cdot b_2)], \tag{7.3}$$

$$\left(\tilde{V}_{F_1^{(n)} b_2 C^{(n+1)}} \right)_{\nu_1 \dots \nu_{n+1}} = -2i\kappa \frac{1}{(n+1)!} k^\mu b_{2\mu[\nu_1} F_{1\nu_2 \dots \nu_{n+1}]}, \tag{7.4}$$

$$\tilde{V}_{F_1^{(n)} F_2^{(n)} b_3 b_4} = -\frac{2i\kappa^2 (n+2)(n+1)}{n!} b_{3[\mu\nu} F_{1\mu_1 \dots \mu_n]} b_4^{\mu\nu} F_2^{\mu_1 \dots \mu_n} + (3 \leftrightarrow 4). \tag{7.5}$$

Putting these contributions together, one finds the following couplings between two RR 1-form field strengths and two B-field strengths:

$$S_{(\partial F^{(1)})^2(\partial H)^2} = \frac{\gamma}{32 \cdot 2^6 \kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(12F_a^{a,b} F^{c,d} H_a{}^{ef}{}_{,c} H_{bef,d} + 6F_a{}^c F^{a,b} H_b{}^{de,f} H_{cde,f} - F_{a,b} F^{a,b} H_{cde,f} H^{cde,f} \right), \tag{7.6}$$

which is written in its minimal-term form. In a similar way, the couplings between two RR 2-form field strengths and two B-field strengths have appeared in the action,

$$S_{(\partial F^{(2)})^2(\partial H)^2} = -\frac{\gamma}{32 \cdot 2^5 \kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(3F^{ab,c} F^{de}{}_{,c} H_{ad}{}^{f,g} H_{beg,f} + 6F^{ab,c} F^{de,f} H_{ac}{}^g{}_{,b} H_{dfg,e} - 3F_a{}^{d,e} F^{ab,c} H_b{}^{fg}{}_{,c} H_{dfg,e} + 3F^{ab,c} F_c{}^{d,e} H_d{}^{fg}{}_{,a} H_{efg,b} + 3F^{ab,c} F^{de,f} H_{ab}{}^g{}_{,d} H_{efg,c} - 3F_a{}^{d,e} F^{ab,c} H_b{}^{fg}{}_{,d} H_{efg,c} - 3F_a{}^{d,e} F^{ab,c} H_b{}^{fg}{}_{,c} H_{efg,d} + 3F_a{}^{d,e} F^{ab,c} H_c{}^{fg}{}_{,b} H_{efg,d} + 2F_{ac}{}^d F^{ab,c} H_{efg,d} H^{efg}{}_{,b} - F_a{}^d{}_{,c} F^{ab,c} H_{efg,d} H^{efg}{}_{,b} \right). \tag{7.7}$$

It is also straightforward to find the couplings between two RR 3-form field strengths and two B-field strengths as

$$S_{(\partial F^{(3)})^2(\partial H)^2} = \frac{\gamma}{3^3 \cdot 2^6 \kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(18F^{abc,d} F^{efg,h} H_{abd,h} H_{cef,g} + 27F_a{}^{ef,g} F^{abc,d} H_{bcg}{}^h H_{def,h} + 36F_{ab}{}^{e,f} F^{abc,d} H_{cf}{}^g{}_{,h} H_{deg,h} - 36F_a{}^{ef,g} F^{abc,d} H_{bg}{}^h{}_{,c} H_{deh,f} - 9F^{abc,d} F^{efg,h} H_{abh,e} H_{dfg,c} \right)$$

$$\begin{aligned}
 & -36F_a^{ef,g} F^{abc,d} H_{bc}{}^h{}_e H_{dfg,h} \\
 & +36F_a^{ef,g} F^{abc,d} H_{bg}{}^h{}_e H_{dfh,c} \\
 & +72F^{abc,d} F^{efg,h} H_{abe,c} H_{dfh,g} \\
 & -36F_a^{ef,g} F^{abc,d} H_{bc}{}^h{}_e H_{dfh,g} \\
 & +54F_{ab}{}^{e,f} F^{abcd} H_{cf}{}^{g,h} H_{dgh,e} \\
 & -9F_{ab}{}^{e,f} F^{abcd} H_c{}^{gh}{}_e H_{dgh,e} \\
 & +216F_a^{ef,g} F^{abc,d} H_{be}{}^h{}_c H_{dgh,f} \\
 & +72F_{ab}{}^{e,f} F^{abcd} H_{ce}{}^{g,h} H_{dgh,f} \\
 & +18F_{ab}{}^{e,f} F^{abcd} H_c{}^{gh}{}_e H_{dgh,f} \\
 & -F^{abc,d} F^{efg,h} H_{abc,d} H_{efg,h} \\
 & +27F_a^{ef,g} F^{abc,d} H_{bcd}{}^h H_{efg,h} \\
 & -18F^{abc,d} F_d{}^{ef}{}_a H_{bc}{}^{g,h} H_{efg,h} \\
 & +108F_{ad}{}^{e,f} F^{abc,d} H_b{}^{gh}{}_c H_{efg,h} \\
 & -72F_{ab}{}^{e,f} F^{abc,d} H_{cd}{}^{g,h} H_{efg,h} \\
 & -9F_{ab}{}^e{}_d F^{abcd} H_c{}^{fg,h} H_{efg,h} \\
 & +36F_{ad}{}^e{}_b F^{abcd} H_c{}^{fg,h} H_{efg,h} \\
 & +9F_{ab}{}^{e,f} F^{abc,d} H_d{}^{gh}{}_c H_{fgh,e} \Big). \tag{7.8}
 \end{aligned}$$

For $n = 4$, the contribution of the u channel in the amplitude is replaced by

$$\begin{aligned}
 A_u = & \left(\tilde{V}_{F_1^{(n)} b_3 C^{(n-3)}} \right)^{\mu_1 \dots \mu_{n-3}} \left(\tilde{G}_{C^{(n-3)}} \right)_{\mu_1 \dots \mu_{n-3}}{}^{\nu_1 \dots \nu_{n-3}} \\
 & \times \left(\tilde{V}_{C^{(n-3)} F_2^{(n)} b_4} \right)_{\nu_1 \dots \nu_{n-3}} \\
 & + \left(\tilde{V}_{F_1^{(n)} b_3 C^{(n+1)}} \right)^{\mu_1 \dots \mu_{n+1}} \\
 & \times \left(\tilde{G}_{C^{(n+1)}} \right)_{\mu_1 \dots \mu_{n+1}}{}^{\nu_1 \dots \nu_{n+1}} \left(\tilde{V}_{C^{(n+1)} F_2^{(n)} b_4} \right)_{\nu_1 \dots \nu_{n+1}}, \tag{7.9}
 \end{aligned}$$

where

$$\left(\tilde{V}_{F_1^{(n)} b_2 C^{(n-3)}} \right)_{\nu_1 \dots \nu_{n-3}} = -i\kappa \frac{1}{(n-3)!} F_{1\mu\nu\rho\nu_1 \dots \nu_{n-3}} b_2^{\mu\nu} k^\rho. \tag{7.10}$$

Inserting this contribution into the amplitude, one finds the following couplings between two RR 4-form field strengths and two B-field strengths:

$$\begin{aligned}
 S_{(\partial F^{(4)})^2(\partial H)^2} = & -\frac{\gamma}{32 \cdot 2^8 \kappa^2} \\
 & \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(F_{fghi,e} F^{fghi}{}_d H_{abc}{}^e H^{abc,d} \right. \\
 & -2F_{efgi,h} F^{efgh,i} H_{abd,c} H^{abc,d} \\
 & -8F_d{}^{ghi}{}_c F_{fghi,e} H_{ab}{}^e{}_f H^{abc,d} \\
 & +12F_{bc}{}^{gh,i} F_{efgi,h} H_a{}^{ef}{}_d H^{abc,d} \\
 & -8F_b{}^{ghi}{}_c F_{eghi,f} H_a{}^{ef}{}_d H^{abc,d} \\
 & +48F_{bd}{}^{hi}{}_e F_{cfhi,g} H_a{}^{ef,g} H^{abc,d} \\
 & \left. -48F_{bg}{}^{hi}{}_c F_{dehi,f} H_a{}^{ef,g} H^{abc,d} \right)
 \end{aligned}$$

$$\begin{aligned}
 & -24F_{bc}{}^{hi}{}_e F_{dfhi,g} H_a{}^{ef,g} H^{abc,d} \\
 & -48F_{be}{}^{hi}{}_c F_{dfhi,g} H_a{}^{ef,g} H^{abc,d} \\
 & -48F_{be}{}^{hi}{}_c F_{dghi,f} H_a{}^{ef,g} H^{abc,d} \\
 & -24F_{bcd}{}^{h,i} F_{efgi,h} H_a{}^{ef,g} H^{abc,d} \\
 & -6F_{bc}{}^{hi}{}_g F_{efhi,d} H_a{}^{ef,g} H^{abc,d} \\
 & +24F_{bcd}{}^{h,i} F_{efhi,g} H_a{}^{ef,g} H^{abc,d} \\
 & +6F_{bc}{}^{hi}{}_d F_{efhi,g} H_a{}^{ef,g} H^{abc,d} \\
 & \left. -96F_{bcd}{}^{h,i} F_{eghi,f} H_a{}^{ef,g} H^{abc,d} \right). \tag{7.11}
 \end{aligned}$$

For $n = 5$, in addition to the u channel amplitude, which is given by (7.9), the s channel amplitude is also replaced by

$$A_s = \left(\tilde{V}_{F_1^{(n)} F_2^{(n)} h} \right)^{\mu\nu} \left(\tilde{G}_h \right)_{\mu\nu,\lambda\rho} \left(\tilde{V}_{hb_3 b_4} \right)^{\lambda\rho}. \tag{7.12}$$

Applying these changes to the amplitude along with imposing the self-duality condition, after some simplifications, one finds the following couplings between two RR 5-form field strengths and two B-field strengths:

$$\begin{aligned}
 S_{(\partial F^{(5)})^2(\partial H)^2} = & -\frac{\gamma}{5 \cdot 3^2 \cdot 2^9 \kappa^2} \\
 & \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(2F_{fghij,e} F^{fghij}{}_d H_{abc}{}^e H^{abc,d} \right. \\
 & -5F_{efghj,i} F^{efghi,j} H_{abd,c} H^{abc,d} \\
 & -20F_d{}^{ghij}{}_c F_{fghij,e} H_{ab}{}^e{}_f H^{abc,d} \\
 & +60F_{bc}{}^{ghi,j} F_{efghj,i} H_a{}^{ef}{}_d H^{abc,d} \\
 & -20F_b{}^{ghij}{}_c F_{eghij,f} H_a{}^{ef}{}_d H^{abc,d} \\
 & +160F_{bd}{}^{hij}{}_e F_{cfhij,g} H_a{}^{ef,g} H^{abc,d} \\
 & -160F_{bg}{}^{hij}{}_c F_{dehij,f} H_a{}^{ef,g} H^{abc,d} \\
 & -80F_{bc}{}^{hij}{}_e F_{dfhij,g} H_a{}^{ef,g} H^{abc,d} \\
 & -160F_{be}{}^{hij}{}_c F_{dfhij,g} H_a{}^{ef,g} H^{abc,d} \\
 & -160F_{be}{}^{hij}{}_c F_{dghij,f} H_a{}^{ef,g} H^{abc,d} \\
 & -240F_{bcd}{}^{hi,j} F_{efghj,i} H_a{}^{ef,g} H^{abc,d} \\
 & -20F_{bc}{}^{hij}{}_g F_{efhij,d} H_a{}^{ef,g} H^{abc,d} \\
 & -120F_{bcd}{}^{hi,j} F_{efhij,g} H_a{}^{ef,g} H^{abc,d} \\
 & +20F_{bc}{}^{hij}{}_d F_{efhij,g} H_a{}^{ef,g} H^{abc,d} \\
 & \left. +480F_{bcd}{}^{hi,j} F_{eghij,f} H_a{}^{ef,g} H^{abc,d} \right). \tag{7.13}
 \end{aligned}$$

Imposing the self-duality constraint also leads to a new coupling with structure $\epsilon_{10}(\partial F^{(5)})^2(\partial H)^2$, which has been neglected here.

8 $\partial F^{(n)} \partial F^{(n-4)} (\partial H)^2$ couplings

Since the minimum rank of RR field strength is 1, there is only one type of couplings in this section, *i.e.*, with $n = 5$. The effective action (2.2) shows that the amplitude in the s channel and contact terms vanishes. It also produces the

following u channel amplitude:

$$A_u = \left(\tilde{V}_{F_1^{(n)} b_3 C^{(n-3)}} \right)^{\mu_1 \dots \mu_{n-3}} \left(\tilde{G}_{C^{(n-3)}} \right)_{\mu_1 \dots \mu_{n-3}}^{\nu_1 \dots \nu_{n-3}} \times \left(\tilde{V}_{C^{(n-3)} F_2^{(n-4)} b_4} \right)_{\nu_1 \dots \nu_{n-3}}, \tag{8.1}$$

where the vertices and propagator are given in (7.10), (7.4) and (4.6), respectively. The amplitude in the t channel is the same as A_u in which the particle labels of the external B-fields are interchanged which means that $A_t = A_u (3 \leftrightarrow 4)$. Gathering these two contributions, imposing the self-duality constraint, one finds the coupling

$$S_{\partial F^{(5)} \partial F^{(1)} (\partial H)^2} = -\frac{\gamma}{3 \cdot 2^5 \kappa^2} \times \int d^{10} x e^{2\phi_0} \sqrt{-G} (3 F_{cdefg,h} F^{a,b} H_a^{cd,e} H_b^{fg,h} + 3 F_{cdfgh,b} F^{a,b} H_a^{cd,e} H_e^{fg,h} - F_{cdefg,h} F^{a,b} H_a^{cd,b} H^{efg,h}) \tag{8.2}$$

plus a coupling of the form $\epsilon_{10} \partial F^{(5)} \partial F^{(1)} (\partial H)^2$, which we drop here. The above coupling is written in its minimal-term form.

9 $\partial F^{(n)} \partial F^{(n-2)} (\partial H) R$ couplings

Here, we will calculate the couplings with structure $\partial F^{(n)} \partial F^{(n-2)} (\partial H) R$. There are three possibilities in this case, with $n = 3, 4, 5$. The supergravity actions (2.1) and (2.2) dictate that the amplitudes in the s channel and contact terms are given by

$$A_s = \left(\tilde{V}_{F_1^{(n)} F_2^{(n-2)} b} \right)^{\mu\nu} \left(\tilde{G}_b \right)_{\mu\nu, \lambda\rho} \left(\tilde{V}_{bb_3 h_4} \right)^{\lambda\rho}, \tag{9.1}$$

$$A_c = \tilde{V}_{F_1^{(n)} F_2^{(n-2)} b_3 h_4},$$

where the vertex operators and B-field propagator are given by

$$\left(\tilde{V}_{F_1^{(n)} F_2^{(n-2)} b} \right)^{\mu\nu} = -\frac{i\kappa}{(n-2)!} F_1^{\mu\nu}{}_{\nu_1 \dots \nu_{n-2}} F_2^{\nu_1 \dots \nu_{n-2}}, \tag{9.2}$$

$$\left(\tilde{G}_b \right)_{\mu\nu, \lambda\rho} = -\frac{i}{2k^2} (\eta_{\mu\lambda} \eta_{\nu\rho} - \eta_{\mu\rho} \eta_{\nu\lambda}), \tag{9.3}$$

$$\left(\tilde{V}_{b_1 h_2 b} \right)^{\mu\nu} = -2i\kappa (k_2 \cdot b_1 \cdot k_2 h_2^{\mu\nu} + 2k_2 \cdot b_1^{[\mu} h_2^{\nu]} \cdot k_1 + 2k_2 \cdot b_1 \cdot h_2^{[\mu} k_2^{\nu]} + 2k_1 \cdot k_2 h_2^{[\mu} \cdot b_1^{\nu]} + k_2^{[\mu} b_1^{\nu]} \cdot h_2 \cdot k_1 + k_1 \cdot h_2 \cdot b_1^{[\mu} k_2^{\nu]}), \tag{9.4}$$

$$\tilde{V}_{F_1^{(n)} F_2^{(n-2)} b_3 h_4} = \frac{i\kappa^2}{n!} \left(F_{1\mu_1 \dots \mu_n} b_3^{[\mu_1 \mu_2} F_2^{\mu_3 \dots \mu_n]} h_4^{\lambda} \right. \\ \left. - 2n F_{1\lambda \mu_2 \dots \mu_n} b_3^{[\rho \mu_2} F_2^{\mu_3 \dots \mu_n]} h_4^{\lambda} \right). \tag{9.5}$$

The amplitude in the u channel is the same as A_s in which the particle labels of the RR $(n-2)$ -form field strength and B-field are interchanged, *i.e.*, $A_u = A_s (2 \leftrightarrow 3)$. We have

$$A_u = \left(\tilde{V}_{F_1^{(n)} b_3 C^{(n-3)}} \right)^{\mu_1 \dots \mu_{n-3}} \left(\tilde{G}_{C^{(n-3)}} \right)_{\mu_1 \dots \mu_{n-3}}^{\nu_1 \dots \nu_{n-3}} \times \left(\tilde{V}_{C^{(n-3)} F_2^{(n-2)} h_4} \right)_{\nu_1 \dots \nu_{n-3}}. \tag{9.6}$$

Similarly, the amplitude in the t channel is the same as A_u in which the particle labels of the external B-field and graviton are interchanged, *i.e.*, $A_t = A_u (3 \leftrightarrow 4)$. We have

$$A_t = \left(\tilde{V}_{F_1^{(n)} h_4 C^{(n+1)}} \right)^{\mu_1 \dots \mu_{n+1}} \left(\tilde{G}_{C^{(n+1)}} \right)_{\mu_1 \dots \mu_{n+1}}^{\nu_1 \dots \nu_{n+1}} \times \left(\tilde{V}_{C^{(n+1)} F_2^{(n-2)} b_3} \right)_{\nu_1 \dots \nu_{n+1}}. \tag{9.7}$$

Replacing the vertices and propagators in the above amplitudes and then summing them yields the following result for the couplings of one RR 3-form field strength, one RR 1-form field strength, one B-field strength and one Riemann tensor:

$$S_{\partial F^{(3)} \partial F^{(1)} (\partial H) R} = -\frac{\gamma}{3 \cdot 2^5 \kappa^2} \times \int d^{10} x e^{2\phi_0} \sqrt{-G} \left(2 F^{cde,f} F^{a,b} H_{ac}{}^g{}_{,b} R_{dfeg} + 4 F^{cde,f} F^{a,b} H_{ac}{}^g{}_{,f} R_{bdeg} - 10 F^{cde,f} F^{a,b} H_{af}{}^g{}_{,c} R_{bdeg} - 2 F_a{}^{cd,e} F^{a,b} H_{bc}{}^{f,g} R_{dgef} + 2 F_a{}^{cd,e} F^{a,b} H_{cd}{}^{f,g} R_{bgef} - 2 F^{cde}{}_{,a} F^{a,b} H_{cd}{}^{f,g} R_{bgef} - F_a{}^{cd,b} F^{a,b} H_c{}^{ef,g} R_{dgef} + 4 F^{cde,f} F^{a,b} H_{cf}{}^g{}_{,a} R_{bdeg} + 2 F^{cde,f} F^{a,b} H_{cf}{}^g{}_{,d} R_{aebg} + F^{cde}{}_{,a} F^{a,b} H_c{}^{fg}{}_{,b} R_{dfeg} \right). \tag{9.8}$$

The couplings of one RR 4-form field strength, one RR 2-form field strength, one B-field strength and one Riemann curvature can be written in the string frame as

$$S_{\partial F^{(4)} \partial F^{(2)} (\partial H) R} = -\frac{\gamma}{3 \cdot 2^5 \kappa^2} \times \int d^{10} x e^{2\phi_0} \sqrt{-G} \left(12 F_{dfgh,e} F^{ab,c} H_a{}^{de,f} R_{bc}{}^{gh} - 6 F_{bdeh,g} F^{ab,c} H_c{}^{de,f} R_a{}^g{}_{,f}{}^h - 12 F_{bdgh,e} F^{ab,c} H_c{}^{de,f} R_a{}^g{}_{,f}{}^h + 6 F_{befh,g} F^{ab,c} H^{def}{}_{,c} R_a{}^g{}_{,d}{}^h + 12 F_{begh,f} F^{ab,c} H^{def}{}_{,c} R_a{}^g{}_{,d}{}^h + 6 F_{efgh,b} F^{ab,c} H^{def}{}_{,c} R_a{}^g{}_{,d}{}^h - 12 F_{bcgh,f} F^{ab,c} H^{def,g} R_a{}^h{}_{,de} - 6 F_{befh,c} F^{ab,c} H^{def,g} R_a{}^h{}_{,dg} - 12 F_{begh,f} F^{ab,c} H^{def,g} R_a{}^h{}_{,cd} - F_{cdef,h} F^{ab,c} H^{def,g} R_{abg}{}^h + 3 F_{cdeh,f} F^{ab,c} H^{def,g} R_{abg}{}^h - 6 F_{cefh,b} F^{ab,c} H^{def,g} R_{agd}{}^h + 12 F_{cefh,b} F^{ab,c} H^{def,g} R_a{}^h{}_{,dg} \right)$$

$$-2F_{defh,b}F^{ab,c}H^{def,g}R_a{}^h{}_{cg}). \tag{9.9}$$

Finally, we find the couplings of one RR 5-form field strength, one RR 3-form field strength, one B-field strength and one Riemann curvature in the type IIB theory as

$$S_{\partial F^{(5)}\partial F^{(3)}(\partial H)R} = -\frac{\gamma}{3^2 \cdot 2^5 \kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(3F_{bcefi,h}F^{abc,d}H_d{}^{ef,g}R_a{}^h{}_{g}{}^i + 6F_{bcehi,f}F^{abc,d}H_d{}^{ef,g}R_a{}^h{}_{g}{}^i - 12F_{bcghi,f}F^{abc,d}H_d{}^{ef,g}R_a{}^h{}_{e}{}^i - 12F_{ceghi,f}F^{abc,d}H_d{}^{ef,g}R_a{}^h{}_{b}{}^i - 3F_{bcfgi,h}F^{abc,d}H_d{}^{ef,g}R_a{}^h{}_{e}{}^i - 6F_{bcfhi,g}F^{abc,d}H_d{}^{ef,g}R_a{}^h{}_{e}{}^i - 6F_{bfgih,c}F^{abc,d}H_d{}^{ef,g}R_a{}^h{}_{e}{}^i - 3F_{bcfgi,d}F^{abc,d}H_d{}^{ef,g}R_{aeh}{}^i + 3F_{bcfgi,d}F^{abc,d}H_d{}^{ef,g}R_{ahe}{}^i + 6F_{bcfhi,g}F^{abc,d}H_d{}^{ef,g}R_a{}^i{}_{de} + 12F_{bdfgi,c}F^{abc,d}H_d{}^{ef,g}R_{aeh}{}^i - 6F_{bdfgi,c}F^{abc,d}H_d{}^{ef,g}R_{ahe}{}^i - 2F_{befgi,c}F^{abc,d}H_d{}^{ef,g}R_{ahd}{}^i + 12F_{bfgih,c}F^{abc,d}H_d{}^{ef,g}R_a{}^i{}_{de} + 4F_{defgi,c}F^{abc,d}H_d{}^{ef,g}R_{ahb}{}^i \right). \tag{9.10}$$

Similar to the previous cases, imposing the self-duality constraint in the above action leads to a new coupling with structure $\epsilon_{10}\partial F^{(5)}\partial F^{(3)}(\partial H)R$. We also find that there is another coupling with structure $\epsilon_{10}\partial F^{(n)}\partial F^{(n-2)}(\partial H)R$, which has not been considered here.

10 $\partial F^{(n)}\partial F^{(n-2)}\partial\phi\partial H$ couplings

The last coupling that we aim to obtain is $\partial F^{(n)}\partial F^{(n-2)}\partial\phi\partial H$. There are three types of couplings in this case and those are with $n = 3, 4, 5$. When $n = 3, 4$, from the supergravity actions (2.1) and (2.2), one can observe that the field-theory amplitudes in the s channel and contact terms are given by

$$A_s = \left(\tilde{V}_{F_1^{(n)}F_2^{(n-2)}b} \right)^{\mu\nu} \left(\tilde{G}_b \right)_{\mu\nu,\lambda\rho} \left(\tilde{V}_b\phi_3b_4 \right)^{\lambda\rho}, \tag{10.1}$$

$$A_c = \tilde{V}_{F_1^{(n)}F_2^{(n-2)}\phi_3b_4},$$

respectively. Here, the vertices, which have not been previously introduced, are given by

$$\left(\tilde{V}_{\phi_1b_2b} \right)^{\mu\nu} = -\sqrt{2}i\kappa \left(2k_1 \cdot b_2^{[\nu} k_2^{\mu]} - k_1 \cdot k_2 b_2^{\mu\nu} \right), \tag{10.2}$$

$$\tilde{V}_{F_1^{(n)}F_2^{(n-2)}\phi_3b_4} = \frac{\sqrt{2}i\kappa^2(5-n)}{n!} F_{1\mu_1\dots\mu_n} b_4^{[\mu_1\mu_2} F_2^{\mu_3\dots\mu_n]}. \tag{10.3}$$

The u channel amplitude A_u can now be obtained from A_s by permuting the particles lines 2 and 3. Similarly, the t channel amplitude A_t is obtained by permuting the external lines 3 and 4:

$$A_u = \left(\tilde{V}_{F_1^{(n)}\phi_3C^{(n-1)}} \right)^{\mu_1\dots\mu_{n-1}} \left(\tilde{G}_{C^{(n-1)}} \right)_{\mu_1\dots\mu_{n-1}}{}^{\nu_1\dots\nu_{n-1}} \times \left(\tilde{V}_{C^{(n-1)}F_2^{(n-2)}b_4} \right)_{\nu_1\dots\nu_{n-1}}. \tag{10.4}$$

Similarly, the t channel amplitude A_t is obtained by permuting the external lines 3 and 4:

$$A_t = \left(\tilde{V}_{F_1^{(n)}b_4C^{(n-3)}} \right)^{\mu_1\dots\mu_{n-3}} \left(\tilde{G}_{C^{(n-3)}} \right)_{\mu_1\dots\mu_{n-3}}{}^{\nu_1\dots\nu_{n-3}} \times \left(\tilde{V}_{C^{(n-3)}F_2^{(n-2)}\phi_3} \right)_{\nu_1\dots\nu_{n-3}}. \tag{10.5}$$

The sum of the pole diagrams, after a simple calculation, leads to the following couplings for one RR 3-form field strength, one RR 1-form field strength, one dilaton and one B-field strength in the type IIB supergravity:

$$S_{\partial F^{(3)}\partial F^{(1)}\partial\phi\partial H} = \frac{\gamma}{3 \cdot 2^{\frac{9}{2}} \kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(F^{cde,f}F^{a,b}\phi_{a,b}H_{cdf,e} - 3F^{def}{}_{,b}F^{a,b}\phi_a{}^{,c}H_{cde,f} - F_c{}^{de,f}F^{a,b}\phi_a{}^{,c}H_{def,b} + 2F_a{}^{ef}{}_{,b}F^{a,b}\phi^c{}_{,d}H_{cef,d} - 2F_a{}^{ef}{}_{,c}F^{a,b}\phi^c{}_{,d}H_{def,b} \right). \tag{10.6}$$

The amplitude of one RR 4-form field strength, one RR 2-form field strength, one dilaton and one B-field strength produces the couplings

$$S_{\partial F^{(4)}\partial F^{(2)}\partial\phi\partial H} = \frac{\gamma}{3^2 \cdot 2^{\frac{13}{2}} \kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(12F_{cefg,d}F^{ab,c}\phi_a{}^{,d}H_b{}^{ef,g} + 12F_{defg,c}F^{ab,c}\phi_a{}^{,d}H_b{}^{ef,g} - 12F_{bdef,g}F^{ab,c}\phi_a{}^{,d}H_c{}^{ef,g} + 12F_{bdeg,f}F^{ab,c}\phi_a{}^{,d}H_c{}^{ef,g} - 2F_{aefg,b}F^{ab,c}\phi_c{}^{,d}H_d{}^{ef,g} + 12F_{bdfg,e}F^{ab,c}\phi^d{}_{,e}H_c{}^{fg}{}_{,a} - 9F_{abfg,e}F^{ab,c}\phi^d{}_{,e}H_d{}^{fg}{}_{,c} + 24F_{aefg,b}F^{ab,c}\phi^d{}_{,e}H_d{}^{fg}{}_{,c} + 6F_{acfg,b}F^{ab,c}\phi^d{}_{,e}H_d{}^{fg}{}_{,e} \right) \tag{10.7}$$

in the string frame. In particular, when $n = 5$, as can be seen from the type IIB supergravity action (2.2), the u channel amplitude as well as contact terms vanishes. In other words, the s and t channels are the only two diagrams that contribute to the process. Putting these two contributions together, one

can obtain the action which reproduces the corresponding amplitude as

$$S_{\partial F^{(5)}\partial F^{(3)}\partial\phi\partial H} = \frac{\gamma}{3^2 \cdot 2^{\frac{11}{2}} \kappa^2} \times \int d^{10}x e^{2\phi_0} \sqrt{-G} \left(2F_{cdegh,b} F^{cde,f} \phi^{a,b} H_{af}{}^{g,h} + F_{cdegh,b} F^{cde,f} \phi^{a,b} H_f{}^{gh}{}_{,a} - 3F_{adegh,b} F^{cde,f} \phi^{a,b} H_f{}^{gh}{}_{,c} + 4F_{bcfgh,d} F_a{}^{cd,e} \phi^{a,b} H^{fgh}{}_{,e} + F_{cdfgh,b} F_a{}^{cd,e} \phi^{a,b} H^{fgh}{}_{,e} \right). \quad (10.8)$$

The number of terms in (10.6) and (10.8) has been reduced as much as possible. Here also imposing the self-duality condition yields a coupling with structure $\epsilon_{10}\partial F^{(5)}\partial F^{(3)}\partial\phi\partial H$, which is not considered here because the number of indices is too large to find all possible contractions of this structure.

We have compared the results of this paper with the corresponding couplings which have been obtained from string amplitude calculations in the RNS [23] and pure-spinor formalisms [28] as well as the T-duality approach [18,23], and we find exact agreement when we write both couplings in terms of independent variables.

It is interesting to find the remaining couplings by improving the computer program. One can also apply the S- and T-duality approaches to get these couplings.

Acknowledgements This work has been financially supported by the research deputy of Sirjan University of Technology.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

References

- D.J. Gross, E. Witten, Nucl. Phys. B **277**, 1 (1986). [https://doi.org/10.1016/0550-3213\(86\)90429-3](https://doi.org/10.1016/0550-3213(86)90429-3)
- E. Kiritsis, B. Pioline, Nucl. Phys. B **508**, 509 (1997). [https://doi.org/10.1016/S0550-3213\(97\)00645-7](https://doi.org/10.1016/S0550-3213(97)00645-7). arXiv:hep-th/9707018
- J.G. Russo, A.A. Tseytlin, Nucl. Phys. B **508**, 245 (1997). [https://doi.org/10.1016/S0550-3213\(97\)00631-7](https://doi.org/10.1016/S0550-3213(97)00631-7), [https://doi.org/10.1016/S0550-3213\(97\)80012-0](https://doi.org/10.1016/S0550-3213(97)80012-0). arXiv:hep-th/9707134
- M.B. Green, P. Vanhove, Phys. Lett. B **408**, 122 (1997). [https://doi.org/10.1016/S0370-2693\(97\)00785-5](https://doi.org/10.1016/S0370-2693(97)00785-5). arXiv:hep-th/9704145
- M.B. Green, M. Gutperle, P. Vanhove, Phys. Lett. B **409**, 177 (1997). [https://doi.org/10.1016/S0370-2693\(97\)00931-3](https://doi.org/10.1016/S0370-2693(97)00931-3). arXiv:hep-th/9706175
- L. Anguelova, P.A. Grassi, P. Vanhove, Nucl. Phys. B **702**, 269 (2004). <https://doi.org/10.1016/j.nuclphysb.2004.09.024>. arXiv:hep-th/0408171
- S. Stieberger. arXiv:0907.2211 [hep-th]
- D.J. Gross, J.H. Sloan, Nucl. Phys. B **291**, 41 (1987). [https://doi.org/10.1016/0550-3213\(87\)90465-2](https://doi.org/10.1016/0550-3213(87)90465-2)
- M.T. Grisaru, A.E.M. van de Ven, D. Zanon, Phys. Lett. B **173**, 423 (1986). [https://doi.org/10.1016/0370-2693\(86\)90408-9](https://doi.org/10.1016/0370-2693(86)90408-9)
- M. de Roo, H. Suelmann, A. Wiedemann, Nucl. Phys. B **405**, 326 (1993). [https://doi.org/10.1016/0550-3213\(93\)90550-9](https://doi.org/10.1016/0550-3213(93)90550-9). arXiv:hep-th/9210099
- M. de Roo, H. Suelmann, A. Wiedemann, Phys. Lett. B **280**, 39 (1992). [https://doi.org/10.1016/0370-2693\(92\)90769-Z](https://doi.org/10.1016/0370-2693(92)90769-Z)
- A. Rajaraman, Phys. Rev. D **72**, 125008 (2005). <https://doi.org/10.1103/PhysRevD.74.085018>. arXiv:hep-th/0505155
- M.F. Paulos, JHEP **0810**, 047 (2008). <https://doi.org/10.1088/1126-6708/2008/10/047>. arXiv:0804.0763 [hep-th]
- E. Cremmer, B. Julia, J. Scherk, Phys. Lett. **76B**, 409 (1978). [https://doi.org/10.1016/0370-2693\(78\)90894-8](https://doi.org/10.1016/0370-2693(78)90894-8)
- Y. Hyakutake, S. Ogushi, JHEP **0602**, 068 (2006). <https://doi.org/10.1088/1126-6708/2006/02/068>. arXiv:hep-th/0601092
- A. Rajaraman, Phys. Rev. D **74**, 085018 (2006). <https://doi.org/10.1103/PhysRevD.72.125008>. arXiv:hep-th/0512333
- K. Peeters, P. Vanhove, A. Westerberg, Class. Quant. Grav. **18**, 843 (2001). <https://doi.org/10.1088/0264-9381/18/5/307>. arXiv:hep-th/0010167
- M.R. Garousi, JHEP **1306**, 030 (2013). [https://doi.org/10.1007/JHEP06\(2013\)030](https://doi.org/10.1007/JHEP06(2013)030). arXiv:1302.7275 [hep-th]
- H. Godazgar, M. Godazgar, JHEP **1309**, 140 (2013). [https://doi.org/10.1007/JHEP09\(2013\)140](https://doi.org/10.1007/JHEP09(2013)140). arXiv:1306.4918 [hep-th]
- J.T. Liu, R. Minasian, Nucl. Phys. B **874**, 413 (2013). <https://doi.org/10.1016/j.nuclphysb.2013.06.002>. arXiv:1304.3137 [hep-th]
- M.R. Garousi. arXiv:1702.00191 [hep-th]
- K. Peeters, A. Westerberg, Class. Quant. Grav. **21**, 1643 (2004). <https://doi.org/10.1088/0264-9381/21/6/022>. arXiv:hep-th/0307298
- H.R. Bakhtiarzadeh, M.R. Garousi, Nucl. Phys. B **884**, 408 (2014). <https://doi.org/10.1016/j.nuclphysb.2014.05.002>. arXiv:1312.4703 [hep-th]
- H.R. Bakhtiarzadeh, M.R. Garousi, Phys. Rev. D **92**, no. 2, 026010 (2015). <https://doi.org/10.1103/PhysRevD.92.026010>. arXiv:1502.02853 [hep-th]
- S. Sannan, Phys. Rev. D **34**, 1749 (1986). <https://doi.org/10.1103/PhysRevD.34.1749>
- B.E.W. Nilsson, Nucl. Phys. B **188**, 176 (1981). [https://doi.org/10.1016/0550-3213\(81\)90111-5](https://doi.org/10.1016/0550-3213(81)90111-5)
- P.S. Howe, P.C. West, Nucl. Phys. B **238**, 181 (1984). [https://doi.org/10.1016/0550-3213\(84\)90472-3](https://doi.org/10.1016/0550-3213(84)90472-3)
- G. Policastro, D. Tsimpis, Class. Quant. Grav. **23**, 4753 (2006). <https://doi.org/10.1088/0264-9381/23/14/012>. arXiv:hep-th/0603165
- I. Antoniadis, R. Minasian, P. Vanhove, Nucl. Phys. B **648**, 69 (2003). [https://doi.org/10.1016/S0550-3213\(02\)00974-4](https://doi.org/10.1016/S0550-3213(02)00974-4). arXiv:hep-th/0209030
- I. Antoniadis, S. Ferrara, R. Minasian, K.S. Narain, Nucl. Phys. B **507**, 571 (1997). [https://doi.org/10.1016/S0550-3213\(97\)00572-5](https://doi.org/10.1016/S0550-3213(97)00572-5). arXiv:hep-th/9707013
- V. Iyer, R.M. Wald, Phys. Rev. D **50**, 846 (1994). <https://doi.org/10.1103/PhysRevD.50.846>. arXiv:gr-qc/9403028
- S. de Haro, A. Sinkovics, K. Skenderis, Phys. Rev. D **68**, 066001 (2003). <https://doi.org/10.1103/PhysRevD.68.066001>. arXiv:hep-th/0302136
- P. Kraus, F. Larsen, JHEP **0509**, 034 (2005). <https://doi.org/10.1088/1126-6708/2005/09/034>. arXiv:hep-th/0506176
- S.S. Gubser, I.R. Klebanov, A.A. Tseytlin, Nucl. Phys. B **534**, 202 (1998). [https://doi.org/10.1016/S0550-3213\(98\)00514-8](https://doi.org/10.1016/S0550-3213(98)00514-8). arXiv:hep-th/9805156
- J. Pawelczyk, S. Theisen, JHEP **9809**, 010 (1998). <https://doi.org/10.1088/1126-6708/1998/09/010>. arXiv:hep-th/9808126
- S. Frolov, I.R. Klebanov, A.A. Tseytlin, Nucl. Phys. B **620**, 84 (2002). [https://doi.org/10.1016/S0550-3213\(01\)00554-5](https://doi.org/10.1016/S0550-3213(01)00554-5). arXiv:hep-th/0108106

37. K. Becker, M. Becker, J.H. Schwarz, *String Theory and M-Theory: A Modern Introduction* (Cambridge University Press, Cambridge, 2007)
38. J.H. Schwarz, Phys. Rept. **89**, 223 (1982). [https://doi.org/10.1016/0370-1573\(82\)90087-4](https://doi.org/10.1016/0370-1573(82)90087-4)
39. M.R. Garousi, R.C. Myers, Nucl. Phys. B **475**, 193 (1996). [https://doi.org/10.1016/0550-3213\(96\)00316-1](https://doi.org/10.1016/0550-3213(96)00316-1). arXiv:hep-th/9603194
40. M.R. Garousi, JHEP **9812**, 008 (1998). <https://doi.org/10.1088/1126-6708/1998/12/008>. arXiv:hep-th/9805078
41. T.Nutma, Comput. Phys. Commun. **185**, 1719 (2014). <https://doi.org/10.1016/j.cpc.2014.02.006>. arXiv:1308.3493 [cs.SC]