

Erratum to: Non-perturbative contribution to the thrust distribution in e^+e^- annihilation

R.A. Davison^{1,2,a}, B.R. Webber^{1,b}

¹Cavendish Laboratory, J.J. Thomson Avenue, Cambridge CB3 0HE, UK

²Present address: Rudolf Peierls Centre for Theoretical Physics, 1 Keble Road, Oxford OX1 3NP, UK

Received: 4 June 2010 / Published online: 6 August 2010

© Springer-Verlag / Società Italiana di Fisica 2010

Erratum to: Eur. Phys. J. C (2009) 59: 13–25
DOI 10.1140/epjc/s10052-008-0836-7

The fits presented here omitted renormalization of the power correction due to higher-order effects, which introduce an overall coefficient of $2\mathcal{M}/\pi \simeq 0.95$ in equation (36), where \mathcal{M} is the ‘Milan factor’ [1]. We have redone the fits including this factor and confirmed that the results do not change significantly. Uncertainty in the value of \mathcal{M} , usually take to be $\sim 20\%$, gives rise to a correspondingly increased un-

certainty in α_0 , but the fitted combination $2\mathcal{M}\alpha_0/\pi$ and its error remain unchanged within the quoted values.

References

1. Y.L. Dokshitzer, A. Lucenti, G. Marchesini, G.P. Salam, Nucl. Phys. B **511**, 396 (1998) [Erratum-ibid. B 593, 729 (2001)].
[arXiv:hep-ph/9707532](https://arxiv.org/abs/hep-ph/9707532)

The online version of the original article can be found under
doi:[10.1140/epjc/s10052-008-0836-7](https://doi.org/10.1140/epjc/s10052-008-0836-7).

^a e-mail: richard.davison@chch.ox.ac.uk

^b e-mail: webber@hep.phy.cam.ac.uk