

Bounds on LFV Higgs decays in a vector-like lepton model and searching for doubly charged leptons at the LHC

Chuan-Hung Chen^{1,a}, Takaaki Nomura^{2,b}

¹ Department of Physics, National Cheng-Kung University, Tainan 70101, Taiwan

² School of Physics, Korea Institute for Advanced Study, Seoul 130-722, Republic of Korea

Received: 15 April 2016 / Accepted: 10 June 2016 / Published online: 24 June 2016

© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract The Higgs-portal lepton flavor violation is studied in a vector-like lepton model. To avoid the constraints from rare $Z \rightarrow \ell_i^\pm \ell_j^\mp$ decays, we introduce two triplet vector-like leptons, $(1, 3)_{-1}$ and $(1, 3)_0$. The resultant branching ratio for $h \rightarrow \mu\tau$ can be up to 10^{-4} when the constraints from the invisible Z decays are applied. As a result, the signal strength for the $\tau\tau$ channel has a 12 % deviation from the standard model prediction, while the muon $g - 2$ is two orders of magnitude smaller than the data, and $\text{BR}(\tau \rightarrow \mu\gamma)$ is of the order of 10^{-12} . A predicted doubly charged lepton in pp collisions at $\sqrt{s} = 13$ TeV is analyzed, and it is found that the interesting production channels are $pp \rightarrow (\Psi_1^- \Psi_1^{++}, \Psi_1^{\pm\pm} \Psi_1^\mp)$. Both single and pair production cross sections of Ψ_1^{++} are comparable, and can be a few hundred fb. The main decay channels for the doubly charged lepton are $\Psi^{\pm\pm} \rightarrow \ell^\pm W^\pm$, and for the heavy singly charged lepton they are $\Psi_1^\pm \rightarrow \nu W^\pm, \ell^\pm Z$. The numerical analysis is carried out with regard to 13 TeV LHC with 100 fb^{-1} luminosity.

Due to a number of unsolved issues, such as the origin of neutrino mass, dark matter (DM), and matter–antimatter asymmetry, it is believed that the standard model (SM) of particle physics is an effective theory at the electroweak scale. Since rare decays are important in the development of new physics, the loop-induced flavor-changing neutral current (FCNC) processes are generally used to examine the SM. However, most hadronic processes involve very uncertain non-perturbative quantum chromodynamic (QCD) effects, and thus, even if new physics exists, it is not easy to distinguish this from the SM results due to QCD uncertainty.

Leptons do not carry a color charge, and QCD uncertainty is thus much smaller in this case. However, due to the Glashow–Iliopoulos–Maiani (GIM) mechanism, lepton

FCNC processes in the SM (e.g., $\mu \rightarrow e\gamma$ and $\tau \rightarrow (e, \mu)\gamma$) are highly suppressed; if any signal of lepton flavor violation (LFV) is observed, it is certainly strong evidence for new physics. It is thus important to search for new physics through the lepton sector [1–3].

With the discovery of the SM Higgs in the ATLAS [4] and CMS [5] experiments, we are moving toward better understanding the process of electroweak symmetry breaking (EWSB) through the spontaneous symmetry breaking (SSB) mechanism in the scalar sector. The next mission for the High Luminosity Large Hadron Collider (LHC) is to explore not only the detailed properties of the SM Higgs, but also the new physics effects.

Since the SM Higgs has been discovered, it is of interest to search for new physics through the Higgs portal. For instance, an excess of events with a significance of 2.4σ in $h \rightarrow \mu\tau$ decay was reported by CMS in pp collisions at $\sqrt{s} = 8$ TeV, where the branching ratio (BR) with the best fit is given by [6]:

$$\text{BR}(h \rightarrow \mu\tau) = (0.84_{-0.37}^{+0.39})\% \quad [\text{CMS}]. \quad (1)$$

ATLAS also reported the same measurement and found no significant excess, where the best fit is [7]:

$$\text{BR}(h \rightarrow \mu\tau) = (0.77 \pm 0.62)\% \quad [\text{ATLAS}]. \quad (2)$$

Although the measurements of $\text{BR}(h \rightarrow \mu\tau)$ are not conclusive yet, inspired by the Higgs-portal events, a number of the possible new physics effects that could explain the large BR for $h \rightarrow \mu\tau$ decay have been studied [8–34]. In this study, we explore the LFV in the framework of a vector-like lepton model.

FCNCs are quite a common phenomenon in the quark sector, such as neutral meson oscillations and $b \rightarrow s\gamma$. However, with the exceptions of the neutrino oscillations we have no concrete and solid signals to verify the LFV in the lepton sector, thus limiting our knowledge about this. In this context, the measurements from ATLAS and CMS of the Higgs-portal

^a e-mail: physchen@mail.ncku.edu.tw

^b e-mail: nomura@kias.re.kr

LFV provide a good chance to better understand the lepton sector. Following the hint of the SM with regard to whether the Higgs couplings to the fermions appear in the Yukawa sector, a possible minimal extension of the SM for Higgs-mediated LFV is to include exotic heavy leptons or to add a new Higgs doublet without imposing any symmetry [30]. In this work, we study the implications of adding heavy leptons. In order to avoid the gauge anomaly, we investigate the model with vector-like leptons (VLLs).

To achieve mixing with the SM leptons, the representations of VLL under $SU(2) \times U(1)_Y$ gauge symmetry can be singlet, doublet [35–43], and triplet [44]. The VLLs from a composite model are discussed in earlier work [45,46]. In order to avoid the constraints from rare $Z \rightarrow \ell_i^\pm \ell_j^\mp$ decays, we study the triplet representations $(1, 3)_{-1}$ and $(1, 3)_0$ with hypercharges $Y = -1$ and $Y = 0$, respectively. The new Yukawa couplings are thus written

$$-\mathcal{L}_Y = \bar{L}\mathbf{Y}_1\Psi_{1R}H + \bar{L}\mathbf{Y}_2\Psi_{2R}\tilde{H} + m_{\psi_1}Tr\bar{\Psi}_{1L}\Psi_{1R} + m_{\psi_2}Tr\bar{\Psi}_{2L}\Psi_{2R} + H.c., \tag{3}$$

where we have suppressed the flavor indices; H is the SM Higgs doublet, $\tilde{H} = i\tau_2 H^*$, the neutral component of Higgs field is $H^0 = (v + h)/\sqrt{2}$, and the representations of two VLLs are:

$$\Psi_1 = \begin{pmatrix} \Psi_1^-/\sqrt{2} & \Psi_1^0 \\ \Psi_1^{--} & -\Psi_1^-/\sqrt{2} \end{pmatrix}, \tag{4}$$

$$\Psi_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \Psi_2^0/\sqrt{2} & \Psi_2^+ \\ \Psi_2^- & -\Psi_2^0/\sqrt{2} \end{pmatrix}$$

with $\Psi_2^+ = C\bar{\Psi}_2^-$ and $\Psi_2^0 = C\bar{\Psi}_2^0$. Since Ψ_2 is a real representation of $SU(2)$, the factor of $1/\sqrt{2}$ in Ψ_2 is to obtain the correct mass term for Majorana fermion Ψ_2^0 . Due to the new Yukawa terms of $\mathbf{Y}_{1,2}$, the heavy neutral and charged leptons mix with the SM leptons; after EWSB, the lepton mass matrices become 5×5 matrices and are expressed by

$$M_\ell = \begin{pmatrix} \mathbf{m}_\ell & \mathbf{Y}^\ell v \\ 0 & \mathbf{m}_\Psi \end{pmatrix}, \quad M_\nu = \begin{pmatrix} \mathbf{m}_\nu & \mathbf{Y}^\nu v \\ 0 & \mathbf{m}_\Psi \end{pmatrix}, \tag{5}$$

where we have chosen the basis such that the SM leptons are in diagonalized states, \mathbf{m}_ℓ is the SM charged lepton mass matrix, $\mathbf{m}_\Psi = \text{diag}(m_{\psi_1}, m_{\psi_2})$, and

$$\mathbf{Y}^\ell = \frac{1}{2} \begin{pmatrix} -Y_{11} & Y_{21} \\ -Y_{12} & Y_{22} \\ -Y_{13} & Y_{23} \end{pmatrix}, \quad \mathbf{Y}^\nu = \sqrt{2} \begin{pmatrix} Y_{11} & Y_{21}/2 \\ Y_{12} & Y_{22}/2 \\ Y_{13} & Y_{23}/2 \end{pmatrix}. \tag{6}$$

We note that the elements of \mathbf{Y}^χ should be read as $Y_{ij} = (\mathbf{Y}_i)_j$, where the index $i = 1, 2$ distinguishes the Yukawa couplings of the different VLLs and the index $j = 1, 2, 3$ stands for the flavors of the SM leptons. Since the origin of neutrino mass has not been concluded yet and is still model dependent, we directly put the Majorana type of neutrino mass term to the Yukawa sector. Since the detailed effects of

neutrino physics are irrelevant to this study, we do not further pursue issues related to this and $\mathbf{m}_\nu = 0$.

To diagonalize M_ℓ and M_ν , we introduce the unitary matrices $V_{R,L}^\chi$ with $\chi = \ell, \nu$ so that $M_\chi^{\text{dia}} = V_L^\chi M_\chi V_R^{\chi\dagger}$. The information of V_L^χ and V_R^χ can be obtained through $M_\chi M_\chi^\dagger$ and $M_\chi^\dagger M_\chi$, respectively. According to Eq. (5), it can easily be found that the flavor mixings between heavy and light leptons in V_R^χ are proportional to the lepton masses. Since the neutrino masses are small, it is a good approximation to take $V_R^\nu \approx 1$. If we further set $m_e = m_\mu = 0$ in the phenomenological analysis, only τ -related processes have significant contributions. Unlike V_R^χ , the off-diagonal elements in flavor-mixing matrices V_L^χ are associated with $\mathbf{Y}_{1,2}v/\mathbf{m}_\Psi$; in principle, the mixing effects can be of the order of 0.1. In this study, we examine these effects on $h \rightarrow \tau\mu$ and rare tau related decays. To be more specific, we parametrize the unitary matrices in terms of $\mathbf{Y}_{1,2}$ as:

$$V_L^\chi \approx \begin{pmatrix} \mathbb{1}_{3 \times 3} - \boldsymbol{\varepsilon}_L^\chi \boldsymbol{\varepsilon}_L^{\chi\dagger}/2 & -\boldsymbol{\varepsilon}_L^\chi \\ \boldsymbol{\varepsilon}_L^{\chi\dagger} & \mathbb{1}_{3 \times 3} - \boldsymbol{\varepsilon}_L^{\chi\dagger} \boldsymbol{\varepsilon}_L^\chi/2 \end{pmatrix},$$

$$V_R^\ell \approx \begin{pmatrix} \mathbb{1}_{3 \times 3} & -\boldsymbol{\varepsilon}_R^\ell \\ \boldsymbol{\varepsilon}_R^{\ell\dagger} & \mathbb{1}_{3 \times 3} \end{pmatrix}, \tag{7}$$

where $V_R^\nu \approx 1$ is implied, $\boldsymbol{\varepsilon}_L^\chi \approx v\mathbf{Y}^\chi/\mathbf{m}_\Psi$, and $\boldsymbol{\varepsilon}_R^\ell \approx v\mathbf{m}_\ell^\dagger \mathbf{Y}^\ell/\mathbf{m}_\Psi^2$.

Combining the SM Higgs couplings and the new Yukawa couplings of Eq. (3), the Higgs couplings to all singly charged leptons are given by

$$-\mathcal{L}_{h\ell'\ell''} = h\bar{\ell}'_L V_L^\ell \begin{pmatrix} \mathbf{m}_\ell/v & Y^\ell \\ 0 & 0 \end{pmatrix} V_R^{\ell\dagger} \ell''_R + H.c., \tag{8}$$

where $\ell'^T = (e, \mu, \tau, \tau', \tau'')$ is the state of a physical charged lepton in lepton flavor space. We use the notations of τ' and τ'' to denote the heavy charged VLLs. Using the parametrization of Eq. (7), the Higgs couplings to the SM leptons can be formulated by

$$-\mathcal{L}_{h\ell\ell} = C_{ij}^h \bar{\ell}_i \ell_j h + H.c.,$$

$$C_{ij}^h = \frac{m_{\ell j}}{v} \left[\delta_{ij} - \frac{3}{8} \left(\frac{v^2 Y_{1i} Y_{1j}}{m_{\psi_1}^2} + \frac{v^2 Y_{2i} Y_{2j}}{m_{\psi_2}^2} \right) \right]. \tag{9}$$

If we set $m_e = m_\mu = 0$, it is clear that in addition to the $h\tau\tau$ coupling being modified, the tree-level flavor-changing $h\tau\mu$ and $h\tau e$ couplings are induced, and the couplings are proportional to $m_\tau/v \approx 7.2 \times 10^{-3}$. In order to study the VLL contributions to $h \rightarrow \gamma\gamma$, the couplings for $h\tau'\tau'$ and $h\tau''\tau''$ are expressed as:

$$-\mathcal{L}_{h\Psi\Psi} = \frac{v \sum_i Y_{1i}^2}{2m_{\psi_1}} h\tau'\tau' + \frac{v \sum_i Y_{2i}^2}{2m_{\psi_2}} h\tau''\tau''. \tag{10}$$

Due to the mixture between VLLs and the SM leptons, the same effects also influence the gauge couplings of Z and W to the SM leptons. To understand the modifications, we discuss the gauge interactions below. With the covariant derivative for triplet VLLs [51], we first write the Z -boson gauge interactions with VLLs,

$$\begin{aligned} \mathcal{L}_Z = & -\frac{g}{c_W} Z_\mu \left[\bar{\Psi}_L \gamma^\mu (I_3^\Psi - s_W^2 Q_\Psi) \Psi_L + \bar{\Psi}_R \gamma^\mu (-s_W^2 Q_\Psi) \Psi_R \right. \\ & + \bar{\Psi}_1^- \gamma^\mu (-1 + 2s_W^2) \Psi_1^- + \bar{\Psi}_L \gamma^\mu \begin{pmatrix} 1/2 & 0 \\ 0 & -1/2 \end{pmatrix} \Psi_L \\ & \left. + \bar{N}_R \gamma^\mu \begin{pmatrix} 1/2 & 0 \\ 0 & 0 \end{pmatrix} N_R + \bar{L}'_R \gamma^\mu \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix} L'_R \right], \end{aligned} \tag{11}$$

where we have expressed the forms of couplings to be the same as those in the SM, $I_3^\Psi = 1/2(-1/2)$ and $Q_\Psi = 0(-1)$ for neutral (charged) VLLs, $\Psi^T = (\Psi_1^0, \Psi_2^0)$ or (Ψ_1^-, Ψ_2^-) , $N_R^T = (\Psi_{1R}^0, \Psi_{2R}^0)$, and $L_R^T = (\Psi_{1R}^-, \Psi_{2R}^-)$. It is clear that the first two terms in Eq. (11) provide the flavor-conserving effects; however, the last three terms lead to Z -mediated FCNC couplings at the tree level. As mentioned earlier, the flavor mixings in V_R^χ are associated with the lepton masses; if we ignore the small effects, the gauge couplings of the Z -boson to the neutral and singly charged leptons can be expressed as

$$\begin{aligned} \mathcal{L}_{Z\ell'\ell'} = & -\frac{g}{c_W} C_{ij}^{\ell'L} \bar{\ell}'_L \gamma^\mu \ell'_L Z_\mu - \frac{g}{c_W} C_{ij}^{\ell'R} \bar{\ell}'_R \gamma^\mu \ell'_R Z_\mu, \tag{12} \\ C_{ij}^{\ell'L} = & (I_3^{\ell'} - s_W^2 Q_{\ell'}) \delta_{ij} + \frac{1}{2} \left(V_{Li4}^{\ell'*} V_{Lj4}^{\ell'} - V_{Li5}^{\ell'} V_{Lj5}^{\ell'*} \right), \\ C_{ij}^{\ell'R} \approx & -s_W^2 Q_{\ell'} \delta_{ij} + \begin{cases} \frac{1}{2} \delta_{i4} \delta_{j4} & \text{for } Q_{\ell'} = 0 \\ -\delta_{i5} \delta_{j5} & \text{for } Q_{\ell'} = -1 \end{cases} \end{aligned}$$

with $\ell'^T = (\nu_1, \nu_2, \nu_3, \nu_4, \nu_5)$ or $(e, \mu, \tau, \tau', \tau'')$. As a result, the tree-level Z -mediated FCNCs only occur in the left-handed currents.

The new gauge interactions of the W -boson with VLLs are given by

$$\begin{aligned} \mathcal{L}_W = & -g \left(\bar{\Psi}_1^0 \gamma^\mu \Psi_1^- + \bar{\Psi}_1^- \gamma^\mu \Psi_1^- \right) W_\mu^+ \\ & -g \left(\bar{\Psi}_2^0 \gamma^\mu \Psi_2^- \right) W_\mu^+ + H.c.. \end{aligned} \tag{13}$$

With $V_R^\chi \approx 1$, the W -mediated interactions of neutral and singly charged leptons are expressed as:

$$\begin{aligned} \mathcal{L}_{W\nu'\ell'} = & -\frac{g}{\sqrt{2}} \bar{N}_L \gamma^\mu V_L^\nu \begin{pmatrix} V'_{PMNS} & 0 \\ 0 & \sqrt{2} \end{pmatrix} \\ & \times V_L^{\ell'\dagger} \ell'_L W_\mu^+ + H.c., \end{aligned} \tag{14}$$

where V'_{PMNS} is the Pontecorvo–Maki–Nakagawa–Sakata (PMNS) matrix without triplet VLLs, which can be regarded as a model-dependent result, and is uncertain. Since $\mathbf{m}_\nu = 0$, in this study $V'_{PMNS} = 1$. With the flavor mixings in Eq. (7), from Eq. (14) the W -boson interactions with the SM leptons are formulated by

$$\mathcal{L}_{W\nu\ell} = -\frac{g}{\sqrt{2}} (\bar{\nu}_1, \bar{\nu}_2, \bar{\nu}_3)_L \gamma^\mu V_{PMNS} \begin{pmatrix} e \\ \mu \\ \tau \end{pmatrix}_L W_\mu^+ + H.c., \tag{15}$$

$$V_{PMNS} = V'_{PMNS} - V'_{PMNS} \frac{\boldsymbol{\varepsilon}_L^\ell \boldsymbol{\varepsilon}_L^{\ell\dagger}}{2} - \frac{\boldsymbol{\varepsilon}_L^\nu \boldsymbol{\varepsilon}_L^{\nu\dagger}}{2} V'_{PMNS} + \sqrt{2} \boldsymbol{\varepsilon}_L^\nu \boldsymbol{\varepsilon}_L^{\ell\dagger}. \tag{16}$$

The V_{PMNS} is the 3×3 PMNS matrix, which can be extracted from the matrix product of $V_L^\nu(\dots)V_L^{\ell\dagger}$ in Eq. (14). Since the minimal value of the PMNS matrix element is around 0.15 [47], the limits of Y_{1i} and Y_{2i} from the charged current interactions may not be as clear as those from the rare Z decays. Therefore, to constrain the free parameters, we focus on the rare Z decays, such as $Z \rightarrow \ell_i^\pm \ell_j^\mp$, invisible Z decays, and so on.

Before studying the LFV-related phenomenologies, we discuss the possible constraints on the free parameters $Y_{1i,2i}$. From Eq. (12), it can be seen that the Z -mediated lepton flavor-violating effects can contribute to $Z \rightarrow (e\mu, e\tau, \mu\tau)$, where the current upper limits of the data are [47]

$$\begin{aligned} Br(Z \rightarrow e^\pm \mu^\mp) & < 7.5 \times 10^{-7}, \\ Br(Z \rightarrow e^\pm \tau^\mp) & < 9.8 \times 10^{-6}, \\ Br(Z \rightarrow \mu^\pm \tau^\mp) & < 1.2 \times 10^{-5}. \end{aligned} \tag{17}$$

The severe constraints make the BR of $h \rightarrow \tau\mu$ decay far smaller than the CMS measurements. In order to avoid the bounds from the rare Z decays, we set $Y_{1i} = Y_{2i}$ and $m_{\psi_1} = m_{\psi_2}$. As a result, the second term of $C_{ij}^{\ell'L}$ in Eq. (12) for charged leptons vanishes. However, the cancellations are not complete in the $Z \rightarrow \bar{\nu}_i \nu_j$ decays, due to the structure of \mathbf{Y}^ν in Eq. (6). That is, the invisible Z -boson decays can directly constrain the parameters, where the current measurement is $\Gamma_{\text{inv}}^Z = 499 \pm 1.5 \text{ MeV}$ [47] and the SM prediction is around 500 MeV. With $Y_{1i} = Y_{2i} = Y_i$ and $m_{\psi_1} = m_{\psi_2} = m_\psi$, the partial decay rates for $Z \rightarrow \bar{\nu}_i \nu_j$ and $h \rightarrow \mu\tau$ are given as:

$$\Gamma(Z \rightarrow \bar{\nu}_i \nu_j) \approx \frac{m_Z}{24\pi} \frac{g^2}{c_W^2} \sum_{ij} |C_{ij}^\nu|^2, \tag{18}$$

$$\Gamma(h \rightarrow \mu\tau) \approx \frac{m_h}{16\pi} \frac{m_\tau^2}{v^2} \left| \frac{3v^2 Y_2 Y_3}{2m_\psi^2} \right|^2. \tag{19}$$

Accordingly, we present the contours for $\Gamma(Z \rightarrow \bar{\nu}_i \nu_j)$ and $BR(h \rightarrow \tau\mu)$ as a function of Y_i and m_ψ in Fig. 1a, where numerically we adopt $Y_2 = Y_3 = Y$ and $Y_1 \ll 1$, the solid line stands for $Z \rightarrow \bar{\nu}_i \nu_j$, the dashed line is for $h \rightarrow \mu\tau$ and the values on the plot are in units of 10^{-5} , and $\Gamma_h \approx 4.21 \text{ MeV}$ is used. The results clearly show that the lepton flavor-




Fig. 1 Contours as a function of Y and m_ψ : **a** for $\Gamma(Z \rightarrow \nu_i \nu_j)$ and $\text{BR}(h \rightarrow \mu\tau)$ and **b** for signal strength $\mu_{\gamma\gamma}$, where the numbers on the plot **a** for $\text{BR}(h \rightarrow \mu\tau)$ are in units of 10^{-5}

violating Higgs decay can only be up to 10^{-4} when the data for invisible Z decays are applied.

Next, we discuss the influence of new flavor mixings on other Higgs decays. From Eq. (10), it can be seen that the induced couplings of Higgs to VLLs can contribute to $h \rightarrow \gamma\gamma$ through the loop diagrams; the decay rate can be formulated as:

$$\Gamma(h \rightarrow \gamma\gamma) \approx \Gamma^{\text{SM}}(h \rightarrow \gamma\gamma) \left| 1 + \frac{C_{VLL}}{A_W + N_C Q_t^2 A_t} \right|^2, \tag{20}$$

$$C_{LLV} = \frac{v^2 \sum_i Y_i^2}{2m_\psi^2} A_{1/2}(\tau_\psi) \tag{21}$$

where $N_C = 3$, $Q_t = 2/3$, $A_W \approx 8.3$, $A_t \approx -1.38$, and the loop integral from VLL is [48]

$$A_{1/2}(\tau) = -2\tau[1 + (1 - \tau)f(\tau)^2]$$

with $\tau_\psi = 4m_\psi^2/m_h^2$ and $f(x) = \sin^{-1}(1/\sqrt{x})$. We note that although the coupling $h\tau\tau$ is modified by the new flavor-mixing effects, we ignore its small contribution to the loop-induced $h \rightarrow \gamma\gamma$ decay. The signal strength, which is used to show the Higgs measurement, is defined as:

$$\mu_{\gamma\gamma} = \frac{\sigma(pp \rightarrow h)}{\sigma(pp \rightarrow h)_{\text{SM}}} \times \frac{\text{BR}(h \rightarrow f)}{\text{BR}(h \rightarrow f)_{\text{SM}}}, \tag{22}$$

where f denotes the possible decay channel. Taking $\Gamma_h \approx 4.21$ MeV and $\sigma(pp \rightarrow h) = \sigma(pp \rightarrow h)_{\text{SM}}$, we plot the contours for $\mu_{\gamma\gamma}$ as a function of Y and m_ψ in Fig. 1b. For comparison, we also show the constraint from Γ_{inv}^Z in the same plot. In these results we see that the deviation from the SM prediction is about 4 % and is consistent with $\mu_{\gamma\gamma} = 1.17 \pm 0.27$ and 1.13 ± 0.24 , as measured by ATLAS [52] and CMS [53], respectively.

From Eq. (9), it can be seen that the modified Higgs couplings to the SM leptons are still proportional lepton masses.

By comparison with other lepton channels, it can be seen that the $\tau\tau$ mode is more significant, and thus we study the influence on $h \rightarrow \tau^+\tau^-$. Using the values that satisfy $\text{BR}(h \rightarrow \mu\tau) \approx 10^{-4}$, the deviation of $\Gamma(h \rightarrow \tau^+\tau^-)$ from the SM results can be obtained:

$$\kappa_{\tau\tau} \equiv \frac{\Gamma(h \rightarrow \tau^+\tau^-)}{\Gamma^{\text{SM}}(h \rightarrow \tau^+\tau^-)} = \left| 1 - \frac{6v^2 Y_3^2}{8m_\psi^2} \right|^2 \approx 0.88. \tag{23}$$

If the SM Higgs production cross section is not changed, the signal strength for $pp \rightarrow h \rightarrow \tau^+\tau^-$ in this model is $\mu_{\tau\tau} \approx 0.88$, where the measurements from ATLAS and CMS are $1.44_{-0.37}^{+0.42}$ [52] and 0.91 ± 0.27 [53], respectively. Although the current data errors for the $\tau\tau$ channel are still large, the precision measurement of $\mu_{\tau\tau}$ can test the model or give strict limits on the parameters.

In the following text we investigate the contributions of new couplings in Eq. (9) to the rare tau decays and to the flavor-conserving muon anomalous magnetic moment. We first examine the muon $g - 2$, denoted by Δa_μ . The lepton flavor-changing coupling $h\mu\tau$ can contribute to the Δa_μ through the Higgs-mediated loop diagrams. However, as shown in Eq. (9), the induced couplings are associated with $m_{\ell j}/v \bar{\ell}_L \ell_R$; that is, only the right-handed tau-lepton has a significant contribution. The induced Δa_μ is thus suppressed by $m_\mu^2 m_\tau / (vm_h^2)$ so that the value of Δa_μ is two orders of magnitude smaller than current data $\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (28.8 \pm 8.0) \times 10^{-10}$ [47]. A similar situation happens in $\tau \rightarrow 3\mu$ decay. Since the couplings are suppressed by m_τ/v and m_μ/v , the BR for $\tau \rightarrow 3\mu$ is of the order of 10^{-14} . We also examine the process $\tau \rightarrow \mu\gamma$ via the h -mediation. The effective interaction for $\tau \rightarrow \mu\gamma$ is expressed by

$$\mathcal{L}_{\tau \rightarrow \mu\gamma} = \frac{e}{16\pi^2} m_\tau \bar{\mu} \sigma_{\mu\nu} (C_L P_L + C_R P_R) \tau F^{\mu\nu}, \tag{24}$$

where $C_L = 0$ and the Wilson coefficient C_R from the one-loop is formulated as:




Fig. 2 Contours for $BR(\tau \rightarrow \mu\gamma)$ (dashed) as a function of Y and m_ψ , where the constraint from Γ_{inv}^Z (solid) is included

$$C_R \approx \frac{C_{23}^h C_{33}^h}{2m_h^2} \left(\ln \frac{m_h^2}{m_\tau^2} - \frac{4}{3} \right). \tag{25}$$

Accordingly, the BR for $\tau \rightarrow \mu\gamma$ is expressed as

$$\frac{BR(\tau \rightarrow \mu\gamma)}{BR(\tau \rightarrow e\bar{\nu}_e\nu_\tau)} = \frac{3\alpha_e}{4\pi G_F^2} |C_R|^2. \tag{26}$$

We present the contours for $BR(\tau \rightarrow \mu\gamma)$ as a function of Y and m_ψ in Fig. 2, where the numbers on the plots are in units of 10^{-12} . It can be seen that the resultant $BR(\tau \rightarrow \mu\gamma)$ can be only up to 10^{-12} , where the current experimental upper bound is $BR(\tau \rightarrow \mu\gamma) < 4.4 \times 10^{-8}$ [47].

In this model, we have two new neutral leptons, two new singly charged leptons, and one doubly charged lepton. Since a particle carrying an electrical charge of 2 can have less background and a clearer signature in colliders, we discuss the potential for discovering the doubly charged lepton Ψ_1^{--} . By electroweak interactions, Ψ_1^{--} can be produced singly and in pairs through the channels $\Psi_1^{\mp\mp}\Psi_1^\pm$ and $\Psi^{--}\Psi^{++}$, where the former is from charged weak interactions while the latter is from Z and electromagnetic interactions. In addition, due to the flavor-mixing effects, the W couplings to Ψ_1^{--} and the SM leptons can be written

$$\mathcal{L}_{W\Psi_1^{--}\ell} = -g\bar{\ell}_i\gamma^\mu \left(\epsilon_{Li4}^\ell P_L + \epsilon_{Ri4}^\ell P_R \right) \Psi_1^{--} W^+ + H.c. \tag{27}$$

By the induced gauge couplings, the doubly charged lepton can be produced through the $\Psi_1^{--}\ell$ channels. We note that due to $\epsilon_R^\ell \propto \mathbf{m}_\ell$, the right-handed current contributions can be neglected. In order to discuss the production cross section in pp collisions, we implement our model in CalcHEP [49] and use CTEQ6L PDF [50] to do the numerical calculations. In Fig. 3, we show the single and pair production cross sections of $\Psi_1^{\pm\pm}$ as a function of m_ψ in pp collisions at $\sqrt{s} = 13$ TeV, where $vY/m_\psi = 0.3$ is used for the $\Psi^{++}\ell$ production.




Fig. 3 Doubly charged lepton production cross section (in units of fb) as a function of m_ψ in pp collisions at $\sqrt{s} = 13$ TeV, where X in the y axis denotes the possible channel

It can be seen that the production cross sections for $\Psi^{\pm\pm}\ell$ modes are one order of magnitude smaller than those for other modes. For $m_\psi < 400$ GeV, the production cross sections for $\Psi^{--}\Psi^{++}$ and $\Psi^{++}\Psi^{--}$, which only depend on gauge couplings, can be over 50 fb.

Next, we discuss the decays of Ψ_1^{--} and Ψ_1^- . From Eq. (3), we see that before EWSB, the triplet VLLs in Ψ_1 are degenerate; however, the masses are split when the \mathbf{Y}_1 effects are involved. Since Ψ_1^{--} cannot mix with other leptons, the mass splittings occur in neutral and singly charged leptons. According to Eqs. (5) and (7), the mass of Ψ_1^- shifted from Ψ_1^{--} is:

$$m_{\Psi_1^-} \approx m_{\Psi_1^{--}} \left(1 + \frac{v^2 \sum_i Y_{1i}^2}{8m_{\Psi_1^{--}}^2} \right). \tag{28}$$

With the bound from Γ_{inv}^Z , the mass splitting is only 3%. Therefore, we still take $m_{\Psi_1^-} \approx m_{\Psi_1^{--}} \approx \Psi_1^0$ in this study.

From Eq. (27), we see that the Ψ_1^{--} decay pattern depends on each value of Yukawa coupling Y_{1i} . If we adopt $Y_{11} \approx Y_{12} \approx Y_{13} = Y$, the better channels to search for the doubly charged lepton are $\Psi_1^{--} \rightarrow (e, \mu, \tau)W^-$ and the corresponding BRs are fixed as:

$$BR(\Psi_1^{--} \rightarrow (e, \mu, \tau)W^-) \approx 1/3, \tag{29}$$

where the lepton masses are ignored. Since τ has hadronic and leptonic decays and accompanies the neutrino when it decays, the clear signal for probing the doubly charged lepton should be $\Psi_1^{--} \rightarrow \ell W^- \rightarrow \ell(\ell\nu)$, where $\ell = e, \mu$ and the final states are the same-sign dilepton. The Higgs and gauge boson couplings of Ψ_1^- to the SM leptons are given by

$$\begin{aligned} \mathcal{L}_{\Psi_1^-} = & \frac{vY_{1i}}{2} \bar{\ell}_{iL} \Psi_{1R}^- h + \frac{g}{c_W} \frac{vY_{1i}}{2m_\psi} \bar{\ell}_{iL} \gamma^\mu \Psi_{1L}^- Z_\mu \\ & + g \frac{3vY_{1i}}{2m_\psi} \bar{\nu}_{iL} \gamma^\mu \Psi_{1L}^- W_\mu^+ + H.c. \end{aligned} \tag{30}$$

Table 1 Number of events for the processes in Eqs. (32) and (33), where a luminosity of 100 fb^{-1} and the center-of-mass energy of 13 TeV are used

Final state (GeV)	$\ell^- W^- \ell^+ W^+$	$\ell^+ \nu W^+ W^-$	$\ell^- \nu W^- W^+$	$\ell^+ \ell^- W^+ Z$	$\ell^- \ell^+ W^- Z$
$m_\Psi = 300$	7600	12,200	5800	900	400
400	2400	4000	1760	297	121
500	928	1620	648	119	44.7
600	405	727	273	53.6	18.9
700	192	355	126	26.2	8.67
800	97.2	183	61.8	13.5	4.26

It is found that the BRs for $\Psi_1^- \rightarrow (\ell_i h, \ell_i Z, \nu W)$ are insensitive to $Y = Y_{1i}$ and m_Ψ , and their ratios are

$$\ell_i h : \ell_i Z : \nu W \approx 0.02/3 : 0.1/3 : 0.89, \tag{31}$$

where ℓ_i denotes one of the SM leptons and the νW channel includes all SM neutrinos. It is clear that the BR for νW is one order of magnitude larger than other decay modes.

According to the analysis, there are two ways to search for the doubly charged lepton in the model. In pair production, the search channel is

$$pp \rightarrow \Psi_1^- \Psi_1^{++} \rightarrow (\ell^- W^-)(\ell^+ W^+), \tag{32}$$

where $\ell = e, \mu$, the W -boson can decay to leptons or jets, and their corresponding cross section with $m_\Psi = 300 \text{ GeV}$ is 76 fb at $\sqrt{s} = 13 \text{ TeV}$. The expected events with a luminosity of 100 fb^{-1} are shown in Table 1. In single production, the search channels are

$$pp \rightarrow \Psi_1^{\pm\pm} \Psi_1^\mp \rightarrow \begin{cases} (\ell^\pm W^\pm)(\nu W^\mp), \\ (\ell^\pm W^\pm)(\ell^\mp Z), \end{cases} \tag{33}$$

where the associated cross sections for $m_\Psi = 300 \text{ GeV}$ at $\sqrt{s} = 13 \text{ TeV}$ are $\sigma(\ell^+ \nu W^+ W^-) = 122 \text{ fb}$, $\sigma(\ell^- \nu W^- W^+) = 58 \text{ fb}$, $\sigma(\ell^+ \ell^- W^+ Z) = 9 \text{ fb}$, and $\sigma(\ell^- \ell^+ W^- Z) = 4 \text{ fb}$. Without event selection criteria and event kinematic cuts, we naively show the expected number of events with 100 fb^{-1} in Table 1. In this work, we just show the potential for discovering the VLLs, and the detailed event simulation with kinematic cuts will be given elsewhere.

We also examine $\Psi^{\pm\pm}$ production via flavor-changing interactions: $pp \rightarrow \Psi_1^{\pm\pm} \ell^\mp$. Although the cross section is $O(1) - O(10) \text{ fb}$ for $m_\Psi = 300 - 400 \text{ GeV}$, as shown in Fig. 3, it would be possible to find the signal with a sufficiently large luminosity. This process will thus also be important, since we could test the flavor-changing coupling $\nu Y/m_\Psi$ in collider experiments. However, detailed analysis of this is left for future work.

In summary, we investigated Higgs-portal lepton flavor violation by introducing two triplet vector-like leptons to the SM; one is the hypercharge $Y = -1$ and the other is $Y = 0$. The model has the Higgs-mediated and Z -mediated flavor-changing neutral currents at the tree level. When the bounds

from rare $Z \rightarrow \bar{\ell}_i \ell_j$ decays are smeared out, the invisible Z decays become the dominant constraints. As a result, the branching ratio for $h \rightarrow \mu\tau$ can be up to 10^{-4} , muon $g - 2$ is two orders of magnitude smaller than the current data, and $\text{BR}(\tau \rightarrow \mu\gamma)$ is of $O(10^{-12})$. The deviation of signal strength from the SM prediction in $\tau\tau$ mode is 12 %. We analyze the production channels for the predicted doubly charged lepton. We find that the interesting production channels in pp collisions are $pp \rightarrow (\Psi_1^- \Psi_1^{++}, \Psi_1^{\pm\pm} \Psi_1^\mp)$. Both single and pair production cross sections of Ψ_1^{++} are comparable and can be a few hundred fb. The main decay channel for the doubly charged lepton is $\Psi^{\pm\pm} \rightarrow \ell^\pm W^\pm$, while the heavy singly charged lepton is $\Psi_1^\pm \rightarrow \nu W^\pm, \ell^\pm Z$. We then summarize possible signatures of our model with the expected number of events for 100 fb^{-1} luminosity at the 13 TeV LHC.

Acknowledgments The work of C.H.C. was supported by the Ministry of Science and Technology of Taiwan, Republic of China, under Grant No. MOST-103-2112-M-006-004-MY3.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

References

1. A. Crivellin, A. Kokulu, C. Greub, Phys. Rev. D **87**(9), 094031 (2013). [arXiv:1303.5877](https://arxiv.org/abs/1303.5877) [hep-ph]
2. M.E. Gomez, T. Hahn, S. Heinemeyer, M. Rehman, Lepton-Flavor-Violating MSSM, Phys. Rev. D **90**(7), 074016 (2014). [arXiv:1408.0663](https://arxiv.org/abs/1408.0663) [hep-ph]
3. A. Crivellin, J. Heeck, P. Stoffer. [arXiv:1507.07567](https://arxiv.org/abs/1507.07567) [hep-ph]
4. G. Aad et al. [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012). [arXiv:1207.7214](https://arxiv.org/abs/1207.7214) [hep-ex]
5. S. Chatrchyan et al. [CMS Collaboration], Phys. Lett. B **716**, 30 (2012). [arXiv:1207.7235](https://arxiv.org/abs/1207.7235) [hep-ex]
6. V. Khachatryan et al. [CMS Collaboration], Phys. Lett. B **749**, 337 (2015). [arXiv:1502.07400](https://arxiv.org/abs/1502.07400) [hep-ex]
7. G. Aad et al. [ATLAS Collaboration], JHEP **1511**, 211 (2015). [arXiv:1508.03372](https://arxiv.org/abs/1508.03372) [hep-ex]

8. M.D. Campos, A.E.C. Hernandez, H. Pas, E. Schumacher, Phys. Rev. D **91**(11), 116011 (2015). [arXiv:1408.1652](#) [hep-ph]
9. D. Aristizabal Sierra, A. Vicente, Phys. Rev. D **90**(11), 115004 (2014). [arXiv:1409.7690](#) [hep-ph]
10. C.J. Lee, J. Tandean, JHEP **1504**, 174 (2015). [arXiv:1410.6803](#) [hep-ph]
11. J. Heeck, M. Holthausen, W. Rodejohann, Y. Shimizu, Nucl. Phys. B **896**, 281 (2015). [arXiv:1412.3671](#) [hep-ph]
12. A. Crivellin, G. D'Ambrosio, J. Heeck, Phys. Rev. Lett. **114**, 151801 (2015). [arXiv:1501.00993](#) [hep-ph]
13. I. Dorsner, S. Fajfer, A. Greljo, J.F. Kamenik, N. Kosnik, I. Nisandzic, JHEP **1506**, 108 (2015). [arXiv:1502.07784](#) [hep-ph]
14. Y. Omura, E. Senaha, K. Tobe, JHEP **1505**, 028 (2015). [arXiv:1502.07824](#) [hep-ph]
15. A. Crivellin, G.D'Ambrosio, J. Heeck, Phys. Rev. D **91**(7), 075006 (2015). [arXiv:1503.03477](#) [hep-ph]
16. D. Das, A. Kundu, Phys. Rev. D **92**(1), 015009 (2015). [arXiv:1504.01125](#) [hep-ph]
17. F. Bishara, J. Brod, P. Uttayaraj, J. Zupan, JHEP **1601**, 010 (2016). [arXiv:1504.04022](#) [hep-ph]
18. I. de Medeiros Varzielas, O. Fischer, V. Maurer, JHEP **1508**, 080 (2015). [arXiv:1504.03955](#) [hep-ph]
19. X.G. He, J. Tandean, Y.J. Zheng, JHEP **1509**, 093 (2015). [arXiv:1507.02673](#) [hep-ph]
20. C.W. Chiang, H. Fukuda, M. Takeuchi, T.T. Yanagida, JHEP **1511**, 057 (2015). [arXiv:1507.04354](#) [hep-ph]
21. W. Altmannshofer, S. Gori, A.L. Kagan, L. Silvestrini, J. Zupan. [arXiv:1507.07927](#) [hep-ph]
22. K. Cheung, W.Y. Keung, P.Y. Tseng, Phys. Rev. D **93**(1), 015010 (2016). [arXiv:1508.01897](#) [hep-ph]
23. E. Arganda, M.J. Herrero, X. Marcano, C. Weiland. [arXiv:1508.04623](#) [hep-ph]
24. F.J. Botella, G.C. Branco, M. Nebot, M.N. Rebelo. [arXiv:1508.05101](#) [hep-ph]
25. S. Baek, K. Nishiwaki, Phys. Rev. D **93**(1), 015002 (2016). [arXiv:1509.07410](#) [hep-ph]
26. W. Huang, Y.L. Tang, Phys. Rev. D **92**, 094015 (2015). [arXiv:1509.08599](#) [hep-ph]
27. S. Baek, Z.F. Kang. [arXiv:1510.00100](#) [hep-ph]
28. E. Arganda, M.J. Herrero, R. Morales, A. Szykman. [arXiv:1510.04685](#) [hep-ph]
29. D. Aloni, Y. Nir, E. Stamou. [arXiv:1511.00979](#) [hep-ph]
30. R. Benbrik, C.H. Chen, T. Nomura. [arXiv:1511.08544](#) [hep-ph]
31. M. Buschmann, J. Kopp, J. Liu, X.P. Wang. [arXiv:1601.02616](#) [hep-ph]
32. M. Sher, K. Thrasher. [arXiv:1601.03973](#) [hep-ph]
33. C.F. Chang, C.H.V. Chang, C.S. Nugroho, T.C. Yuan. [arXiv:1602.00680](#) [hep-ph]
34. X.F. Han, L. Wang, J.M. Yang. [arXiv:1601.04954](#) [hep-ph]
35. W. Altmannshofer, M. Bauer, M. Carena, JHEP **1401**, 060 (2014). [arXiv:1308.1987](#) [hep-ph]
36. A. Falkowski, D.M. Straub, A. Vicente, JHEP **1405**, 092 (2014). [arXiv:1312.5329](#) [hep-ph]
37. T. Ma, B. Zhang, G. Cacciapaglia, Phys. Rev. D **89**(9), 093022 (2014). [arXiv:1404.2375](#) [hep-ph]
38. R. Dermisek, A. Raval, S. Shin, Phys. Rev. D **90**(3), 034023 (2014). [arXiv:1406.7018](#) [hep-ph]
39. R. Dermisek, J.P. Hall, E. Lunghi, S. Shin, JHEP **1412**, 013 (2014). [arXiv:1408.3123](#) [hep-ph]
40. R. Dermisek, E. Lunghi, S. Shin, JHEP **1508**, 126 (2015). [arXiv:1503.08829](#) [hep-ph]
41. R. Dermisek, E. Lunghi, S. Shin, JHEP **1602**, 119 (2016). [arXiv:1509.04292](#) [hep-ph]
42. N. Kumar, S.P. Martin, Phys. Rev. D **92**(11), 115018 (2015). [arXiv:1510.03456](#) [hep-ph]
43. R. Dermisek, E. Lunghi, S. Shin. [arXiv:1512.07837](#) [hep-ph]
44. T. Ma, B. Zhang, G. Cacciapaglia, Phys. Rev. D **89**(1), 015020 (2014). [arXiv:1309.7396](#) [hep-ph]
45. S. Biondini, O. Panella, G. Pancheri, Y.N. Srivastava, L. Fano, Phys. Rev. D **85**, 095018 (2012). [arXiv:1201.3764](#) [hep-ph]
46. R. Leonardi, O. Panella, L. Fano, Phys. Rev. D **90**(3), 035001 (2014). [arXiv:1405.3911](#) [hep-ph]
47. K.A. Olive et al., Particle Data Group. Chin. Phys. C **38**, 090001 (2014)
48. J.F. Gunion, H.E. Haber, G.L. Kane, S. Dawson, Front. Phys. **80**, 1 (2000)
49. A. Belyaev, N.D. Christensen, A. Pukhov, Comput. Phys. Commun. **184**, 1729 (2013). [arXiv:1207.6082](#) [hep-ph]
50. P.M. Nadolsky, H.L. Lai, Q.H. Cao, J. Huston, J. Pumplin, D. Stump, W.K. Tung, C.-P. Yuan, Phys. Rev. D **78**, 013004 (2008). [arXiv:0802.0007](#) [hep-ph]
51. C.H. Chen, T. Nomura, Phys. Rev. D **92**(11), 115021 (2015). [arXiv:1509.02039](#) [hep-ph]
52. G. Aad et al. [ATLAS Collaboration]. [arXiv:1507.04548](#) [hep-ex]
53. V. Khachatryan et al. [CMS Collaboration], Eur. Phys. J. C **75**(5), 212 (2015). [arXiv:1412.8662](#) [hep-ex]