Production of $Z_c(3900)$ and $Z_c(4020)$ in B_c decay

Qi Wu¹, Dian-Yong Chen^{1,a}, Xue-Jia Fan², Gang Li^{2,b}

¹ School of Physics, Southeast University, Nanjing 210094, China

² College of Physics and Engineering, Qufu Normal University, Qufu 273165, China

Received: 19 February 2019 / Accepted: 13 March 2019 / Published online: 22 March 2019
© The Author(s) 2019

Abstract In the present work, we propose to search the charmonium-like states $Z_c(3900)$ and $Z_c(4020)$ in the B_c decay. In an effective Lagrangian approach, the branching ratios of $B_c^+ \rightarrow Z_c(3900)^+\pi^0$ and $B_c^+ \rightarrow Z_c(4020)^+\pi^0$ are estimated to be of order of 10^{-4} and 10^{-7} , respectively. The large production rate of $Z_c(3900)$ could provide an important source of the production of $Z_c(3900)$ from the semi-exclusive decay of b -flavored hadrons reported by D0 Collaboration, which can be tested by the exclusive measurements in LHCb.

1 Introduction

The charmonium-like state $Z_c^\pm(3900)$ (abbreviate to Z_c here and after) was first observed in the $J/\psi\pi^\pm$ invariant mass spectrum of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ by BESIII and Belle Collaborations [1,2], and then confirmed in $D^*\bar{D}$ invariant mass spectrum of $e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$ process [3]. The spin and parity quantum numbers of the Z_c had been determined to be $J^P = 1^+$ by the partial wave analysis of the process $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ [4]. As a partner of Z_c , $Z_c(4020)$ (abbreviate to Z'_c here and after), was discovered in the $h_c\pi^\pm$ invariant mass spectrum of $e^+e^- \rightarrow \pi^+\pi^-h_c$ [5] and confirmed in the $D^*\bar{D}^*$ invariant mass spectrum of $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$ process [6] by the BESIII Collaboration.

The decays $Z_c^\pm(3900) \rightarrow J/\psi\pi^\pm$ and $Z_c^\pm(4020) \rightarrow h_c\pi^\pm$ indicates Z_c^+ and Z_c^+ most likely contains at least four constituent quarks. Thus these two states can be good candidates of tetraquark states. In Ref. [7], Z_c was considered as the charm counterpart of $Z_b(10610)$ with $[cq][c\bar{q}]$ tetraquark configuration, and the mass estimated by using QCD sum rule (QSR) was consistent with the observed one of Z_c within the errors. The authors in Refs. [8–12] assigned both Z_c and Z'_c as

a diquark-antidiquark tetraquark state with $J^P = 1^+$, but the estimation in Ref. [12] indicated that the lowest axial-vector tetraquark states dominantly decay into $J/\psi\pi$, while their open charm decay modes were strongly suppressed, which is contrast with the experimental measurements. By reanalysing the experimental data of $J/\psi\pi$ invariant mass spectrum from the BESIII and Belle collaboration, the authors indicated that the data may contain another resonance and this resonance as well as $Z_c(3900)$ could be tetraquark states [13]. There are also some estimation based on constituent quark model, such as, QCD confining model based on SU(3) flavor symmetry [14], color flux-tube model with a four-body confinement potential [15], relativized diquark model [16], and in Ref. [17], the structures observed in $J/\psi\pi$ and $D^*\bar{D}$ were not the same, the former one could be interpreted as a $[Q\bar{q}][\bar{Q}q]$ tetraquark state, while the latter one was considered as a $[Qq][\bar{Q}\bar{q}]$ tetraquark state.

It should be notice that the observed masses of Z_c and Z'_c are in the vicinity of $D^*\bar{D}$ and $D^*\bar{D}^*$ thresholds, respectively, thus, they have been considered as the molecular or resonance states resulted from the $D^*\bar{D}^{(*)}$ interactions. The estimations in a one-boson-exchange model indicated that Z_c and Z'_c could be molecular states composed of $D\bar{D}^* + c.c$ and $D^*\bar{D}^*$, respectively [18–21], where the long range pion exchange plays an important role. By using a $D^*\bar{D}^*$ interpolating current within QSR, the mass of the molecular state was consistent with Z'_c [22]. The decay behaviors of Z_c and Z'_c were estimated in a phenomenological Lagrangian approach [23,24]. In Refs. [25,26], the hidden charm decays of Z_c and Z'_c were estimated based on heavy quark symmetry. The estimations in Ref. [27] indicated that the hidden charm decays of Z_c and Z'_c could be understood by the final states interactions, which indicated that these two charmonium-like states could be molecular candidates. In Ref. [28], the charmless decays of Z_c and Z'_c were investigated in an effective Lagrangian approach. With final states interaction effect, a pole corresponding to Z_c in the complex energy plane in different Riemann sheets could be found, which supported

^a e-mail: chendy@seu.edu.cn

^b e-mail: gli@qfnu.edu.cn

Z_c as a $D^*\bar{D} + c.c$ molecular state [29]. By considering the couple channel effect of $\pi J/\psi - \bar{D}^*D$ interaction, Z_c was found to be a virtual state [30].

Besides the tetraquark and molecular scenarios, there are also some nonresonance interpretations, such as cusp effect [31, 32] and initial single pion emission mechanism [33, 34], in which Z_c and Z'_c were considered as the kinetic effects due to $D^*\bar{D} + h.c$ and $D^*\bar{D}^*$ interactions, respectively. In addition, the Lattice QCD estimations also indicated that Z_c is not a conventional resonance but a threshold cusp [35, 36].

In addition to the mass spectrum and decay behaviors, the production process can also provide some important information of the internal structures of $Z_c^{(\prime)}$, thus investigating the productions of $Z_c^{(\prime)}$ in different processes are very critical. The observed process $Y(4260) \rightarrow Z_c\pi$ were investigated in Ref. [37], where $Y(4260)$ and Z_c were treated as $D\bar{D}_1 + c.c$ and $D^*\bar{D} + c.c$ molecular states, respectively. In Ref. [38], a large cross sections for $\gamma p \rightarrow Z_c^+ n$ was predicted, however, the COMPASS Collaboration found no evidence of Z_c , and an upper limit for the ratio $\mathcal{B}(Z_c \rightarrow J/\psi\pi) \times \sigma(\gamma N \rightarrow Z_c N)/\sigma(\gamma N \rightarrow J/\psi N)$ were measured to be 3.7×10^{-3} at 90% credit level. The cross section for the Pion-induced production of Z_c off a nuclear target was estimated in Ref. [39], where the nuclear target will enhance the meson productions. Recently, Z_c is observed in the $J/\psi\pi$ invariant mass spectrum of the semi-inclusive weak decays of b-flavored hadron [40]. However, in the exclusive B decay process, such as $B \rightarrow K J/\psi\pi\pi$ [41] and $B \rightarrow K D^*\bar{D}$ [42], no signal of Z_c is observed. Then, searching for the source of Z_c production in the semi-inclusive b-flavor hadron decay will be interesting. In addition, the production of Z_c and Z'_c are similar in electron-positron annihilation process, whether such kind of similarity still holds in b -flavored hadron decay is a question needed to be answer. Comparing to B_s^0 , B_c^+ should more easily be detected experimentally, thus, in the present work, we investigate $Z_c^{(\prime)}$ production in the B_c^+ decays.

The experimental measurements indicate that $Z_c^{(\prime)}$ dominantly decay into a pair of charmed meson or a charmonium and a pion. Thus, we have two choice to connect $Z_c^{(\prime)}$ with the mother particle B_c^+ , one is through the charmed meson loop and the other is via the loop formed by a charmonia and a light meson. Theoretical estimations of the branching ratios of two charm meson decays of B_c^+ are of order of 10^{-6} [43–46], and the LHCb collaboration have measured the decays of $B_c^+ \rightarrow D_s^{(*)+} \bar{D}^{(*)0}$ and $B_c^+ \rightarrow D_s^{(*)+} D^{(*)0}$ and no signals was found [47]. Thus, the scenario of $Z_c^{(\prime)}$ productions from B_c^+ decay via a charmed meson loop should be rule out. As for the second choice, one should notice that $B_c^+ \rightarrow c\bar{c} + \rho^+$ is a CKM enhanced mode with $(\Delta b = 1, \Delta C = 1, \Delta S = 0)$ and $\mathcal{B}(B_c^+ \rightarrow J/\psi/h_c + \rho^+)$ were predicted to be of order of 10^{-3} [46, 48]. The ρ meson dominantly decays into a pair of pion, thus, there should exists meson loops composed of

Fig. 1 Sketch diagrams for $B_c \rightarrow Z_c\pi$ (diagram a) and $B_c \rightarrow Z'_c\pi$ (diagram b)

$J/\psi\rho/h_c\rho$ in the decay channel $B_c \rightarrow Z_c\pi/Z'_c\pi$ as presented in Fig. 1. Furthermore, all the involved mesons in the loops could be on-shell and the contributions from such loop will be further enhanced, thus $B_c \rightarrow Z_c^{(\prime)}\pi$ might be a gold channel of $Z_c^{(\prime)}$ production from b flavor mesons.

This work is organized as follows. After introduction, we present formula of decays of $B_c \rightarrow c\bar{c}\rho$ and the amplitudes of the meson loop contributions to $B_c \rightarrow c\bar{c}\rho \rightarrow Z_c^{(\prime)}\pi$ in Sect. 2. The numerical results and discussions are given in Sect. 3, and Sect. 4 is devoted to a short summary.

2 Theoretical framework and the decay amplitudes

The weak decays of $B_c \rightarrow (c\bar{c})\rho$ have been well investigated in the literatures [46, 48]. For the completeness of this paper, we just present a short review of the decay amplitudes of $B_c \rightarrow (c\bar{c})\rho$, which will be used in our following estimations. At the quark level, the process happened in $B_c^+ \rightarrow c\bar{c} + \rho^+$ is the weak decay $\bar{b} \rightarrow \bar{c}u\bar{d}$ and the charm quark as a spectator. The effective Hamiltonian related to $B_c^+ \rightarrow c\bar{c} + \rho^+$ is [46]

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^\dagger [c_1 (\bar{c}b)_{V-A} (\bar{d}u)_{V-A} + c_2 (\bar{d}b)_{V-A} (\bar{c}u)_{V-A}] + \text{h.c.}, \quad (1)$$

where the subscript $V - A$ denotes the left-chiral current $\gamma^\mu (1 - \gamma^5)$, c_1 and c_2 is the Wilson coefficients.

The decay amplitudes for the non-leptonic decays can be formulated into the three factors: the non-leptonic decay constants, the weak current matrix elements and the relevant coefficients in the combinations. In particular, the non-leptonic decay constant are defined by hadronic matrix elements

$$\langle 0 | \mathcal{J}_\mu | \rho(k, \varepsilon) \rangle = f_\rho \varepsilon_\mu m_\rho, \quad (2)$$

where f_ρ stands for leptonic decay constant of ρ meson, ε_μ denotes the polarization of the vector meson, and the current $\mathcal{J}_\mu$ is $\mathcal{J}_\mu = \bar{q}_1 \gamma_\mu (1 - \gamma^5) \bar{q}_2$. The transition matrix element between B_c and (pseudo-)vector meson can be expressed in the combination of four form factors, $V, A_{0,\pm}$, which are the function of the square of the transfer momentum between B_c

and vector meson. The transition matrix element could be expressed as [49–53]

$$\langle (c\bar{c})_{J=1}(p_2, \epsilon) | \mathcal{J}^\mu | B_c^+(p) \rangle = \epsilon_v^* [i\epsilon^{\mu\nu\alpha\beta} P_\alpha Q_\beta V(Q^2) - g^{\mu\nu}(P \cdot Q)A_0(Q^2) + P^\mu P^\nu A_+(Q^2) + Q^\mu P^\nu A_-(Q^2)], \quad (3)$$

where $(c\bar{c})_{J=1}$ indicates J/ψ and h_c here and after and $P_\mu = (p+p_2)_\mu$ and $Q_\mu = (p-p_2)_\mu$.¹ The concrete expression of the form factors V , $A_{0,\pm}$ will be discussed in the following section.

With Eqs. (1)–(3), one can get the decay amplitude of $B_c \rightarrow \rho(c\bar{c})_{J=1}$,

$$\mathcal{M}[B_c^+(p) \rightarrow \rho^+(p_1)(c\bar{c})_{J=1}(p_2)] \equiv \mathcal{A}^{\mu\nu}(P, Q)\epsilon_\mu(p_1)\epsilon_\nu(p_2), \quad (6)$$

where $\mathcal{A}^{\mu\nu}(P, Q)$ is defined as

$$\mathcal{A}^{\mu\nu}(P, Q) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^\dagger a_1 f_\rho m_\rho [i\epsilon^{\mu\nu\alpha\beta} P_\alpha Q_\beta V(Q^2) - g^{\mu\nu} P \cdot Q A_0(Q^2) + P^\mu P^\nu A_+(Q^2) + Q^\mu P^\nu A_-(Q^2)], \quad (7)$$

where the Fermi coupling constant $G_F = 0.16637 \times 10^{-5} \text{ GeV}^{-2}$, the CKM matrix elements $|V_{cb}| = 0.0405$, $|V_{ud}| = 0.974$ [54], and $a_1 = c_1 + \frac{1}{N_c} c_2$ with $N_c = 3$ to be the number of color. In the present work, we take $a_1 = 1.14$ [46]. The decay constant of ρ meson is taken as $f_\rho = 221 \text{ MeV}$ [55,56].

As shown in Fig. 1, the initial B_c and final $Z_c^{(\prime)}\pi$ could be connected by meson loops composed by $(c\bar{c})_{J=1}\rho$. In the present work, we estimate the meson loop contribution in an effective Lagrangian approach. Besides the effective

¹ Actually, the expressions of the transition matrix in Refs. [49,50,53] is not exactly the same as Eq. (3). As for $B_c \rightarrow J/\psi$, the transition matrix defined in Refs. [49,50] is,

$$\langle J/\psi(p_2, \epsilon^*) | \mathcal{J}^\mu | B_c(p) \rangle = \frac{\epsilon_v^*}{m_{B_c} + m_{J/\psi}} [i\epsilon^{\mu\nu\alpha\beta} P_\alpha Q_\beta V(Q^2) - g^{\mu\nu}(P \cdot Q)A_0(Q^2) + P^\mu P^\nu A_+(Q^2) + Q^\mu P^\nu A_-(Q^2)], \quad (4)$$

and the one of $B_c \rightarrow h_c$ presented in Ref. [53] is,

$$\langle h_c(p_2, \epsilon^*) | \mathcal{J}^\mu | B_c(p) \rangle = \frac{1}{m_{B_c} - m_{h_c}} \epsilon^{\mu\nu\alpha\beta} \epsilon_v^* P_\alpha Q_\beta V(Q^2) - i(m_{B_c} - m_{h_c}) \left(\epsilon^{\mu\nu} - \frac{\epsilon^* \cdot P}{Q^2} Q^\mu \right) A_+(Q^2) - i \left[-\frac{\epsilon^* \cdot P}{m_{B_c} - m_{h_c}} P^\mu + (m_{B_c} + m_{h_c}) \frac{\epsilon^* \cdot P}{Q^2} Q^\mu \right] A_-(Q^2) - i 2 m_{h_c} \frac{\epsilon^* \cdot P}{Q^2} Q^\mu A_0(Q^2). \quad (5)$$

One can find the expressions of the transition matrix in Eqs. (4) and (5) can be reduced to the form of Eq. (3) by absorbing some constant factors into the form factors.

interactions of $B_c \rightarrow (c\bar{c})_{J=1}\rho$ as shown in Eq. (7), we still need the effective Lagrangian for $\rho\pi\pi$, which is [57,58]

$$\mathcal{L}_{\rho\pi\pi} = -i g_{\rho\pi\pi} \rho_\mu^+ [\pi^0 \partial^\mu \pi^+ - \partial^\mu \pi^0 \pi^+], \quad (8)$$

where the coupling constants $g_{\rho\pi\pi} = 6.05$ [55], which is determined from the $\rho \rightarrow \pi\pi$ partial width.

The J^P quantum numbers of $Z_c^{(\prime)}$ are 1^+ , thus Z_c couples to $J/\psi\pi$ via S -wave, while Z_c' couples to $h_c\pi$ via P -wave. Considering the chiral symmetry, the effective Lagrangian related to $Z_c^{(\prime)}$ are [39,59,60]

$$\begin{aligned} \mathcal{L}_{Z_c\psi\pi} &= g_{Z_c\psi\pi} \partial_\mu \psi \partial^\mu \pi Z_c^\nu - \partial^\nu \pi Z_c^\mu, \\ \mathcal{L}_{Z_c'h_c\pi} &= g_{Z_c'h_c\pi} \epsilon_{\mu\nu\alpha\beta} \partial^\mu h_c^\nu Z_c'^{\alpha} \partial^\beta \pi + \text{H.c.}, \end{aligned} \quad (9)$$

respectively. The coupling constants in the Lagrangian will be discussed in the next section.

The amplitude of $B_c^+(p) \rightarrow \rho^+(p_1)J/\psi(p_2)[\pi^+(q)] \rightarrow \pi^0(p_3)Z_c^+(p_4)$ corresponding to Fig. 1a is

$$\begin{aligned} \mathcal{M}_a &= (i)^3 \int \frac{d^4 q}{(2\pi)^4} \mathcal{A}^{\mu\nu}(P, Q) [g_{\rho\pi\pi}(q - p_3)^\xi] \\ &\quad \left[-g_{Z_c\psi\pi} p_{2\rho} (q^\rho \epsilon_4^\sigma - q^\sigma \epsilon_4^\rho) \right] \frac{-g_{\mu\xi} + p_{1\mu} p_{1\xi} / m_\rho^2}{p_1^2 - m_\rho^2} \\ &\quad \frac{-g_{\nu\sigma} + p_{2\nu} p_{2\sigma} / m_\psi^2}{p_2^2 - m_\psi^2} \frac{1}{q^2 - m_\pi^2} \mathcal{F}(q^2, m_\pi^2), \end{aligned} \quad (10)$$

and in a similar way, one can get the amplitude of $B_c^+(p) \rightarrow \rho^+(p_1)h_c(p_2)[\pi^+(q)] \rightarrow \pi^0(p_3)Z_c'^+(p_4)$ corresponding to corresponding to Fig. 1b, which is,

$$\begin{aligned} \mathcal{M}_b &= (i)^3 \int \frac{d^4 q}{(2\pi)^4} \mathcal{A}^{\mu\nu}(P, Q) [g_{\rho\pi\pi}(q - p_3)^\xi] \\ &\quad \left[-g_{Z_c'h_c\pi} \epsilon^{\rho\sigma\tau\lambda} \epsilon_{4\rho} p_{2\tau} q_\lambda \right] \frac{-g_{\mu\xi} + p_{1\mu} p_{1\xi} / m_\rho^2}{p_1^2 - m_\rho^2} \\ &\quad \frac{-g_{\nu\sigma} + p_{2\nu} p_{2\sigma} / m_{h_c}^2}{p_2^2 - m_{h_c}^2} \frac{1}{q^2 - m_\pi^2} \mathcal{F}(q^2, m_\pi^2), \end{aligned} \quad (11)$$

where $\mathcal{F}(q^2, m_\pi^2)$ is a form factor introduced to depict the structure effects of the exchanged pion meson. In the present work, a monopole form factor is adopted, which is,

$$\mathcal{F}(q^2, m_\pi^2) = \frac{m_\pi^2 - \Lambda^2}{q^2 - \Lambda^2}, \quad (12)$$

where Λ is a parameter, which is of order 1 GeV [61–64].

3 Numerical results and discussion

Before we estimate the partial width of $B_c^+ \rightarrow Z_c^{(\prime)+}\pi^0$, we have to discuss the coupling constants $g_{Z_c J/\psi\pi}$ and $g_{Z_c' h_c\pi}$. Experimentally, Z_c dominantly decay into $D^* \bar{D} + c.c$ and $J/\psi\pi$, and the ratio of these two channels are measured to be $6.2 \pm 1.1 \pm 2.7$. With the center value of Z_c width

Table 1 The values of the parameters $F(0)$, a and b in the form factors of $B_c \rightarrow J/\psi$ and $B_c \rightarrow h_c$ [50, 53]. Here, one should notice that these parameters correspond to the definitions in Eqs. (4) and (5)

$(c\bar{c})$	Parameters	A_0	A_+	A_-	V
J/ψ	$F(0)$	1.65	0.55	-0.87	0.78
	a	1.19	1.68	1.85	1.82
	b	0.17	0.70	0.91	0.87
h_c	$F(0)$	0.64	0.50	-0.32	0.07
	a	1.92	1.54	2.63	2.32
	b	0.39	0.24	0.63	0.49

and the assumption that the width of Z_c comes from the partial widths of $D^*\bar{D}$ and $\pi J/\psi$, we can get the partial width of $Z_c \rightarrow \pi J/\psi$, and then the estimated coupling constant $g_{Z_c J/\psi \pi}$ is

$$g_{Z_c J/\psi \pi} = 0.47 \text{ GeV}^{-1}. \quad (13)$$

In a similar way, one can estimate the coupling constant $g_{Z_c' h_c \pi}$. We suppose Z_c' dominantly decay into $D^*\bar{D}^*$ and $h_c \pi$ and the ratio of the partial widths of these two decay modes were measured to be 12.0 ± 3.3 [5, 6], then with the center values of the width and the ratio, we can get the partial width of $Z_c' \rightarrow h_c \pi$ and the corresponding coupling constant $g_{Z_c' h_c \pi}$, which is

$$g_{Z_c' h_c \pi} = 0.65 \text{ GeV}^{-1}. \quad (14)$$

In the weak interaction vertex, there are four form factors, which are $V(Q^2)$, $A_{0,\pm}(Q^2)$. As indicated in Ref. [65], the form factors are usually estimated in the quark model, thus, the form factors are known only in spacelike region. Thus, one need to analytically continue the form factors to the timelike region, where the physical decay processes are relevant. In Ref. [50], the form factors for $B_c \rightarrow J/\psi$ are parameterized as the form

$$F(Q^2) = \frac{F(0)}{1 - a\zeta + b\zeta^2}. \quad (15)$$

As for $B_c \rightarrow h_c$, the form factors are parameterized as [53].

$$F(Q^2) = F(0) \exp(a\zeta + b\zeta^2), \quad (16)$$

with $\zeta = Q^2/m_{B_c}^2$ and $F(0)$, a and b are parameters. For completeness of the present work, we collect the values of these parameters in Table 1.

In order to avoid ultraviolet divergence in the loop integrals and evaluate the loop integrals with Feynman parameterization methods, we further parameterize the form factors in the form

$$F(Q^2) = -F(0) \frac{\Lambda_1^2}{Q^2 - \Lambda_1^2} \frac{\Lambda_2^2}{Q^2 - \Lambda_2^2} \frac{\Lambda_3^2}{Q^2 - \Lambda_3^2}, \quad (17)$$

Table 2 Values of parameters Λ_1 , Λ_2 and Λ_3

$(c\bar{c})$	Parameter	A_0	A_+	A_-	V
J/ψ	Λ_1	10.0	9.05	7.35	7.69
	Λ_2	7.30	6.13	6.34	6.24
	Λ_3	17.0	15.9	16.3	16.2
h_c	Λ_1	7.02	7.88	6.05	6.43
	Λ_2	7.02	7.88	6.05	6.41
	Λ_3	14.0	14.0	14.0	14.0

Fig. 2 Branching ratios of $B_c \rightarrow Z_c \pi$ (right panel) in unit of 10^{-4} and $B_c \rightarrow Z_c' \pi$ (left panel) in unit of 10^{-7} depending on parameter Λ

where the values of Λ_1 , Λ_2 and Λ_3 are obtained by fitting Eqs. (15) and (16) with Eq. (17) and the fitted parameter values are list in Table 2.

In the present work, there is only one model parameter, Λ , which is introduced by the form factor in the amplitudes and generally, the magnitude of Λ is of order 1 GeV. In Fig. 2, we present the Λ dependence of the branching ratio of $B_c^+ \rightarrow Z_c^{(\prime)+} \pi^0$. Here, we varies Λ from 1 GeV to 1.5 GeV, and in this parameter range, we find the branching ratio of $B_c^+ \rightarrow Z_c^+ \pi^0$ monotonously increases with the increasing of Λ . In particular, the branching ratio is predicted to be $(1.71 - 6.37) \times 10^{-4}$, which is large enough to be detected in further exclusive B_c decays. This is comparable to the branching ratio of $B_c \rightarrow X(3872)\pi$ in a model calculation [66, 67]. As for $B_c^+ \rightarrow Z_c'^+ \pi^0$, the branching ratio is estimated to be $(2.34 - 5.92) \times 10^{-7}$, which is about 3 order smaller than the one of $B_c^+ \rightarrow Z_c^+ \pi^0$.

Our estimation indicates a strong suppression of Z_c' production from B_c decay comparing to the case of Z_c production. This suppression is partially resulted from the P -wave coupling of $Z_c' h_c \pi$. Further more, in the estimation of $B_c \rightarrow Z_c \pi$, the term related to $V(Q^2)$ in Eq. (7) vanishes after performing the loop integral in Eq. (10). While for $B_c \rightarrow Z_c' \pi$, only the term related to $V(Q^2)$

survive, which further suppress Z'_c production from B_c^+ decay.

4 Summary

After the observations of $Z_c^{(\prime)}$ by BES III and Belle Collaborations, the specific properties of these two charmonium-like states have stimulated great interests of theorist and experimentist to reveal their nature. Addition to mass spectrum and decays of the charmonium-like states, searching more production process are also interesting. Besides the electron-positron annihilation, the signal of Z_c have been observed in the semi-exclusive b -flavored hadron. However, the charmonium-like state Z_c were not observed in the B meson decay, such as $B \rightarrow KJ/\psi\pi\pi$ and $B \rightarrow KD^*\bar{D}$. Thus, searching for the source of Z_c production in the semi-inclusive b -flavor hadron decay will be intriguing.

In present work, we investigate $Z_c^{(\prime)}$ production from B_c^+ decay. We find that production process can occur via $(c\bar{c})_{J=1\rho}$ meson loop, which could enhance the rates of $B_c \rightarrow Z_c^{(\prime)}\pi$ since all the internal mesons could be on-shell. Our estimation shows that the branching ratio of $B_c^+ \rightarrow Z_c^+\pi^0$ is of order of 10^{-4} , which indicates that Z_c production from B_c decay could be a important source of Z_c production from the b -flavored hadron semi-exclusive decay. Such a large branching fraction can also be tested by the exclusive decay of B_c^+ by LHCb. As for Z'_c , we find the branching ratio of $B_c^+ \rightarrow Z_c'^+\pi^0$ is three orders smaller than the one of $B_c^+ \rightarrow Z_c^+\pi^0$, which indicate strong suppression of Z'_c production from B_c decay.

Acknowledgements This work is supported in part by the National Natural Science Foundation of China (NSFC) under Grant nos. 11775050, 11675091, and 11835015.

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: This is a theoretical research work, so no additional data are associated with this work.]

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

References

1. M. Ablikim et al. [BESIII Collaboration], Phys. Rev. Lett. **110**, 252001 (2013). <https://doi.org/10.1103/PhysRevLett.110.252001>. arXiv:1303.5949 [hep-ex]
2. Z.Q. Liu et al. [Belle Collaboration], Phys. Rev. Lett. **110**, 252002 (2013). <https://doi.org/10.1103/PhysRevLett.110.252002>. arXiv:1304.0121 [hep-ex]
3. M. Ablikim et al. [BESIII Collaboration], Phys. Rev. Lett. **112**(2), 022001 (2014). <https://doi.org/10.1103/PhysRevLett.112.022001>. arXiv:1310.1163 [hep-ex]
4. M. Ablikim et al. [BESIII Collaboration], Phys. Rev. Lett. **119**(7), 072001 (2017). <https://doi.org/10.1103/PhysRevLett.119.072001>. arXiv:1706.04100 [hep-ex]
5. M. Ablikim et al. [BESIII Collaboration], Phys. Rev. Lett. **111**(24), 242001 (2013). <https://doi.org/10.1103/PhysRevLett.111.242001>. arXiv:1309.1896 [hep-ex]
6. M. Ablikim et al. [BESIII Collaboration], Phys. Rev. Lett. **112**(13), 132001 (2014). <https://doi.org/10.1103/PhysRevLett.112.132001>. arXiv:1308.2760 [hep-ex]
7. C.F. Qiao, L. Tang, Eur. Phys. J. C **74**, 2810 (2014). <https://doi.org/10.1140/epjc/s10052-014-2810-x>. arXiv:1308.3439 [hep-ph]
8. Z.G. Wang, T. Huang, Phys. Rev. D **89**(5), 054019 (2014). <https://doi.org/10.1103/PhysRevD.89.054019>. arXiv:1310.2422 [hep-ph]
9. Z.G. Wang, Commun. Theor. Phys. **63**(4), 466 (2015). <https://doi.org/10.1088/0253-6102/63/4/466>. arXiv:1312.1537 [hep-ph]
10. Z.G. Wang, Eur. Phys. J. C **74**(5), 2874 (2014). <https://doi.org/10.1140/epjc/s10052-014-2874-7>. arXiv:1311.1046 [hep-ph]
11. S.S. Agaev, K. Azizi, H. Sundu, Phys. Rev. D **96**(3), 034026 (2017). <https://doi.org/10.1103/PhysRevD.96.034026>. arXiv:1706.01216 [hep-ph]
12. L. Zhao, W.Z. Deng, S.L. Zhu, Phys. Rev. D **90**(9), 094031 (2014). <https://doi.org/10.1103/PhysRevD.90.094031>. arXiv:1408.3924 [hep-ph]
13. L. Maiani, V. Riquer, R. Faccini, F. Piccinini, A. Pilloni, A.D. Polosa, Phys. Rev. D **87**(11), 111102 (2013). <https://doi.org/10.1103/PhysRevD.87.111102>. arXiv:1303.6857 [hep-ph]
14. R. Zhu, Phys. Rev. D **94**(5), 054009 (2016). <https://doi.org/10.1103/PhysRevD.94.054009>. arXiv:1607.02799 [hep-ph]
15. C. Deng, J. Ping, F. Wang, Phys. Rev. D **90**, 054009 (2014). <https://doi.org/10.1103/PhysRevD.90.054009>. arXiv:1402.0777 [hep-ph]
16. M.N. Anwar, J. Ferretti, E. Santopinto, Phys. Rev. D **98**(9), 094015 (2018). <https://doi.org/10.1103/PhysRevD.98.094015>. arXiv:1805.06276 [hep-ph]
17. S. Patel, M. Shah, P.C. Vinodkumar, Eur. Phys. J. A **50**, 131 (2014). <https://doi.org/10.1140/epja/i2014-14131-9>. arXiv:1402.3974 [hep-ph]
18. X. Liu, Z.G. Luo, Y.R. Liu, S.L. Zhu, Eur. Phys. J. C **61**, 411 (2009). <https://doi.org/10.1140/epjc/s10052-009-1020-4>. arXiv:0808.0073 [hep-ph]
19. Y.R. Liu, X. Liu, W.Z. Deng, S.L. Zhu, Eur. Phys. J. C **56**, 63 (2008). <https://doi.org/10.1140/epjc/s10052-008-0640-4>. arXiv:0801.3540 [hep-ph]
20. Z.F. Sun, Z.G. Luo, J. He, X. Liu, S.L. Zhu, Chin. Phys. C **36**, 194 (2012). <https://doi.org/10.1088/1674-1137/36/3/002>
21. J. He, X. Liu, Z.F. Sun, S.L. Zhu, Eur. Phys. J. C **73**(11), 2635 (2013). <https://doi.org/10.1140/epjc/s10052-013-2635-z>. arXiv:1308.2999 [hep-ph]
22. W. Chen, T.G. Steele, M.L. Du, S.L. Zhu, Eur. Phys. J. C **74**(2), 2773 (2014). <https://doi.org/10.1140/epjc/s10052-014-2773-y>. arXiv:1308.5060 [hep-ph]
23. T. Gutsche, M. Kesenheimer, V.E. Lyubovitskij, Phys. Rev. D **90**(9), 094013 (2014). <https://doi.org/10.1103/PhysRevD.90.094013>. arXiv:1410.0259 [hep-ph]
24. D.Y. Chen, Y.B. Dong, Phys. Rev. D **93**(1), 014003 (2016). <https://doi.org/10.1103/PhysRevD.93.014003>. arXiv:1510.00829 [hep-ph]
25. G. Li, Eur. Phys. J. C **73**(11), 2621 (2013). <https://doi.org/10.1140/epjc/s10052-013-2621-5>. arXiv:1304.4458 [hep-ph]

26. G. Li, X.H. Liu, Z. Zhou, Phys. Rev. D **90**(5), 054006 (2014). <https://doi.org/10.1103/PhysRevD.90.054006>. arXiv:1409.0754 [hep-ph]
27. C.J. Xiao, D.Y. Chen, Y.B. Dong, W. Zuo, T. Matsuki. arXiv:1811.04688 [hep-ph]
28. Q. Wu, G. Li, F. Shao, R. Wang, Phys. Rev. D **94**(1), 014015 (2016). <https://doi.org/10.1103/PhysRevD.94.014015>
29. Q.R. Gong, Z.H. Guo, C. Meng, G.Y. Tang, Y.F. Wang, H.Q. Zheng, Phys. Rev. D **94**(11), 114019 (2016). <https://doi.org/10.1103/PhysRevD.94.114019>. arXiv:1604.08836 [hep-ph]
30. J. He, D.Y. Chen, Eur. Phys. J. C **78**(2), 94 (2018). <https://doi.org/10.1140/epjc/s10052-018-5580-z>. arXiv:1712.05653 [hep-ph]
31. X.H. Liu, G. Li, Phys. Rev. D **88**, 014013 (2013). <https://doi.org/10.1103/PhysRevD.88.014013>. arXiv:1306.1384 [hep-ph]
32. E.S. Swanson, Phys. Rev. D **91**(3), 034009 (2015). <https://doi.org/10.1103/PhysRevD.91.034009>. arXiv:1409.3291 [hep-ph]
33. D.Y. Chen, X. Liu, Phys. Rev. D **84**, 034032 (2011). <https://doi.org/10.1103/PhysRevD.84.034032>. arXiv:1106.5290 [hep-ph]
34. D.Y. Chen, X. Liu, T. Matsuki, Phys. Rev. D **88**(3), 036008 (2013). <https://doi.org/10.1103/PhysRevD.88.036008>. arXiv:1304.5845 [hep-ph]
35. Y. Ikeda [HAL QCD Collaboration], J. Phys. G **45**(2), 024002 (2018). <https://doi.org/10.1088/1361-6471/aa9afd>. arXiv:1706.07300 [hep-lat]
36. Y. Ikeda et al. [HAL QCD Collaboration], Phys. Rev. Lett. **117**(24), 242001 (2016). <https://doi.org/10.1103/PhysRevLett.117.242001>. arXiv:1602.03465 [hep-lat]
37. D.Y. Chen, Y.B. Dong, M.T. Li, W.L. Wang, Eur. Phys. J. A **52**(10), 310 (2016). <https://doi.org/10.1140/epja/i2016-16310-0>
38. Q.Y. Lin, X. Liu, H.S. Xu, Phys. Rev. D **88**, 114009 (2013). <https://doi.org/10.1103/PhysRevD.88.114009>. arXiv:1308.6345 [hep-ph]
39. Y. Huang, J. He, X. Liu, H.F. Zhang, J.J. Xie, X.R. Chen, Phys. Rev. D **93**(3), 034022 (2016). <https://doi.org/10.1103/PhysRevD.93.034022>. arXiv:1512.00981 [hep-ph]
40. V.M. Abazov et al. [D0 Collaboration], Phys. Rev. D **98**(5), 052010 (2018). <https://doi.org/10.1103/PhysRevD.98.052010>. arXiv:1807.00183 [hep-ex]
41. S.K. Choi et al. [Belle Collaboration], Phys. Rev. Lett. **91**, 262001 (2003). <https://doi.org/10.1103/PhysRevLett.91.262001>. arXiv:hep-ex/0309032
42. T. Aushev et al. [Belle Collaboration], Phys. Rev. D **81**, 031103 (2010). <https://doi.org/10.1103/PhysRevD.81.031103>. arXiv:0810.0358 [hep-ex]
43. Z. Rui, Z. Zhitian, C.D. Lu, Phys. Rev. D **86**, 074019 (2012). <https://doi.org/10.1103/PhysRevD.86.074019>. arXiv:1203.2303 [hep-ph]
44. V.V. Kiselev. arXiv:hep-ph/0211021
45. M.A. Ivanov, J.G. Korner, O.N. Pakhomova, Phys. Lett. B **555**, 189 (2003). [https://doi.org/10.1016/S0370-2693\(03\)00052-2](https://doi.org/10.1016/S0370-2693(03)00052-2). arXiv:hep-ph/0212291
46. M.A. Ivanov, J.G. Korner, P. Santorelli, Phys. Rev. D **73**, 054024 (2006). <https://doi.org/10.1103/PhysRevD.73.054024>. arXiv:hep-ph/0602050
47. R. Aaij et al. [LHCb Collaboration], Nucl. Phys. B **930**, 563 (2018). <https://doi.org/10.1016/j.nuclphysb.2018.03.015>. arXiv:1712.04702 [hep-ex]
48. C.H. Chang, Y.Q. Chen, G.L. Wang, H.S. Zong, Phys. Rev. D **65**, 014017 (2002). <https://doi.org/10.1103/PhysRevD.65.014017>. arXiv:hep-ph/0103036
49. W. Wang, Y.L. Shen, C.D. Lu, Phys. Rev. D **79**, 054012 (2009). <https://doi.org/10.1103/PhysRevD.79.054012>. arXiv:0811.3748 [hep-ph]
50. A. Issadykov, M.A. Ivanov, Phys. Lett. B **783**, 178 (2018). <https://doi.org/10.1016/j.physletb.2018.06.056>. arXiv:1804.00472 [hep-ph]
51. Z. Rui, Phys. Rev. D **97**(3), 033001 (2018). <https://doi.org/10.1103/PhysRevD.97.033001>. arXiv:1712.08928 [hep-ph]
52. Z. Rui, Z.T. Zou, Phys. Rev. D **90**(11), 114030 (2014). <https://doi.org/10.1103/PhysRevD.90.114030>. arXiv:1407.5550 [hep-ph]
53. X.X. Wang, W. Wang, C.D. Lu, Phys. Rev. D **79**, 114018 (2009). <https://doi.org/10.1103/PhysRevD.79.114018>. arXiv:0901.1934 [hep-ph]
54. C. Patrignani et al. [Particle Data Group], Chin. Phys. C **40**(10), 100001 (2016). <https://doi.org/10.1088/1674-1137/40/10/100001>
55. H.Y. Cheng, C.K. Chua, A. Soni, Phys. Rev. D **71**, 014030 (2005). <https://doi.org/10.1103/PhysRevD.71.014030>. arXiv:hep-ph/0409317
56. C.D. Lu, Y.L. Shen, W. Wang, Phys. Rev. D **73**, 034005 (2006). <https://doi.org/10.1103/PhysRevD.73.034005>. arXiv:hep-ph/0511255
57. T.M. Yan, H.Y. Cheng, C.Y. Cheung, G.L. Lin, Y.C. Lin, H.L. Yu, Phys. Rev. D **46**, 1148 (1992). Erratum: [Phys. Rev. D **55**, 5851 (1997)]. <https://doi.org/10.1103/PhysRevD.46.1148>, <https://doi.org/10.1103/PhysRevD.55.5851>
58. R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, G. Nardulli, Phys. Rep. **281**, 145 (1997). [https://doi.org/10.1016/S0370-1573\(96\)00027-0](https://doi.org/10.1016/S0370-1573(96)00027-0). arXiv:hep-ph/9605342
59. X.H. Liu, Q. Zhao, F.E. Close, Phys. Rev. D **77**, 094005 (2008). <https://doi.org/10.1103/PhysRevD.77.094005>. arXiv:0802.2648 [hep-ph]
60. D.Y. Chen, C.J. Xiao, J. He, Phys. Rev. D **96**(5), 054017 (2017). <https://doi.org/10.1103/PhysRevD.96.054017>
61. N.A. Tornqvist, Nuovo Cim. A **107**, 2471 (1994)
62. N.A. Tornqvist, Z. Phys. C **61**, 525 (1994)
63. M.P. Locher, Y. Lu, B.S. Zou, Z. Phys. A **347**, 281 (1994)
64. X.Q. Li, D.V. Bugg, B.S. Zou, Phys. Rev. D **55**, 1421 (1997)
65. H.Y. Cheng, C.K. Chua, C.W. Hwang, Phys. Rev. D **69**, 074025 (2004). <https://doi.org/10.1103/PhysRevD.69.074025>. arXiv:hep-ph/0310359
66. W. Wang, Y.L. Shen, C.D. Lu, Eur. Phys. J. C **51**, 841 (2007). <https://doi.org/10.1140/epjc/s10052-007-0334-3>. arXiv:0704.2493 [hep-ph]
67. W. Wang, Q. Zhao, Phys. Lett. B **755**, 261 (2016). <https://doi.org/10.1016/j.physletb.2016.02.012>. arXiv:1512.03123 [hep-ph]