



NNLO final-state quark-pair corrections in four dimensions

B. Page^{1,2,a}, R. Pittau^{3,b}

¹ Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, 79104 Freiburg, Germany

² Present Address: Institut de Physique Théorique, CEA, CNRS, Université Paris-Saclay, 91191 Gif-sur-Yvette Cedex, France

³ Departamento de Física Teórica y del Cosmos and CAFPE, Universidad de Granada, Campus Fuentenueva s.n., 18071 Granada, Spain

Received: 21 January 2019 / Accepted: 8 April 2019 / Published online: 25 April 2019
© The Author(s) 2019

Abstract We describe how NNLO final state quark-pair corrections are computed in FDR by directly enforcing gauge invariance and unitarity in the definition of the regularized divergent integrals. We give details of our approach and show how virtual and real contributions can be merged together without relying, explicitly or implicitly, on dimensional regularization. As an example, we recompute the $H \rightarrow b\bar{b} + jets$ and $\gamma^* \rightarrow jets$ inclusive rates at the NNLO accuracy in the large N_F limit of QCD. This demonstrates, for the first time, that physical results with intermediate infrared singularities can be obtained at NNLO using a fully four-dimensional procedure.

1 Introduction

It is well known that calculations of observables in quantum field theory (QFT) are complicated by divergences in intermediate steps. On general grounds these divergences fall into two categories – ultraviolet (UV) divergences, associated with short wavelengths, and infrared (IR) divergences, associated with large wavelengths and/or collinear configurations. They show up in the form of integrals which do not exist in the four-dimensional physical space.

The customary way to deal with UV infinities is a two step procedure. Firstly, one regularizes the divergent integrals. Secondly, one reabsorbs the UV pieces into the free parameters of the Lagrangian. As for the IR infinities, they cancel when considering sufficiently inclusive observables, or can be reabsorbed in the parton densities. Depending on the approach, this cancellation can be achieved before or after integration. In the latter case, IR divergent integrals also need to be regulated.

When performing calculations of observables in QFT, the exact method of regulating the UV/IR divergences is arbi-

trary. Nevertheless, this freedom is not absolute as the chosen method should not interfere with two core tenets of QFT, that are

- gauge invariance; (1a)
- unitarity. (1b)

These general principles have concrete consequences in perturbative calculations. Gauge invariance implies a set of graphical identities (see e.g. [1]) that need to be fulfilled to all perturbative orders by the Feynman diagrams of the QFT. Unitarity, meanwhile, demands the validity of the cutting equations [2,3] corresponding to the relation

$$i(T - T^\dagger) = -T^\dagger T \quad (2)$$

for the T matrix [4]. Both (1) are essential to preserve the Ward/Slavnov-Taylor identities (WI) at the regularized level, known as the regularized quantum action principle [5].

The most commonly used technique is dimensional regularization [6,7] (DReg). DReg exploits the fact that gauge invariance and unitarity are naturally preserved as they hold for the theory in all values of the dimensionality d of the space-time. Hence, the divergent integrals are analytically evaluated in d dimensions, and the asymptotic $d \rightarrow 4$ limit is eventually taken. By doing so, UV/IR divergences are parametrized in terms of negative powers of a Laurent expansion in $(d - 4)$. In this framework, one still has some freedom to define objects in intermediate steps. Hence, several variants of DReg exist such as conventional dimensional regularization [8], 'tHooft-Veltman [6], four-dimensional helicity [9] and dimensional reduction [10]. We refer to [11] for an exact definition of all of them.

In recent years, a considerable effort has been pursued by several groups to introduce more four-dimensional ingredients in the definition of regularization. The main motivation being an attempt to simplify analytical and numerical methods, as well as to try to consider different theoretical perspectives. This four-dimensional program has resulted in

^a e-mail: bpage@ipht.fr

^b e-mail: pittau@ugr.es

a number of methods such as the four-dimensional formulation of FDH [12], implicit regularization [13,14], four-dimensional unsubtraction [15–17] and FDR [18–22]. They are described and compared in [23]. Whilst all approaches are different, FDR is the only method that does not rely on the customary UV renormalization procedure. Instead, the result of an FDR calculation is directly a renormalized quantity.

FDR treats UV divergences by performing a subtraction, extracting from the loop integrands the divergent parts which do not contain physical information – the so-called *vacua*. In the case of IR finite amplitudes, the validity of the FDR strategy has been explicitly demonstrated in [22]. In the presence of IR divergent configurations, the IR regulator should not interfere with principles (1). If this is achieved, the correct physical result is obtained for IR safe quantities. At NLO, it is known how to match real and one-loop contributions in the presence of final state IR singularities [21,23], while, so far, no FDR NNLO calculation has been performed involving infrared divergent configurations. In this paper we bridge this gap and give the first example of such a computation. We describe how NNLO final state quark-pair corrections can be computed in FDR in a way that automatically respects the principles (1). In particular, we reproduce the $\overline{\text{MS}}$ results for the N_F part of $H \rightarrow b\bar{b} + jets$ and $\gamma^* \rightarrow jets$. Since DReg is never used, explicitly or implicitly, this represents, to our knowledge, the first example of a realistic fully four-dimensional NNLO calculation.

The structure of the paper is as follows. In Sect. 2 we present our definitions of the virtual and real integrals appearing when computing NNLO quark-pair corrections. In Sect. 3 we discuss the relevant renormalization issues. Sections 4 and 5 give a detailed description of the $H \rightarrow b\bar{b} + jets$ and $\gamma^* \rightarrow jets$ calculations. Finally, in Sect. 6 we summarize our findings and discuss perspectives and directions opened by the procedures introduced in this work.

2 Inclusive quark-pair corrections and FDR

Our aim is to compute the large N_F limit of cross sections including NNLO quark-pair corrections. The relevant contributions are the Born, Virtual and Real reactions given by

$$\begin{aligned} \sigma_B &\propto \int d\Phi_n \sum_{\text{spin}} |A_n^{(0)}|^2, \\ \sigma_V &\propto \int d\Phi_n \sum_{\text{spin}} \left\{ A_n^{(2)} (A_n^{(0)})^* + A_n^{(0)} (A_n^{(2)})^* \right\}, \\ \sigma_R &\propto \int d\Phi_{n+2} \sum_{\text{spin}} \left\{ A_{n+2}^{(0)} (A_{n+2}^{(0)})^* \right\}. \end{aligned} \tag{3}$$




Fig. 1 The lowest order amplitude (a), the IR divergent final-state virtual quark-pair correction (b) and the IR divergent real component (c). The blob stands for the emission of $n - 1$ particles. Additional IR finite corrections are created if the gluons with momenta q_1 and k_{34} are emitted by an off-shell particle in the blob

In (3), $A_n^{(j)}$ represents the amplitude for the emission of n partons computed at the j^{th} perturbative order, while $d\Phi_m$ is the m -particle phase-space

$$d\Phi_m := \delta \left(P - \sum_{i=1}^m p_i \right) \prod_{i=1}^m d^4 p_i \delta_+(p_i^2), \tag{4}$$

with $\delta_+(p_i^2) := \delta(p_i^2) \Theta(p_i^0)$, in which P is the initial state momentum. The amplitude $A_n^{(0)}$ is drawn in Fig. 1a, where the line with momentum p is an on-shell QCD parton and the blob denotes $n-1$ additional final-state particles. The NNLO amplitudes can be split into IR divergent and finite parts

$$\begin{aligned} A_n^{(2)} &:= A_{n,\text{IR}}^{(2)} + A_{n,\text{F}}^{(2)}, \\ A_{n+2}^{(0)} &:= A_{n+2,\text{IR}}^{(0)} + A_{n+2,\text{F}}^{(0)}. \end{aligned} \tag{5}$$


The infrared singular contributions are depicted in Fig. 1b, c, while the finite pieces are created when the splitting gluons with momenta q_1 and k_{34} are emitted by an off-shell particle contained in the blob.

Due to the presence of IR parts in (5), both σ_V and σ_R are IR divergent, although, as is well known, their combination in infrared safe quantities is IR finite. In addition, UV infinities are present in σ_V that are renormalized away when bare parameters are determined at the perturbative order appropriate to match the NNLO accuracy. In conclusion, the fully inclusive sum

$$\sigma^{\text{NNLO}} = \sigma_B + \sigma_V + \sigma_R \tag{6}$$

is a physical quantity, although its parts are separately plagued by IR and UV divergences. Indeed, it is the sim-

Fig. 2 The indices $\hat{\rho}$ and $\hat{\sigma}$ are external to the divergent sub-diagram disconnected from the rest



plest case of an infrared safe observable, which must give the same result when computed in any consistent scheme used to deal with the divergences.

In this section we use the four-dimensional FDR framework, and describe the procedures which allow one to compute σ^{NNLO} . We put particular emphasis on the steps needed to cope with the simultaneous presence of IR and UV infinities at two loops, and on how to merge virtual and real components. Section 2.1 presents the steps needed to define σ_V , while 2.2 deals with σ_R . Section 2.3 contains an explicit example which guides the reader across both real and virtual procedures.

2.1 The NNLO definition of the virtual contribution

A generic two-loop integrand in $A_{n,IR}^{(2)}$ has the form

$$J(q_1, q_2) = \frac{F_{\hat{\rho}\hat{\sigma}}(q_1, q_1^2)}{D} \frac{q_2^\rho q_{12}^\sigma + q_{12}^\rho q_2^\sigma - g^{\rho\sigma} (q_2 \cdot q_{12})}{q_2^2 q_{12}^2} := \frac{N}{Dq_2^2 q_{12}^2}, \tag{7}$$

with internal loop momenta q_1, q_2 and $q_{12} := q_1 + q_2$. The denominator D collects all q_2 -independent propagators

$$D = D_p q_1^4 \left(\prod_{i=1}^k D_i \right), \quad D_p = (q_1 + p)^2, \tag{8}$$

where k is the number of propagators in the blob of Fig. 1b, and N is the numerator of the integrand. The hats on Lorentz indices means that they are external to the divergent sub-diagram, as in Fig. 2. The difference between hatted and un-hatted indices can be ignored until equation (23). Here we just point out that hatted indices can be identified in more general cases as described in [22].

We now analyze all possible divergences generated upon loop integration. We do this to determine the form of the FDR UV subtractions and the structure of the IR regulator needed to define the FDR integration over (7) given in (35). $J(q_1, q_2)$ is quadratically divergent when $q_2 \rightarrow \infty$ at fixed q_1 . Given the presence of many propagators in D , this is the only possible UV sub-divergence. Depending on where the lower gluon reconnects to the blob, it may also generate global UV infinities. In addition, due to the on-shell condition $p^2 = 0$, a double collinear IR singularity arises when both q_1 and q_2 are proportional to p . A second potential IR divergent configuration is $q_1^2 \rightarrow 0$ but $q_2^2 \neq 0$. Nevertheless, this divergence is cancelled by the UV behavior of the q_2 integration.

This can be understood by noticing that a scaleless q_2 -type sub-integral is generated in this case, that vanishes in FDR.¹ An additional double collinear IR singularity is created if the lower gluon is attached to a second external massless parton.

In FDR, a common unphysical scale μ^2 is added to all propagators in order to regulate divergences. As is well known, performing this operation only in the denominator is contrary to the core principle (1a) of gauge invariance. As such, in FDR one performs a *Global Prescription* (GP) [18], also making the replacement in the numerator such that all integrand cancellations between numerator and denominator take place at the regulated level.² These cancellations are called ‘‘gauge cancellations’’. In practice, one first determines the dependence of $J(q_1, q_2)$ on q_1^2, q_2^2 and q_{12}^2 generated by Feynman rules. This is the reason for the explicit q_1^2 as an argument of F in (7). Subsequently, one adds an unphysical scale μ^2 to all such self-contractions

$$q_i^2 \rightarrow q_i^2 - \mu^2 := \bar{q}_i^2 \quad (i = 1, 2, 12). \tag{9}$$

We denote this procedure by the symbol \xrightarrow{GP} . Therefore, the replacement $f \xrightarrow{GP} \bar{f}$, applied to any function $f(q_1^2, q_2^2, q_{12}^2)$, produces a new function \bar{f} with arguments replaced as in (9),

$$\bar{f}(\bar{q}_1^2, \bar{q}_2^2, \bar{q}_{12}^2) := f(\bar{q}_1^2, \bar{q}_2^2, \bar{q}_{12}^2). \tag{10}$$

Note that \bar{f} has the same functional form as f . Nevertheless, we keep the bar to make explicit that GP has been implemented on f . In the case at hand one has

$$J(q_1, q_2) \xrightarrow{GP} \bar{J}(q_1, q_2) := \frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \bar{G}^{\rho\sigma} = \frac{\bar{N}}{\bar{D}\bar{q}_2^2 \bar{q}_{12}^2}, \tag{11}$$

with

$$\bar{D} := \bar{D}_p \bar{q}_1^4 \left(\prod_{i=1}^k \bar{D}_i \right), \quad \bar{D}_{p,i} := D_{p,i} - \mu^2, \tag{12}$$

and

$$\bar{G}^{\rho\sigma} := \frac{q_2^\rho q_{12}^\sigma + q_{12}^\rho q_2^\sigma - g^{\rho\sigma} (\bar{q}_{12}^2 + \bar{q}_2^2 - \bar{q}_1^2)/2}{\bar{q}_2^2 \bar{q}_{12}^2}. \tag{13}$$

In our calculation, we simplify the \bar{q}_i^2 s in \bar{N} with the denominators of (11),

$$\frac{\bar{q}_i^2}{\bar{q}_i^2} = 1, \quad \frac{\bar{q}_1^2}{\bar{D}_p} = 1 - 2 \frac{(q_1 \cdot p)}{\bar{D}_p}. \tag{14}$$

After this is done, μ^2 only appears in the propagators.

The IR singularities of $J(q_1, q_2)$ are now regulated by the addition of μ^2 in the propagators. As we shall see, the asymptotic limit $\mu^2 \rightarrow 0$ will be eventually taken after integration. That generates logarithms of μ^2 of IR origin. Nevertheless

¹ See Appendix B.

² This, combined with the shift invariance of FDR integrals is sufficient to prove all Ward identities graphically [20].

$\bar{J}(q_1, q_2)$ is still UV divergent. The global UV infinities are subtracted by separating physical and non-physical scales in \bar{D}_p . That means using the identity

$$\frac{1}{\bar{D}_p} = \frac{1}{\bar{q}_1^2} - \frac{2(q_1 \cdot p)}{\bar{q}_1^2 \bar{D}_p}, \tag{15}$$

and noticing that the second term is more UV convergent than the original propagator. The same expansion has to be applied to the other propagators in $1/\bar{D}$, until \bar{F}/\bar{D} is written as follows

$$\frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} = \left[\frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \right]_V + \left(\frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \right)_F, \tag{16}$$

where $[\bar{F}/\bar{D}]_V$ do not depend on physical scales. Since also $\bar{G}^{\rho\sigma}$ does not contain physical scales, $[\bar{F}/\bar{D}]_V$ defines the global UV divergent behavior of $\bar{J}(q_1, q_2)$:

$$[\bar{J}(q_1, q_2)]_{GV} := \left[\frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \right]_V \bar{G}^{\rho\sigma}. \tag{17}$$

$[\bar{J}(q_1, q_2)]_{GV}$ is called a *Global Vacuum* (GV) and is written between square brackets, that is the standard FDR notation to indicate the vacuum part of an object. Note that $(\bar{F}/\bar{D})_F$ in (16) gives rise to a subtracted integrand which is globally UV convergent but still divergent when $q_2 \rightarrow \infty$:

$$\left(\frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \right)_F \bar{G}^{\rho\sigma}. \tag{18}$$

This is fixed by subtracting the *Sub-Vacuum* (SV) from $\bar{G}^{\rho\sigma}$ by means of the expansion

$$\frac{1}{\bar{q}_{12}^2} = \frac{1}{\bar{q}_2^2} - \frac{q_1^2 + 2(q_1 \cdot q_2)}{\bar{q}_2^2 \bar{q}_{12}^2}. \tag{19}$$

The final result has the form

$$\bar{G}^{\rho\sigma} = [\bar{G}^{\rho\sigma}]_{SV} + (\bar{G}^{\rho\sigma})_F, \tag{20}$$

so that the fully UV subtracted integrand

$$\left(\frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} - \left[\frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \right]_V \right) (\bar{G}^{\rho\sigma} - [\bar{G}^{\rho\sigma}]_{SV}) \tag{21}$$

is integrable in four dimensions.³ Upon integration, the vacua subtracted in (21) induce the appearance of logarithms of μ^2 of UV origin, so that both IR and UV singularities are regulated by the same regulator.

³ The described procedure is quite general. For example, despite the fact that subtracting sub-divergences via (19) potentially re-introduces UV singularities due to q_1 , the expressions can be arranged in such a way that fully subtracted integrands can be always written down. Indeed, this has been automated in the case of two-loop off-shell QCD amplitudes in [22].

The procedure leading to (21) is conveniently encoded in a linear integral operator

$$\int [d^4 q_1][d^4 q_2], \tag{22}$$

whose action on a two-loop integrand is defined by three subsequent operations:

- subtract the vacua;
- integrate over q_1 and q_2 ;
- take the asymptotic limit $\mu^2 \rightarrow 0$.

The last operation means retaining only the logarithmic pieces in the asymptotic expansion, neglecting $\mathcal{O}(\mu^2)$ terms. Thus, the FDR two-loop integration over $\bar{J}(q_1, q_2)$ in (11) is defined as follows⁴

$$\begin{aligned} \bar{I} &:= \int [d^4 q_1][d^4 q_2] \frac{\bar{N}}{\bar{D}\bar{q}_2^2\bar{q}_{12}^2} \\ &= \int d^4 q_1 d^4 q_2 \left(\frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \right)_F (\bar{G}^{\rho\sigma})_F. \end{aligned} \tag{23}$$

In the following, we will often omit terms that integrate to zero in globally prescribed numerators. Hence, it is convenient to introduce a notation for that, that is $\bar{N}' \simeq \bar{N}$ if both numerators give the same result upon FDR integration.

Equation (23) defines a gauge-invariant object, in which the necessary gauge cancellations are preserved by the GP operation. By ‘‘gauge-invariant object’’ we mean that a calculation of \bar{I} in a different gauge will give the same result. It is instructive to check that no change in \bar{I} is produced if one shifts the numerator of the gluon propagator as

$$g_{\rho\hat{\rho}} \rightarrow g_{\rho\hat{\rho}} + \lambda_1 \frac{q_1^\rho q_1^{\hat{\rho}}}{\bar{q}_1^2}, \quad g_{\sigma\hat{\sigma}} \rightarrow g_{\sigma\hat{\sigma}} + \lambda_2 \frac{q_1^\sigma q_1^{\hat{\sigma}}}{\bar{q}_1^2}, \quad \forall \lambda_{1,2}. \tag{24}$$

Thus, \bar{I} gives the same result when computed in any gauge. Another consequence of the WIs is that the term proportional $q_2^\rho q_1^\sigma + q_1^\rho q_2^\sigma$ in (13) should not contribute to \bar{I} when contracted with $\bar{F}_{\hat{\rho}\hat{\sigma}}$. That is,

$$\int [d^4 q_1][d^4 q_2] \frac{\bar{F}_{\hat{\rho}\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}\bar{q}_2^2\bar{q}_{12}^2} (q_1^\rho q_2^\sigma + q_2^\rho q_1^\sigma) = 0. \tag{25}$$

After taking into account the vanishing of scaleless integrals, this corresponds to the WI depicted in Fig. 3. Nevertheless, we keep this piece in (13), as we will explicitly show in our calculation that it never contributes.

⁴ Due to the vacuum subtraction, $q_i^2/\bar{q}_i^2 \neq 1$ under FDR integration. This is the reason why we perform the explicit gauge cancellations of (14).




Fig. 3 The graphical version of the WI in (25). The scalar gluon, with coupling proportional to the gluon momentum, is denoted by a dashed arrow. The symbol \otimes indicates that it is emitted by the quark loop




Fig. 4 Perturbative expansion of (2) for $H \rightarrow b\bar{b} + jets$ in the large N_F limit

Let us now consider the unitarity properties of this prescription. Equation (2) relates different orders of perturbation theory, so a given divergent subgraph S_G can appear embedded in loop diagrams of different orders ℓ . If the result of S_G depends on ℓ then the relation (2) will in general not survive. For a more concrete example relevant to the calculations of this paper, consider the consequence of (2) shown in Fig. 4. This specifies that the discontinuity of a two-loop graph is given in terms of a one-loop graph. As such, the higher loop regularization must be consistent with the lower loop one. In FDR this places strong constraints on the GP. In our example of Fig. 4 we see that on the left hand side, the momenta associated to the gluon lines take part in the GP, but this is not true for the right hand side. In FDR, this tension is resolved by enforcing “sub-integration consistency” (SIC) [22].

We now consider this procedure in detail for the integrand $J(q_1, q_2)$. We analyze a subset of the full integrand, specifically the numerator of the gluonic self-energy of Fig. 2, $q_2^\rho q_{12}^\sigma + q_{12}^\rho q_2^\sigma - g^{\rho\sigma} (q_2 \cdot q_{12})$, which appears in (7). This is the piece that needs to be treated carefully to maintain SIC. We consider, in particular, the result of the contraction with a $g_{\rho\hat{\sigma}}$ ⁵ (potentially) contained in $F_{\rho\hat{\sigma}}$. It reads

$$N_s(q_1^2, q_2^2, q_{12}^2, \hat{q}_2^2) := q_1^2 - 3q_2^2 - q_{12}^2 + 2\hat{q}_2^2, \tag{26}$$

where we have explicitly separated the contribution $\hat{q}_2^2 := g_{\rho\hat{\sigma}} q_2^\rho q_2^\sigma$. From the point of view of the sub-diagram disconnected from the rest, \hat{q}_2^2 should not be globally prescribed, hence the GP replacement to be performed is

$$N_s(q_1^2, q_2^2, q_{12}^2, \hat{q}_2^2) \rightarrow N_s(q_1^2, \bar{q}_2^2, \bar{q}_{12}^2, q_2^2). \tag{27}$$

⁵ The analysis for terms like $\gamma_\rho \gamma_{\hat{\sigma}}$ is also required, but equivalent.

On the other hand, embedding this in the full diagram requires integrating over q_1 , therefore q_1^2 needs to be barred

$$N_s(q_1^2, q_2^2, q_{12}^2, \hat{q}_2^2) \rightarrow N_s(\bar{q}_1^2, \bar{q}_2^2, \bar{q}_{12}^2, q_2^2). \tag{28}$$

Nevertheless, ρ and σ are internal indices of the two-loop diagrams of Fig. 1b, so that GP is needed for \hat{q}_2^2 as well

$$N_s(q_1^2, q_2^2, q_{12}^2, \hat{q}_2^2) \xrightarrow{\text{GP}} N_s(\bar{q}_1^2, \bar{q}_2^2, \bar{q}_{12}^2, \bar{q}_2^2), \tag{29}$$

which is the prescription used in (23). It is the mismatch among (27)–(29) which violates SIC. Our solution to this problem is modifying the integrand in (23) as follows:

- we do not apply GP to \hat{q}_2^2 terms whose origin is a contraction with indices external to the UV divergent sub-diagram;⁶
- we replace back everywhere $\bar{q}_1^2 \rightarrow q_1^2$ after GV subtraction.

Note that the last operation is possible because barring q_2^2 and q_{12}^2 is sufficient to regulate the IR divergences. This is a consequence of the fact that the only IR divergent configuration is the double collinear limit. Furthermore, this solution does not affect any of the WIs and so still defines a gauge invariant object. In summary, after GV subtraction, (27) is maintained as it is also when embedded in a two-loop calculation. A comparison between this solution and the IR-free case is given in Appendix A.

Let us now return to the matter of enforcing SIC in the entirety of (23), and how one applies our solution. We enforce SIC by rewriting (23) as

$$\bar{I} = \int [d^4 q_1][d^4 q_2] \left(\frac{\bar{F}_{\rho\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \right)_F \bar{G}^{\rho\sigma}(\bar{q}_1^2), \tag{30}$$

where we have used the fact that $[\bar{F}/\bar{D}]_V$ is subtracted by the integral operator. The structure of the expansions needed to extract the GV is such that \bar{D} can be always pulled out from the rest. Thus, it is possible to rewrite

$$\left(\frac{\bar{F}_{\rho\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}} \right)_F = \frac{\bar{H}_{\rho\hat{\sigma}}(q_1, \bar{q}_1^2)}{\bar{D}}. \tag{31}$$

Next, we introduce the numerator function

$$\bar{Z}(\bar{q}_1^2, \hat{q}_2^2) := \bar{H}_{\rho\hat{\sigma}}(q_1, \bar{q}_1^2) \bar{G}^{\rho\sigma}(\bar{q}_1^2) \bar{q}_2^2 \bar{q}_{12}^2, \tag{32}$$

where the explicit dependence on \hat{q}_2^2 is generated by the l.h.s of (27). In practice, $\bar{Z}(\bar{q}_1^2, \hat{q}_2^2)$ is constructed from N in (7) as follows:

- globally prescribe N , $N \xrightarrow{\text{GP}} \bar{N}$, leaving \hat{q}_2^2 unbarred;

⁶ This reproduces the effect of the evanescent operators needed in the four-dimensional helicity scheme and dimensional reduction to restore renormalizability [24].

- perform GV subtraction and determine, for each term T in \bar{N} , the appropriate function \bar{H} to be used in (32). The result of this will always have a factorized form. For instance, if (15) has to be used once to subtract the global vacuum, the contribution of T to $\bar{Z}(\bar{q}_1^2, \hat{q}_2^2)$ is $-2(q_1 \cdot p)/\bar{q}_1^2 \times T$;
- eliminates the bars from the \bar{q}_1^2 s;
- identify \hat{q}_2^2 with q_2^2 .

We denote the last two operations with the symbol $\xrightarrow{\text{SIC}}$. Thus, the change $\bar{f} \xrightarrow{\text{SIC}} \tilde{f}$, applied to any globally prescribed and GV subtracted function $\bar{f}(\bar{q}_1^2, \hat{q}_2^2)$, produces a new function defined as

$$\tilde{f}(q_1^2, q_2^2) := \bar{f}(q_1^2, q_2^2). \tag{33}$$

In the equation above, \tilde{f} has the same functional form as \bar{f} , but we use the tilded notation to denote that the SIC operation has been imposed on \bar{f} . Thus, the SIC compatible version of (32) reads

$$\bar{Z}(\bar{q}_1^2, \hat{q}_2^2) \xrightarrow{\text{SIC}} \tilde{Z}(q_1^2, q_2^2). \tag{34}$$

To continue to ensure gauge cancellations, we unbar also the propagators, that leads to

$$\tilde{I} := \int d^4 q_1 [d^4 q_2] \frac{\tilde{Z}(q_1^2, q_2^2)}{D \bar{q}_2^2 \bar{q}_1^2}. \tag{35}$$

Equation (35) defines the SIC preserving four-dimensional integration over the integrand in (7). Note that, since the GV has been subtracted in it, $[d^4 q_1]$ is replaced by a customary integration $d^4 q_1$. The asymptotic limit $\mu^2 \rightarrow 0$ is understood after taking the two integrations.

A first consequence of this definition is that external wavefunction corrections vanish for massless particles, so that they can always be neglected in actual calculations. The proof is given in Appendix B.

2.2 The NNLO definition of the real component

Given the propagator structure of Fig. 1c, the integrands contributing to σ_R in (3) have the following form

$$J_R = \frac{N_R}{S s_{34}^\alpha s_{134}^\beta}, \quad s_{i\dots j} := (k_i + \dots + k_j)^2, \tag{36}$$

where $0 \leq \alpha, \beta \leq 2$. The symbol S collects all the remaining propagators and N_R is the numerator of the amplitude squared. Depending on the value of the exponents α and β , J_R becomes IR divergent under integration over Φ_{n+2} . These IR singularities must be regulated coherently with our treatment of the virtual component, without violating unitarity and gauge invariance. In this section, we determine a four-dimensional integration that achieves this.




Fig. 5 Virtual and real cuts contributing to the IR divergent parts of σ_V (a, b) and σ_R (c, d). The special indices needed to enforce (27) and (38b) are denoted by $\hat{\rho}$ and $\hat{\sigma}$

Our starting point is the representation of σ_V and σ_R in terms of cut diagrams, in which we put the complex conjugate amplitudes on the right side. With this convention, normal Feynman rules are assumed on the left and complex conjugate ones on the right. The cuts generating IR divergent configurations are obtained by squaring the amplitudes in Fig. 1b, c and are depicted in Fig. 5. IR divergences manifest themselves as pinch singularities of the loop integrals in (a,b) and endpoint phase-space singularities in (c,d). These two kinds of singularities are related to each other by the identity

$$\frac{1}{k^2 + i0^+} = \frac{2\pi}{i} \delta_+(k^2) + \frac{1}{k^2 + ik^0 0^+}, \tag{37}$$

in which the poles of the propagator on the l.h.s. may create a pinch in the complex plane of the loop integration and the $\delta_+(k^2)$ on the r.h.s. may induce a non-integrable end-point configuration. Dubbing Σ_c the sum over all cuts that appears in the r.h.s. of (2), the cutting equations [2, 3] ensure that the last term in (37) does not contribute to the singular part of each cut in Σ_c , and that Σ_c is IR finite. This theorem implies a unitarity-preserving cancellation of the IR singularities if the Cutkosky relation

$$\frac{1}{k^2 + i0^+} \leftrightarrow \frac{2\pi}{i} \delta_+(k^2), \tag{38a}$$

giving the possibility of a one-to-one integrand level identification of the infrared divergent parts contributing to different

cuts in Σ_c , is preserved. However, one should also prove that $\Sigma_c = \sigma_R + \sigma_V$. The reason for this second requirement is the different origin of the potential numerators multiplying the two sides of (38a). In the case of a fermion line, that is the only cut relevant for this paper, the l.h.s. gets multiplied by the numerator of the propagator $k := k_{\text{prop}}$, while the r.h.s. by $\sum_{\text{spin}} u(k)\bar{u}(k) = \sum_{\text{spin}} v(k)\bar{v}(k) := k_{\text{spin}}$. Hence, in addition to (38a), the identity

$$k_{\text{prop}} = k_{\text{spin}} \tag{38b}$$

must hold to guarantee the validity of (2). Equation (38b) states that the numerators of the loop propagators in Fig. 5a, b should be treated consistently with the sum over the spin polarizations of the cut lines in Fig. 5c, d. Note that (38b) also guarantees consistent gauge cancellations in all terms contributing to Σ_c .

Let us consider how to make these relations consistent with the procedure from the previous section where we make two modifications to the integrand:

- adding μ^2 to the propagators with momenta q_2 and q_{12} ;
- SV and GV subtraction from the integrand.

We first study how FDR preserves (38a). We start dealing with the effect of the $q_2^2 \xrightarrow{\text{GP}} \bar{q}_2^2$ and $q_{12}^2 \xrightarrow{\text{GP}} \bar{q}_{12}^2$ replacements in cuts (a,b). Equation (38a) is preserved if

$$\frac{1}{(\bar{q}_2^2 + i0^+)(\bar{q}_{12}^2 + i0^+)} \leftrightarrow \left(\frac{2\pi}{i}\right)^2 \delta_+(\bar{k}_3^2)\delta_+(\bar{k}_4^2), \tag{39}$$

with $\bar{k}_{3,4}^2 := k_{3,4}^2 - \mu^2$. Thus, the \bar{q}_2^2 and \bar{q}_{12}^2 propagators in σ_V must correspond to external particles in σ_R obeying

$$k_{3,4}^2 = \mu^2. \tag{40}$$

Hence, we replace in (3) $\Phi_{n+2} \rightarrow \tilde{\Phi}_{n+2}$, where the phase-space $\tilde{\Phi}_{n+2}$ is such that $k_3^2 = k_4^2 = \mu^2$ and $k_i^2 = 0$ when $i \neq 3, 4$. However, this is not enough. One also needs to show that (39) survives the SV subtraction of (21). We prove this explicitly in the case of the last term in (13). The proof is unchanged for the other contributions. The relevant expansion is

$$\frac{1}{\bar{q}_2^2 \bar{q}_{12}^2} = \left[\frac{1}{\bar{q}_2^4} \right]_{\text{SV}} - \frac{q_1^2 + 2(q_1 \cdot q_2)}{\bar{q}_2^4 \bar{q}_{12}^2}, \tag{41}$$

where SV is the term to be subtracted. We consider a piece of the finite part of (41) as a numerator factor λ

$$-\frac{q_1^2 + 2(q_1 \cdot q_2)}{\bar{q}_2^4 \bar{q}_{12}^2} = \frac{\lambda}{\bar{q}_2^2 \bar{q}_{12}^2}, \quad \lambda := -\frac{q_1^2 + 2(q_1 \cdot q_2)}{\bar{q}_2^2} \tag{42}$$

and observe that $\lambda \rightarrow 1$ when $\bar{q}_{12}^2 \rightarrow 0$. We therefore first put the \bar{q}_{12}^2 propagator on-shell, giving

$$\begin{aligned} \frac{1}{\bar{q}_2^2 \bar{q}_{12}^2} &\leftrightarrow \frac{(2\pi)}{i} \delta_+(\bar{k}_4^2) \left\{ \frac{2(k_3 \cdot k_{34}) - k_{34}^2}{\bar{k}_3^4} \right\} \\ &= \frac{(2\pi)}{i} \delta_+(\bar{k}_4^2) \frac{1}{\bar{k}_3^2}. \end{aligned} \tag{43}$$

When also \bar{k}_3^2 goes on-shell, one obtains the same result as applying (39) before subtracting the vacuum. Hence, the SV subtraction is “invisible” from the point of view of (39), and (38a) is fulfilled if $k_{3,4}$ obey (40).

As for (38b), in order to preserve it, one must treat k_3^2, k_4^2 and $(k_3 + k_4)^2 = s_{34}$ in the numerator N_R of (36) using the same prescriptions imposed on q_2^2, q_{12}^2 and q_1^2 in N of (7). This means replacing in N_R

$$\begin{aligned} k_{3,4}^2 &\rightarrow \bar{k}_{3,4}^2 = 0, \\ (k_3 \cdot k_4) &= \frac{s_{34} - k_3^2 - k_4^2}{2} \rightarrow \frac{s_{34} - \bar{k}_3^2 - \bar{k}_4^2}{2} = \frac{s_{34}}{2}, \end{aligned} \tag{44}$$

where the last equalities are induced by the delta functions in (39). These changes should be performed everywhere in N_R except in contractions induced by the external indices $\hat{\rho}$ and $\hat{\sigma}$ in cuts (c,d). In this case

$$\hat{k}_{3,4}^2 \rightarrow k_{3,4}^2 = \mu^2, \quad (\hat{k}_3 \cdot \hat{k}_4) \rightarrow (k_3 \cdot k_4) = \frac{s_{34} - 2\mu^2}{2}, \tag{45}$$

in accordance to the SIC preserving requirement we have used to construct $\tilde{Z}(q_1^2, q_2^2)$ from N .⁷ We denote all of this by introducing a globally prescribed and SIC preserving version of N_R

$$N_R \xrightarrow{\text{GS}} \tilde{N}_R(\mu^2), \tag{46}$$

where the action of $\xrightarrow{\text{GS}}$ on a function $f(k_3^2, k_4^2, \hat{k}_3^2, \hat{k}_4^2)$ is defined to be

$$\begin{aligned} f(k_3^2, k_4^2, \hat{k}_3^2, \hat{k}_4^2) &\xrightarrow{\text{GS}} \tilde{f}(\bar{k}_3^2, \bar{k}_4^2, k_3^2, k_4^2) \\ &:= f(\bar{k}_3^2, \bar{k}_4^2, k_3^2, k_4^2), \end{aligned} \tag{47}$$

in which the tilde on \tilde{f} expresses the fact that GS has been imposed. Only a dependence on μ^2 is left in the r.h.s. of (46) because of the deltas in $\tilde{\Phi}_{n+2}$.

The only remaining propagator involved in our definition of the virtual component is in the top-left line of cut (a), that

⁷ Although it does not occur in our calculation, it is worth mentioning what happens when loop propagators and cut lines refer to spin-1 particles. In this case (38b) is replaced by $g_{\text{prop}}^{\mu\nu} = g_{\text{spin}}^{\mu\nu}$, which states that the metric tensor in the numerator of a loop propagator should be handled consistently with the one generated by the sum over polarizations. This is again achieved by treating the self contractions $k_3^2, k_4^2, (k_3 + k_4)^2$ in σ_R exactly as their counterparts q_2^2, q_{12}^2, q_1^2 in σ_V , as done in (44) and (45).

must correspond to the cut k_1 propagator in (d) through the relation⁸

$$\frac{1}{(q_1 + p)^2 + i0^+} \leftrightarrow \frac{2\pi}{i} \delta_+(k_1^2). \tag{48}$$

Everything is massless in this case, so that (38b) is fulfilled and it is sufficient to check that GV subtraction does not alter (48). The proof is similar to the one used for the SV. In fact, if m expansions

$$\frac{1}{(q_1 + p)^2 - \mu^2} = \frac{1}{\bar{q}_1^2} - \frac{2(q_1 \cdot p)}{\bar{q}_1^2((q_1 + p)^2 - \mu^2)} \tag{49}$$

are needed to subtract the vacuum in front of a term in $\tilde{Z}(\bar{q}_1^2, \hat{q}_2^2)$, this term gets multiplied by a factor λ^m in $\tilde{Z}(q_1^2, q_2^2)$, where

$$\lambda := \frac{-2(q_1 \cdot p)}{q_1^2}. \tag{50}$$

But $\lambda = 1$ when the propagator goes on-shell. So that (48) survives the GV subtraction.

In summary, we define the four-dimensional integration over the integrand in (36) as follows

$$\tilde{R} := \int d\tilde{\Phi}_{n+2} \frac{\tilde{N}_R(\mu^2)}{S_{34}^\alpha S_{134}^\beta} \equiv \int d\tilde{\Phi}_{n+2} \tilde{J}_R. \tag{51}$$

By doing that, unitarity preserving IR cancellations occur by construction between σ_R and σ_V , without violating gauge invariance.

2.3 An example of cancellation

The integrals in (35) and (51) can be computed independently, and this is the strategy we adopt in this paper. In fact, term by term cancellations in Σ_c are difficult to find. The reason is that one can add to the numerator of the virtual piece arbitrary vanishing terms that nevertheless contribute to the real part, and vice-versa. This is due to the different structure of the deltas contained in $d\Phi_n$ and $d\tilde{\Phi}_{n+2}$. However, if the numerator of a term does not change – modulo a relabelling of the momenta – when multiplied by both phase-spaces, the IR cancellation must occur between integrals constructed one from the other via the replacement in (38a). In this section we illustrate this phenomenon by considering a piece of the full $H \rightarrow b\bar{b} + jets$ calculation presented in the following section. This also serves us as a concrete example on how the procedures of Sects. 2.1 and 2.2 work in practice.

The two-loop contribution to σ_V is depicted in Fig. 6. We focus on the term generated by the $g_{\rho\sigma}$ piece of the internal trace. It reads

$$\tilde{\Gamma}_V = -\frac{1}{(2\pi)^{10}} \int d^4 p_1 d^4 p_2 [d^4 q_1][d^4 q_2] \delta_-(p_1^2) \delta_+(p_2^2)$$

⁸ The complex conjugate of (48) links cuts (b) and (c).




Fig. 6 A cut contributing to $H \rightarrow b\bar{b}$ at NNLO. Only the term proportional to $g_{\rho\sigma}$ is considered in (52)




Fig. 7 A cut contributing to $H \rightarrow b\bar{b}q\bar{q}$. Only the $g_{\rho\sigma}$ term is considered in (61)

$$\times \delta^4(P - p_2 + p_1) \frac{\tilde{N}_V}{\bar{q}_1^4 \bar{D}_1 \bar{D}_2 \bar{q}_2^2 \bar{q}_1^2}, \tag{52}$$

with $D_i := (q_1 + p_i)^2$. The numerator \tilde{N}_V is obtained via GP from its unbarred version

$$N_V = -64(p_1 \cdot p_2)(q_2 \cdot q_{12})(q_1 + p_1) \cdot (q_1 + p_2), \tag{53}$$

where we have neglected couplings and color factors, but not phases needed to compute the overall sign. We rewrite

$$N_V = 16s(q_{12}^2 + q_2^2 - q_1^2) \left(D_1 + (P \cdot q_1) - \frac{s}{2} \right), \tag{54}$$

with $s = P^2$, that gives

$$N_V \xrightarrow{\text{GP}} \bar{N}_V = 16s(\bar{q}_{12}^2 + \bar{q}_2^2 - \bar{q}_1^2) \left(\bar{D}_1 + (P \cdot q_1) - \frac{s}{2} \right). \tag{55}$$

This expression can be simplified by dropping terms which integrate to zero, for example \bar{q}_2^2 and \bar{q}_{12}^2 generate vacua and \bar{D}_1 gives a scale-less integral. Furthermore, $(P \cdot q_1)$ does not contribute because it is antisymmetric when $p_1 \leftrightarrow p_2$, while the denominator in (52) is symmetric. Thus $\tilde{N}_V \simeq \bar{N}'_V$, with

$$\bar{N}'_V(\bar{q}_1^2) = 8s^2 \bar{q}_1^2. \tag{56}$$

The GV in (52) is fully removed by the subtraction of the scale-less integral. Thus, $\tilde{Z}_V(\bar{q}_1^2) = \tilde{N}'_V(\bar{q}_1^2)$ and

$$\tilde{Z}_V(\bar{q}_1^2) \xrightarrow{\text{sic}} \tilde{Z}_V(q_1^2) = \tilde{N}'_V(q_1^2) = 8s^2q_1^2, \tag{57}$$

so that the physically relevant two-loop contribution reads

$$\begin{aligned} \tilde{\Gamma}_V &= \frac{1}{(2\pi)^{10}} \int d^4p_1 d^4p_2 d^4q_1 [d^4q_2] \delta_-(p_1^2) \delta_+(p_2^2) \\ &\times \delta^4(P - p_2 + p_1) \frac{\tilde{Z}_V(q_1^2)}{q_1^4 D_1 D_2 \bar{q}_2^2 \bar{q}_{12}^2}. \end{aligned} \tag{58}$$

$\tilde{\Gamma}_V$ develops IR divergences in the form of powers of

$$L := \ln \frac{\mu^2}{s}. \tag{59}$$

When splitting the result of the integration in a part which collects all terms containing powers of L , dubbed logarithmic part ($L.P.$), plus a remainder, one finds

$$\begin{aligned} L.P.(\mathcal{R}e(\tilde{\Gamma}_V)) &= -\frac{s}{64\pi^5} \left(L \left(1 - \frac{\pi^2}{12} \right) + \frac{L^2}{4} + \frac{L^3}{24} \right). \end{aligned} \tag{60}$$

These logarithms are cancelled by a term contributing to the four-particle cut-diagram in Fig. 7

$$\begin{aligned} \tilde{\Gamma}_R &= \frac{1}{(2\pi)^8} \int d^4k_1 d^4k_2 d^4k_3 d^4k_4 \delta_+(k_1^2) \delta_+(k_2^2) \\ &\times \delta_+(\bar{k}_3^2) \delta_+(\bar{k}_4^2) \delta^4(P - k_{1234}) \frac{\tilde{N}_R}{s_{34}^2 s_{134} s_{234}}, \end{aligned} \tag{61}$$

where the IR behavior is now regulated by the two external massive lines $k_3^2 = k_4^2 = \mu^2$. \tilde{N}_R is obtained by applying the GS operation defined in (47) to the numerator of the diagram

$$N_R = 64(k_3 \cdot k_4)(k_1 \cdot k_{234})(k_2 \cdot k_{134}). \tag{62}$$

the result reads

$$\tilde{N}_R = 8s_{34}(s - s_{234})(s - s_{134}). \tag{63}$$

\tilde{N}_R does not depend on μ^2 because in this case there are no contractions of $k_{3,4}$ vectors with the external indices $\hat{\rho}$ and $\hat{\sigma}$.

We now can check that the integrand in (61) is the correct object to cancel the logarithms in (60). In fact, it can be obtained from (58) by means of the Cutkosky replacement in (38a), together with the relabellings

$$q_1 \rightarrow k_{34}, \quad q_2 \rightarrow -k_3, \quad p_2 \rightarrow k_2, \quad p_1 \rightarrow -k_{134}, \tag{64}$$

inferred by comparing Figs. 6 and 7, and the substitution

$$\frac{1}{(q_1^2 + i0^+)^2} \rightarrow -\frac{1}{(s_{34} + i0^+)(s_{34} - i0^+)}, \tag{65}$$

which is necessary because of the gluon propagator appearing on the r.h.s. of Fig. 7. Therefore, $\tilde{\Gamma}_R$ must contain a contribution with the same singular behavior of $-\tilde{\Gamma}_V$. We dub

$\tilde{\Gamma}'_R$ such a contribution, and \tilde{N}'_R its numerator function. The terms proportional to s_{234} or s_{134} in (63) give zero when evaluated at the two-particle cut: they cannot be “seen” by $\tilde{\Gamma}_V$. This leads us to the conclusion that

$$\tilde{N}'_R = 8s^2s_{34}, \tag{66}$$

which corresponds to $\tilde{Z}_V(q_1^2)$ in (57), modulo the first replacement in (64). Thus, $\tilde{\Gamma}'_R$ is obtained by replacing $\tilde{N}_R \rightarrow \tilde{N}'_R$ in (61). An explicit calculation confirms that

$$L.P.(\mathcal{R}e(\tilde{\Gamma}_V) + \tilde{\Gamma}'_R) = 0. \tag{67}$$

3 Renormalization

In this section, we discuss and implement the FDR renormalization program in the context of our calculation. To do this we need to distinguish, at least conceptually, between UV regulator, IR regulator and renormalization scale. We denote them by μ_{UV} , μ_{IR} and μ_R , respectively.

In the case of IR free observables, the FDR integral operator in (22) subtracts the UV infinities before integration. For this reason, after taking the asymptotic limit $\mu_{UV} \rightarrow 0$, μ_{UV} can be directly interpreted as the finite renormalization scale μ_R [18]. In this sense, FDR directly produces a finite, renormalized result for the loop part: nothing needs to be subtracted from it. However, this result is arbitrary until bare parameters are fixed by experimental measurements. After doing so, if the theory is renormalizable, the scale μ_R gets replaced by physical scales, leading to an unambiguous prediction. This can be understood as a finite renormalization necessary to make the theory predictive [25].

In the presence of IR divergences, no distinction is made in the virtual component between μ_{UV} and μ_{IR} . As a matter of fact, the procedure in Sect. 2.1 assumes $\mu = \mu_{UV} = \mu_{IR}$, preventing one from setting $\mu = \mu_{UV} = \mu_R$, as is possible in the IR free case. Our solution is fixing the bare parameters in terms of physical quantities before combining virtual and real components. After this is done, the μ_R scales get automatically replaced by physical scales, hence the left over μ_S are the μ_{IR} s which cancel the IR behavior of the real counterpart.

The bare parameters in our calculation are α_S^0 and the Yukawa coupling y_b^0 . In order to implement our renormalization program we need relations linking them to measured quantities at the appropriate perturbative order, which is one loop for α_S^0 and two loops for y_b^0 .

We are interested in corrections proportional to N_F . Hence, α_S^0 can be linked to the customary $\alpha_S^{\overline{\text{MS}}}(s)$ by using the fact that the N_F contribution to the running coincides in FDR and $\overline{\text{MS}}$ [21]. As a consequence, we choose our renormalized strong coupling constant to be $\alpha_S = \alpha_S^{\overline{\text{MS}}}(s)$, that gives the relation

$$a^0 = a \left(1 + a\delta_a^{(1)} \right), \tag{68}$$

with

$$a^0 := \frac{\alpha_S^0}{4\pi}, \quad a := \frac{\alpha_S}{4\pi}, \quad \delta_a^{(1)} = \frac{2}{3} N_F L, \tag{69}$$

where L is given in (59).

The Yukawa coupling is renormalized by using its proportionality to the bottom mass. The corrected bottom propagator at the pole is proportional to

$$\frac{1}{\not{p} - m^0 + \Sigma^{(1)} + \Sigma^{(2)}}, \tag{70}$$

where m^0 is the bare mass and the $\Sigma^{(j)}$ s are computed in Appendix C. This gives a relation between m^0 and the pole mass m

$$m^0 = m + \Sigma^{(1)} + \Sigma^{(2)}, \tag{71}$$

which translates into

$$y_b^0 = y_b(1 + a^0\delta_y^{(1)} + a^2\delta_y^{(2)}), \tag{72}$$

with

$$\begin{aligned} \delta_y^{(1)} &= -C_F(3L'' + 5), \\ \delta_y^{(2)} &= C_F N_F \left(L''^2 + \frac{13}{3}L'' + \frac{2}{3}\pi^2 + \frac{151}{18} \right), \end{aligned} \tag{73}$$

where

$$L'' := \ln \frac{\mu^2}{m^2}. \tag{74}$$

Equation (72) contains the bare QCD coupling. Inserting (68) gives the desired two-loop relation between bare and renormalized Yukawa coupling

$$y_b^0 = y_b(1 + a\delta_y^{(1)} + a^2(\delta_y^{(2)} + \delta_a^{(1)}\delta_y^{(1)})). \tag{75}$$

4 $H \rightarrow b\bar{b} + jets$

In this section, we use FDR to reproduce the physical prediction for the inclusive decay width of the Higgs into two b jets up to the NNLO accuracy in the large N_F limit of QCD. That means computing the observable

$$\Gamma^{\text{NNLO}}(y_b) = \Gamma_2^{(0)}(y_b) + \delta\Gamma^{N_F}, \tag{76}$$

where $\Gamma_2^{(0)}(y_b)$ is the tree-level decay width and $\delta\Gamma^{N_F}$ collects all the NNLO terms proportional to $\alpha_S^2 N_F$.

The correction factor $\delta\Gamma^{N_F}$ receives contributions from processes with up to four final-state particles, namely:

- $H \rightarrow b\bar{b}$ up to two loops;
- $H \rightarrow b\bar{b}g$ at the tree level;
- $H \rightarrow b\bar{b}q\bar{q}$ at the tree level.




Fig. 8 The one-loop QCD correction to the $Hb\bar{b}$ vertex

The tree- and one-loop two- and three-body decays in the above list contribute to $\delta\Gamma^{N_F}$ through renormalization. As a matter of fact, due to the scalar nature of the Yukawa coupling, $\Gamma^{\text{NNLO}}(y_b)$ is a simple process in terms of the contributing tensor structures. Nevertheless, it requires the two-loop renormalization of (75).

In the following, we compute all components in the massless limit of QCD, namely with $m \neq 0$ only in y_b . Our notation is as follows. We dub $V_i^{(j)}$ the H decay amplitudes into i final state partons computed at the j^{th} order of the QCD perturbative expansion. We shall omit for brevity the multiplication of the appropriate quark spinors in any expressions for the $V_i^{(j)}$. The decay widths are obtained by squaring the amplitudes and are denoted by $\Gamma_i^{(j)}$.

In this paper we focus on the new aspects of FDR at NNLO, namely the procedures presented in Sects. 2.1 and 2.2. For this reason we do not go into detail of the calculation of the NLO part. The corresponding expressions can be computed as described in references [21,23]. However, we emphasize that FDR NLO formulae stay the same also when they contribute to a NNLO calculation. The same holds true for LO expressions multiplying higher order corrections. This is in contrast to d -dimensional regularization methods, in which higher powers in the $(d - 4)$ expansion must be added.

4.1 $H \rightarrow b\bar{b}$ up to two loops

In our conventions, the lowest order $H \rightarrow b\bar{b}$ vertex is

$$V_2^{(0)} = \delta_{kl} y_b^0, \tag{77}$$

where k and l are the color indices of the bottom quarks. Squaring $V_2^{(0)}$ gives the LO $H \rightarrow b\bar{b}$ decay width

$$\Gamma_2^{(0)}(y_b^0) = (y_b^0)^2 M_H \frac{N_C}{8\pi}, \tag{78}$$

in which N_C is the number of colors.

The one-loop correction is depicted in Fig. 8. Following reference [21] one obtains

$$V_2^{(1)} = -\delta_{kl} \left(\frac{\alpha_S^0}{4\pi} \right) y_b^0 C_F L^2, \tag{79}$$

with

$$L' := \ln \frac{\mu^2}{-s - i0^+}, \quad s = M_H^2, \quad C_F = \frac{N_C^2 - 1}{2N_C}. \quad (80)$$

The globally prescribed integral needed to compute the two-loop correction is given by left part of Fig. 6. It reads

$$\bar{V}_2^{(2)} = \delta_{kl} \frac{y_b}{8} C_F N_F \frac{\alpha_S^2}{\pi^6} \int [d^4 q_1][d^4 q_2] \frac{\bar{N}_A + \bar{N}_B}{\bar{q}_1^4 \bar{D}_1 \bar{D}_2 \bar{q}_2^2 \bar{q}_{12}^2}, \quad (81)$$

with \bar{D}_i given right after (52). Here we have replaced bare quantities with renormalized ones, because the difference is $\mathcal{O}(\alpha_S^3)$. In the following, we compute $\bar{N}_{A,B}$ starting from their unbarred counterparts $N_{A,B}$, which can be obtained from (7) by taking $F_{\hat{\rho}\hat{\sigma}} = \gamma_{\hat{\rho}}(\not{q}_1 + \not{p}_1)(\not{q}_1 + \not{p}_2)\gamma_{\hat{\sigma}}$. By denoting $\hat{q}_2 := \gamma_{\hat{\rho}} q_2^{\rho} = \gamma_{\hat{\sigma}} q_2^{\sigma}$ one finds

$$\begin{aligned} N_A &= \not{q}_1(\not{q}_1 + \not{p}_1)(\not{q}_1 + \not{p}_2)\not{q}_2 + \not{q}_2(\not{q}_1 + \not{p}_1)(\not{q}_1 + \not{p}_2)\not{q}_1, \\ N_B &= 2\not{q}_2(\not{q}_1 + \not{p}_1)(\not{q}_1 + \not{p}_2)\not{q}_2 - \frac{N_V}{8s}, \end{aligned} \quad (82)$$

where the last term originates from the piece we have studied in detail in Sect. 2.3, with N_V given in (53). N_A does not contribute to $\bar{V}_2^{(2)}$. In fact, a $\bar{u}(p_1)(u(p_2))$ is understood on the l.h.s.(r.h.s.), so that, by virtue of the Dirac equation, one can replace $N_A \rightarrow N'_A = D_1(\not{q}_1 + \not{p}_2)\not{q}_2 + \not{q}_2(\not{q}_1 + \not{p}_1)D_2$, giving

$$N'_A \xrightarrow{\text{GP}} \bar{N}'_A = \bar{D}_1(\not{q}_1 + \not{p}_2)\not{q}_2 + \not{q}_2(\not{q}_1 + \not{p}_1)\bar{D}_2, \quad (83)$$

which generates scale-less integrals. That explicitly proves the WI in (25). As for the N_B piece, there are several ways [19,20] to deal with strings of γ -matrices to extract the dependence on $(q_i \cdot q_j)$ and q_i^2 needed to implement GP. They are based on replacements of the type $\not{q}_i \rightarrow \not{q}_i - \mu_i$, where the ‘‘masses’’ μ_i serve as a bookkeeping tool. In this paper we find it more convenient to use Clifford algebra until we reach the configurations⁹

$$\not{q}_i \not{q}_i \xrightarrow{\text{GP}} \bar{q}_i^2, \quad \not{q}_i \not{q}_j \xrightarrow{\text{GP}} \frac{1}{2}(\bar{q}_{ij}^2 - \bar{q}_i^2 - \bar{q}_j^2 + \not{q}_i \not{q}_j - \not{q}_j \not{q}_i). \quad (84)$$

⁹ Note the invariance of the last term under $\not{q}_i \rightarrow \not{q}_i - \mu_i$.




Fig. 9 The LO $Hb\bar{b}g$ vertex

By using this method one finds $N_B \xrightarrow{\text{GP}} \bar{N}_B \simeq \bar{N}'_B$ with

$$\begin{aligned} \bar{N}'_B(\bar{q}_1^2, \hat{q}_2^2) &= -4(q_1 \cdot p_1)\hat{q}_2^2 + 8(q_2 \cdot p_2)((q_2 \cdot q_1) + (q_2 \cdot p_1)) \\ &\quad - \frac{8}{s}(P \cdot q_2)((q_1 \cdot p_2)(q_2 \cdot p_1) - (q_1 \cdot p_1)(q_2 \cdot p_2)) \\ &\quad - s\bar{q}_1^2. \end{aligned} \quad (85)$$

\bar{N}'_B does not induce the appearance of global UV divergences in (81), hence the numerator function directly reads $\bar{Z}(\bar{q}_1^2, \hat{q}_2^2) = \bar{N}'_B(\bar{q}_1^2, \hat{q}_2^2)$. Thus, the SIC preserving numerator function is $\tilde{Z}(q_1^2, q_2^2) = \bar{N}'_B(q_1^2, q_2^2)$, and the two-loop correction is

$$V_2^{(2)} = \delta_{kl} \frac{y_b}{8} C_F N_F \frac{\alpha_S^2}{\pi^6} \int d^4 q_1 [d^4 q_2] \frac{\tilde{Z}(q_1^2, q_2^2)}{q_1^4 D_1 D_2 \bar{q}_2^2 \bar{q}_{12}^2}. \quad (86)$$

In terms of the master integrals listed in Appendix D it reads

$$V_2^{(2)} = \delta_{kl} \frac{4}{3} y_b C_F N_F \left(\frac{\alpha_S}{4\pi}\right)^2 \tilde{I}_1. \quad (87)$$

4.2 $H \rightarrow b\bar{b}g$ and $H \rightarrow b\bar{b}q\bar{q}$ at the tree level

The LO $H \rightarrow b\bar{b}g$ decay width is obtained by squaring the $V_3^{(0)}$ vertex drawn in Fig. 9 and integrating over a phase-space in which all final-state particles acquire a small mass μ , as described in reference [21]. The result is

$$\Gamma_3^{(0)} = \left(\frac{\alpha_S^0}{4\pi}\right) \Gamma_2^{(0)}(y_b^0) C_F (2L^2 + 6L + 19 - 2\pi^2), \quad (88)$$

with L defined in (59).

As for $H \rightarrow b\bar{b}q\bar{q}$, two diagrams contribute to the amplitude $V_4^{(0)}$. They can be read from Fig. 9 by allowing the gluon to split into a $q\bar{q}$ pair. As described in Sect. 2.2, $\Gamma_4^{(0)}$ is obtained by squaring $V_4^{(0)}$ and integrating over a massive 4-particle phase-space $\tilde{\Phi}_4$ such that $k_1^2 = k_2^2 = 0$ and $k_3^2 = k_4^2 = \mu^2$. Prior to integration, the integrand should be modified according to the GP and SIC replacements given in (46). As a result of this, the function to be integrated in (51) is a rational combination of the invariants s_{34}, s_{134}, s_{234} and μ^2 :

$$\tilde{J}_R = \tilde{J}_R(s_{34}, s_{134}, s_{234}, \mu^2), \quad (89)$$

where the μ^2 dependence is induced by (45). It is interesting to note this μ^2 dependence factorizes. In particular, one finds

$$\tilde{J}_R(s_{34}, s_{134}, s_{234}, \mu^2) = \tilde{J}'_R(s_{34}, s_{134}, s_{234})w(\mu^2), \tag{90}$$

with

$$w(\mu^2) = \left(1 + 2\frac{\mu^2}{s_{34}}\right). \tag{91}$$

Therefore, (51) has the form of a massless-like matrix element $\tilde{J}'_R(s_{34}, s_{134}, s_{234})$, obeying all relevant WIs, times a global factor $w(\mu^2)$ integrated over a massive phase-space. We conjecture that a similar gauge-invariant structure should naturally appear also when dealing with more complicated cases involving, for instance, IR divergences induced by vector particles like gluons.

In terms of the integrals reported in Appendix E the result reads

$$\Gamma_4^{(0)} = \frac{64}{3}C_F N_F \Gamma_2^{(0)}(y_b) \times \left(\frac{\alpha_S}{4\pi}\right)^2 (\tilde{R}_8 + \tilde{R}_7 + \tilde{R}_6 - 2\tilde{R}_5 - \tilde{R}_4). \tag{92}$$

4.3 The large N_F limit of the inclusive width

Here we gather all the calculated components and compute $\Gamma^{\text{NNLO}}(y_b)$ in (76). The correction factor $\delta\Gamma^{N_F}$ receives contributions from processes with up to four partons

$$\delta\Gamma^{N_F} = \Gamma_2^{N_F} + \Gamma_3^{N_F} + \Gamma_4^{N_F}, \tag{93}$$

that are obtained by inserting the renormalization equations (68) and (75) in the amplitudes given in the previous sections.

One finds

$$\begin{aligned} \Gamma_2^{N_F} &= \Gamma_2^0(y_b)a^2 \\ &\quad \times 2\Re e \left(\delta V_2^{(2)} + \delta_a^{(1)}\delta V_2^{(1)} + \delta_y^{(2)} + \delta_a^{(1)}\delta_y^{(1)}\right), \\ \Gamma_3^{N_F} &= a^2\delta_a^{(1)}\Gamma_2^0(y_b)C_F(2L^2 + 6L + 19 - 2\pi^2), \\ \Gamma_4^{N_F} &= a^2C_F N_F \Gamma_2^{(0)}(y_b) \\ &\quad \times \frac{4}{9} \left\{-L^3 - \frac{19}{2}L^2 - L\left(\frac{155}{3} - 2\pi^2\right) + 30\zeta_3\right. \\ &\quad \left. + \frac{29}{6}\pi^2 - \frac{4345}{36}\right\}, \end{aligned} \tag{94}$$

where

$$\begin{aligned} \delta V_2^{(1)} &= -C_F(L')^2, \\ \delta V_2^{(2)} &= \frac{2}{9}C_F N_F \left(L'^3 + 5L'^2 + L'\left(\frac{56}{3} + \pi^2\right) - 12\zeta_3\right. \\ &\quad \left. + \frac{5}{3}\pi^2 + \frac{328}{9}\right). \end{aligned} \tag{95}$$

Equations (94) are written in a form that highlights the contributions generated by renormalization. Collecting all the pieces gives the IR finite result

$$\begin{aligned} \Gamma^{\text{NNLO}}(y_b) &= \Gamma_2^{(0)}(y_b) \left\{1 + a^2C_F N_F\right. \\ &\quad \left.\times \left(2\ln^2\frac{m^2}{s} - \frac{26}{3}\ln\frac{m^2}{s} + 8\zeta_3 + 2\pi^2 - \frac{62}{3}\right)\right\}. \end{aligned} \tag{96}$$

Equation (96) is written in terms of the pole mass m . It is possible to reabsorb the large logarithms of the ratio m^2/s in a new Yukawa coupling $y_b^{\overline{\text{MS}}}$ defined through the known two-loop relation between m and the $\overline{\text{MS}}$ mass [26]. Using the N_F part of it gives

$$\begin{aligned} \Gamma_2^{(0)}(y_b) &= \Gamma_2^{(0)}(y_b^{\overline{\text{MS}}}(s)) \left\{1 + a^2C_F N_F\right. \\ &\quad \left.\times \left(-2\ln^2\frac{m^2}{s} + \frac{26}{3}\ln\frac{m^2}{s} - \frac{4}{3}\pi^2 - \frac{71}{6}\right)\right\}, \end{aligned} \tag{97}$$

hence

$$\begin{aligned} \Gamma^{\text{NNLO}}(y_b^{\overline{\text{MS}}}(s)) &= \Gamma_2^{(0)}(y_b^{\overline{\text{MS}}}(s)) \left\{1 + a^2C_F N_F\right. \\ &\quad \left.\times \left(8\zeta_3 + \frac{2}{3}\pi^2 - \frac{65}{2}\right)\right\}. \end{aligned} \tag{98}$$

Equation (98) coincides with the known $\overline{\text{MS}}$ result [27,28].

5 $\gamma^* \rightarrow jets$

In this section we compute the large N_F limit of the inclusive $e^+e^- \rightarrow \gamma^* \rightarrow jets$ production rate up to the NNLO accuracy. That is the observable

$$\sigma^{\text{NNLO}} = \sigma_2^{(0)} + \delta\sigma^{N_F}, \tag{99}$$

where $\sigma_2^{(0)}$ is the tree-level $e^+e^- \rightarrow \gamma^* \rightarrow q\bar{q}$ cross-section and $\delta\sigma^{N_F}$ contains the QCD corrections proportional to $\alpha_S^2 N_F$. QCD renormalization only involves α_S , in this case. Nevertheless, higher rank tensors contribute, so that preserving gauge cancellations and unitarity in such an environment provides a more stringent test for our procedures. In this respect, $\gamma^* \rightarrow jets$ is complementary to $H \rightarrow b\bar{b} + jets$.

The processes which contribute to $\delta\sigma^{N_F}$ are

- $e^+e^- \rightarrow q\bar{q}$ up to two loops;
- $e^+e^- \rightarrow q\bar{q}g$ at the tree level;
- $e^+e^- \rightarrow q\bar{q}q'\bar{q}'$ at the tree level,

where we understand a photon mediating the reactions. We dub $V_i^{(j)\beta}$ the final-state current producing i partons computed at the j^{th} QCD order, where β is the Lorentz index of the virtual photon. The Feynman diagrams representing the vertices are obtained from those in the previous section by replacing the Higgs with a photon. Hence, we do not draw them. Contracting $V_i^{(j)\beta}$ with the initial-state current,

squaring and integrating over the phase-space gives the corresponding cross section, denoted by $\sigma_i^{(j)}$.

In the following, we describe the FDR computation of the various components.

5.1 $e^+e^- \rightarrow q\bar{q}$ up to two loops

The lowest order vertex is

$$V_2^{(0)\beta} = -ieQ_q\delta_{kl}\gamma^\beta, \tag{100}$$

where Q_q is the electric charge of the quark. The corresponding cross section reads

$$\sigma_2^{(0)} = N_C \frac{4}{3}\pi \frac{\alpha^2}{s} Q_q^2, \tag{101}$$

in which α is the fine-structure constant.

The computation of $V_2^{(1)\beta}$ is described in [23]. The result is

$$V_2^{(1)\beta} = -a^0 C_F V_2^{(0)\beta} (L'^2 + 3L' + 7), \tag{102}$$

with a^0 and L' defined in (69) and (80), respectively.

The globally prescribed two-loop integral we need to compute $V_2^{(2)\beta}$ reads

$$\begin{aligned} \bar{V}_2^{(2)\beta} &= (-ieQ_q\delta_{kl}) \frac{C_F N_F \alpha_S^2}{8 \pi^6} \\ &\times \int [d^4 q_1][d^4 q_2] \frac{\bar{N}^\beta}{\bar{q}_1^4 \bar{D}_1 \bar{D}_2 \bar{q}_2^2 \bar{q}_{12}^2}. \end{aligned} \tag{103}$$

The unbarred N^β is obtained from (7) with $F_{\hat{\rho}\hat{\sigma}} = \gamma_{\hat{\rho}}(\not{q}_1 + \not{p}_1)\gamma^{\hat{\beta}}(\not{q}_1 + \not{p}_2)\gamma_{\hat{\sigma}}$ by using the fact that, according to the WI in (25), the term proportional to $q_2^\rho q_1^\sigma + q_1^\rho q_2^\sigma$ in the fermion trace does not contribute.¹⁰ Hence

$$\begin{aligned} N^\beta &= +2(q_2 \cdot q_{12})(\not{q}_1 + \not{p}_2)\gamma^\beta(\not{q}_1 + \not{p}_1) \\ &+ 2\not{q}_2(\not{q}_1 + \not{p}_1)\gamma^\beta(\not{q}_1 + \not{p}_2)\not{q}_2. \end{aligned} \tag{104}$$

Using tensor decomposition and the $p_1 \leftrightarrow p_2$ symmetry gives $N^\beta \xrightarrow{\text{GP}} \bar{N}^\beta \simeq \bar{M}^\beta$ with

$$\begin{aligned} \bar{M}^\beta(\hat{q}_1^2, \hat{q}_2^2) &= +2\gamma^\beta \left\{ 4(q_2 \cdot p_1)(q_2 \cdot p_2) \right. \\ &+ \frac{4}{s}(q_2 \cdot P)((q_1 \cdot p_1)(q_2 \cdot p_2) - (q_1 \cdot p_2)(q_2 \cdot p_1)) \\ &+ 2\hat{q}_1^2(q_1 \cdot p_1) - \frac{s}{2}\hat{q}_1^2 \left. \right\} + 8(q_1 \cdot p_1)\not{q}_2\gamma^\beta \\ &- 4\hat{q}_2^2 \left\{ (q_1 \cdot p_1)\gamma^\beta + q_1^\beta \not{q}_1 \frac{-2(q_1 \cdot p_1)}{\hat{q}_1^2} \right\}, \end{aligned} \tag{105}$$

where the factor $-2(q_1 \cdot p_1)/\hat{q}_1^2$ multiplying the last term subtracts its GV. Thus, the SIC preserving numerator function

¹⁰ The proof is analogous to the one given in Sect. 4.1.

is $\tilde{Z}^\beta(q_1^2, q_2^2) = \bar{M}^\beta(q_1^2, q_2^2)$, giving

$$\begin{aligned} V_2^{(2)\beta} &= (-ieQ_q\delta_{kl}) \frac{C_F N_F \alpha_S^2}{8 \pi^6} \\ &\times \int d^4 q_1 [d^4 q_2] \frac{\tilde{Z}^\beta(q_1^2, q_2^2)}{q_1^4 D_1 D_2 \bar{q}_2^2 \bar{q}_{12}^2}. \end{aligned} \tag{106}$$

In terms of the two-loop integrals in Appendix D one finds

$$V_2^{(2)\beta} = V_2^{(0)\beta} \frac{16C_F N_F a^2}{3} \left(\tilde{I}_3 - \tilde{I}_2 + \frac{\tilde{I}_1}{4} \right). \tag{107}$$

5.2 $e^+e^- \rightarrow q\bar{q}g$ and $e^+e^- \rightarrow q\bar{q}q'\bar{q}'$ at the tree level

A NLO computation produces

$$\sigma_3^{(0)} = a^0 \sigma_2^{(0)} C_F (2L^2 + 6L + 17 - 2\pi^2), \tag{108}$$

with L is given in (59).

As for $\sigma_4^{(0)}$, it is obtained by computing the amplitude squared, modifying it according to the prescription in (46) and integrating over the $\hat{\Phi}_4$ phase-space. In terms of the integrals in Appendix E the result reads

$$\begin{aligned} \sigma_4^{(0)} &= \frac{64}{3} C_F N_F \sigma_2^{(0)} a^2 \\ &\times (\tilde{R}_7 + \tilde{R}_6 - 2\tilde{R}_5 - \tilde{R}_4 + \tilde{R}_3 + 2\tilde{R}_2 - 2\tilde{R}_1). \end{aligned} \tag{109}$$

5.3 The large N_F limit of the inclusive jet production rate

Here we collect all components needed to compute σ^{NNLO} . The correction can be split as follows

$$\delta\sigma^{N_F} = \sigma_2^{N_F} + \sigma_3^{N_F} + \sigma_4^{N_F}, \tag{110}$$

where the various contributions are obtained by inserting (68) in the results of the previous sections. One has

$$\begin{aligned} \sigma_2^{N_F} &= \sigma_2^{(0)} a^2 2\Re e \left(\delta\hat{V}_2^{(2)} + \delta_a^{(1)} \delta\hat{V}_2^{(1)} \right), \\ \sigma_3^{N_F} &= a^2 \delta_a^{(1)} \sigma_2^{(0)} C_F (2L^2 + 6L + 17 - 2\pi^2), \\ \sigma_4^{N_F} &= a^2 C_F N_F \sigma_2^{(0)} \frac{4}{9} \left\{ -L^3 - \frac{19}{2}L^2 \right. \\ &- L \left(\frac{146}{3} - 2\pi^2 \right) \\ &\left. + 30\zeta_3 + \frac{19}{3}\pi^2 - \frac{2123}{18} \right\}, \end{aligned} \tag{111}$$

with

$$\begin{aligned} \delta\hat{V}_2^{(1)} &= -C_F (L'^2 + 3L' + 7), \\ \delta\hat{V}_2^{(2)} &= \frac{2}{9} C_F N_F \left\{ L'^3 + \frac{19}{2}L'^2 + L' \left(\frac{265}{6} + \pi^2 \right) \right. \\ &\left. - 12\zeta_3 + \frac{19}{6}\pi^2 + \frac{3355}{36} \right\}. \end{aligned} \tag{112}$$

Gathering all the pieces gives

$$\sigma^{\text{NNLO}} = \sigma_2^{(0)} \{1 + a^2 C_F N_F (8\zeta_3 - 11)\}, \quad (113)$$

which reproduces the $\overline{\text{MS}}$ result [29,30].

6 Conclusion and outlook

In this paper we have demonstrated that a fully four-dimensional framework to compute NNLO quark-pair corrections can be constructed based on the requirement of preserving the two principles given in (1). The FDR idea of enforcing gauge invariance and unitarity at the level of the UV subtracted integrands is at the base of the procedures we have used to define UV and IR divergent integrals.

A few advantages of such an approach that have appeared in our calculation are, for the UV part

- no (explicit or implicit) UV counterterms have to be included in the Lagrangian;
- lower-order substructures are used in higher-order calculations without any modification (see e.g. (79) and (102)).

For the IR sector

- infrared divergences in the real component directly show up in terms of logarithms of a small cut-off parameter μ_{IR} , with no need for a prior subtraction of $1/(d-4)$ poles (see, for instance, the four-parton rates in (94) and (111));
- one-to-one integrand correspondences can be written down between virtual and real contributions (see Sect. 2.3).

In this paper we have focused our attention on a special class of NNLO corrections. However, we believe that the basic principles that have guided us towards a consistent treatment of all the pieces contributing to the final NNLO answer will remain valid also when considering more complicated environments, with the final aim of constructing a completely general procedure including also initial state IR singularities. This is certainly the main subject of our future investigations. Other possible directions are: using μ_{IR} as a separation parameter in slicing-based subtraction methods at NNLO [31], or exploiting the virtual/real integrand correspondence to construct four-dimensional local counter-terms directly from the virtuals,¹¹ in the same spirit of the procedure presented at NLO in the FDR section of [23].

¹¹ A two-loop DReg algorithm along these lines has been recently proposed in [32].

On a more general ground, we envisage that the intrinsic four-dimensionality of FDR can pave the way to new numerical methods and that there is room for fully exploiting its potential in NNLO calculations.

Acknowledgements R.P. acknowledges the financial support of the MINECO Project FPA2016-78220-C3-3-P and the hospitality of the CERN TH department during the completion of this work. Our figures are prepared with AxoDraw [33].

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: This is a theoretical paper with no external associated data.]

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

Appendix A: Sub-integration consistency with and w/o IR divergences

When no IR infinities are present, the mismatch between Eqs. (28) and (29) is cured by adding the so called *extra-extra integrals* (EEI) introduced in [22]. Their exact definition is not needed here. It suffices to say that terms proportional to the difference

$$q_2^2 - \bar{q}_2^2 = \mu^2 \quad (\text{A.1})$$

are included. They multiply UV $1/\mu^2$ poles and generate logarithms of μ^2 that restore the correct renormalization properties of the two-loop amplitude. Such contributions are missed by (29).

In the presence of IR divergences an additional complication is generated by the GP $q_1^2 \rightarrow \bar{q}_1^2$ in (28) and (29). After GV subtraction, the difference

$$\mu^2 = q_1^2 - \bar{q}_1^2 \quad (\text{A.2})$$

also hits $1/\mu^2$ poles of IR origin. This gives rise to different renormalization constants for processes with or without IR divergences, which is unacceptable. This leads to the choice of letting q_1^2 unbarred, as discussed in Sect. 2.1. For the sake of consistency, also the EEIs part needs to be modified accordingly. The problem is that the EEIs become unregulated when unbarred \bar{q}_1^2 at the integrand level. The solution to this is replacing EEIs with the difference of two ordinary FDR integrals, generated by the combination

$$q_2^2 - \bar{q}_2^2, \quad (\text{A.3})$$

which is sometimes referred as an *extra-integral* (EI). One shows that EEIs and EIs share the same logarithmic content,

which fixes the correct UV behavior. In addition, EIs admit the $\bar{q}_1^2 \rightarrow q_1^2$ limit that matches the rest of the calculation.

In summary, the solution presented in Sect. 2.1 is equivalent to the following procedure:

- apply GP;
- subtract GV;
- downgrade $\bar{q}_1^2 \rightarrow q_1^2$ in the result;
- identify the EEIs to be added (using the same algorithm as in the IR-free case);
- replace each EEI with the corresponding EI.

It would be interesting to establish whether this strategy works also for IR finite two-loop calculations. That would make unnecessary the use of the EEIs. We leave this to further investigations.

Appendix B: Massless wave-function corrections

Wave function corrections are generated when the lower gluon in Fig. 1b reconnects to the emitting massless parton. In this appendix, we use the results of Sect. 2.1 to demonstrate that they vanish.

The relevant integrand is obtained by taking $F_{\hat{\rho}\hat{\sigma}} = \gamma_{\hat{\rho}}(\not{q}_1 + \not{p})\gamma_{\hat{\sigma}}$ in (7), that gives $N = N_A + N_B$ with

$$\begin{aligned} N_A &= \not{q}_1(\not{q}_1 + \not{p})\not{q}_2 + \not{q}_2(\not{q}_1 + \not{p})\not{q}_1, \\ N_B &= 2\hat{q}_2(\not{q}_1 + \not{p})\hat{q}_2 + 2(q_2 \cdot q_{12})(\not{q}_1 + \not{p}). \end{aligned} \tag{B.4}$$

Furthermore, $D = q_1^4 D_p$, so that the integrals we have to consider are

$$\bar{I}_{A,B} = \int [d^4 q_1][d^4 q_2] \frac{\bar{N}_{A,B}}{\bar{q}_1^4 \bar{D}_p \bar{q}_2^2 \bar{q}_{12}^2}. \tag{B.5}$$

One finds

$$\begin{aligned} N_A \xrightarrow{\text{GP}} \bar{N}_A &= 2\not{q}_2 \bar{D}_p - \not{p}(\bar{q}_{12}^2 - \bar{q}_1^2 - \bar{q}_2^2) \\ &\quad - \frac{1}{2}\not{p}(\not{q}_1\not{q}_2 - \not{q}_2\not{q}_1) - \frac{1}{2}(\not{q}_2\not{q}_1 - \not{q}_1\not{q}_2)\not{p}. \end{aligned} \tag{B.6}$$

Only the third term contributes to \bar{I}_A . All the others generate vacua or result from contractions of antisymmetric combinations of γ -matrices with symmetric integrals. Thus

$$\bar{I}_A = \not{p} \int [d^4 q_1][d^4 q_2] \frac{1}{\bar{q}_1^2 \bar{D}_p \bar{q}_2^2 \bar{q}_{12}^2}. \tag{B.7}$$

\bar{I}_A only depends on $p^2 = 0$. In addition, it is both UV divergent and logarithmically IR divergent, so that it is a *scale-less* integral. Such integrals vanish in FDR as a consequence of an exact cancellation between UV and IR singularities,

therefore $\bar{I}_A = 0$, as required by (25). In the same way, \bar{I}_B is fully scale-less

$$\begin{aligned} \bar{I}_B &= \int [d^4 q_1][d^4 q_2] \frac{1}{\bar{q}_1^4 \bar{D}_p \bar{q}_2^2 \bar{q}_{12}^2} (-2\hat{q}_2^2(\not{q}_1 + \not{p}) \\ &\quad - \bar{q}_1^2(\not{q}_1 + \not{p}) - 2\bar{q}_1^2\not{q}_2 + 4(q_2 \cdot p)\not{q}_2) = 0, \end{aligned} \tag{B.8}$$

so that self-energy corrections $\bar{I}_A + \bar{I}_B$ vanish.

The proof that scale-less integrals do not contribute can be found in [20]. Here we prove that they vanish also when q_1^2 is unbarred, as in (35). We concentrate on first term of (B.8)

$$\bar{I}_C := \int [d^4 q_1][d^4 q_2] \frac{\bar{N}_C}{\bar{q}_1^4 \bar{D}_p \bar{q}_2^2 \bar{q}_{12}^2}, \quad \bar{N}_C := \hat{q}_2^2(\not{q}_1 + \not{p}). \tag{B.9}$$

The proof is unchanged for all the other contributions. A GV subtraction is needed in front of $\hat{q}_2^2\not{q}_1$. This is achieved by using twice the identity in (15)

$$\frac{1}{\bar{D}_p} = \left[\frac{1}{\bar{q}_1^2} - \frac{2(q_1 \cdot p)}{\bar{q}_1^4} \right]_V + 4 \frac{(q_1 \cdot p)^2}{\bar{q}_1^4 \bar{D}_p}. \tag{B.10}$$

The $\hat{q}_2^2\not{p}$ piece is less UV divergent, so that a single subtraction is sufficient

$$\frac{1}{\bar{D}_p} = \left[\frac{1}{\bar{q}_1^2} \right]_V - \frac{2(q_1 \cdot p)}{\bar{q}_1^2 \bar{D}_p}. \tag{B.11}$$

The vacua are subtracted by the integral operator. That defines the numerator function associated with \bar{N}_C

$$\bar{Z}_C(\bar{q}_1^2, \hat{q}_2^2) = 4(\hat{q}_2^2\not{q}_1) \frac{(q_1 \cdot p)^2}{\bar{q}_1^4} - 2(\hat{q}_2^2\not{p}) \frac{(q_1 \cdot p)}{\bar{q}_1^2}. \tag{B.12}$$

Hence

$$\begin{aligned} \bar{Z}_C(\bar{q}_1^2, \hat{q}_2^2) \xrightarrow{\text{SIC}} \bar{Z}_C(q_1^2, q_2^2) &= +4(q_2^2\not{q}_1) \frac{(q_1 \cdot p)^2}{q_1^4} \\ &\quad - 2(q_2^2\not{p}) \frac{(q_1 \cdot p)}{q_1^2}, \end{aligned} \tag{B.13}$$

which produces

$$\begin{aligned} \bar{I}_C &= +4 \int d^4 q_1 [d^4 q_2] \frac{q_2^2\not{q}_1(q_1 \cdot p)^2}{q_1^8 D_p \bar{q}_2^2 \bar{q}_{12}^2} \\ &\quad - 2\not{p} \int d^4 q_1 [d^4 q_2] \frac{q_2^2(q_1 \cdot p)}{q_1^6 D_p \bar{q}_2^2 \bar{q}_{12}^2}. \end{aligned} \tag{B.14}$$

\bar{I}_C diverges logarithmically in the double collinear configuration in the absence of regulator. The barred q_2 -type denominators are sufficient to regulate this. That is a consequence of the fact that (B.10) and (B.11) do not alter the IR power counting. By tensor decomposition $\bar{I}_C \sim \not{p}(p^2/\mu^2 + \mathcal{O}(p^4/\mu^4))$, so that it vanishes on-shell. In summary, the GV subtraction does not leave finite pieces in scale-less integrals.




Fig. 10 The one- and two-loop QCD corrections to the massive bottom propagator. N_F quarks run in the loop

Appendix C: Correcting the bottom propagator

To renormalize the Yukawa coupling, we need the one- and two-loop QCD corrections of Fig. 10 computed at the value $\not{p} = m$. We dub them $\Sigma^{(j)} := \Sigma^{(j)}(\not{p} = m)$.

One finds

$$\Sigma^{(1)} = -m \left(\frac{\alpha_S^0}{4\pi} \right) C_F (3L'' + 5), \tag{C.15}$$

with L'' in (74).

As for the second order contribution, one has

$$i\bar{\Sigma}^{(2)} = -i \frac{2}{\pi^4} C_F N_F \left(\frac{\alpha_S}{4\pi} \right)^2 \times \int [d^4 q_1][d^4 q_2] \frac{\bar{N}(\not{p} = m)}{\bar{q}_1^4 \bar{D}_p \bar{q}_2^2 \bar{q}_{12}^2} \Big|_{p^2=m^2}, \tag{C.16}$$

with $D_p = q_1^2 + 2(q_1 \cdot p)$. The unbarred $N(\not{p})$ is given by (7) with $F_{\hat{\rho}\hat{\sigma}} = \gamma_{\hat{\rho}}(\not{q}_1 + \not{p} + m)\gamma_{\hat{\sigma}}$. The result reads $N(\not{p}) = N_A(\not{p}) + N_B(\not{p})$, where

$$\begin{aligned} N_A(\not{p}) &= \not{q}_1(\not{q}_1 + \not{p} + m)\not{q}_2 + \not{q}_2(\not{q}_1 + \not{p} + m)\not{q}_1 \\ &= 2\not{q}_2 D_p - (\not{p} - m)\not{q}_1\not{q}_2 - \not{q}_2\not{q}_1(\not{p} - m), \\ N_B(\not{p}) &= 2(q_2 \cdot q_{12})(\not{q}_1 + \not{p} - 2m) + 2\not{q}_2(\not{q}_1 + \not{p} + m)\not{q}_2. \end{aligned} \tag{C.17}$$

When barring N_A one obtains a vanishing contribution to $i\bar{\Sigma}^{(2)}$. As for N_B , one computes $N_B(\not{p} = m) = 2(q_2 \cdot q_{12})(\not{q}_1 - m) + 4(q_1 \cdot q_2)\not{q}_2 - 2\hat{q}_2^2(\not{q}_1 - 2m)$. Using tensor decomposition gives

$$N_B(\not{p} = m) \xrightarrow{\text{GR}} \bar{N}_B(\not{p} = m) \simeq \bar{N}'_B(\not{p} = m), \tag{C.18}$$

with

$$\bar{N}'_B(\not{p} = m) = m(\bar{q}_1^2 - 2\hat{q}_2^2((q_1 \cdot p)/m^2 - 2)). \tag{C.19}$$

To subtract the GV from \bar{N}'_B we expand $1/\bar{D}_p = 1/\bar{q}_1^2 + \bar{\lambda}/\bar{D}_p$ with $\bar{\lambda} = -2(q_1 \cdot p)/\bar{q}_1^2$, that gives the numerator function

$$\bar{\mathcal{Z}}(\hat{q}_1^2, \hat{q}_2^2) = m(\bar{q}_1^2 \bar{\lambda} - 2\hat{q}_2^2((q_1 \cdot p)/m^2 \bar{\lambda}^2 - 2\bar{\lambda})). \tag{C.20}$$

Hence

$$\begin{aligned} \bar{\mathcal{Z}}(\hat{q}_1^2, \hat{q}_2^2) &\xrightarrow{\text{SIC}} \bar{\mathcal{Z}}(q_1^2, q_2^2) \\ &= m(q_1^2 \lambda - 2q_2^2((q_1 \cdot p)/m^2 \lambda^2 - 2\lambda)), \end{aligned} \tag{C.21}$$

where $\lambda = -2\frac{(q_1 \cdot p)}{q_1^2}$. In summary, the two-loop correction is

$$\begin{aligned} \Sigma^{(2)} &= -\frac{2}{\pi^4} C_F N_F \left(\frac{\alpha_S}{4\pi} \right)^2 \\ &\times \int d^4 q_1 [d^4 q_2] \frac{\bar{\mathcal{Z}}(q_1^2, q_2^2)}{q_1^4 D_p q_2^2 \bar{q}_{12}^2} \Big|_{p^2=m^2} \\ &= 4m \left(\frac{\alpha_S}{4\pi} \right)^2 C_F N_F (2\tilde{I}_5 - \tilde{I}_4), \end{aligned} \tag{C.22}$$

with $\tilde{I}_{4,5}$ written in Appendix D.

Appendix D: The virtual master integrals

In this appendix we sketch out the computation of the two-loop integrals appearing in our calculation.

The q_2 integration is performed first.¹² This means computing

$$B^{:\alpha:\alpha\beta} := \int [d^4 q_2] \frac{1; q_2^\alpha; q_2^\alpha q_2^\beta}{\bar{q}_2^2 \bar{q}_{12}^2}. \tag{D.23}$$

As for B , we use the expansion in (41) to subtract its sub-vacuum. Then we use Feynman parametrization and integrate over the UV finite remainder. The result is

$$B = -i\pi^2 q_1^2 \int_0^1 dx \left(\frac{1}{x} - 2 \right) \frac{1}{D_0}, \tag{D.24}$$

with

$$D_0 := q_1^2 - \mu_0^2, \quad \mu_0^2 := \frac{\mu^2}{x(1-x)}. \tag{D.25}$$

To determine B^α we use tensor decomposition

$$B^\alpha = \frac{q_1^\alpha}{2} \int [d^4 q_2] \frac{q_{12}^2 - q_2^2 - q_1^2}{q_1^2 \bar{q}_2^2 \bar{q}_{12}^2}. \tag{D.26}$$

The first two terms cancel each other due to the $q_2^2 \leftrightarrow q_{12}^2$ symmetry of the integral. Thus

$$B^\alpha = -\frac{q_1^\alpha}{2} B. \tag{D.27}$$

Finally, tensor decomposition gives

$$\begin{aligned} B^{\alpha\beta} &= \frac{1}{3} \int [d^4 q_2] \frac{1}{\bar{q}_2^2 \bar{q}_{12}^2} \left\{ q_2^2 \left(g^{\alpha\beta} - \frac{q_1^\alpha q_1^\beta}{q_1^2} \right) \right. \\ &\quad \left. - \frac{(q_1 \cdot q_2)^2}{q_1^2} \left(g^{\alpha\beta} - 4 \frac{q_1^\alpha q_1^\beta}{q_1^2} \right) \right\}. \end{aligned} \tag{D.28}$$

¹² Assuming the appropriate GV subtraction in the rest of the integral.

The coefficients are obtained by subtracting the SV by means of (19), and integrating over the finite part. The result reads

$$\int [d^4 q_2] \frac{q_2^2}{\bar{q}_2^2 \bar{q}_{12}^2} = -\frac{i\pi^2}{2} q_1^4 \int_0^1 dx (4x^2 - 1) \frac{1}{D_0},$$

$$\int [d^4 q_2] \frac{(q_1 \cdot q_2)^2}{\bar{q}_2^2 \bar{q}_{12}^2} = \frac{q_1^4}{4} B. \tag{D.29}$$

When inserting these results in Eqs. (86) and (106) one finds that the two-loop vertex corrections can be expressed in terms of three master integrals

$$\tilde{I}_j := s^{2-j} \frac{i}{\pi^2} \int_0^1 dx \left(\frac{1}{x} - 3 + 4x^2 \right) \times \int d^4 q_1 \frac{(q_1 \cdot p_1)^{j-1}}{D_0 D_1 D_2}, \quad j = 1, 2, 3, \tag{D.30}$$

where \tilde{I}_2 is UV finite because $p_1^2 = 0$. Integrating over q_1 and x and neglecting $\mathcal{O}(\mu^2)$ terms gives

$$\tilde{I}_1 = \frac{1}{6} \left(L'^3 + 5L'^2 + L' \left(\frac{56}{3} + \pi^2 \right) - 12\zeta_3 + \frac{5}{3}\pi^2 + \frac{328}{9} \right),$$

$$\tilde{I}_2 = -\frac{1}{4} \left(L'^2 + \frac{16}{3}L' + \frac{\pi^2}{3} + \frac{104}{9} \right),$$

$$\tilde{I}_3 = -\frac{1}{16} \left(L'^2 + \frac{13}{3}L' + \frac{\pi^2}{3} + \frac{151}{18} \right), \tag{D.31}$$

with L' given in (80).

Finally, the two-loop integrals in (C.22) are

$$\tilde{I}_4 := \frac{i}{\pi^2} \int_0^1 dx \left(\frac{1}{x} - 4 + 8x^2 \right) \times \int d^4 q_1 \frac{(q_1 \cdot p_1)}{q_1^2 D_p D_0} \Big|_{p^2=m^2},$$

$$\tilde{I}_5 := \frac{i}{\pi^2 m^2} \int_0^1 dx \left(1 - 4x^2 \right) \times \int d^4 q_1 \frac{(q_1 \cdot p_1)^3}{q_1^4 D_p D_0} \Big|_{p^2=m^2}. \tag{D.32}$$

Their asymptotic expansions read

$$\tilde{I}_4 = -\frac{1}{4}L''^2 - \frac{7}{6}L'' - \frac{\pi^2}{6} - \frac{41}{18},$$

$$\tilde{I}_5 = -\frac{1}{24}L'' - \frac{13}{144}, \tag{D.33}$$

with L'' in (74).

Appendix E: The real integrals

The real component of the NNLO corrections computed in this paper can be expressed in terms of the following eight integrals

$$\tilde{R}_1 := \frac{1}{s\pi^3} \int d^4 \tilde{\Phi}_4 w(\mu^2) \frac{1}{s_{134}},$$

$$\tilde{R}_2 := \frac{1}{\pi^3} \int d^4 \tilde{\Phi}_4 w(\mu^2) \frac{1}{s_{134}s_{234}},$$

$$\tilde{R}_3 := \frac{1}{s\pi^3} \int d^4 \tilde{\Phi}_4 w(\mu^2) \frac{s_{34}}{s_{134}s_{234}},$$

$$\tilde{R}_4 := \frac{1}{\pi^3} \int d^4 \tilde{\Phi}_4 w(\mu^2) \frac{1}{s_{134}^2},$$

$$\tilde{R}_5 := \frac{1}{\pi^3} \int d^4 \tilde{\Phi}_4 w(\mu^2) \frac{1}{s_{134}s_{34}},$$

$$\tilde{R}_6 := \frac{1}{s\pi^3} \int d^4 \tilde{\Phi}_4 w(\mu^2) \frac{s_{234}}{s_{134}s_{34}},$$

$$\tilde{R}_7 := \frac{s}{\pi^3} \int d^4 \tilde{\Phi}_4 w(\mu^2) \frac{1}{s_{34}s_{134}s_{234}},$$

$$\tilde{R}_8 := \frac{1}{s\pi^3} \int d^4 \tilde{\Phi}_4 w(\mu^2) \frac{1}{s_{34}}, \tag{E.34}$$

with $w(\mu^2)$ given in (91).

To compute the \tilde{R}_i s it is convenient to use the following phase-space parametrization

$$\int d\tilde{\Phi}_4 = \frac{s^2\pi^3}{8} \int_{4\epsilon^2}^{(1-2\epsilon)^2} dz \sqrt{1 - 4\epsilon^2/z} \times \int_z^1 dy \int_{\frac{z}{y}}^{1-y+z} dx, \tag{E.35}$$

where $z = s_{34}/s$, $y = s_{234}/s$, $x = s_{134}/s$, and $\epsilon^2 = \mu^2/s$. The asymptotic $\mu^2 \rightarrow 0$ behavior can be extracted with the change of variable $\tau = 4\epsilon^2/z$, to be used when $\lim_{\epsilon \rightarrow 0}$ cannot be taken before integration.

The first three integrals are IR finite

$$\tilde{R}_1 = \frac{1}{32},$$

$$\tilde{R}_2 = \frac{\pi^2}{48} - \frac{1}{8},$$

$$\tilde{R}_3 = -\frac{\pi^2}{96} + \frac{7}{64}. \tag{E.36}$$

As for the remaining ones, (E.35) gives

$$\tilde{R}_4 = -\frac{L}{16} - \frac{25}{96},$$

$$\tilde{R}_5 = \frac{L^2}{16} + \frac{11}{24}L - \frac{\pi^2}{48} + \frac{85}{72},$$

$$\tilde{R}_6 = \frac{L^2}{32} + \frac{11}{48}L - \frac{\pi^2}{96} + \frac{349}{576},$$

$$\begin{aligned}\tilde{R}_7 &= -\frac{L^3}{48} - \frac{5}{48}L^2 + L\left(\frac{\pi^2}{24} - \frac{7}{18}\right) + \frac{5}{8}\zeta_3 \\ &\quad + \frac{5}{72}\pi^2 - \frac{41}{54}, \\ \tilde{R}_8 &= \tilde{R}_4,\end{aligned}\tag{E.37}$$

with L written in (59).

References

- G.F. Sterman, *An Introduction to Quantum Field Theory* (Cambridge University Press, Cambridge, 1994)
- R.E. Cutkosky, *J. Math. Phys.* **1**, 429 (1960). <https://doi.org/10.1063/1.1703676>
- M.J.G. Veltman, *Physica* **29**, 186 (1963). [https://doi.org/10.1016/S0031-8914\(63\)80277-3](https://doi.org/10.1016/S0031-8914(63)80277-3)
- G. 't Hooft, M.J.G. Veltman, *NATO Sci. Ser. B* **4**, 177 (1974)
- D. Stöckinger, *JHEP* **0503**, 076 (2005). <https://doi.org/10.1088/1126-6708/2005/03/076>
- G. 't Hooft, M. Veltman, *Nucl. Phys. B* **44**, 189 (1972). [https://doi.org/10.1016/0550-3213\(72\)90279-9](https://doi.org/10.1016/0550-3213(72)90279-9)
- C.G. Bollini, J.J. Giambiagi, *Nuovo Cim. B* **12**, 20 (1972). <https://doi.org/10.1007/BF02895558>
- J.C. Collins, *Renormalization* (Cambridge University Press, Cambridge, 1984)
- Z. Bern, D.A. Kosower, *Nucl. Phys. B* **379**, 451 (1992). [https://doi.org/10.1016/0550-3213\(92\)90134-W](https://doi.org/10.1016/0550-3213(92)90134-W)
- W. Siegel, *Phys. Lett. B* **84**, 193 (1979). [https://doi.org/10.1016/0370-2693\(79\)90282-X](https://doi.org/10.1016/0370-2693(79)90282-X)
- A. Signer, D. Stöckinger, *Nucl. Phys. B* **808**, 88 (2009). <https://doi.org/10.1016/j.nuclphysb.2008.09.016>
- R.A. Fazio, P. Mastrolia, E. Mirabella, W.J. Torres Bobadilla, *Eur. Phys. J. C* **74**(12), 3197 (2014). <https://doi.org/10.1140/epjc/s10052-014-3197-4>
- O.A. Battistel, A.L. Mota, M.C. Nemes, *Mod. Phys. Lett. A* **13**, 1597 (1998). <https://doi.org/10.1142/S0217732398001686>
- A.L. Cherchiglia, M. Sampaio, M.C. Nemes, *Int. J. Mod. Phys. A* **26**, 2591 (2011). <https://doi.org/10.1142/S0217751X11053419>
- R.J. Hernandez-Pinto, G.F.R. Sborlini, G. Rodrigo, *JHEP* **02**, 044 (2016). [https://doi.org/10.1007/JHEP02\(2016\)044](https://doi.org/10.1007/JHEP02(2016)044)
- G.F.R. Sborlini, F. Driencourt-Mangin, R. Hernandez-Pinto, G. Rodrigo, *JHEP* **08**, 160 (2016). [https://doi.org/10.1007/JHEP08\(2016\)160](https://doi.org/10.1007/JHEP08(2016)160)
- G.F.R. Sborlini, F. Driencourt-Mangin, G. Rodrigo, *JHEP* **10**, 162 (2016). [https://doi.org/10.1007/JHEP10\(2016\)162](https://doi.org/10.1007/JHEP10(2016)162)
- R. Pittau, *JHEP* **11**, 151 (2012). [https://doi.org/10.1007/JHEP11\(2012\)151](https://doi.org/10.1007/JHEP11(2012)151)
- A.M. Donati, R. Pittau, *JHEP* **04**, 167 (2013). [https://doi.org/10.1007/JHEP04\(2013\)167](https://doi.org/10.1007/JHEP04(2013)167)
- A.M. Donati, R. Pittau, *Eur. Phys. J. C* **74**, 2864 (2014). <https://doi.org/10.1140/epjc/s10052-014-2864-9>
- R. Pittau, *Eur. Phys. J. C* **74**(1), 2686 (2014). <https://doi.org/10.1140/epjc/s10052-013-2686-1>
- B. Page, R. Pittau, *JHEP* **11**, 183 (2015). [https://doi.org/10.1007/JHEP11\(2015\)183](https://doi.org/10.1007/JHEP11(2015)183)
- C. Gnendiger et al., *Eur. Phys. J. C* **77**(7), 471 (2017). <https://doi.org/10.1140/epjc/s10052-017-5023-2>
- C. Gnendiger, A. Signer, Private communication
- R. Pittau, *Fortschr. Phys.* **63**, 132 (2015). <https://doi.org/10.1002/prop.201400079>
- A.V. Bednyakov, B.A. Kniehl, A.F. Pikelner, O.L. Veretin, *Nucl. Phys. B* **916**, 463 (2017). <https://doi.org/10.1016/j.nuclphysb.2017.01.004>
- S.G. Gorishnii, A.L. Kataev, S.A. Larin, L.R. Surguladze, *Mod. Phys. Lett. A* **5**, 2703 (1990). <https://doi.org/10.1142/S0217732390003152>
- S.G. Gorishnii, A.L. Kataev, S.A. Larin, L.R. Surguladze, *Phys. Rev. D* **43**, 1633 (1991). <https://doi.org/10.1103/PhysRevD.43.1633>
- K.G. Chetyrkin, J.H. Kuhn, A. Kwiatkowski, *Phys. Rep.* **277**, 189 (1996). [https://doi.org/10.1016/S0370-1573\(96\)00012-9](https://doi.org/10.1016/S0370-1573(96)00012-9)
- A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover, *Nucl. Phys. B* **691**, 195 (2004). <https://doi.org/10.1016/j.nuclphysb.2004.05.017>
- R. Boughezal, J.M. Campbell, R.K. Ellis, C. Focke, W.T. Giele, X. Liu, F. Petriello, *Phys. Rev. Lett.* **116**(15), 152001 (2016). <https://doi.org/10.1103/PhysRevLett.116.152001>
- L. Magnea, E. Maina, G. Pelliccioli, C. Signorile-Signorile, P. Torrielli, S. Uccirati, *JHEP* **12**, 062 (2018). [https://doi.org/10.1007/JHEP12\(2018\)062](https://doi.org/10.1007/JHEP12(2018)062)
- J.A.M. Vermaseren, *Comput. Phys. Commun.* **83**, 45 (1994). [https://doi.org/10.1016/0010-4655\(94\)90034-5](https://doi.org/10.1016/0010-4655(94)90034-5)