



Cosmological perturbations and dynamical analysis for interacting quintessence

Ricardo G. Landim^a

Physik Department T70, Technische Universität München, James-Franck-Strasse, 85748 Garching, Germany

Received: 26 August 2019 / Accepted: 19 October 2019 / Published online: 5 November 2019
© The Author(s) 2019

Abstract We present the dynamical analysis for interacting quintessence, considering linear cosmological perturbations. Matter perturbations improve the background analysis and viable critical points describing the transition of the three cosmological eras are found. The stability of those fixed points are similar to previous studies in the literature, for both coupled and uncoupled cases, leading to a late-time attractor.

1 Introduction

Observations of Type IA Supernova indicate that the Universe undergoes an accelerated expansion [1, 2], which is dominant today ($\sim 68\%$) [3]. Ordinary matter represents only 5% of the energy content of the Universe, and the remaining 27% is the still unknown dark matter (DM). The nature of the dark sector is one of the biggest challenges in the modern cosmology, whose plethora of dark energy (DE) candidates include scalar fields [4–14], vector fields [15–21], holographic dark energy [22–38], models of false vacuum decay [39–45], modifications of gravity and different kinds of cosmological fluids [46–48]. In addition, the two components of the dark sector may interact with each other [25–28, 43, 48–69], since their densities are comparable and the interaction can eventually alleviate the coincidence problem [70, 71].

When a scalar field is in the presence of a barotropic fluid the relevant evolution equations can be converted into an autonomous system and the asymptotic states of the cosmological models can be analysed. Such approach is well-known, at the background level, for uncoupled dark energy (quintessence, tachyon field and phantom field for instance [72–76]) and coupled dark energy [21, 50, 55, 61, 77–82]. On the other hand, cosmological perturbations were only studied using dynamical analysis for Λ CDM [83, 84] and quintessence [84]. The role of cosmological perturbations in uncoupled and coupled quintessence (with diverse forms of

interactions between the dark sector) has been investigated in several works [85–96], whose aim is to constrain the free parameters of the model using sets of observations. Results from dynamical analysis are also usually employed in such works, as in [97], for instance. Therefore, it is interesting to improve the background analysis in order to understand whether the fixed points are viable to describe each one of the cosmological eras of the Universe or not. This can be done if one takes cosmological perturbations into account. In this paper we go in this direction, analysing interacting quintessence with cosmological perturbations, in the light of dynamical systems theory. Our findings mostly agree with previous results in the literature, where it was used only background equations, for coupled (and uncoupled) quintessence, including the stability of the fixed points. One of those critical points, however, no longer can describe a DE-dominated Universe, when one uses cosmological perturbations.

The rest of the paper is organized as follows. In Sect. 2 we present the basics of the interacting DE and the dynamical analysis theory. In Sect. 3 we present the dynamics of the canonical scalar field, with the correspondent equations for the background and for linear perturbations. We use the dynamical system theory in Sect. 4 to study interacting quintessence at cosmological perturbation level, analysing the critical points and their stabilities. Section 5 is reserved for conclusions. We use Planck units ($\hbar = c = M_{pl} = 1$) throughout the text.

2 Interacting dark energy and dynamical analysis

We will consider that DE is described by single real scalar field (quintessence) with energy density ρ_ϕ and pressure p_ϕ , whose equation of state is $w_\phi = p_\phi/\rho_\phi$. DE is interacting with DM through a transfer of energy-momentum between them, such that that total energy-momentum is conserved. In the flat Friedmann–Lamaître–Robertson–Walker (FLRW)

^ae-mail: ricardo.landim@tum.de

background with a scale factor a , the continuity equations for both components and for radiation are

$$\begin{aligned} \dot{\rho}_\phi + 3H(\rho_\phi + p_\phi) &= -\mathcal{Q}, \\ \dot{\rho}_m + 3H\rho_m &= \mathcal{Q}, \\ \dot{\rho}_r + 4H\rho_r &= 0, \end{aligned} \tag{1}$$

respectively, where $H = \dot{a}/a$ is the Hubble rate, \mathcal{Q} is the coupling between DM and DE and the dot is a derivative with respect to the cosmic time. A positive \mathcal{Q} corresponds to DE being transformed into DM, while negative \mathcal{Q} means the transformation in the opposite direction. In principle, the coupling can depend upon several variables $\mathcal{Q} = \mathcal{Q}(\rho_m, \rho_\phi, \dot{\phi}, H, t, \dots)$, thus we assume here the first form used in the literature $\mathcal{Q} = Q\rho_m\dot{\phi}$ [49,50], where Q is a positive constant (a negative constant would give similar results). A coupling of the form $Q\rho_\phi\dot{\phi}$ would have no cosmological perturbations, because as we will point out later, DE is expected not to cluster at sub-horizon scales [98]. On the other hand, a coupling $Q(\rho_\phi + \rho_m)\dot{\phi}$ was shown not to be viable to describe all three cosmological eras [61].

To deal with the dynamics of the system, we will define dimensionless variables. The new variables are going to characterize a system of differential equations in the form

$$X' = f[X], \tag{2}$$

where X is a column vector of dimensionless variables and the prime is the derivative with respect to $\log a$, where we set the present scale factor a_0 to be one. The critical points X_c are those ones that satisfy $X' = 0$. In order to study stability of the fixed points, we consider linear perturbations Y around them, thus $X = X_c + Y$. At the critical point the perturbations Y satisfy the following equation

$$Y' = \mathcal{J}Y, \tag{3}$$

where \mathcal{J} is the Jacobian matrix. The eigenvalues of \mathcal{J} determine if the critical points are stable (if all eigenvalues are negative), unstable (if all eigenvalues are positive) or saddle points (if at least one eigenvalue is positive and the others are negative, or vice-versa).

3 Quintessence field dynamics

The real canonical scalar field ϕ is described by the Lagrangian

$$\mathcal{L} = -\sqrt{-g} \left(\frac{1}{2} \partial^\mu \phi \partial_\mu \phi + V(\phi) \right), \tag{4}$$

where $V(\phi) = V_0 e^{-\lambda\phi}$ is the potential and V_0 and $\lambda > 0$ are constants. A negative λ is obtained if the field is replaced by $\phi \rightarrow -\phi$, thus we may restrict our attention to a positive λ . For a homogeneous field $\phi \equiv \phi(t)$ in an expanding Universe

with FLRW metric and scale factor $a \equiv a(t)$, the equation of motion becomes

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = -Q\rho_m. \tag{5}$$

In the presence of matter and radiation, the Friedmann equations are

$$H^2 = \frac{1}{3} \left(\frac{\dot{\phi}^2}{2} + V(\phi) + \rho_m + \rho_r \right), \tag{6}$$

$$\dot{H} = -\frac{1}{2} \left(\dot{\phi}^2 + \rho_m + \frac{4}{3}\rho_r \right), \tag{7}$$

and the equation of state becomes

$$w_\phi = \frac{p_\phi}{\rho_\phi} = \frac{\dot{\phi}^2 - 2V(\phi)}{\dot{\phi}^2 + 2V(\phi)}. \tag{8}$$

Cosmological perturbations are the roots of structure formation, and they are reached perturbing the energy-momentum tensor and the metric. It is convenient to work in the conformal (Newtonian) gauge, where the density perturbation $\delta \equiv \delta\rho/\rho$ and the divergence of the velocity perturbation in Fourier space $\theta \equiv a^{-1}ik^j\delta u_j$ obey the following equations for a general interacting DE model [60]

$$\begin{aligned} \dot{\delta} + \left[3H(c_s^2 - w) - \frac{\mathcal{Q}}{\rho} \right] \delta + (1+w)(\theta - 3\dot{\phi}) &= -\frac{\delta\mathcal{Q}}{\rho}, \\ \dot{\theta} + \left[H(1-3w) - \frac{\mathcal{Q}}{\rho} + \frac{\dot{w}}{1+w} \right] \theta - k^2\phi - \frac{c_s^2}{1+w}k^2\delta &= 0, \end{aligned} \tag{9}$$

where ϕ is the metric perturbation in Newtonian gauge, $c_s \equiv \delta p/\delta\rho$ is the sound speed and k^i are the components of the wave-vector in Fourier space. For radiation the density fluctuations do not cluster, and for quintessence $c_s = 1$ and DE perturbations are expected to be negligible at sub-horizon scales [98], thus they can be neglected. It is interesting, therefore, to analyse only DM perturbations, and to do so it is more convenient to merge Eq. (9) into a second-order differential equation. This is done using the Poisson equation

$$k^2\phi = -\frac{3}{2}H^2\Omega_m\delta_m, \tag{10}$$

whose result gives

$$\ddot{\delta}_m + (2H - Q\dot{\phi})\dot{\delta}_m - \frac{3}{2}H^2\Omega_m\delta_m = 0. \tag{11}$$

Now we may proceed to the dynamical analysis of the system.

4 Autonomous system

The new dimensionless variables are defined as

$$\begin{aligned} x &\equiv \frac{\dot{\phi}}{\sqrt{6}H}, & y &\equiv \frac{\sqrt{V(\phi)}}{\sqrt{3}H}, & z &\equiv \frac{\sqrt{\rho_r}}{\sqrt{3}H}, & \lambda &\equiv -\frac{V'}{V}, \\ \Gamma &\equiv \frac{V V''}{V^2}, & U_m &\equiv \frac{\delta'_m}{\delta_m}, \end{aligned} \tag{12}$$

Table 1 Critical points (x, y, z, U_m) for quintessence field. U_m was found only for the viable points that may describe one of the three cosmological eras. The table shows the correspondent equation of state for

DE (15), the effective equation of state (16) and the density parameter for DE (13)

Point	x	y	z	U_m	w_ϕ	Ω_ϕ	w_{eff}
(a1)	$-\frac{\sqrt{6}Q}{3}$	0	0	$-\frac{1}{4} \left(1 + 2Q^2 + \sqrt{4Q^4 - 12Q^2 + 25} \right)$	1	$\frac{2Q^2}{3}$	$\frac{2Q^2}{3}$
(a2)	$-\frac{\sqrt{6}Q}{3}$	0	0	$-\frac{1}{4} \left(1 + 2Q^2 - \sqrt{4Q^4 - 12Q^2 + 25} \right)$	1	$\frac{2Q^2}{3}$	$\frac{2Q^2}{3}$
(b)	0	0	1	0	-	0	$\frac{1}{3}$
(c1)	$\frac{\sqrt{6}}{2(\lambda+Q)}$	$\sqrt{\frac{2Q(Q+\lambda)+3}{2(\lambda+Q)^2}}$	0	$\frac{Q \left(2 - \sqrt{\frac{25\lambda^2+4Q^2+20\lambda Q-72}{(\lambda+Q)^2}} \right) - \lambda \left(1 + \sqrt{\frac{25\lambda^2+4Q^2+20\lambda Q-72}{(\lambda+Q)^2}} \right)}{4(\lambda+Q)}$	$-\frac{Q(Q+\lambda)}{Q(Q+\lambda)+3}$	$\frac{Q(Q+\lambda)+3}{(\lambda+Q)^2}$	$-\frac{Q}{\lambda+Q}$
(c2)	$\frac{\sqrt{6}}{2(\lambda+Q)}$	$\sqrt{\frac{2Q(Q+\lambda)+3}{2(\lambda+Q)^2}}$	0	$\frac{Q \left(2 + \sqrt{\frac{25\lambda^2+4Q^2+20\lambda Q-72}{(\lambda+Q)^2}} \right) - \lambda \left(1 - \sqrt{\frac{25\lambda^2+4Q^2+20\lambda Q-72}{(\lambda+Q)^2}} \right)}{4(\lambda+Q)}$	$-\frac{Q(Q+\lambda)}{Q(Q+\lambda)+3}$	$\frac{Q(Q+\lambda)+3}{(\lambda+Q)^2}$	$-\frac{Q}{\lambda+Q}$
(d1)	$\frac{\lambda}{\sqrt{6}}$	$\sqrt{1 - \frac{\lambda^2}{6}}$	0	0	$-1 + \frac{\lambda^2}{3}$	1	$-1 + \frac{\lambda^2}{3}$
(d2)	$\frac{\lambda}{\sqrt{6}}$	$\sqrt{1 - \frac{\lambda^2}{6}}$	0	$\frac{1}{4} (2\lambda^2 + 4\lambda Q - 8)$	$-1 + \frac{\lambda^2}{3}$	1	$-1 + \frac{\lambda^2}{3}$

where the prime is the derivative with respect to $N \equiv \ln a$.

The DE density parameter is written in terms of these new variables as

$$\Omega_\phi \equiv \frac{\rho_\phi}{3H^2} = x^2 + y^2, \tag{13}$$

thus the first Friedmann Eq. (6) becomes

$$\Omega_\phi + \Omega_m + \Omega_r = 1, \tag{14}$$

where the matter and radiation density parameter are defined by $\Omega_i = \rho_i/(3H^2)$, with $i = m, r$. From Eqs. (13) and (14) x and y are restricted in the phase plane $x^2 + y^2 \leq 1$.

The equation of state w_ϕ is written in terms of the dimensionless variables as

$$w_\phi = \frac{\epsilon x^2 - y^2}{\epsilon x^2 + y^2}, \tag{15}$$

and the total effective equation of state is

$$w_{eff} = \frac{p_\phi + p_r}{\rho_\phi + \rho_m + \rho_r} = x^2 - y^2 + \frac{z^2}{3}, \tag{16}$$

with an accelerated expansion for $w_{eff} < -1/3$. The dynamical system for the variables x, y, z, λ and U_m are

$$\frac{dx}{dN} = -3x + \frac{\sqrt{6}}{2}y^2\lambda - \frac{\sqrt{6}}{2}Q(1 - x^2 - y^2 - z^2) - xH^{-1}\frac{dH}{dN}, \tag{17}$$

$$\frac{dy}{dN} = -\frac{\sqrt{6}}{2}xy\lambda - yH^{-1}\frac{dH}{dN}, \tag{18}$$

$$\frac{dz}{dN} = -2z - zH^{-1}\frac{dH}{dN}, \tag{19}$$

$$\frac{d\lambda}{dN} = -\sqrt{6}\lambda^2x(\Gamma - 1), \tag{20}$$

$$\frac{dU_m}{dN} = -U_m(U_m + 2) + \frac{3}{2}(1 - x^2 - y^2 - z^2)$$

$$+ \sqrt{6}QxU_m - U_mH^{-1}\frac{dH}{dN}, \tag{21}$$

where

$$H^{-1}\frac{dH}{dN} = -\frac{3}{2}(1 + x^2 - y^2) - \frac{z^2}{2}. \tag{22}$$

4.1 Critical points

The fixed points of the system are obtained by setting $dx/dN = 0, dy/dN = 0, dz/dN, d\lambda/dN = 0$ and $dU_m/dN = 0$ in Eqs. (17)–(21). When $\Gamma = 1, \lambda$ is constant the potential is $V(\phi) = V_0e^{-\lambda\phi}$ [72,73]. The fixed points for coupled [50] or uncoupled quintessence [72] are well-known in the literature, and only the critical points that may satisfactorily represent one of the three cosmological eras (radiation-dominated, matter-dominated or DE-dominated) are shown in Table 1 (see [99] for a review). For those points, the additional critical point U_m was found.

The eigenvalues of the Jacobian matrix were found for each fixed point in Table 1 and the results are shown in Table 2. The eigenvalues μ_{3c} and μ_{4c} are

$$\begin{aligned} \mu_{3c,4c} = & -\frac{3\lambda^2 + 9\lambda Q + 6Q^2}{4(\lambda + Q)^2} \\ & \pm \frac{\sqrt{3}}{4(\lambda + Q)} \left[72 - 21\lambda^2 - 16\lambda Q^3 \right. \\ & \left. - 4Q^2(8\lambda^2 - 15) - 4\lambda(4\lambda^2 - 9)Q \right]^{1/2}. \end{aligned} \tag{23}$$

The points (a1) and (a2) are the so-called “ ϕ -matter-dominated epoch” (ϕ MDE) [50] and they may describe a matter-dominated universe if $\Omega_\phi = 2Q^2/3 \ll 1$. Thus μ_1 and μ_2 are negative, while μ_3 and μ_4 are always positive. Therefore (a1) and (a2) are saddle points. In the small Q limit, described above, we have $U_m = 1 - \frac{4Q^2}{5}$ for (a1) and

Table 2 Eigenvalues and stability of the fixed points

Point	μ_1	μ_2	μ_3	μ_4	Stability
(a1)	$\frac{1}{2}(2Q^2 - 3)$	$\frac{1}{2}(2Q^2 - 1)$	$\frac{1}{2}\sqrt{4Q^4 - 12Q^2 + 25}$	$\frac{1}{2}(2Q^2 + 2\lambda Q + 3)$	Saddle
(a2)	$\frac{1}{2}(2Q^2 - 3)$	$\frac{1}{2}(2Q^2 - 1)$	$-\frac{1}{2}\sqrt{4Q^4 - 12Q^2 + 25}$	$\frac{1}{2}(2Q^2 + 2\lambda Q + 3)$	Saddle
(b)	2	-1	1	0	Unstable
(c1)	$-\frac{\lambda+4Q}{2(\lambda+Q)}$	$\frac{1}{2}\sqrt{\frac{25\lambda^2+4Q^2+20\lambda Q-72}{(\lambda+Q)^2}}$	μ_{3c}	μ_{4c}	Saddle
(c2)	$-\frac{\lambda+4Q}{2(\lambda+Q)}$	$-\frac{1}{2}\sqrt{\frac{25\lambda^2+4Q^2+20\lambda Q-72}{(\lambda+Q)^2}}$	μ_{3c}	μ_{4c}	Saddle
(d1)	$\frac{1}{2}(\lambda^2 - 6)$	$\frac{1}{2}(\lambda^2 - 4)$	$\lambda^2 + \lambda Q - 3$	$\frac{\lambda^2}{2} + \lambda Q - 2$	Saddle or stable
(d2)	$\frac{1}{2}(\lambda^2 - 6)$	$\frac{1}{2}(\lambda^2 - 4)$	$\lambda^2 + \lambda Q - 3$	$-\frac{\lambda^2}{2} - \lambda Q + 2$	Saddle or stable

$U_m = -\frac{3}{2} - \frac{Q^2}{5}$ for (a2). The point (a1) correctly describes the growth of the perturbation $\delta_m \sim a$, with a small correction due to the coupling with DE. On the other hand, the point (a2) does not describe the expected growth of structures.

The radiation-dominated Universe is described by the critical point (b) and matter perturbations do not increase during this epoch ($U_m = 0$). For this case, the fixed point is unstable, because it has one positive, one negative and one zero eigenvalue [100]. At the background level (without the variable U_m) (b) is a saddle point, which indicates that the presence of linear cosmological perturbations drives the point away from the unstable equilibrium (*i.e.* saddle).

At first glance one might think that the points (c1) and (c2) could describe a matter-dominated universe, when $Q \ll \lambda$. In this situation, we would have $U_m = -\frac{1}{4} \left(1 \pm \sqrt{25 - \frac{72}{\lambda^2}} \right)$, which is real for $\lambda > \frac{6\sqrt{2}}{5}$. This value for the parameter λ was excluded by cosmological observations more than one decade ago [97, 101], being $\lambda > 1$ ruled out by at least the 3σ level [102]. Without cosmological perturbations the fixed points (c) might represent a DE-dominated Universe, although the match between the coupling constant in this case and the points (a) is difficult. In our case, both points cannot describe the late accelerated expansion of the Universe. The reason is that (c1) is a saddle point, because μ_2 is always positive while the other eigenvalues can be negative. On the other hand, (c2) could be a stable point if μ_4 were negative. This condition would be satisfied for the set of values Q and λ shown in Fig. 1, but as we said before, these values of λ are already excluded by current observations. Therefore (c2) is also a saddle point.

Points (d1) and (d2) exist for $\lambda^2 < 6$, they can describe the accelerated expansion of the Universe if $\lambda^2 < 2$ (because $w_{eff} < -1/3$) and they are either saddle or stable, depending on the values of Q and λ . The point (d1) is an attractor if $Q < \frac{4-\lambda^2}{2\lambda}$, while (d2) is attractor for $\frac{4-\lambda^2}{2\lambda} < Q < \frac{3-\lambda^2}{\lambda}$. The matter perturbation is constant ($U_m = 0$) for (d1) or decrease for (d2), indicating that the formation of structures does not happen in the DE-dominated Universe.




Fig. 1 Allowed parameter space for Q and λ in order for (c2) to be a stable point. These possible values of λ are ruled out by current cosmological observations [102]

5 Conclusions

In this paper we have used dynamical system theory to analyse the evolution of cosmological (matter) perturbations for interacting quintessence. Previous results in the literature [50], regarding the possible fixed points that represent one of each cosmological eras, are maintained (with the exception of point (c), which no longer can describe a DE-dominated Universe) and the viable cosmological transition radiation \rightarrow matter \rightarrow DE is achieved considering the sequence of critical points (a1) \rightarrow (b) \rightarrow (d1) or (d2). The stability of these points remain similar to previous studies, *i.e.*, to those ones considering only background evolution, for both coupled and uncoupled cases. Future constraints on the parameter λ will

elucidate whether quintessence can still be a DE candidate with an exponential potential or not.

Acknowledgements This work was supported by CAPES under the process 88881.162206/2017-01 and Alexander von Humboldt Foundation.

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: The paper has no associated data. All calculations were done using the software Mathematica.]

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

References

1. A.G. Riess et al., Observational evidence from supernovae for an accelerating universe and a cosmological constant. *Astron. J.* **116**, 1009–1038 (1998)
2. S. Perlmutter et al., Measurements of Omega and Lambda from 42 high redshift supernovae. *Astrophys. J.* **517**, 565–586 (1999)
3. N. Aghanim et al., Planck 2018 results. VI. Cosmological parameters (2018)
4. P.J.E. Peebles, B. Ratra, Cosmology with a time variable cosmological constant. *Astrophys. J.* **325**, L17 (1988)
5. B. Ratra, P.J.E. Peebles, Cosmological consequences of a rolling homogeneous scalar field. *Phys. Rev. D* **37**, 3406 (1988)
6. J.A. Frieman, C.T. Hill, R. Watkins, Late time cosmological phase transitions. I. Particle physics models and cosmic evolution. *Phys. Rev. D* **46**, 1226–1238 (1992)
7. J.A. Frieman, C.T. Hill, A. Stebbins, I. Waga, Cosmology with ultralight pseudo Nambu–Goldstone bosons. *Phys. Rev. Lett.* **75**, 2077 (1995)
8. R.R. Caldwell, R. Dave, P.J. Steinhardt, Cosmological imprint of an energy component with general equation of state. *Phys. Rev. Lett.* **80**, 1582 (1998)
9. T. Padmanabhan, Accelerated expansion of the universe driven by tachyonic matter. *Phys. Rev. D* **66**, 021301 (2002)
10. J.S. Bagla, H.K. Jassal, T. Padmanabhan, Cosmology with tachyon field as dark energy. *Phys. Rev. D* **67**, 063504 (2003)
11. C. Armendariz-Picon, V.F. Mukhanov, P.J. Steinhardt, A dynamical solution to the problem of a small cosmological constant and late time cosmic acceleration. *Phys. Rev. Lett.* **85**, 4438–4441 (2000)
12. P. Brax, J. Martin, Quintessence and supergravity. *Phys. Lett. B* **468**, 40–45 (1999)
13. E.J. Copeland, N.J. Nunes, F. Rosati, Quintessence models in supergravity. *Phys. Rev. D* **62**, 123503 (2000)
14. S. Vagnozzi, S. Dhawan, M. Gerbino, K. Freese, A. Goobar, O. Mena, Constraints on the sum of the neutrino masses in dynamical dark energy models with $w(z) \geq -1$ are tighter than those obtained in Λ CDM. *Phys. Rev. D* **98**(8), 083501 (2018)
15. T. Koivisto, D.F. Mota, Vector field models of inflation and dark energy. *JCAP* **0808**, 021 (2008)
16. K. Bamba, S.D. Odintsov, Inflation and late-time cosmic acceleration in non-minimal Maxwell- $F(R)$ gravity and the generation of large-scale magnetic fields. *JCAP* **0804**, 024 (2008)
17. V. Emelyanov, F.R. Klinkhamer, Possible solution to the main cosmological constant problem. *Phys. Rev. D* **85**, 103508 (2012)
18. V. Emelyanov, F.R. Klinkhamer, Reconsidering a higher-spin-field solution to the main cosmological constant problem. *Phys. Rev. D* **85**, 063522 (2012)
19. V. Emelyanov, F.R. Klinkhamer, Vector-field model with compensated cosmological constant and radiation-dominated FRW phase. *Int. J. Mod. Phys. D* **21**, 1250025 (2012)
20. S. Kouwn, P. Oh, C.-G. Park, Massive photon and dark energy. *Phys. Rev. D* **93**(8), 083012 (2016)
21. R.C.G. Landim, Dynamical analysis for a vector-like dark energy. *Eur. Phys. J. C* **76**, 480 (2016)
22. S.D.H. Hsu, Entropy bounds and dark energy. *Phys. Lett. B* **594**, 13–16 (2004)
23. M. Li, A model of holographic dark energy. *Phys. Lett. B* **603**, 1 (2004)
24. D. Pavon, W. Zimdahl, Holographic dark energy and cosmic coincidence. *Phys. Lett. B* **628**, 206–210 (2005)
25. B. Wang, Y.-G. Gong, E. Abdalla, Transition of the dark energy equation of state in an interacting holographic dark energy model. *Phys. Lett. B* **624**, 141–146 (2005)
26. B. Wang, Y. Gong, E. Abdalla, Thermodynamics of an accelerated expanding universe. *Phys. Rev. D* **74**, 083520 (2006)
27. B. Wang, C.-Y. Lin, E. Abdalla, Constraints on the interacting holographic dark energy model. *Phys. Lett. B* **637**, 357–361 (2006)
28. B. Wang, C.-Y. Lin, D. Pavon, E. Abdalla, Thermodynamical description of the interaction between dark energy and dark matter. *Phys. Lett. B* **662**, 1–6 (2008)
29. R.C.G. Landim, Holographic dark energy from minimal supergravity. *Int. J. Mod. Phys. D* **25**(4), 1650050 (2016)
30. M. Li, X.-D. Li, S. Wang, X. Zhang, Holographic dark energy models: a comparison from the latest observational data. *JCAP* **0906**, 036 (2009)
31. M. Li, X.-D. Li, S. Wang, Y. Wang, X. Zhang, Probing interaction and spatial curvature in the holographic dark energy model. *JCAP* **0912**, 014 (2009)
32. M. Li, X.-D. Li, S. Wang, Y. Wang, Dark energy. *Commun. Theor. Phys.* **56**, 525–604 (2011)
33. E.N. Saridakis, Ricci-Gauss-Bonnet holographic dark energy. *Phys. Rev. D* **97**(6), 064035 (2018)
34. A. Al Mamon, Reconstruction of interaction rate in holographic dark energy model with Hubble horizon as the infrared cut-off. *Int. J. Mod. Phys. D* **26**(11), 1750136 (2017)
35. A. Mukherjee, Reconstruction of interaction rate in Holographic dark energy. *JCAP* **1611**, 055 (2016)
36. L. Feng, X. Zhang, Revisit of the interacting holographic dark energy model after Planck 2015. *JCAP* **1608**(08), 072 (2016)
37. R. Herrera, W.S. Hipolito-Ricaldi, N. Videla, Instability in interacting dark sector: an appropriate holographic Ricci dark energy model. *JCAP* **1608**, 065 (2016)
38. M. Forte, Holographik, the k-essential approach to interactive models with modified holographic Ricci dark energy. *Eur. Phys. J. C* **76**(12), 707 (2016)
39. M. Szydlowski, A. Stachowski, K. Urbanowski, Quantum mechanical look at the radioactive-like decay of metastable dark energy. *Eur. Phys. J. C* **77**(12), 902 (2017)
40. A. Stachowski, M. Szydlowski, Urbanowski, Cosmological implications of the transition from the false vacuum to the true vacuum state. *Eur. Phys. J. C* **77**(6), 357 (2017)
41. D. Stojkovic, G.D. Starkman, R. Matsuo, Dark energy, the colored anti-de Sitter vacuum, and LHC phenomenology. *Phys. Rev. D* **77**, 063006 (2008)
42. E. Greenwood, E. Halstead, R. Poltis, D. Stojkovic, Dark energy, the electroweak vacua and collider phenomenology. *Phys. Rev. D* **79**, 103003 (2009)

43. E. Abdalla, L.L. Graef, B. Wang, A model for dark energy decay. *Phys. Lett. B* **726**, 786–790 (2013)
44. A. Shafieloo, D.K. Hazra, V. Sahni, A.A. Starobinsky, Metastable dark energy with radioactive-like decay. *Mon. Not. R. Astron. Soc.* **473**, 2760–2770 (2018)
45. R.G. Landim, E. Abdalla, Metastable dark energy. *Phys. Lett. B.* **764**, 271 (2017)
46. E.J. Copeland, M. Sami, S. Tsujikawa, Dynamics of dark energy. *Int. J. Mod. Phys. D* **15**, 1753–1936 (2006)
47. G. Dvali, G. Gabadadze, M. Porrati, 4D gravity on a brane in 5D Minkowski space. *Phys. Lett. B* **485**, 208 (2000)
48. S. Yin, B. Wang, E. Abdalla, C. Lin, Transition of equation of state of effective dark energy in the Dvali-Gabadadze-Porrati model with bulk contents. *Phys. Rev. D* **76**, 124026 (2007)
49. C. Wetterich, The Cosmon model for an asymptotically vanishing time dependent cosmological 'constant'. *Astron. Astrophys.* **301**, 321–328 (1995)
50. L. Amendola, Coupled quintessence. *Phys. Rev. D* **62**, 043511 (2000)
51. Z.-K. Guo, Y.-Z. Zhang, Interacting phantom energy. *Phys. Rev. D.* **71**, 023501 (2005)
52. R.-G. Cai, A. Wang, Cosmology with interaction between phantom dark energy and dark matter and the coincidence problem. *JCAP* **0503**, 002 (2005)
53. Z.-K. Guo, R.-G. Cai, Y.-Z. Zhang, Cosmological evolution of interacting phantom energy with dark matter. *JCAP* **0505**, 002 (2005)
54. X.-J. Bi, B. Feng, H. Li, X. Zhang, Cosmological evolution of interacting dark energy models with mass varying neutrinos. *Phys. Rev. D.* **72**, 123523 (2005)
55. B. Gumjudpai, T. Naskar, M. Sami, S. Tsujikawa, Coupled dark energy: towards a general description of the dynamics. *JCAP* **0506**, 007 (2005)
56. A.A. Costa, X.-D. Xu, B. Wang, E.G.M. Ferreira, E. Abdalla, Testing the Interaction between dark energy and dark matter with Planck data. *Phys. Rev. D* **89**(10), 103531 (2014)
57. E.G.M. Ferreira, J. Quintin, A.A. Costa, E. Abdalla, B. Wang, Evidence for interacting dark energy from BOSS. *Phys. Rev. D* **95**(4), 043520 (2017)
58. A.A. Costa, L.C. Olivari, E. Abdalla, Quintessence with Yukawa interaction. *Phys. Rev. D* **92**(10), 103501 (2015)
59. A.A. Costa, X.-D. Xu, B. Wang, E. Abdalla, Constraints on interacting dark energy models from Planck 2015 and redshift-space distortion data. *JCAP* **1701**(01), 028 (2017)
60. R.J.F. Marcondes, R.C.G. Landim, A.A. Costa, B. Wang, E. Abdalla, Analytic study of the effect of dark energy-dark matter interaction on the growth of structures. *JCAP* **1612**(12), 009 (2016)
61. F.F. Bernardi, R.G. Landim, Coupled quintessence and the impossibility of an interaction: a dynamical analysis study. *Eur. Phys. J. C* **77**(5), 290 (2017)
62. B. Wang, E. Abdalla, F. Atrio-Barandela, D. Pavon, Dark matter and dark energy interactions: theoretical challenges, cosmological implications and observational signatures. *Rep. Prog. Phys.* **79**(9), 096901 (2016)
63. G.R. Farrar, P.J.E. Peebles, Interacting dark matter and dark energy. *Astrophys. J.* **604**, 1–11 (2004)
64. S. Micheletti, E. Abdalla, B. Wang, A Field Theory Model for Dark Matter and Dark Energy in Interaction. *Phys. Rev. D* **79**, 123506 (2009)
65. W. Yang, N. Banerjee, S. Pan, Constraining a dark matter and dark energy interaction scenario with a dynamical equation of state. *Phys. Rev. D* **95**(12), 123527 (2017)
66. R.F. vom Marttens, L. Casarini, W.S. Hipólito-Ricaldi, W. Zimdahl, CMB and matter power spectra with non-linear dark-sector interactions. *JCAP* **1701**(01), 050 (2017)
67. W. Yang, S. Pan, J.D. Barrow, Large-scale stability and astronomical constraints for coupled dark-energy models. *Phys. Rev. D* **97**(4), 043529 (2018)
68. A.A. Costa, R.C.G. Landim, B. Wang, E. Abdalla, Interacting dark energy: possible explanation for 21-cm absorption at cosmic dawn. *Eur. Phys. J. C* **78**(9), 746 (2018)
69. W. Yang, S. Pan, E. Di Valentino, R.C. Nunes, S. Vagnozzi, D.F. Mota, Tale of stable interacting dark energy, observational signatures, and the H_0 tension. *JCAP* **1809**(09), 019 (2018)
70. W. Zimdahl, D. Pavon, L.P. Chimento, Interacting quintessence. *Phys. Lett. B* **521**, 133–138 (2001)
71. L.P. Chimento, A.S. Jakubi, D. Pavon, W. Zimdahl, Interacting quintessence solution to the coincidence problem. *Phys. Rev. D* **67**, 083513 (2003)
72. E.J. Copeland, A.R. Liddle, D. Wands, Exponential potentials and cosmological scaling solutions. *Phys. Rev. D* **57**, 4686–4690 (1998)
73. S.C.C. Ng, N.J. Nunes, F. Rosati, Applications of scalar attractor solutions to cosmology. *Phys. Rev. D* **64**, 083510 (2001)
74. E.J. Copeland, M.R. Garousi, M. Sami, S. Tsujikawa, What is needed of a tachyon if it is to be the dark energy? *Phys. Rev. D* **71**, 043003 (2005)
75. X.-H. Zhai, Y.-B. Zhao, A cosmological model with complex scalar field. *Nuovo Cim. B* **120**, 1007–1016 (2005)
76. J. De-Santiago, J.L. Cervantes-Cota, D. Wands, Cosmological phase space analysis of the $F(X)$ - $V(\phi)$ scalar field and bouncing solutions. *Phys. Rev. D* **87**(2), 023502 (2013)
77. S. Tsujikawa, General analytic formulae for attractor solutions of scalar-field dark energy models and their multi-field generalizations. *Phys. Rev. D* **73**, 103504 (2006)
78. L. Amendola, M. Quartin, S. Tsujikawa, I. Waga, Challenges for scaling cosmologies. *Phys. Rev. D* **74**, 023525 (2006)
79. X.-M. Chen, Y.-G. Gong, E.N. Saridakis, Phase-space analysis of interacting phantom cosmology. *JCAP* **0904**, 001 (2009)
80. N. Mahata, S. Chakraborty, Dynamical system analysis for DBI dark energy interacting with dark matter. *Mod. Phys. Lett. A* **30**(02), 1550009 (2015)
81. R.C.G. Landim, Coupled tachyonic dark energy: a dynamical analysis. *Int. J. Mod. Phys. D* **24**, 1550085 (2015)
82. R.C.G. Landim, Coupled dark energy: a dynamical analysis with complex scalar field. *Eur. Phys. J. C* **76**(1), 31 (2016)
83. A. Alho, C. Uggla, J. Wainwright, Perturbations of the Lambda-CDM model in a dynamical systems perspective. *JCAP* **1909**(09), 045 (2019)
84. S. Basilakos, G. Leon, G. Papagiannopoulos, E.N. Saridakis, Dynamical system analysis at background and perturbation levels: quintessence in severe disadvantage comparing to Λ CDM. *Phys. Rev. D* **100**(4), 043524 (2019)
85. O. Bertolami, P.J. Martins, Nonminimal coupling and quintessence. *Phys. Rev. D* **61**, 064007 (2000)
86. C. Baccigalupi, A. Balbi, S. Matarrese, F. Perrotta, N. Vittorio, Constraints on flat cosmologies with tracking quintessence from cosmic microwave background observations. *Phys. Rev. D* **65**, 063520 (2002)
87. R. Dave, R.R. Caldwell, P.J. Steinhardt, Sensitivity of the cosmic microwave background anisotropy to initial conditions in quintessence cosmology. *Phys. Rev. D* **66**, 023516 (2002)
88. V. Pettorino, C. Baccigalupi, G. Mangano, Extended quintessence with an exponential coupling. *JCAP* **0501**, 014 (2005)
89. A. W. Brookfield, C. van de Bruck, D. F. Mota, D. Tocchini-Valentini, Cosmology of mass-varying neutrinos driven by quintessence: theory and observations. *Phys. Rev. D* **73**:083515 (2006). [Erratum: *Phys. Rev. D* **76**, 049901 (2007)]
90. T. Koivisto, Growth of perturbations in dark matter coupled with quintessence. *Phys. Rev. D* **72**, 043516 (2005)

91. S. Lee, G.-C. Liu, K.-W. Ng, Constraints on the coupled quintessence from cosmic microwave background anisotropy and matter power spectrum. *Phys. Rev. D* **73**, 083516 (2006)
92. G. Olivares, F. Atrio-Barandela, D. Pavon, Matter density perturbations in interacting quintessence models. *Phys. Rev. D* **74**, 043521 (2006)
93. V. Pettorino, C. Baccigalupi, Coupled and Extended Quintessence: theoretical differences and structure formation. *Phys. Rev. D* **77**, 103003 (2008)
94. E.R.M. Tarrant, C. van de Bruck, E.J. Copeland, A.M. Green, Coupled quintessence and the halo mass function. *Phys. Rev. D* **85**, 023503 (2012)
95. E. Sefusatti, F. Vernizzi, Cosmological structure formation with clustering quintessence. *JCAP* **1103**, 047 (2011)
96. X.W.Liu, C. Heneka, L. Amendola, Constraining coupled quintessence with the 21cm signal (2019)
97. U. França, R. Rosenfeld, Fine tuning in quintessence models with exponential potentials. *JHEP* **10**, 015 (2002)
98. D. Duniya, D. Bertacca, R. Maartens, Clustering of quintessence on horizon scales and its imprint on HI intensity mapping. *JCAP* **1310**, 015 (2013)
99. S. Bahamonde, C.G. Boehmer, S. Carloni, E.J. Copeland, W. Fang, N. Tamanini, Dynamical systems applied to cosmology: dark energy and modified gravity. *Phys. Rep.* **775–777**, 1–122 (2018)
100. C.G. Boehmer, N. Chan, R. Lazkoz, Dynamics of dark energy models and centre manifolds. *Phys. Lett. B* **714**, 11–17 (2012)
101. R. Kallosh, A.D. Linde, S. Prokushkin, M. Shmakova, Supergravity, dark energy and the fate of the universe. *Phys. Rev. D* **66**, 123503 (2002)
102. Y. Akrami, R. Kallosh, A. Linde, V. Vardanyan, The landscape, the swampland and the era of precision cosmology. *Fortschr. Phys.* **67**(1–2), 1800075 (2019)