Scalar fully-heavy tetraquark states $Q\bar{Q}'\bar{Q}\bar{Q}'$ in QCD sum rules

Bo-Cheng Yang¹, Liang Tang^{1,2,a}, Cong-Feng Qiao^{3,4,b}

¹ College of Physics, Hebei Normal University, Shijiazhuang 050024, China

² Hebei Key Laboratory of Photophysics Research and Application, Hebei Normal University, Shijiazhuang 050024, China

³ School of Physics, University of Chinese Academy of Sciences, Beijing 100049, China

⁴ CAS Center for Excellence in Particle Physics, Beijing 100049, China

Received: 17 February 2021 / Accepted: 27 March 2021 / Published online: 16 April 2021

© The Author(s) 2021

Abstract Very recently, the LHCb Collaboration observed distinct structures with the $cc\bar{c}\bar{c}$ in the J/ψ -pair mass spectrum. In this work, we construct four scalar ($J^{PC} = 0^{++}$) $[8_c]_{Q\bar{Q}'} \otimes [8_c]_{\bar{Q}\bar{Q}'}$ type currents to investigate the fully-heavy tetraquark state $Q\bar{Q}'\bar{Q}\bar{Q}'$ in the framework of QCD sum rules, where $Q = c, b$ and $Q' = c, b$. Our results suggest that the broad structure around 6.2–6.8 GeV can be interpreted as the 0^{++} octet–octet tetraquark states with masses 6.44 ± 0.11 GeV and 6.52 ± 0.10 GeV, and the narrow structure around 6.9 GeV can be interpreted as the 0^{++} octet–octet tetraquark states with masses 6.87 ± 0.11 GeV and 6.96 ± 0.11 GeV, respectively. Extending to the b-quark sector, the masses of their fully-bottom partners are found to be around 18.38–18.59 GeV. Additionally, we also analyze the spectra of the $[8_c]_{c\bar{c}} \otimes [8_c]_{b\bar{b}}$ and $[8_c]_{c\bar{b}} \otimes [8_c]_{b\bar{c}}$ tetraquark states, which lie in the range of 12.51–12.74 GeV and 12.49–12.81 GeV, respectively.

1 Introduction

Hadrons with more than the minimal quark content ($q\bar{q}$ or qqq) was proposed by Gell-Mann [1] and Zweig [2] in 1964, which were named as multi-quark exotic states and were also allowed by Quantum Chromodynamics (QCD). Since the discovery of X(3872) in 2003 [3], benefited from the accumulation of more and more experimental data, many multi-quark states have been discovered at B factories, LHC and BESIII experiments. Needless to say, understanding the nature of the multi-quark exotic states is one of the most intriguing research topics of hadronic physics, which attracts more and more interests from theorists and experimentalists.

Very recently, the LHCb Collaboration reported resonant structures in the double- J/ψ invariant mass distribu-

tion using data for pp collisions at centre-of-mass energies of 7, 8 and 13 TeV collected by the LHCb experiment at the Large Hadron Collider, corresponding to an integrated luminosity of 9fb^{-1} [4]. They observed a broad structure above the threshold ranging from 6.2 to 6.8 GeV and a narrow structure at around 6.9 GeV, referred to as X(6900). Such structures are naturally assigned to have the constituent quarks $cc\bar{c}\bar{c}$, making them the first fully-heavy multi-quark exotic candidates claimed to date in the experimental literature. In the fully-bottom tetraquark sector, the CMS Collaboration observed the $\Upsilon(1S)$ pair production and indicated a $bb\bar{b}\bar{b}$ signal around 18.4 GeV with a global significance of 3.6σ [5] in 2017. However, the LHCb did not observe this tetraquark state by searching the invariant mass distribution of $\Upsilon(1S)\mu^+\mu^-$ [6, 7]. The contradictory information from CMS and LHCb Collaborations on fully-bottom tetraquark state urge more researches on its corresponding fully-charm tetraquark partner for both theorists and experimentalists.

Theoretical studies of $cc\bar{c}\bar{c}$ states have a much longer history [8]. As early as in 1981, the exotic hadrons composed of $cc\bar{c}\bar{c}$ have been systematically studied in quark–gluon model by Chao for the first time [9], where their masses were predicted to lie in the range 6.4–6.8 GeV and were all above the threshold of two charmonia. Later, the fully-charm tetraquark states were investigated in Refs. [10–12], which have predicted that fully-charm tetraquark states mainly decay into two charmonia. In recent years, the fully heavy tetraquark states attracted much attention and have been studied extensively in various model schemes [13–52]. Among these methods, the QCD sum rules technique has some peculiar advantages in exploring hadron properties involving nonperturbative QCD, and has been used to analyze X(6900) in recent works [44–52].

Since the broad structure and X(6900) respectively lie well above the $\eta_c\eta_c$ threshold and $J/\psi J/\psi$ threshold and they don't contain light flavor quark, the four-charm structures

^a e-mail: tangl@hebtu.edu.cn (corresponding author)

^b e-mail: qiaocf@ucas.ac.cn

are unlikely to be the hadronic molecules, which are usually formed by light meson exchanges with small binding energies. Then the color binding diquark–antidiquark structure is widely used to interpret the newly reported X(6900). It should be noted that according to Quantum Chromodynamics (QCD), there exists another possible tetraquark configuration $[8_c]_{Q\bar{Q}} \otimes [8_c]_{Q\bar{Q}}$, which is composed of two color-octet parts [53–58]. Since there exists a QCD interaction, it is different from molecular state with two color-singlet mesons. That is to say, it could decay to two charmonia via exchanging one or more gluons. Therefore, the study of the color-octet tetraquark state is very important for possible new exotic hadrons. In this paper, we endeavor to analyze in the framework of QCD sum rules whether there exists stable $J^{PC} = 0^{++}$ tetraquark states with color octet–octet configuration or not, and compare the results with the above-mentioned new structures reported by the LHCb Collaboration.

The rest of the paper is arranged as follows. After the introduction, some primary formulas of the QCD sum rules in our calculation are presented in Sect. 2. The numerical analysis and results are given in Sect. 3. The last part is left for conclusions and discussion of the results.

2 Formalism

The starting point of the QCD sum rules [59–63] is the two-point correlation function constructed from two hadronic currents. For a scalar state considered in this work, the two-point correlation function is expressed as the following form:

$$\Pi(q) = i \int d^4x e^{iq \cdot x} \langle 0 | T \{ j(x), j^\dagger(0) \} | 0 \rangle, \quad (1)$$

where $j(x)$ and $j(0)$ are the interpolating currents with $J^{PC} = 0^{++}$.

The interpolating currents of fully-charm or fully-bottom tetraquark states with $J^{PC} = 0^{++}$ are, respectively, constructed as

$$j_A^{0^{++}}(x) = [i \bar{Q}^j(x) \gamma^5 (t^a)_{jk} Q^k(x)] [i \bar{Q}^m(x) \gamma^5 (t^a)_{mn} Q^n(x)], \quad (2)$$

$$j_B^{0^{++}}(x) = [\bar{Q}^j(x) \gamma_\mu (t^a)_{jk} Q^k(x)] [\bar{Q}^m(x) \gamma^\mu (t^a)_{mn} Q^n(x)], \quad (3)$$

$$j_C^{0^{++}}(x) = [i \bar{Q}^j(x) (t^a)_{jk} Q^k(x)] [i \bar{Q}^m(x) (t^a)_{mn} Q^n(x)], \quad (4)$$

$$j_D^{0^{++}}(x) = [\bar{Q}^j(x) \gamma_\mu \gamma_5 (t^a)_{jk} Q^k(x)] [\bar{Q}^m(x) \gamma^\mu \gamma_5 (t^a)_{mn} Q^n(x)], \quad (5)$$

where j, k, m , and n are the color indices, the t^a is the Gell-Mann matrix, and Q represents the heavy-quark c or b . Here, the subscripts A to D of $J^{PC} = 0^{++}$ indicate the currents composed of two 0^- color-octet $Q\bar{Q}$ system, two 1^- color-octet $Q\bar{Q}$ system, two 0^+ color-octet $Q\bar{Q}$ system, and two 1^+ color-octet $Q\bar{Q}$ system, respectively.

At the quark–gluon level, the correlation function can be calculated with the operator product expansion (OPE). In our

evaluation, the heavy-quark ($Q = c$ or b) propagator $S_{ij}^Q(p)$ is considered in momentum space, which can be expanded as

$$\begin{aligned} S_{jk}^Q(p) = & \frac{i \delta_{jk} (\not{p} + m_Q)}{p^2 - m_Q^2} - \frac{i}{4} \frac{t_{jk}^a G_{a\beta}^a}{(p^2 - m_Q^2)^2} [\sigma^{\alpha\beta} (\not{p} + m_Q) \\ & + (\not{p} + m_Q) \sigma^{\alpha\beta}] \\ & + \frac{i \delta_{jk} m_Q \langle g_s^2 G^2 \rangle}{12(p^2 - m_Q^2)^3} \left[1 + \frac{m_Q (\not{p} + m_Q)}{p^2 - m_Q^2} \right] \\ & + \frac{i \delta_{jk}}{48} \left\{ \frac{(\not{p} + m_Q) [\not{p}(p^2 - 3m_Q^2) + 2m_Q(2p^2 - m_Q^2)]}{(p^2 - m_Q^2)^6} \right. \\ & \left. \times (\not{p} + m_Q) \right\} \langle g_s^3 G^3 \rangle. \end{aligned} \quad (6)$$

Here, the vacuum condensates are clearly displayed. For more interpretation on above propagators, the reader is referred to Refs. [64, 65].

Based on the dispersion relation, the correlation function $\Pi(q^2)$ of the quark–gluon side can be expressed as

$$\Pi^{OPE}(q^2) = \int_{(4m_Q)^2}^{\infty} ds \frac{\rho^{OPE}(s)}{s - q^2} + \Pi^{(GG)}(q^2), \quad (7)$$

where $\rho^{OPE}(s) = \text{Im}[\Pi^{OPE}(s)]/\pi$ and

$$\rho^{OPE}(s) = \rho^{\text{pert}}(s) + \rho^{(GG)}(s). \quad (8)$$

The second term $\Pi^{(GG)}(q^2)$ in Eq. (7) represents the contribution in the correlation function that have no imaginary part but have nontrivial magnitudes after the Borel transformation. After making the Borel transformation to Eq. (7), we can obtain

$$\Pi^{\text{OPE}}(M_B^2) = \int_{(4m_Q)^2}^{\infty} ds \rho^{\text{OPE}}(s) e^{-s/M_B^2} + \Pi^{(GG)}(M_B^2). \quad (9)$$

For all the tetraquark states considered in this work, we put the lengthy expressions of spectral densities $\rho^{OPE}(s)$ and $\Pi^{(GG)}(M_B^2)$ in Eq. (9) into the Appendix.

On the phenomenological side, after separating the ground state contribution from the pole term, the correlation function $\Pi(q^2)$ can be expressed as the dispersion integral over the physical region, i.e.,

$$\Pi(q^2) = \frac{(\lambda_X)^2}{(M_X)^2 - q^2} + \frac{1}{\pi} \int_{s_0}^{\infty} ds \frac{\rho(s)}{s - q^2}, \quad (10)$$

where the subscript X means the lowest lying tetraquark state, M_X denotes its mass, and $\rho(s)$ is the spectral density that contains the contribution from higher excited states and the continuum states above the threshold s_0 . As in Refs. [61, 63], in order to work out the phenomenological side of the QCD sum rules, a complete set of intermediate states should be inserted into the two color-octet–octet tetraquark interpolating currents, where the summation goes over all possible

hadronic states created by the color-octet–octet tetraquark current. The coupling constant λ_X is defined through

$$\langle 0 | j^{0^{++}}(0) | X \rangle = \lambda_X, \quad (11)$$

where X stands for the tetraquark state.

By performing the Borel transform on the phenomenological side, Eq. (10), and matching it to Eq. (9), we can then obtain the mass of the tetraquark state,

$$M_X^i(s_0, M_B^2) = \sqrt{-\frac{L_1(s_0, M_B^2)}{L_0(s_0, M_B^2)}}, \quad (12)$$

where the superscript i runs from A to D , respectively. The moments L_1 and L_0 are, respectively, defined as

$$L_0(s_0, M_B^2) = \int_{(4m_Q)^2}^{\infty} ds \rho^{OPE}(s) e^{-s/M_B^2} + \Pi^{(GG)}(M_B^2), \quad (13)$$

$$L_1(s_0, M_B^2) = \frac{\partial}{\partial (M_B^2)^{-1}} L_0(s_0, M_B^2). \quad (14)$$

3 Numerical evaluation

In order to yield meaningful physical results in QCD sum rules, as in any practical theory, one needs to give certain inputs. To perform numerical analyses, we use the following values for various condensate and heavy quark masses [59–63]: $m_c(m_c) = \bar{m}_c = (1.27 \pm 0.03) \text{ GeV}$, $m_b(m_b) = \bar{m}_b = (4.18 \pm 0.03) \text{ GeV}$, $\langle g_s^2 G^2 \rangle = 0.48 \pm 0.14 \text{ GeV}^4$, where, $\bar{m}_c$ and $\bar{m}_b$ represent heavy-quark running masses in $\overline{MS}$ scheme.

Moreover, there exist two additional parameters M_B^2 and s_0 introduced in establishing the sum rule, which will be constrained by three criteria [59–61, 63]. First, to extract the information on ground state tetraquark state, one should guarantee pole contribution (PC) is bigger than the continuum contribution [63, 66], which can be determined by the formula

$$R_i^{PC} = \frac{L_0(s_0, M_B^2)}{L_0(\infty, M_B^2)}, \quad (15)$$

where the subscript i runs from A to D . Under this prerequisite, the most part of the contribution in the mass equation (12) comes directly from the ground state, and the critical value of M_B^2 is the upper limit $(M_B^2)_{max}$.

The second one asks for the convergence of the OPE, which fixes the lower limit on M_B^2 , that is, $(M_B^2)_{min}$. In general, one can determine the $(M_B^2)_{min}$ value by the following ratio

$$R_i^{OPE} = \frac{L_0^{dim}(s_0, M_B^2)}{L_0(s_0, M_B^2)}, \quad (16)$$

where the contribution of the higher dimension condensate in the OPE side is smaller than 10% to 25% of the total contribution [63, 65]. Here the superscript dim means the dimension of relevant condensate in the OPE of Eq. (13), the subscript i runs from A to D . As in Refs. [44, 45, 48, 49], for the fully heavy tetraquark systems, the three-gluon condensate is not only lower than the two-gluon contribution but also too tiny in our calculations. Then we conclude that if M_B^2 are larger than 4.0 GeV^2 , the operator product expansion is well convergent for the scalar fully-charm tetraquark states. Therefore, for simplicity, we neglect three-gluon condensate in our numerical analyses in the following.

The third criterion is to require the dependence of the mass of the tetraquark state M_X on the parameter s_0 to be weak. To find a proper value for continuum threshold s_0 , we perform a similar analysis as in Refs. [67–69]. Notice that the s_0 relates to the mass of the ground state by $\sqrt{s_0} \sim (M_X + \delta) \text{ GeV}$, in which δ lies in the scope of $0.4 \sim 0.8 \text{ GeV}$. Therefore, various $\sqrt{s_0}$ satisfying this constraint should be taken into account in the numerical analysis. Among these values, we need to pick up the one which yields an optimal window for Borel parameter M_B^2 . That is to say, in the optimal window, the fully heavy tetraquark mass M_X is somehow independent of the Borel parameter M_B^2 . Eventually, the value of $\sqrt{s_0}$ corresponding to the optimal mass curve will be taken as its central value. In practice, in order to estimate the uncertainties stemming from s_0 , we may vary $\sqrt{s_0}$ by 0.20 GeV in numerical calculation [68], which set the upper and lower bounds on $\sqrt{s_0}$.

For 0^{++} fully-charm tetraquark states, we plot the mass curves as functions of the Borel parameter M_B^2 for different $\sqrt{s_0}$ in Fig. 1, where A to D stand for cases A to D , and the two vertical lines in each case indicate the upper and lower bounds of the valid Borel window with the central value of $\sqrt{s_0}$. We search for the Borel parameter M_B^2 and continuum threshold parameters s_0 to satisfy the two criteria of the QCD sum rules: pole dominance at the phenomenological side and convergence of the operator product expansion at the QCD side. Furthermore, we take the relation $\sqrt{s_0} = M_X + (0.4 - 0.8) \text{ GeV}$ as an additional constraint to obey.

The resulting Borel parameters, continuum threshold parameters, pole contributions are shown explicitly in Table 1. From the table, we can see that the pole dominance at the phenomenological side is well satisfied. In the Borel windows, the contributions of the two-gluon condensate are smaller than the total contribution. Moreover, the mass curves in these Borel windows have the optimal platform. Now the three criteria of the QCD sum rules are all satisfied, we expect to make reasonable predictions.

After the above evaluation, we can then determine the masses of the scalar fully-charm tetraquark states with cur-

Fig. 1 The $[8_c]_{c\bar{c}} \otimes [8_c]_{c\bar{c}}$ tetraquark masses M_X as a function of the Borel parameter M_B^2 with $J^{PC} = 0^{++}$ and different values of $\sqrt{s_0}$, where A to D stand for cases A to D, and the two vertical lines in each case indicate the upper and lower bounds of the valid Borel window with the central value of $\sqrt{s_0}$

Table 1 The windows of the Borel parameter, continuum thresholds, pole contributions, two-gluon contributions, and predicted masses for $[8_c]_{c\bar{c}} \otimes [8_c]_{c\bar{c}}$ tetraquark states

	$M_B^2 (\text{GeV}^2)$	$\sqrt{s_0} (\text{GeV})$	PC	$R_i^{(GG)}$	$M_X (\text{GeV})$
0^{++} case A	4.7–5.3	7.0 ± 0.2	61–50%	0.37–0.24%	6.44 ± 0.11
0^{++} case B	4.9–5.6	7.1 ± 0.2	60–50%	0.05–0.02%	6.52 ± 0.11
0^{++} case C	4.0–4.6	7.3 ± 0.2	62–50%	28.98–23.60%	6.87 ± 0.10
0^{++} case D	6.1–7.0	7.7 ± 0.2	62–50%	2.77–2.48%	6.96 ± 0.11

Table 2 Predicted masses of the ground states for $[8_c]_{c\bar{c}} \otimes [8_c]_{b\bar{b}}$, $[8_c]_{c\bar{b}} \otimes [8_c]_{b\bar{c}}$, and $[8_c]_{b\bar{b}} \otimes [8_c]_{b\bar{b}}$, where cases A to D correspond to the currents in Eqs. (2–5)

	Case A	Case B	Case C	Case D
$[8_c]_{c\bar{c}} \otimes [8_c]_{b\bar{b}}$	12.51 ± 0.10	12.58 ± 0.10	12.67 ± 0.10	12.74 ± 0.11
$[8_c]_{c\bar{b}} \otimes [8_c]_{b\bar{c}}$	12.49 ± 0.11	12.58 ± 0.10	12.75 ± 0.11	12.81 ± 0.10
$[8_c]_{b\bar{b}} \otimes [8_c]_{b\bar{b}}$	18.38 ± 0.11	18.44 ± 0.10	18.50 ± 0.10	18.59 ± 0.11

rents A to D, that is

$$M_X^{A,0^{++}} = (6.44 \pm 0.11) \text{ GeV}, \quad (17)$$

$$M_X^{B,0^{++}} = (6.52 \pm 0.11) \text{ GeV}, \quad (18)$$

$$M_X^{C,0^{++}} = (6.87 \pm 0.10) \text{ GeV}, \quad (19)$$

$$M_X^{D,0^{++}} = (6.96 \pm 0.11) \text{ GeV}, \quad (20)$$

where the central values correspond to the results with the optimal stability of M_B^2 , and the errors stem from the uncertainties of the condensates, the quark mass, the threshold parameter $\sqrt{s_0}$, and the Borel parameter M_B^2 .

By swapping the flavor of the heavy quarks in the interpolating currents and performing the same calculation and analysis, we can obtain the corresponding results of the fully-heavy tetraquark states for $[8_c]_{c\bar{c}} \otimes [8_c]_{b\bar{b}}$, $[8_c]_{c\bar{b}} \otimes [8_c]_{b\bar{c}}$, and $[8_c]_{b\bar{b}} \otimes [8_c]_{b\bar{b}}$, whose masses are shown in Table 2, respectively.

4 Conclusions

Very recently, the LHCb Collaboration reported their discovery of resonance-like structure in the di- J/ψ mass spectrum, which opens a whole new arena for multiquark exotic hadron. The narrow structure X(6900) is the first clear candidate for a multiquark exotic state that are composed of four charm quarks $cc\bar{c}\bar{c}$. This fully-heavy sector is particularly interesting from a theoretical point of view, since the molecular structure popular for multiquark states that contain light flavor is much less viable, leaving the color binding structures (such as the diquark–antidiquark structure and the octet–octet tetraquark structure) as the leading candidates.

In this work, we study the $J^{PC} = 0^{++}$ fully-heavy tetraquark states via the QCD sum rules by constructing the octet–octet type currents. In the calculation, we consider the nonperturbative condensate contributions up to dimension 4 in the operator product expansion. Using the interpolating currents in Eqs. (2–5), we investigate all these four currents and collect the masses of these $[8_c]_{c\bar{c}} \otimes [8_c]_{c\bar{c}}$ in Table 1. While comparing numerically to the recent LHCb results, we find the masses of currents (2) and (3) are apparently lower than X(6900), but in the scope of the broad structure from 6.2 to 6.8 GeV, this implies that these two configurations should make contributions to the broad structure as well as other possible configurations, for example, the $J/\psi - J/\psi$ molecule [50]. Therefore the mixing effect is the reason that the structure in the range (6.2, 6.8) GeV observed by the LHCb Collaboration is a broad one. Particularly, since the sum rules of the currents (4) and (5) can really produce the mass of X(6900), we may probably conclude that the best interpretation is assigning X(6900) to the 0^{++} octet–octet tetraquark states with the configurations as currents (4) and (5).

Extending to the b-quark sector, the masses of their fully-bottom partners are found to lie in the region 18.38–18.59 GeV. Additionally, we also analyze the spectra of the $[8_c]_{c\bar{c}} \otimes [8_c]_{b\bar{b}}$ and $[8_c]_{c\bar{b}} \otimes [8_c]_{b\bar{c}}$ tetraquark states, which

lie in the range of 12.51–12.74 GeV and 12.49–12.81 GeV, respectively.

Acknowledgements This work was supported in part by the National Natural Science Foundation of China (NSFC) under the Grants 11975236 and 11635009; Science Foundation of Hebei Normal University under Contract No. L2016B08; and National Key Research and Development Program of China under Contracts no. 2020YFA0406400.

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: There are no associated data available.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.
Funded by SCOAP³.

Appendix

We list in this appendix the explicit expressions of the QCD spectral densities $\rho(s)$ and $\Pi(M_B^2)$ in Eq. (9) for all currents shown in Eqs. (2–5).

For the $[Q\bar{Q}][Q\bar{Q}]$ octet–octet tetraquark states with $Q = c, b$, the spectral densities $\rho(s)$ and $\Pi(M_B^2)$ are

$$\begin{aligned} \rho_{j,\text{pert}}^{0^{++}}(s) &= \frac{1}{27 \times \pi^5} \int_{16m_Q^2}^{s_0} ds \int_{x_i}^{x_f} dx \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \\ &\times dz \left\{ \frac{\mathcal{F}_{xyz}^4 x y z \mathcal{A}_{xyz}}{2} \right. \\ &- \frac{\mathcal{F}_{xyz}^3 m_Q^2 [6\hat{s}xyz\mathcal{A}_{xyz} + \mathcal{N}_j 2z\mathcal{A}_{xyz} + \mathcal{N}_j 2xy]}{3} \\ &\left. + \mathcal{F}_{xyz}^2 m_Q^4 [\hat{s}^2 xyz\mathcal{A}_{xyz} + \mathcal{N}_j \hat{s}z\mathcal{A}_{xyz} + \mathcal{N}_j \hat{s}xy + 1] \right\}, \quad (21) \\ \rho_{j,\langle GG \rangle}^{0^{++}}(s) &= \frac{\langle g_s^2 GG \rangle}{2^{13} \times 3^2 \times \pi^5 x^3 y^3 z^3 \mathcal{A}_{xyz}^3} \int_{16m_Q^2}^{s_0} ds \int_{x_i}^{x_f} dx \\ &\times \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \\ &\times \left\{ - \frac{\mathcal{F}_{xyz}^2 [18x^4 z^3 y^4 \mathcal{A}_{xyz}^3 + 18x^3 y^3 z^4 \mathcal{A}_{xyz}^4]}{2} \right. \\ &+ \mathcal{F}_{xyz} m_Q^2 [\mathcal{A}_{xyz}^4 [48xy^3z^4 \\ &\times (y - \mathcal{N}_j) + 48x^4 yz(y^3 + z^3) + \mathcal{N}_j 12x^2 y^2 z^4 \\ &- \mathcal{N}_j 48x^3 yz^4 + 18\hat{s}x^3 y^3 z^4] \\ &\left. + \mathcal{A}_{xyz}^3 (18\hat{s}x^4 y^4 z^3 - \mathcal{N}_j 48x^4 y^4 z + \mathcal{N}_j 12x^3 y^3 z^3) \right\} \end{aligned}$$

$$\begin{aligned}
& +12x^4y^4z^2\mathcal{A}_{xyz}^2 \\
& +48x^4y^4z^3(z-\mathcal{N}_j)\mathcal{A}_{xyz}] - m_Q^4[\mathcal{A}_{xyz}^4[\mathcal{N}_j16x^3z^4 \\
& +3\hat{s}^2x^3y^3z^4 + \mathcal{N}_j16x^3y^3z \\
& + \mathcal{N}_j16y^3z^4 + 6\mathcal{N}_j\hat{s}x^2y^2z^4 - \mathcal{N}_j24\hat{s}x^3yz^4 \\
& +24\hat{s}xy^3z^4(2y-\mathcal{N}_j) \\
& +48\hat{s}x^4yz(y^3+z^3)] + \mathcal{A}_{xyz}^3[\mathcal{N}_j16x^4y^4 + 3\hat{s}^2x^4y^4z^3 \\
& + \mathcal{N}_j6\hat{s}x^3y^3z^3 \\
& +6x^2y^2z^3 + \mathcal{N}_j16x^4yz^3 - 24x^3yz^3 \\
& + \mathcal{N}_j8xy^3z^3(2y-\mathcal{N}_j) \\
& -24x^3y^3z - \mathcal{N}_j24\hat{s}x^4y^4z] + \mathcal{A}_{xyz}^2[6x^3y^3z^2 \\
& + \mathcal{N}_j6\hat{s}x^4y^4z^2] \\
& + \mathcal{A}_{xyz}[\mathcal{N}_j8x^3y^3z^3(2z-\mathcal{N}_j) + 24\hat{s}x^4y^4z^3(2z-\mathcal{N}_j) \\
& + \mathcal{N}_j16x^4y^4z^3] \Big\}, \quad (22)
\end{aligned}$$

$$\begin{aligned}
\Pi_{j,(GG)}^{0++}(M_B^2) = & -\frac{\langle g_s^2 GG \rangle}{2^9 \times 3^2 \pi^6} \int_0^1 \\
& \times dx \int_0^{1-x} dy \int_0^{1-x-y} dz e^{-\frac{f_{xyz}m_Q^2}{M_B^2}} \\
& \times \frac{1}{x^3y^3z^3\mathcal{A}_{xyz}^3} m_Q^6 (f_{xyz}z\mathcal{A}_{xyz} + \mathcal{N}_j)(f_{xyz}xy + \mathcal{N}_j) \\
& \times \left\{ \mathcal{A}_{xyz}^3 [x^3(y^3+z^3) + y^3z^3] + x^3y^3z^3 \right\}, \quad (23)
\end{aligned}$$

$$\begin{aligned}
\rho_{k,pert}^{0++}(s) = & \frac{1}{2^6 \times \pi^5} \int_{16m_Q^2}^{s_0} ds \int_{x_i}^{x_f} dx \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \\
& \times \left\{ \mathcal{F}_{xyz}^4 x y z \mathcal{A}_{xyz} \right. \\
& - \frac{\mathcal{F}_{xyz}^3 m_Q^2 [12\hat{s}xyz\mathcal{A}_{xyz} + \mathcal{N}_k 2z\mathcal{A}_{xyz} + \mathcal{N}_k 2xy]}{3} \\
& \left. + \mathcal{F}_{xyz}^2 m_Q^4 [2\hat{s}^2xyz\mathcal{A}_{xyz} + \mathcal{N}_k \hat{s}z\mathcal{A}_{xyz} + \mathcal{N}_k \hat{s}xy + 2] \right\}, \quad (24)
\end{aligned}$$

$$\begin{aligned}
\rho_{k,(GG)}^{0++}(s) = & \frac{\langle g_s^2 GG \rangle m_Q^2}{2^{10} \times 3^2 \times \pi^5} \int_{16m_Q^2}^{s_0} ds \int_{x_i}^{x_f} dx \int_{y_i}^{y_f} dy \\
& \times dy \int_{z_i}^{z_f} dz \frac{1}{x^3y^3z^3\mathcal{A}_{xyz}^3} \\
& \times \left\{ 6\mathcal{F}_{xyz}xyz\mathcal{A}_{xyz}[\mathcal{A}_{xyz}^3[8x^3(y^3+z^3) \right. \\
& - \mathcal{N}_k 4x^2z^3 + 4y^2z^3(2y-\mathcal{N}_k)] \\
& - \mathcal{N}_k x^2y^2(4xy+z^2)\mathcal{A}_{xyz}^2 + 4x^3y^3z^2 \\
& \times (2z-\mathcal{N}_k)] - m_Q^2[4z\mathcal{A}_{xyz}^4[12\hat{s}x^4y \\
& \times (y^3+z^3) + \mathcal{N}_k x^3(-3\hat{s}yz^3 \\
& + 2y^3+2z^3) + 3\hat{s}xy^3z^3(4y-\mathcal{N}_k) + \mathcal{N}_k 2y^3z^3] \\
& - xy\mathcal{A}_{xyz}^3[4x^3\mathcal{N}_k(y^3(3\hat{s}-2)-2z^3) + \mathcal{N}_k 3x^2(y^2z(\hat{s}z^2 \\
& + 8\mathcal{N}_k) + \mathcal{N}_k 8z^3) \\
& - \mathcal{N}_k 8y^2z^3(y-3\mathcal{N}_k)] + 4x^3y^3z^3\mathcal{A}_{xyz}(3\hat{s}xy(4z-\mathcal{N}_k) \\
& + 2\mathcal{N}_k(z-3\mathcal{N}_k))] \Big\}
\end{aligned}$$

$$+ \mathcal{N}_k 8x^4y^4z^3] \Big\}, \quad (25)$$

$$\begin{aligned}
\Pi_{k,(GG)}^{0++}(M_B^2) = & -\frac{\langle g_s^2 GG \rangle}{2^8 \times 3^2 \times \pi^6} \int_0^1 dx \int_0^{1-x} dy \\
& \times \int_0^{1-x-y} dz e^{-\frac{f_{xyz}m_Q^2}{M_B^2}} \\
& \times \frac{1}{x^3y^3z^3\mathcal{A}_{xyz}^3} m_Q^6 \{ \mathcal{A}_{xyz}^3 [x^3(y^3+z^3) \\
& + y^3z^3] + x^3y^3z^3 \} \\
& \times [2f_{xyz}^2xyz\mathcal{A}_{xyz} \\
& + \mathcal{N}_k f_{xyz}(z\mathcal{A}_{xyz} + xy) + 2], \quad (26)
\end{aligned}$$

where the subscript j represents A and C , and k denotes B and D , and the factors $\mathcal{N}_j$ and $\mathcal{N}_k$ have the following definition: $\mathcal{N}_A = \mathcal{N}_B = 1$ and $\mathcal{N}_C = \mathcal{N}_D = -1$.

For the $[c\bar{c}][b\bar{b}]$ octet–octet tetraquark states, the spectral densities $\rho(s)$ and $\Pi(M_B^2)$ read

$$\begin{aligned}
\rho_{j,pert}^{0++}(s) = & \frac{1}{2^8 \times \pi^5} \int_{(2m_c+2m_b)^2}^{s_0} ds \int_{x_i}^{x_f} dx \\
& \times dy \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \left\{ \frac{\mathcal{F}_{xyz}^4 xyz\mathcal{A}_{xyz}}{2} \right. \\
& - \frac{\mathcal{F}_{xyz}^3 [6m_b^2\hat{s}xyz\mathcal{A}_{xyz} + \mathcal{N}_j 2m_c^2z\mathcal{A}_{xyz} + \mathcal{N}_j 2m_b^2xy]}{3} \\
& \left. + \mathcal{F}_{xyz}^2 [m_b^4\hat{s}^2xyz\mathcal{A}_{xyz} + \mathcal{N}_j m_b^2m_c^2\hat{s}z\mathcal{A}_{xyz} + \right. \\
& \left. \times \mathcal{N}_j m_b^4\hat{s}xy + m_b^2m_c^2] \right\}, \quad (27)
\end{aligned}$$

$$\begin{aligned}
\rho_{j,(GG)}^{0++}(s) = & \frac{\langle g_s^2 GG \rangle}{2^{13} \times 3^2 \times \pi^5} \int_{(2m_c+2m_b)^2}^{s_0} ds \int_{x_i}^{x_f} dx \int_{y_i}^{y_f} dy \\
& \times dy \int_{z_i}^{z_f} dz \frac{1}{x^3y^3z^3\mathcal{A}_{xyz}^3} \\
& \times \left\{ -\frac{\mathcal{F}_{xyz}^2 [18x^3z^3y^4\mathcal{A}_{xyz}^4 + 18x^4y^4z^3\mathcal{A}_{xyz}^3]}{2} \right. \\
& + \mathcal{F}_{xyz}[\mathcal{A}_{xyz}^4[48x^4yz(m_b^2y^3 + m_c^2z^3) \\
& + 18m_b^2\hat{s}x^3y^3z^4 - \mathcal{N}_j 48m_c^2x^3yz^4 + \mathcal{N}_j 12m_c^2x^2y^2z^4 \\
& + 48m_c^2xy^3z^4(y-\mathcal{N}_j)] \\
& + \mathcal{A}_{xyz}^3[18m_b^2\hat{s}x^4y^4z^3 - \mathcal{N}_j 48m_b^2x^4y^4z \\
& + \mathcal{N}_j 6m_b^2x^3y^3z^3 + \mathcal{N}_j 6m_c^2x^3y^3z^3] \\
& + \mathcal{A}_{xyz}^2[\mathcal{N}_j 12m_b^2x^4y^4z^2 + 48m_b^2x^4y^4z^3(z-\mathcal{N}_j)\mathcal{A}_{xyz}] \\
& - [\mathcal{A}_{xyz}^4[3m_b^4\hat{s}^2x^3y^3z^4 \\
& + 48m_b^2\hat{s}x^4yz(m_b^2y^3 + m_c^2z^3) - \mathcal{N}_j 24m_b^2m_c^2\hat{s}x^3yz^4 \\
& + \mathcal{N}_j 6m_b^2m_c^2\hat{s}x^2y^2z^4 \\
& + 24m_b^2m_c^2\hat{s}xy^3z^4(2y-\mathcal{N}_j) + \mathcal{N}_j 16\hat{s}xy^3z^4(2y-\mathcal{N}_j) \\
& + \mathcal{N}_j 16m_b^2m_c^2x^3y^3z] + \mathcal{A}_{xyz}^3[3m_b^4\hat{s}^2x^4y^4z^3 \\
& + 6m_b^2m_c^2x^2y^2z^3 - 24m_b^2m_c^2x^3yz^3 \\
& - \mathcal{N}_j 24m_b^4\hat{s}x^4y^4z + \mathcal{N}_j 16m_b^4x^4y^4] \Big\}
\end{aligned}$$

$$\begin{aligned}
& +\mathcal{N}_j 3m_b^2(m_b^2 + m_c^2)\hat{s}x^3y^3z^3 \\
& +\mathcal{N}_j 16m_b^2m_c^2x^4yz^3 - 24m_b^2m_c^2x^3y^3z \\
& +\mathcal{N}_j 8m_b^2m_c^2xy^3z^3(2y - 3\mathcal{N}_j)] \\
& +\mathcal{A}_{xyz}^2[\mathcal{N}_j 6m_b^4\hat{s}x^4y^4z^2 + 6m_b^2m_c^2x^3y^3z^2] \\
& +\mathcal{A}_{xyz}[24m_b^4\hat{s}x^4y^4z^3(2z - \mathcal{N}_j) \\
& +\mathcal{N}_j 8m_b^2m_c^2x^3y^3z^3(2z - 3\mathcal{N}_j)] + \mathcal{N}_j 16m_b^4x^4y^4z^3 \\
& +\mathcal{N}_j 16m_c^4x^3z^4 + \mathcal{N}_j 16m_c^4y^3z^4] \Big\}, \quad (28)
\end{aligned}$$

$$\begin{aligned}
\Pi_{j,(GG)}^{0++}(M_B^2) &= -\frac{\langle g_s^2 GG \rangle}{2^{10} \times 3^2 \times \pi^6} \int_0^1 \\
&\times dx \int_0^{1-x} dy \int_0^{1-x-y} dz e^{-\frac{f_{xyz}m_b^2}{M_B^2}} \\
&\times \frac{1}{x^3y^3z^3\mathcal{A}_{xyz}^3} m_b^2(f_{xyz}z\mathcal{A}_{xyz} + \mathcal{N}_j)(f_{xyz}m_b^2 \\
&xy + \mathcal{N}_jm_c^2) \\
&\times [m_b^2x^3y^3(\mathcal{A}_{xyz}^3 + z^3) + \mathcal{A}_{xyz}^3m_c^2z^3(x^3 + y^3)], \quad (29)
\end{aligned}$$

$$\begin{aligned}
\rho_{k,pert}^{0++}(s) &= \frac{1}{2^7 \times \pi^5} \int_{(2m_c+2m_b)^2}^{s_0} ds \int_{x_i}^{x_f} \\
&\times dx \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \left\{ \mathcal{F}_{xyz}^4 xy z \mathcal{A}_{xyz} \right. \\
&- \frac{\mathcal{F}_{xyz}^3 [12m_b^2\hat{s}xyz\mathcal{A}_{xyz} + \mathcal{N}_k 2m_c^2z\mathcal{A}_{xyz} + \mathcal{N}_k 2m_b^2xy]}{3} \\
&+ \mathcal{F}_{xyz}^2 [2m_b^4\hat{s}^2xyz\mathcal{A}_{xyz} + \mathcal{N}_k m_b^2m_c^2\hat{s}z\mathcal{A}_{xyz} \\
&+ \mathcal{N}_k m_b^4\hat{s}xy + m_b^2m_c^2] \Big\}, \quad (30)
\end{aligned}$$

$$\begin{aligned}
\rho_{k,(GG)}^{0++}(s) &= \frac{\langle g_s^2 GG \rangle}{2^{12} \times 3^2 \pi^5} \int_{(2m_c+2m_b)^2}^{s_0} ds \int_{x_i}^{x_f} \\
&\times dx \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \frac{1}{x^3y^3z^3\mathcal{A}_{xyz}^3} \\
&\times \left\{ \mathcal{F}_{xyz} [\mathcal{A}_{xyz}^4 [96m_b^2x^4y^4z + 96m_c^2x^4yz^4 \right. \\
&- \mathcal{N}_k 48m_c^2x^3yz^4 \\
&+ 48m_c^2xy^3z^4(2y - \mathcal{N}_k)] - \mathcal{A}_{xyz}^3 (\mathcal{N}_k 48m_b^2x^4y^4 \\
&\times z - \mathcal{N}_k 6m_b^2x^3y^3z^3 \\
&- \mathcal{N}_k 6m_c^2x^3y^3z^3) + 48m_b^2x^4y^4z^3(2z - \mathcal{N}_k)\mathcal{A}_{xyz}] \\
&- [\mathcal{A}_{xyz}^4 [96m_b^4\hat{s}x^4y^4z \\
&+ 96m_b^2m_c^2\hat{s}x^4yz^4 - \mathcal{N}_k 24m_b^2m_c^2\hat{s}x^3yz^4 \\
&+ 24m_b^2m_c^2\hat{s}xy^3z^4(4y - \mathcal{N}_k) \\
&+ \mathcal{N}_k 16m_b^2m_c^2x^3y^3z + \mathcal{N}_k 16m_c^4x^3z^4 + \mathcal{N}_k 16m_c^4y^3z^4] \\
&- \mathcal{A}_{xyz}^3 (\mathcal{N}_k 24m_b^4\hat{s}x^4y^4z - \mathcal{N}_k 3m_b^4\hat{s}x^3y^3z^3) \\
&+ \mathcal{N}_k 16m_b^4y^4(\mathcal{A}_{xyz}^3 + z^3) \\
&- \mathcal{A}_{xyz}^3 [\mathcal{N}_k 3m_b^2m_c^2\hat{s}x^3y^3z^3 \\
&+ \mathcal{N}_k 16m_b^2m_c^2x^4yz^3 - 48m_b^2m_c^2x^3y^3z \\
&- 48m_b^2m_c^2x^3yz^3 + \mathcal{N}_k 16m_b^2m_c^2xy^3z^3(y - 3\mathcal{N}_k)]
\end{aligned}$$

$$\begin{aligned}
& +\mathcal{A}_{xyz}[24m_b^4\hat{s}x^4y^4z^3(4z - \mathcal{N}_k) \\
& +\mathcal{N}_k 16m_b^2m_c^2x^3y^3z^3(z - 3\mathcal{N}_k)] \Big\}, \quad (31)
\end{aligned}$$

$$\begin{aligned}
\Pi_{k,(GG)}^{0++}(M_B^2) &= -\frac{\langle g_s^2 GG \rangle}{2^9 \times 3^2 \times \pi^6} \int_0^1 \\
&\times dx \int_0^{1-x} dy \int_0^{1-x-y} dz e^{-\frac{f_{xyz}m_b^2}{M_B^2}} \\
&\times \frac{m_b^2}{x^3y^3z^3\mathcal{A}_{xyz}^3} [f_{xyz}m_b^2xy(2\mathcal{A}_{xyz}f_{xyz}z + \mathcal{N}_k) \\
&+ m_c^2(\mathcal{N}_k\mathcal{A}_{xyz}f_{xyz}z + 2)] \\
&\times [m_b^2x^3y^3(\mathcal{A}_{xyz}^3 + z^3) \\
&+ \mathcal{A}_{xyz}^3m_c^2z^3(x^3 + y^3)]. \quad (32)
\end{aligned}$$

For the $[c\bar{b}][b\bar{c}]$ octet–octet tetraquark states, the spectral densities $\rho(s)$ and $\Pi(M_B^2)$ can be written as

$$\begin{aligned}
\rho_{j,pert}^{0++}(s) &= \frac{1}{2^7 \times \pi^5} \int_{(2m_c+2m_b)^2}^{s_0} ds \int_{x_i}^{x_f} \\
&\times dx \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \left\{ \frac{\mathcal{F}_{xyz}^4 xy z \mathcal{A}_{xyz}}{2} \right. \\
&- \frac{\mathcal{F}_{xyz}^3 (6m_b^2\hat{s}xyz\mathcal{A}_{xyz} + \mathcal{N}_j 2m_b m_c x \mathcal{A}_{xyz} + \mathcal{N}_j 2m_b m_c y z)}{3} \\
&+ \mathcal{F}_{xyz}^2 [m_b^4\hat{s}^2xyz\mathcal{A}_{xyz} + \mathcal{N}_j m_b^3m_c\hat{s}x\mathcal{A}_{xyz} \\
&+ \mathcal{N}_j m_b^3m_c\hat{s}yz + m_b^2m_c^2] \Big\}, \quad (33)
\end{aligned}$$

$$\begin{aligned}
\rho_{j,(GG)}^{0++}(s) &= \frac{\langle g_s^2 GG \rangle}{2^{12} \times 3^2 \times \pi^5} \int_{(2m_c+2m_b)^2}^{s_0} ds \int_{x_i}^{x_f} \\
&\times dx \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \frac{1}{x^3y^3z^3\mathcal{A}_{xyz}^3} \\
&\times \left\{ -\frac{\mathcal{F}_{xyz}^2 [18x^4z^3y^3\mathcal{A}_{xyz}^4 + 18x^3y^4z^4\mathcal{A}_{xyz}^3]}{2} \right. \\
&+ \mathcal{F}_{xyz} [\mathcal{A}_{xyz}^4 [18m_b^2\hat{s}x^4y^3z^3 \\
&+ 48m_b^2x^4y^4z - \mathcal{N}_j 48m_b m_c x^4y^3z + \mathcal{N}_j 12m_b m_c x^4y^2z^2 \\
&+ 48m_c x^4yz^3(m_c z - \mathcal{N}_j m_b) + 48m_c^2xy^4z^4] \\
&+ \mathcal{A}_{xyz}^3 [18m_b^2\hat{s}x^3y^4z^4 \\
&+ \mathcal{N}_j 12m_b m_c x^3y^3z^3 - \mathcal{N}_j 48m_b m_c xy^4z^4] \\
&+ \mathcal{N}_j 12m_b m_c x^2y^4z^4\mathcal{A}_{xyz}^2 \\
&+ 48m_b^2x^4y^4z^4\mathcal{A}_{xyz} - \mathcal{N}_j 48m_b m_c x^3y^4z^4] \\
&- [\mathcal{A}_{xyz}^4 [3m_b^4\hat{s}^2x^4y^3z^3 \\
&+ 48m_b^4\hat{s}x^4y^4z - \mathcal{N}_j 24m_b^3m_c\hat{s}x^4y^3z \\
&+ \mathcal{N}_j 6m_b^3m_c\hat{s}x^4y^2z^2 \\
&+ \mathcal{N}_j 16m_b^3m_cx^4y^3 + 48m_b^2m_c^2\hat{s}xy^4z^4 - \mathcal{N}_j \\
&\times 24m_b^2m_c\hat{s}x^4yz^3(m_b - \mathcal{N}_j 2m_c z) \\
&+ \mathcal{N}_j 16m_b m_c^3x^4z^3 + \mathcal{N}_j 16m_b m_c^3xy^3z^3] \\
&+ \mathcal{A}_{xyz}^3 [3m_b^4\hat{s}^2x^3y^4z^4
\end{aligned}$$

$$\begin{aligned}
& +\mathcal{N}_j 6 m_b^3 m_c \hat{s} x^3 y^3 z^3 - \mathcal{N}_j 24 m_b^3 m_c \hat{s} x y^4 z^4 \\
& +\mathcal{N}_j 16 m_b^3 m_c \hat{s} x^3 y^4 z \\
& -24 m_b^2 m_c^2 x^3 y^3 z + 6 m_b^2 m_c^2 x^3 y^2 z^2 - 24 m_b^2 m_c^2 x y^3 z^3 \\
& +\mathcal{N}_j 16 m_b m_c^3 y^4 z^4 \\
& +\mathcal{N}_j 8 m_b m_c^2 x^3 y z^3 (2 m_c z - \mathcal{N}_j 3 m_b) \\
& +\mathcal{A}_{xyz}^2 [\mathcal{N}_j 6 m_b^3 m_c \hat{s} x^2 y^4 z^4 \\
& +6 m_b^2 m_c^2 x^2 y^3 z^3] + \mathcal{A}_{xyz} [48 m_b^4 \hat{s} x^4 y^4 z^4 \\
& -\mathcal{N}_j 24 m_b^3 m_c \hat{s} x^3 y^4 z^4 \\
& +\mathcal{N}_j 16 m_b^3 m_c x^4 y^3 z^3 - 24 m_b^2 m_c^2 x^3 y^3 z^3] \\
& +\mathcal{N}_j 16 m_b^3 m_c x^3 y^4 z^4 \Big\}, \quad (34)
\end{aligned}$$

$$\begin{aligned}
\Pi_{j,(GG)}^{0++}(M_B^2) &= -\frac{\langle g_s^2 GG \rangle}{2^9 \times 3^2 \times \pi^6} \int_0^1 \\
&\times dx \int_0^{1-x} dy \int_0^{1-x-y} dz e^{-\frac{f_{xyz} m_b^2}{M_B^2}} \\
&\times \frac{1}{x^3 y^3 z^3 \mathcal{A}_{xyz}^3} m_b^2 (\mathcal{N}_j f_{xyz} m_b x \mathcal{A}_{xyz} + m_c) \\
&\times (\mathcal{N}_j f_{xyz} m_b y z + m_c) \\
&\times [m_b^2 x^3 y^3 (\mathcal{A}_{xyz}^3 + z^3) + \mathcal{A}_{xyz}^3 m_c^2 z^3 (x^3 + y^3)], \quad (35)
\end{aligned}$$

$$\begin{aligned}
\rho_{k,pert}^{0++}(s) &= \frac{1}{2^6 \times \pi^5} \int_{(2m_c+2m_b)^2}^{s_0} ds \int_{x_i}^{x_f} \\
&\times dx \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \left\{ \mathcal{F}_{xyz}^4 x y z \mathcal{A}_{xyz} \right. \\
&- \frac{\mathcal{F}_{xyz}^3 [12 m_b^2 \hat{s} x y z \mathcal{A}_{xyz} + \mathcal{N}_k 2 m_b m_c x \mathcal{A}_{xyz} + \mathcal{N}_k 2 m_b m_c y z]}{3} \\
&+ \mathcal{F}_{xyz}^2 [2 m_b^4 \hat{s}^2 x y z \mathcal{A}_{xyz} + \mathcal{N}_k m_b^3 m_c \hat{s} x \mathcal{A}_{xyz} \\
&+ \mathcal{N}_k m_b^3 m_c \hat{s} y z + 2 m_b^2 m_c^2] \Big\}, \quad (36)
\end{aligned}$$

$$\begin{aligned}
\rho_{k,(GG)}^{0++}(s) &= \frac{\langle g_s^2 GG \rangle}{2^{10} \times 3^2 \times \pi^5} \int_{(2m_c+2m_b)^2}^{s_0} ds \int_{x_i}^{x_f} \\
&\times dx \int_{y_i}^{y_f} dy \int_{z_i}^{z_f} dz \frac{1}{x^3 y^3 z^3 \mathcal{A}_{xyz}^3} \\
&\times \left\{ \mathcal{F}_{xyz} [\mathcal{A}_{xyz}^4 [48 m_b^2 x^4 y^4 z - \mathcal{N}_k 24 m_b m_c x^4 y^3 \right. \\
&\times z - \mathcal{N}_k 24 m_b m_c x^4 y z^3 \\
&+ 48 m_c^2 x y z^4 (x^3 + y^3)] - \mathcal{A}_{xyz}^3 [\mathcal{N}_k 6 m_b m_c x^3 y^3 z^3 \\
&+ \mathcal{N}_k 24 m_b m_c x y^4 z^4] \\
&+ \mathcal{A}_{xyz} [48 m_b^2 x^4 y^4 z^4 - \mathcal{N}_k 24 m_b m_c x^3 y^4 z^4] \\
&- [\mathcal{A}_{xyz}^4 [-\mathcal{N}_k 12 m_b^3 m_c \hat{s} x^4 y^3 z \\
&- \mathcal{N}_k 12 m_b^3 m_c \hat{s} x^4 y z^3 + 48 m_b^2 m_c^2 \hat{s} x y z^4 (x^3 + y^3) \\
&+ \mathcal{N}_k 8 m_b m_c^3 x^4 z^3 \\
&+ \mathcal{N}_k 8 m_b m_c^3 x y^3 z^3] + 48 m_b^4 \hat{s} x^4 y^4 z (\mathcal{A}_{xyz}^3 + z^3) \mathcal{A}_{xyz} \\
&- \mathcal{A}_{xyz}^3 [\mathcal{N}_k 3 m_b^3 m_c \hat{s} \\
&\times x^3 y^3 z^3 + \mathcal{N}_k 12 m_b^3 m_c \hat{s} x y^4 z^4 + 24 m_b^2 m_c^2 x^3 y^3 z \\
&+ 24 m_b^2 m_c^2 x^3 y z^3]
\end{aligned}$$

$$\begin{aligned}
& +24 m_b^2 m_c^2 x y^3 z^3 - \mathcal{N}_k 8 m_b m_c^3 x^3 y z^4 \\
& -\mathcal{N}_k 8 m_b m_c^3 y^4 z^4] \\
& -\mathcal{A}_{xyz} [\mathcal{N}_k 12 m_b^3 m_c \hat{s} x^3 y^4 z^4 + 24 m_b^2 m_c^2 x^3 y^3 z^3] \\
& +\mathcal{N}_k 8 m_b^3 m_c x^3 y^3 (\mathcal{A}_{xyz}^3 + z^3) (\mathcal{A}_{xyz} x + y z) \Big\}, \quad (37)
\end{aligned}$$

$$\begin{aligned}
\Pi_{k,(GG)}^{0++}(M_B^2) &= -\frac{\langle g_s^2 GG \rangle}{2^8 \times 3^2 \times \pi^6} \int_0^1 \\
&\times dx \int_0^{1-x} dy \int_0^{1-x-y} dz e^{-\frac{f_{xyz} m_b^2}{M_B^2}} \\
&\times \frac{m_b^2}{x^3 y^3 z^3 \mathcal{A}_{xyz}^3} [2 f_{xyz}^2 m_b^2 x y z \mathcal{A}_{xyz} + 2 m_c^2 \\
&+ \mathcal{N}_k f_{xyz} m_b m_c (\mathcal{A}_{xyz} x + y z) \\
&\times [m_b^2 x^3 y^3 (\mathcal{A}_{xyz}^3 + z^3) \\
&+ \mathcal{A}_{xyz}^3 m_c^2 z^3 (x^3 + y^3)]. \quad (38)
\end{aligned}$$

Here, we also have the following definitions:

$$\begin{aligned}
\mathcal{A}_x &= (1-x), \quad \mathcal{A}_{xy} = (1-x-y), \\
\mathcal{A}_{xyz} &= (1-x-y-z), \quad (39)
\end{aligned}$$

$$f_{xyz} = \left[\frac{r_m^2}{x} + \frac{r_m^2}{y} + \frac{1}{z} + \frac{1}{(1-x-y-z)} \right], \quad (40)$$

$$x_{f/i} = \frac{\hat{s} - (4r_m + 1) \pm \sqrt{(\hat{s} - 4)(\hat{s} - 4(r_m + 1)^2)}}{2\hat{s}}, \quad (41)$$

$$\begin{aligned}
y_{f/i} &= \frac{1}{2(\hat{s}x - r_m^2)} \left\{ x(\hat{s}\mathcal{A}_x + r_m^2 - 4) - r_m^2 \mathcal{A}_x \right. \\
&\pm \sqrt{4r_m^2 x \mathcal{A}_x (r_m^2 - \hat{s}x) + [r_m^2 \mathcal{A}_x - x(\hat{s}\mathcal{A}_x + r_m^2 - 4)]} \Big\}, \quad (42)
\end{aligned}$$

$$\begin{aligned}
z_{f/i} &= \frac{1}{2[\hat{s}xy - r_m^2(x+y)]} \left\{ \mathcal{A}_{xy} [\hat{s}xy - r_m^2(x+y)] \right. \\
&\pm \sqrt{\mathcal{A}_{xy} [\mathcal{A}_{xy} (\hat{s}xy - r_m^2(x+y))^2 - 4xy(\hat{s}xy - r_m^2(x+y))]} \Big\}. \quad (43)
\end{aligned}$$

It should be noted that for $[Q\bar{Q}][Q\bar{Q}]$ case with $Q = c$ or b , we have the following definitions: $\hat{s} = \frac{s}{m_Q^2}$, $\mathcal{F}_{xyz} = m_Q^2 f_{xyz} - s$, and $r_m = 1$; for $[c\bar{c}][b\bar{b}]$ and $[c\bar{b}][b\bar{c}]$ cases, $\hat{s} = \frac{s}{m_b^2}$, $\mathcal{F}_{xyz} = m_b^2 f_{xyz} - s$, and $r_m = m_c/m_b$.

References

1. M. Gell-Mann, Phys. Lett. **8**, 214 (1964)
2. G. Zweig, Report No. CERN-TH-401
3. S.K. Choi et al., [Belle], Phys. Rev. Lett. **91**, 262001 (2003). [arXiv:hep-ex/0309032](https://arxiv.org/abs/hep-ex/0309032)
4. R. Aaij et al., [LHCb], Sci. Bull. **65** (2020). [arXiv:2006.16957](https://arxiv.org/abs/2006.16957) [hep-ex]
5. V. Khachatryan et al., [CMS], JHEP **05**, 013 (2017). [arXiv:1610.07095](https://arxiv.org/abs/1610.07095) [hep-ex]
6. R. Aaij et al., [LHCb], JHEP **10**, 086 (2018). [arXiv:1806.09707](https://arxiv.org/abs/1806.09707) [hep-ex]

7. A.M. Sirunyan et al., [CMS], Phys. Lett. B **808**, 135578 (2020). [arXiv:2002.06393](#) [hep-ex]
8. Y. Iwasaki, Phys. Rev. Lett. **36**, 1266 (1976)
9. K.T. Chao, Z. Phys. C **7**, 317 (1981)
10. J.P. Ader, J.M. Richard, P. Taxil, Phys. Rev. D **25**, 2370 (1982)
11. A.M. Badalian, B.L. Ioffe, A.V. Smilga, Nucl. Phys. B **281**, 85 (1987)
12. L. Heller, J.A. Tjon, Phys. Rev. D **32**, 755 (1985)
13. M.N. Anwar, J. Ferretti, F.K. Guo, E. Santopinto, B.S. Zou, Eur. Phys. J. C **78**(8), 647 (2018). [arXiv:1710.02540](#) [hep-ph]
14. M.A. Bedolla, J. Ferretti, C.D. Roberts, E. Santopinto, Eur. Phys. J. C **80**(11), 1004 (2020)
15. R.J. Lloyd, J.P. Vary, Phys. Rev. D **70**, 014009 (2004). [arXiv:hep-ph/0311179](#)
16. N. Barnea, J. Vijande, A. Valcarce, Phys. Rev. D **73**, 054004 (2006). [arXiv:hep-ph/0604010](#)
17. V.R. Debastiani, F.S. Navarra, Chin. Phys. C **43**(1), 013105 (2019). [arXiv:1706.07553](#) [hep-ph]
18. J. Wu, Y.R. Liu, K. Chen, X. Liu, S.L. Zhu, Phys. Rev. D **97**(9), 094015 (2018). [arXiv:1605.01134](#) [hep-ph]
19. G.J. Wang, L. Meng, S.L. Zhu, Phys. Rev. D **100**(9), 096013 (2019). [arXiv:1907.05177](#) [hep-ph]
20. M.S. Liu, Q.F. Lü, X.H. Zhong, Q. Zhao, Phys. Rev. D **100**(1), 016006 (2019). [arXiv:1901.02564](#) [hep-ph]
21. R.N. Faustov, V.O. Galkin, E.M. Savchenko, [arXiv:2009.13237](#) [hep-ph]
22. Q.F. Lü, D.Y. Chen, Y.B. Dong, Eur. Phys. J. C **80**(9), 871 (2020). [arXiv:2006.14445](#) [hep-ph]
23. W. Heupel, G. Eichmann, C.S. Fischer, Phys. Lett. B **718**, 545–549 (2012). [arXiv:1206.5129](#) [hep-ph]
24. X.Z. Weng, X.L. Chen, W.Z. Deng, S.L. Zhu, [arXiv:2010.05163](#) [hep-ph]
25. A.V. Berezhnoy, A.V. Luchinsky, A.A. Novoselov, Phys. Rev. D **86**, 034004 (2012). [arXiv:1111.1867](#) [hep-ph]
26. M. Karliner, S. Nussinov, J.L. Rosner, Phys. Rev. D **95**(3), 034011 (2017). [arXiv:1611.00348](#) [hep-ph]
27. A.V. Berezhnoy, A.K. Likhoded, A.V. Luchinsky, A.A. Novoselov, Phys. Rev. D **84**, 094023 (2011). [arXiv:1101.5881](#) [hep-ph]
28. F. Feng, Y. Huang, Y. Jia, W.L. Sang, X. Xiong, J.Y. Zhang, [arXiv:2009.08450](#) [hep-ph]
29. Y.Q. Ma, H.F. Zhang, [arXiv:2009.08376](#) [hep-ph]
30. M. Karliner, J.L. Rosner, [arXiv:2009.04429](#) [hep-ph]
31. J.Z. Wang, D.Y. Chen, X. Liu, T. Matsuki, [arXiv:2008.07430](#) [hep-ph]
32. J.F. Giron, R.F. Lebed, Phys. Rev. D **102**(7), 074003 (2020). [arXiv:2008.01631](#) [hep-ph]
33. K.T. Chao, S.L. Zhu, Sci. Bull. **65**, 1952–1953 (2020)
34. L. Maiani, [arXiv:2008.01637](#) [hep-ph]
35. J.M. Richard, [arXiv:2008.01962](#) [hep-ph]
36. R. Zhu, [arXiv:2010.09082](#) [hep-ph]
37. Z.H. Guo, J.A. Oller, [arXiv:2011.00978](#) [hep-ph]
38. R. Maciula, W. Schäfer, A. Szczurek, [arXiv:2009.02100](#) [hep-ph]
39. J.W. Zhu, X.D. Guo, R.Y. Zhang, W.G. Ma, X.Q. Li, [arXiv:2011.07799](#) [hep-ph]
40. G. Eichmann, C.S. Fischer, W. Heupel, N. Santowsky, P.C. Wallbott, Few Body Syst. **61**(4), 38 (2020)
41. C. Gong, M.C. Du, B. Zhou, Q. Zhao, X.H. Zhong, [arXiv:2011.11374](#) [hep-ph]
42. C. Becchi, J. Ferretti, A. Giachino, L. Maiani, E. Santopinto, Phys. Lett. B **811**, 135952 (2020)
43. X.K. Dong, V. Baru, F.K. Guo, C. Hanhart, A. Nefediev, [arXiv:2009.07795](#) [hep-ph]
44. W. Chen, H.X. Chen, X. Liu, T.G. Steele, S.L. Zhu, Phys. Lett. B **773**, 247–251 (2017). [arXiv:1605.01647](#) [hep-ph]
45. Z.G. Wang, Eur. Phys. J. C **77**(7), 432 (2017). [arXiv:1701.04285](#) [hep-ph]
46. W. Chen, H.X. Chen, X. Liu, T.G. Steele, S.L. Zhu, EPJ Web Conf. **182**, 02028 (2018). [arXiv:1803.02522](#) [hep-ph]
47. Z.G. Wang, Z.Y. Di, Acta Phys. Pol. B **50**, 1335 (2019). [arXiv:1807.08520](#) [hep-ph]
48. J.R. Zhang, Phys. Rev. D **103**(1), 014018 (2021). [arXiv:2010.07719](#) [hep-ph]
49. Z.G. Wang, Int. J. Mod. Phys. A **36**, 2150014 (2021). [arXiv:2009.05371](#) [hep-ph]
50. R.M. Albuquerque, S. Narison, A. Rabemananjara, D. Rabetiarivony, G. Randriamanatrika, Phys. Rev. D **102**(9), 094001 (2020). [arXiv:2008.01569](#) [hep-ph]
51. H.X. Chen, W. Chen, X. Liu, S.L. Zhu, Sci. Bull. **65**, 1994–2000 (2020). [arXiv:2006.16027](#) [hep-ph]
52. B.D. Wan, C.F. Qiao, [arXiv:2012.00454](#) [hep-ph]
53. J.I. Latorre, P. Pascual, J. Phys. G **11**, L231 (1985)
54. S. Narison, Phys. Lett. B **175**, 88 (1986)
55. Z.G. Wang, Nucl. Phys. A **791**, 106–116 (2007). [arXiv:hep-ph/0610171](#)
56. Z.G. Wang, Int. J. Mod. Phys. A **30**(30), 1550168 (2015). [arXiv:1502.01459](#) [hep-ph]
57. L. Tang, C.F. Qiao, Eur. Phys. J. C **76**(10), 558 (2016). [arXiv:1603.04761](#) [hep-ph]
58. L. Tang, B.D. Wan, K. Maltman, C.F. Qiao, Phys. Rev. D **101**(9), 094032 (2020). [arXiv:1911.10951](#) [hep-ph]
59. M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, Nucl. Phys. B **147**, 385 (1979)
60. M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, Nucl. Phys. B **147**, 448 (1979)
61. L.J. Reinders, H. Rubinstein, S. Yazaki, Phys. Rep. **127**, 1 (1985)
62. S. Narison, World Sci. Lect. Notes Phys. **26**, 1 (1989)
63. P. Colangelo, A. Khodjamirian, in *At the frontier of particle physics / Handbook of QCD*, edited by M. (Singapore, Shifman (World Scientific, 2001). [arXiv:hep-ph/0010175](#)
64. Z.G. Wang, T. Huang, Phys. Rev. D **89**, 054019 (2014)
65. R.M. Albuquerque, (n.d.) [arXiv:1306.4671](#) [hep-ph]
66. R.D.E. Matheus, S. Narison, M. Nielsen, J.M. Richard, Phys. Rev. D **75**, 014005 (2007)
67. S.I. Finazzo, M. Nielsen, X. Liu, Phys. Lett. B **701**, 101 (2011). [arXiv:1102.2347](#) [hep-ph]
68. C.F. Qiao, L. Tang, Eur. Phys. J. C **74**(10), 3122 (2014). [arXiv:1307.6654](#) [hep-ph]
69. C.F. Qiao, L. Tang, Eur. Phys. J. C **74**, 2810 (2014)