



Hawking evaporation of Einstein–Gauss–Bonnet AdS black holes in $D \geq 4$ dimensions

Chen-Hao Wu^{1,2,4,a}, Ya-Peng Hu^{1,4,b}, Hao Xu^{2,3,c} 

¹ College of Science, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, China

² Center for Gravitation and Cosmology, College of Physical Science and Technology, Yangzhou University, Yangzhou 225009, China

³ School of Aeronautics and Astronautics, Shanghai Jiao Tong University, Shanghai 200240, China

⁴ MIT Key Laboratory of Aerospace Information Materials and Physics, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, China

Received: 6 March 2021 / Accepted: 13 April 2021 / Published online: 23 April 2021
© The Author(s) 2021

Abstract Einstein–Gauss–Bonnet theory is a string-generated gravity theory when approaching the low energy limit. By introducing the higher order curvature terms, this theory is supposed to help to solve the black hole singularity problem. In this work, we investigate the evaporation of the static spherically symmetric neutral AdS black holes in Einstein–Gauss–Bonnet gravity in various spacetime dimensions with both positive and negative coupling constant α . By summarizing the asymptotic behavior of the evaporation process, we find the lifetime of the black holes is dimensional dependent. For $\alpha > 0$, in $D \geq 6$ cases, the black holes will be completely evaporated in a finite time, which resembles the Schwarzschild-AdS case in Einstein gravity. While in $D = 4, 5$ cases, the black hole lifetime is always infinite, which means the black hole becomes a remnant in the late time. Remarkably, the cases of $\alpha > 0$, $D = 4, 5$ will solve the terminal temperature divergent problem of the Schwarzschild-AdS case. For $\alpha < 0$, in all dimensions, the black hole will always spend a finite time to a minimal mass corresponding to the smallest horizon radius $r_{min} = \sqrt{2|\alpha|}$ which coincide with an additional singularity. This implies that there may exist constraint conditions to the choice of coupling constant.

1 Introduction

In 1974, by applying quantum field theory in curved spacetime, Hawking showed that quantum mechanical effects allow black holes to emit particles, now known as Hawking radiation [1, 2]. Hawking radiation connects quantum physics

and gravity, and it is viewed as a window to help us to understand the quantum gravity. Due to the radiation, the black hole will keep losing mass and thermal entropy, which leaves us the controversial information loss paradox [3, 4].

The spectrum of the Hawking radiation allows us to calculate the particles emission power and to estimate the lifetime of the black hole. For a static spherically symmetric black hole in four-dimensional asymptotically flat spacetime in Einstein gravity, its lifetime obeys the relationship $t \sim M_0^3$ associated with the initial black hole mass M_0 [5], which means the black hole lifetime will be divergent if we take an infinite initial mass. A similar analysis can also be carried out in rotating black holes and charged black holes. A rotating black hole will lose its angular momentum much faster than its mass and eventually it becomes a Schwarzschild black hole and keeps going on its evaporation [6] (unless the case with a lot of scalar particle species [7, 8]). A charged black hole will keep losing its charge and mass, however, the charge-over-mass ratio may increase and gets close to the extreme black hole, so the lifetime of these black holes may be extended by a huge factor, but finally, it also evolves to a Schwarzschild state [9, 10].

The spectrum of the Hawking radiation allows us to calculate the particles emission power and to estimate the lifetime of the black hole. For a static spherically symmetric black hole in four-dimensional asymptotically flat Based on Hawking radiation, the study on the evaporation of the black hole in asymptotically flat spacetime has been a fruitful area. The evaporation in anti-de Sitter (AdS) spacetimes was, however, largely overlooked. This is due to the specific asymptotical behavior of the AdS spacetime. Although AdS has an infinite volume, the gravitational potential acts effectively like a finite confining box for the black hole, and the emitted massive particles reflect inward in a finite time thus preventing

^a e-mail: chenhao_wu@nuaa.edu.cn

^b e-mail: huyup@nuaa.edu.cn

^c e-mail: haoxu_phys@163.com (corresponding author)

the energy loss. Thanks to the AdS/CFT correspondence [11–14], a great mass of attention has been paid on black holes in AdS spacetimes, especially AdS black hole thermodynamics [15], and the evaporation problem of AdS black hole also comes back to our view in recent years. By imposing an absorbing AdS boundary condition [16–19] and considering only massless emitted particle, Page calculated the lifetime of static spherically symmetric AdS black hole in Einstein gravity [20], and surprisingly, the lifetime did not become divergent with any arbitrarily large initial mass and the total time to evaporate from infinite mass to zero was finite, of the order ℓ^3 , where ℓ is the AdS radius. This is because although the black hole may have infinite initial mass, the temperature, and also the particles' emission power, are also divergent, thus the infinite amount of mass can be evaporated away in a finite time. The study on black holes evaporation in asymptotically AdS spacetimes have been extended to many different backgrounds, such as Lovelock gravity [21], Conformal (Weyl) Gravity [22], Hořava–Lifshitz gravity [23], and dRGT massive gravity [24], etc.

From a modern perspective, Einstein gravity could just be an effective field theory in the low energy limit of unknown more fundamental theory. Although we have not known its correct and precise form, there are some attempts by physicists. Inspired by the quadratic term in heterotic string theory and six-dimensional Calabi–Yau compactifications of M-theory, the Gauss–Bonnet quadratic term was added into the Einstein–Hilbert action to analyze the properties of the effective gravity theory [25–29]. This gravity can be derived by string theories when approaching the low energy limit. Studies on the Gauss–Bonnet gravity and its generalizations have attracted lots of attention [30–40], however, this theory just exists in higher spacetime dimensions because the Gauss–Bonnet term is a topological invariant in four dimensions thus it cannot give any non-trivial gravitational dynamics in $D \leq 4$, where D is the dimensions of the spacetime. In [41], Glavan and Lin try to extend this theory to four dimensions by rescaling the coupling constant of the Gauss–Bonnet term by a factor of $1/(D-4)$ and taking the limit $D \rightarrow 4$. Remarkably, from their point of view, this theory can bypass the Lovelock's theorem and be free from Ostrogradsky instability. Interestingly, its black hole solution was also found in other contexts, such as gravity with conformal anomaly [42, 43] and quantum corrections [44]. This processing method in [41] has led to a great deal of discussion, see e.g. [45–60].

In the present work, we investigate the Hawking evaporation process of the static spherically symmetric neutral AdS black holes in Einstein–Gauss–Bonnet gravity. The thermodynamical properties of these black holes in $D \geq 5$ and $D = 4$ are provided in [30, 46]. Unlike the Schwarzschild–AdS case, which always has a divergent temperature in the late period of the evaporation process [61], the Einstein–Gauss–Bonnet

theory reveals richer phenomena by choosing different coupling constant. By testing the Hawking evaporation processes of different Einstein–Gauss–Bonnet AdS black holes, we give some clues to constrain the choice of coupling constant. Especially, the extreme behavior of the cases with negative coupling constant also resembles the Garfinkle–Horowitz–Strominger dilation black hole [62].

This paper is organized as follows: in Sect. 2, we briefly review the thermodynamics of $D \geq 5$ -dimensional neutral Einstein–Gauss–Bonnet AdS black holes. By investigating the Hawking evaporation process of the $D = 5$ and $D \geq 6$ cases, we show that the positive coupling constant may help to solve divergent temperature, while the negative one may make this problem worse. In Sect. 3, our investigation is extended to the $D = 4$ cases. Summary remarks are given in Sect. 4. We adopt the natural unit system, setting the speed of light in vacuum c , the gravitational constant G_N , the Planck constant h , and the Boltzmann constant k_B equal to one.

2 Evaporation of Einstein–Gauss–Bonnet AdS Black Holes in $D \geq 5$

In this section, first, we will give a brief review of Einstein–Gauss–Bonnet AdS black holes for $D \geq 5$ dimensions, then we investigate the black hole evaporation process.

2.1 Black hole thermodynamics in $D \geq 5$ Einstein–Gauss–Bonnet AdS spacetimes

For the $D \geq 5$ dimensional AdS spacetime, the Einstein–Hilbert action with a Gauss–Bonnet term can be written as

$$S = \frac{1}{16\pi} \int d^D x \sqrt{-g} \left(\mathcal{R} + \frac{(D-1)(D-2)}{\ell^2} + \alpha_{\text{GB}} \mathcal{L}_{\text{GB}} \right), \quad (1)$$

where α_{GB} is the Gauss–Bonnet coupling coefficient with dimension $(length)^2$ and the Gauss–Bonnet term $\mathcal{L}_{\text{GB}} = \mathcal{R}_{\mu\nu\gamma\delta} \mathcal{R}^{\mu\nu\gamma\delta} - 4\mathcal{R}_{\mu\nu} \mathcal{R}^{\mu\nu} + \mathcal{R}^2$. The Gauss–Bonnet coefficient α_{GB} is supposed to be a positive value in string theory [25], but for generality we consider both cases of $\alpha_{\text{GB}} > 0$ and $\alpha_{\text{GB}} < 0$ in the present work. The static spherically symmetric black hole metric in D dimensions can be written in the form of

$$ds^2 = -f(r)dt^2 + f^{-1}(r)dr^2 + r^2 d\Omega_{D-2}^2, \quad (2)$$

with the the metric function [30]

$$f(r) = 1 + \frac{r^2}{2\alpha} \left(1 - \sqrt{1 + \frac{64\pi M\alpha}{(D-2)\Omega_{D-2} r^{D-1}} - \frac{4\alpha}{\ell^2}} \right), \quad (3)$$

where $\alpha = (D - 3)(D - 4)\alpha_{\text{GB}}$ and the parameter M is the gravitational mass of the solution. The $\Omega_{D-2} = \frac{(D-1)\pi^{D/2}}{\Gamma(\frac{D+1}{2})}$ is the area of an $(D - 1)$ -dimensional sphere with unit radius. When $\alpha \rightarrow 0$, black hole solution reduces to the Schwarzschild AdS case in [63]. In the limit $M \rightarrow 0$, one can obtain the vacuum solution $f(r) = 1 + r^2/\ell_{\text{eff}}^2$ with the effective AdS radius $\ell_{\text{eff}}^2 = \frac{\ell^2}{2} \left(1 + \sqrt{1 - \frac{4\alpha}{\ell^2}}\right)$, which implies the coupling constant α must obey $\alpha \leq \ell^2/4$, beyond that this theory is undefined. The event horizon of the black hole is determined by $f(r_+) = 0$, and the mass of black hole M can be obtained as [30]

$$M = \frac{(D - 2)\Omega_{D-2}r_+^{D-3}}{16\pi} \left(1 + \frac{\alpha}{r_+^2} + \frac{r_+^2}{l^2}\right), \tag{4}$$

and the Hawking temperature of the black hole can be written as

$$T = \frac{f'(r)}{4\pi} \Big|_{r=r_+} = \frac{(D - 1)r_+^4 + (D - 3)\ell^2r_+^2 + (D - 5)\alpha\ell^2}{4\pi\ell^2r_+(r_+^2 + 2\alpha)}. \tag{5}$$

In $\alpha > 0$, it is worth noting that there is a mass gap $M_c \equiv M(r_+ \rightarrow 0) = 3\pi\alpha/8$ in $D = 5$ that all black holes must satisfy $M \geq M_c$, while in $D \geq 6$, there exists no mass gap. We can also find the last term in (5) vanishes when $D = 5$. Consequently, the temperature T vanishes as $r_+ \rightarrow 0$ when $D = 5$, while it becomes divergent as $r_+ \rightarrow 0$ in higher dimensions. Among various choices for the spacetime dimension D , the particular case $D = 5$ is qualitatively different from other choices. This is perhaps a consequence of $D = 5$ is the lowest dimension in which the Gauss–Bonnet term can affect the local geometry. In $\alpha < 0$, there is a extreme horizon radius $r_{\text{min}} = \sqrt{2|\alpha|}$ corresponding to a divergent temperature, below that the theory is undefined. The black hole entropy can also be calculated by using the first law of black hole thermodynamics, as well as other thermodynamical quantities. We omit them here.

2.2 Black hole evaporation in $D \geq 5$ Einstein–Gauss–Bonnet AdS spacetimes

Since we have obtained all the necessary thermodynamical quantities, we are ready to investigate the evaporation of Einstein–Gauss–Bonnet AdS black holes. By imposing an absorbing boundary condition, the black hole mass M should be monotonically-decreasing functions of the time t because of the Hawking radiation. Applying geometric optics approximation, we assume all the emitted massless particles move along null geodesics. If we orient the extra $(D - 3)$ angular coordinates in $d\Omega_{D-2}^2$ and normalize the affine parameter λ , the geodesic equation of the massless particles reads

$$\left(\frac{dr}{d\lambda}\right)^2 = E^2 - J^2\frac{f(r)}{r^2}, \tag{6}$$

where $E = f(r)\frac{dt}{d\lambda}$ is the energy and $J = r^2\frac{d\theta}{d\lambda}$ is angular momentum of the emitted particle. Consider an emitted particle coming from just outside the black hole horizon, this particle cannot be detected by the observer on the AdS boundary if there is a turning point satisfying $(\frac{dr}{d\lambda})^2 = 0$. Defining an impact parameter $b \equiv J/E$, the massless particle can reach infinity only if

$$\frac{1}{b^2} > \frac{f(r)}{r^2}, \tag{7}$$

for all $r > r_+$. The impact factor b_c is defined by the maximal value of $f(r)/r^2$, which depends on the exact form of metric function $f(r)$.

If we obtain the impact factor b_c , we can get the Hawking emission rate in D -dimensional spacetime according to Stefan–Boltzmann law, yields [10, 21, 23, 64, 65]

$$\frac{dM}{dt} = -Cb_c^{D-2}T^D, \tag{8}$$

with the constant $C = (D - 2)\pi^{\frac{D}{2}-1}\Omega_{D-2}\frac{k^D}{h^{D-1}c^{D-1}}\frac{\Gamma(D)}{\Gamma(\frac{D}{2})}\zeta(D)$. Without loss of generality, we set the constant $C = 1$. The Stefan–Boltzmann law implies that in D -dimensional spacetime the emission power is proportional to the $(D - 2)$ -dimensional cross section b_c^{D-2} and the $(D - 1)$ -dimensional space (spatial dimension only) photon energy T^D . Notice that the photon energy term T^D has the higher order, so the behavior of temperature T , especially the asymptotical behavior, will play a leading role in black hole evaporation process. Next, we are ready to study the black hole evaporation for various features of T and b_c .

Using scaling analysis we know $M \sim l^{D-3}$, $T \sim l^{-1}$, $b_c \sim l$, $\ell \sim l$ and $\alpha \sim l^2$, where l has the dimension (length). In order to get dimensionless parts of M , T and b_c , by defining dimensionless variables $x \equiv r_+/l$ and $y \equiv \alpha/\ell^2$, we can express M and T as

$$M = \frac{(D - 2)\Omega_{D-2}}{16\pi} (1 + y + x^2)x^{D-3}\ell^{D-3}, \tag{9}$$

$$T = \frac{(D - 1)x^4 + (D - 3)x^2 + (D - 5)y}{4\pi x(x^2 + 2y)}\ell^{-1}. \tag{10}$$

The b_c depends on the exact form of metric function $f(r)$, so we cannot get the universal solution for various dimensions.

Firstly we start from the case of $D = 5$, which is qualitatively different from the cases of $D \geq 6$. When $D = 5$, the temperature (5) can be written as

$$T = \frac{2r_+^3 + \ell^2r_+}{2\pi\ell^2(r_+^2 + 2\alpha)}. \tag{11}$$

In Fig. 1 we present the temperature of black holes in 5 dimensions with $\alpha > 0$ and $\alpha < 0$ as the function of r_+ respectively. For $\alpha > 0$, the temperature starts from zero at $r_+ = 0$ and




Fig. 1 The temperature of black holes in 5 dimensions with $\alpha > 0$ and $\alpha < 0$ respectively. The left figure we set $\alpha = 0.15, \ell = 1$, while in the right figure $\alpha = -0.15, \ell = 1$




Fig. 2 The numerical results of the black hole mass M as function of the lifetime t with setting $y > 0$ and $y < 0$ (corresponding to $\alpha > 0$ and $\alpha < 0$, respectively). In the left figure, we set $y = 0.1$ with $\ell = 1, \ell = 1.5$ and $\ell = 2$ from left to right respectively. Note that there is a mass gap $M_c = \frac{3\pi\alpha}{8}$ for $y > 0$ ($\alpha > 0$). In the right figure, we set $y = -0.1$ with $\ell = 1, \ell = 1.5$ and $\ell = 2$ from left to right, corresponding to minimum mass $M_{min1} = 0.164934, M_{min2} = 0.371101$ and $M_{min3} = 0.659734$, respectively

goes to infinity when $r_+ \rightarrow \infty$. The behavior of temperature in $\alpha < 0$ is quite different from the case of $\alpha > 0$. There is an extreme horizon radius $r_{min} = \sqrt{2|\alpha|}$, which coincides with an additional singularity. As shown in the figure, the temperature is divergent at r_{min} when $\alpha < 0$, and as a result, the thermodynamical properties of case $\alpha < 0$ resembles the case of $\alpha \rightarrow 0$ that without Gauss–Bonnet term.

In order to get the impact factor b_c , solving the equation $\frac{\partial}{\partial r} \frac{f(r)}{r^2} = 0$, we can find the root

$$r_p = \frac{2}{\sqrt{3\pi}} \left(\frac{6\pi \ell^2 M \alpha - 16\ell^2 M^2}{4\alpha - \ell^2} \right)^{1/4}. \tag{12}$$

This radius corresponds to the unstable photon orbit, and the impact factor is given by $b_c = r_p / \sqrt{f(r_p)}$. Similarly, taking $x \equiv r_+ / \ell$ and $y \equiv \alpha / \ell^2$, one can obtain the b_c as

$$b_c = \left(\frac{\sqrt{4y - 1}}{2\sqrt{-(x^2 + x^4)(x^2 + x^4 + y)}} - \frac{-1 + \sqrt{\frac{(x^2 + x^4 + y)(4y - 1)}{x^2 + x^4}}}{2y} \right)^{-\frac{1}{2}} \ell \tag{13}$$

For $D = 5$, inserting the black hole mass M (9), temperature T (10) and impact factor b_c (13) into Stefan–Boltzmann law, we can obtain

$$dt = \ell^4 F_1(x, y) dx, \tag{14}$$

where $F_1(x, y)$ is a complicated function which is not worth explicitly written. Setting y to be a constant and integrating the equation from ∞ to x_{min} , where $x_{min} = 0$ for $\alpha > 0$ and $x_{min} = r_{min} / \ell = \sqrt{2|y|}$ for $\alpha < 0$, one can obtain the black hole lifetime is divergent in $\alpha > 0$ and of the order ℓ^4 in $\alpha < 0$.

In Fig. 2, we plot some numerical results of the black hole mass M as function of the lifetime t by setting $y > 0$ and $y < 0$ (corresponding to $\alpha > 0$ and $\alpha < 0$ respectively). In the case of $\alpha > 0$, we set $y = 0.1$, and the initial mass is taken to infinity. The black holes lose their mass in a “short” time, but the evaporation gets harder when the black hole mass and temperature become smaller near the $T = 0$ state, thus the black hole will have an infinite lifetime, satisfying the third law of black hole thermodynamics. This means that the black hole becomes a remnant, which may help us to ameliorate the information paradox [66]. In the case $\alpha < 0$, there is an extreme horizon radius at $r_{min} = \sqrt{2|\alpha|}$, where the temperature of black hole is divergent coinciding with an additional singularity [30]. This r_{min} corresponds to a mass $M_{min} = M(r_{min})$, and the black hole will take a finite time to decrease the mass to M_{min} . Interestingly, this phenomenon resembles the extreme case of Garfinkle-Horowitz-Strominger dilation black hole [62].

Now we extend our study to $D \geq 6$. The temperature behavior in $D \geq 6$ is quite different from the case $D = 5$ when $\alpha > 0$. For $D \geq 6$, the temperature is divergent at $r_+ = 0$ and goes to infinity again when $r_+ \rightarrow \infty$ [30], which resembles the case of Schwarzschild AdS. On the other hand, the temperature behavior of case $\alpha < 0$ in $D \geq 6$ spacetimes is similarly with the case $\alpha < 0$ in 5 dimensions, where there is also the smallest horizon radius at $r_{min} = \sqrt{2|\alpha|}$. Although $\frac{\partial}{\partial r} \frac{f(r)}{r^2} = 0$ is a equation of higher order in $D \geq 6$ dimensions and we cannot get the explicit form of b_c , we can still conclude $b_c \sim \ell$ by scaling analysis. As we have emphasized earlier in the present work, the photon energy term T^D plays the leading role in the black hole evaporation process. Similarly, inserting the black hole mass $M \sim \ell^{D-3}$, temperature $T \sim \ell^{-1}$ and impact factor $b_c \sim \ell$ into Stefan-Boltzmann law, we can also obtain the D -dimensional Stefan-Boltzmann law as

$$dt = \ell^{D-1} F_2(x, y) dx, \tag{15}$$

where $F_2(x, y)$ is still a complicated function. We can conclude this evaporation process qualitatively by analyzing the temperature asymptotic behavior of the black hole. When $\alpha > 0$, the temperature T is divergent in both $r_+ \rightarrow \infty$ and $r_+ \rightarrow 0$, which means that $\int_{\infty}^0 F_2(x, y) dx$ is convergent and just depends on the value of y . Hence, the lifetime of the black hole will be a finite value of the order ℓ^{D-1} with any arbitrarily large initial mass, which resembles the Schwarzschild AdS black hole in Einstein gravity [20]. For $\alpha < 0$, the temperature of black hole is still divergent at $r_{min} = \sqrt{2|\alpha|}$, thus the qualitative features of the evaporation process in $D \geq 6$ dimensions are consistent with the case of $\alpha < 0$ in 5 dimensions.

3 Evaporation of novel 4D Einstein-Gauss-Bonnet AdS black holes

Since Gauss-Bonnet term is a topological invariant in four dimensions, previously the investigation of Gauss-Bonnet gravity is limited to $D \geq 5$ dimensions. This situation is changed when Glavan & Lin declare that rescaling the coupling coefficient α_{GB} of the Gauss-Bonnet term by a factor of $1/(D - 4)$ can lead to non-trivial dynamics when taking the limit $D \rightarrow 4$, thus extend the classic Einstein-Gauss-Bonnet theory to 4 dimensions [41]. Remarkably, in their point of view, this theory can bypass Lovelock’s theorem and be free from Ostrogradsky instability. The black hole solution in AdS space has been provided in [46]. It is worth noting that this black hole solution has been found in other backgrounds, such as gravity with conformal anomaly [42,43] and with quantum corrections [44]. In this section, we will take the metric in [46] and investigate the qualitative evaporation process. We hope our result may help inspiring the researches on 4D Einstein-Gauss-Bonnet gravity.

3.1 Black hole thermodynamics in 4D Einstein-Gauss-Bonnet AdS spacetimes

In this section we give a brief review on thermodynamics of 4D Einstein-Gauss-Bonnet AdS black holes. The static spherically symmetric black hole solution takes the form

$$ds^2 = -f(r)dt^2 + f^{-1}(r)dr^2 + r^2d\Omega_2^2 \tag{16}$$

with the the metric function [46]

$$f(r) = 1 + \frac{r^2}{2\alpha} \left(1 - \sqrt{1 + 4\alpha \left(\frac{2M}{r^3} - \frac{1}{\ell^2} \right)} \right), \tag{17}$$

where α is coupling constant and the parameter M is the mass of black hole. Similarly, in the vanishing limit of mass $M \rightarrow 0$, one can obtain the vacuum solution $f(r) = 1 + r^2/\ell_{eff}^2$ with the effective AdS radius $\ell_{eff}^2 = \frac{\ell^2}{2} \left(1 + \sqrt{1 - \frac{4\alpha}{\ell^2}} \right)$, thus the coupling constant α must obey $\alpha \leq \ell^2/4$. The solution still keeps asymptotically AdS when $\alpha < 0$. Notably, the black hole solution will recover the solution in Einstein gravity when $\alpha \rightarrow 0$. Solving $f(r_+) = 0$ at the event horizon r_+ , we can obtain the mass of black hole

$$M = \frac{r_+}{2} \left(1 + \frac{r_+^2}{\ell^2} + \frac{\alpha}{r_+^2} \right). \tag{18}$$

The Hawking temperature of the black hole can be obtain as

$$T = \frac{f'(r)}{4\pi} \Big|_{r=r_+} = \frac{3r_+^4 + \ell^2(r_+^2 - \alpha)}{4\pi \ell^2 r_+ (r_+^2 + 2\alpha)}, \tag{19}$$

For $\alpha > 0$, solving $T = 0$, there is a critical horizon radius $r_c = \sqrt{(-\ell^2 + \sqrt{\ell^2(\ell^2 + 12\alpha)})}/6$ corresponding to a vanishing temperature. Similarly there will also be a critical mass

$$M_c = \frac{-\ell^2 + 12\alpha + \sqrt{\ell^4 + 12\ell^2\alpha}}{3\sqrt{6}\sqrt{-\ell^2 + \sqrt{\ell^4 + 12\ell^2\alpha}}}, \tag{20}$$

which depends on the AdS radius ℓ and the coupling constant α . For $M < M_c$, there is no black hole. The black hole entropy can also be calculated by the first law of black hole thermodynamics, see [46].

3.2 Black hole evaporation in 4D Einstein–Gauss–Bonnet AdS spacetimes

After the review of the thermodynamics of 4D Einstein–Gauss–Bonnet AdS black holes, we are ready to explore the black holes evaporation. Imposing the same assumptions that we take in the case $D \geq 5$, we can obtain the Stefan–Boltzmann law in 4D, reads

$$\frac{dM}{dt} = -b_c^2 T^4. \tag{21}$$

Note that the behavior of temperature T still plays a leading role in black hole evaporation process since it occupies the highest order. In order to analyze the qualitative behavior of the black hole temperature T , we provide the examples with $\alpha > 0$ and $\alpha < 0$ in Fig. 3. It is clear that $T = 0$ at a critical horizon radius $r_c = \sqrt{(-\ell^2 + \sqrt{\ell^2(\ell^2 + 12\alpha)})}/6$, which means that exists a critical mass M_c . When $\alpha < 0$, this situation resembles the case $\alpha < 0$ in $D \geq 5$ which also have divergent temperature at $r_{min} = \sqrt{2|\alpha|}$.

Solving the equation $\frac{\partial f(r)}{\partial r} = 0$, we can obtain the root corresponding to the photon orbit as

$$r_p = \left(-\frac{4M\alpha}{1 - \frac{4\alpha}{\ell^2}} - \sqrt{-\frac{27M^6}{(1 - \frac{4\alpha}{\ell^2})^3} + \frac{16M^2\alpha^2}{(1 - \frac{4\alpha}{\ell^2})^2}} \right)^{\frac{1}{3}} + \left(-\frac{4M\alpha}{1 - \frac{4\alpha}{\ell^2}} + \sqrt{-\frac{27M^6}{(1 - \frac{4\alpha}{\ell^2})^3} + \frac{16M^2\alpha^2}{(1 - \frac{4\alpha}{\ell^2})^2}} \right)^{\frac{1}{3}}. \tag{22}$$

Similarly, using the dimensionless variables $x \equiv r_+/\ell$ and $y \equiv \alpha/\ell^2$, we can re-express the black hole mass (18) and the temperature (19) as

$$M = \frac{x^2 + x^4 + y}{2x} \ell, \tag{23}$$

$$T = \frac{x^2 + 3x^4 - y}{4\pi x^3 + 8\pi xy} \ell^{-1}. \tag{24}$$

The impact factor $b_c = r_p/\sqrt{f(r_p)}$ can also be expressed in the form of

$$b_c = \left(\frac{4}{(2(A - B)^{1/3} + 2(A + B)^{1/3})^2} - \frac{-1 + \sqrt{1 - 4y + \frac{32y(x^2 + x^4 + y)}{x(2(A - B)^{1/3} + 2(A + B)^{1/3})^3}}}{2y} \right)^{-\frac{1}{2}} \ell$$

$$A = \frac{2y(x^2 + x^4 + y)}{x(-1 + 4y)},$$

$$B = \frac{1}{8} \sqrt{\frac{(x^2 + x^4 + y)^2(256x^4(1 - 4y)y^2 - 27(x^2 + x^4 + y)^4)}{x^6(1 - 4y)^3}}. \tag{25}$$

Inserting the above M , T and b_c into the Stefan–Boltzmann law in 4D spacetimes, we can obtain the lifetime relationship of black hole evaporation, reads

$$dt = \ell^3 F_3(x, y) dx, \tag{26}$$

where $F_3(x, y)$ is a complex function related to the M , T and b_c . Similarly, integrating this equation from ∞ to x_{min} (where the temperature T has divergence or $T = 0$), one can find that the lifetime is of the order $\sim \ell^3$ with $\alpha < 0$ or divergent with $\alpha > 0$, which resembles the case in $D \geq 5$ dimensions. We plot some numerical results of the 4D Einstein–Gauss–Bonnet AdS black hole in Fig. 4. As shown in the figure, the case of $\alpha > 0$ is very similar to the case of 5D with $\alpha > 0$, just has the difference in the value of critical mass which in case 5D is $3\pi\alpha/8$, while in case 4D equals to M_c . While in $\alpha < 0$, the qualitative features of the evaporation process are parallel to the case $\alpha < 0$ in 5D, which also possess a minimal horizon radius at $r_{min} = \sqrt{2|\alpha|}$. The qualitative features of the evaporation process identify what implied from the asymptotical behavior of the black hole temperature T .

4 Summary

By introducing the higher order curvature term, Einstein–Gauss–Bonnet theory is supposed to solve a class of singularity problem of black holes. Because of the modification, the thermodynamic properties of Einstein–Gauss–Bonnet AdS black holes are quite different from the Schwarzschild–AdS case. In the present work, we investigate the evaporation process of Einstein–Gauss–Bonnet AdS black holes in D -dimensional cases with both positive and negative coupling constant. Firstly, for the case of $D = 5$ and $\alpha > 0$, the temperature T is not divergent at a small distance and admits the state of $T = 0$ at $r_+ = 0$, which is different from the Schwarzschild–AdS case. Applying the geometric optics approximation and absorbing AdS boundary condition, we




Fig. 3 The temperature of black holes in 4-dimensions with $\alpha > 0$ and $\alpha < 0$ respectively. The left figure we set $\alpha = 0.15$, $\ell = 1$, while in the right $\alpha = -0.15$, $\ell = 1$




Fig. 4 The numerical results of the 4D Einstein–Gauss–Bonnet AdS black hole mass M as function of the lifetime t with setting $y > 0$ and $y < 0$ (corresponding to $\alpha > 0$ and $\alpha < 0$, respectively). In the left figure, we set $y = 0.1$ with $\ell = 1$, $\ell = 1.5$ and $\ell = 2$ from left to right, corresponding to critical mass $M_{c1} = 0.329509$, $M_{c2} = 0.494264$ and

$M_{c3} = 0.659019$, respectively. In the right figure, we set $y = -0.1$ with $\ell = 1$, $\ell = 1.5$ and $\ell = 2$ from left to right, corresponding to minimum mass $M_{min1} = 0.156525$, $M_{min2} = 0.234787$ and $M_{min3} = 0.313050$, respectively

investigate the evaporation of black holes by using the D -dimensional Stefan–Boltzmann law. The black holes will lose an infinite amount of mass in a “short” time, then the evaporation will become slower and slower near the $T = 0$ state, which means that the black holes with $\alpha > 0$ have a divergent lifetime in 5-dimensional spacetime. This phenomenon also obeys the third law of black hole thermodynamics. When $\alpha < 0$, the temperature is divergent at $r_{min} = \sqrt{2|\alpha|}$, and the black holes will always spend finite time $t \sim \ell^4$ to reach the extreme mass $M(r_{min})$. For the cases of $D \geq 6$, although it is difficult to get the exact form of impact factor b_c , we can still do out the evaporation process by analyzing the asymptotic behavior of the temperature. Since its temperature is always positive and divergent in $r_+ \rightarrow 0$ and $r_+ \rightarrow \infty$, the black hole lifetime is always of the order $t \sim \ell^{D-1}$, which resembles the Schwarzschild–AdS case. Applying the “novel” metric, we extend our investiga-

tion to 4D. Just with a difference in the value of critical mass between case 4D and case 5D when $\alpha > 0$, the evaporation behaviors of both $\alpha > 0$ and $\alpha < 0$ are parallel to the ones in 5D.

In the present work, we consider both $\alpha > 0$ and $\alpha < 0$ in different dimensions and find that some choices of α are not applicable. The case of $\alpha > 0$ may help us solving the problem of the divergent temperature in the terminal of the evaporation process of the Schwarzschild–AdS case, while the $\alpha < 0$ could make it worse. Also, in the cases $D = 4, 5$ there always have remnants, while in the cases $D \geq 6$ that all the black holes will evaporate in a finite time. This phenomenon is quite similar to the Hořava–Lifshitz case [23], which also has the dimensional dependent evaporation behavior.

Finally, we suggest some topics that can be further investigated. In consideration of the lifetime of the black holes in Einstein–Gauss–Bonnet and Hořava–Lifshitz gravity are

dimensional dependent, there may be some relationship between the dimensions D and asymptotic behavior of evaporation, finding this relationship may become an interesting work [67, 68]. Besides, this work can be extended to charged black holes in Einstein–Gauss–Bonnet gravity [46, 69]. By considering the Schwinger effect and the Stefan–Boltzmann law, we could calculate the charged particles emission and the mass loss [10]. How to describe the evolution of charged black holes in Einstein–Gauss–Bonnet gravity will be challenging and meaningful work. Furthermore, modification to the AdS/CFT correspondence can be considered from the perspective of Einstein–Gauss–Bonnet AdS black hole evaporation. In the present work, our derivation relies on the geometric optics approximation and absorbing boundary condition. Nowadays there have been many papers on evaporating black hole in AdS in connection with the formation of islands [70–72]. It would be interesting to study the connection among absorbing boundary conditions, black hole evaporation in AdS spacetime, and the islands. We hope to consider these questions in our future work.

Acknowledgements We would like to thank Yen Chin Ong, Rui-Hong Yue and De-Cheng Zou for useful discussion. This work is supported by National Natural Science Foundation of China (NSFC) under Grant nos. 11575083, 11565017, and the Natural Science Foundation of the Jiangsu Higher Education Institutions of China (Grant no. 20KJD140001).

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: The data used to support the findings of this study are available from the corresponding author upon request.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³.

References

1. S.W. Hawking, Black hole explosions. *Nature* **248**, 30 (1974)
2. S.W. Hawking, Particle creation by black holes. *Commun. Math. Phys.* **43**, 199 (1975). (Erratum: *Commun. Math. Phys.* **46**, 206 (1976))
3. A. Almheiri, D. Marolf, J. Polchinski, J. Sully, Black holes: complementarity or firewalls? *JHEP* **1302**, 062 (2013). [arXiv:1207.3123](https://arxiv.org/abs/1207.3123) [hep-th]
4. S.W. Hawking, M.J. Perry, A. Strominger, Superrotation charge and supertranslation hair on black holes. *JHEP* **1705**, 161 (2017). [arXiv:1611.09175](https://arxiv.org/abs/1611.09175) [hep-th]
5. D.N. Page, Particle emission rates from a black hole: massless particles from an uncharged, nonrotating hole. *Phys. Rev. D* **13**, 198 (1976)
6. D.N. Page, Particle emission rates from a black hole. 2. Massless particles from a rotating hole. *Phys. Rev. D* **14**, 3260 (1976)
7. C.M. Chambers, W.A. Hiscock, B.E. Taylor, The 'Ups' and 'Downs' of a spinning black hole. [arXiv:gr-qc/9710013](https://arxiv.org/abs/gr-qc/9710013)
8. B.E. Taylor, C.M. Chambers, W.A. Hiscock, Evaporation of a Kerr black hole by emission of scalar and higher spin particles. *Phys. Rev. D* **58**, 044012 (1998). [arXiv:gr-qc/9801044](https://arxiv.org/abs/gr-qc/9801044)
9. W.A. Hiscock, L.D. Weems, Evolution of charged evaporating black holes. *Phys. Rev. D* **41**, 1142 (1990)
10. H. Xu, Y.C. Ong, M.H. Yung, Cosmic censorship and the evolution of d-dimensional charged evaporating black holes. *Phys. Rev. D* **101**, 064015 (2020). [arXiv:1911.11990](https://arxiv.org/abs/1911.11990) [gr-qc]
11. J.M. Maldacena, The large N limit of superconformal field theories and supergravity. *Int. J. Theor. Phys.* **38**, 1113–1133 (1999). [arXiv:hep-th/9711200](https://arxiv.org/abs/hep-th/9711200)
12. E. Witten, Anti-de Sitter space, thermal phase transition, and confinement in gauge theories. *Adv. Theor. Math. Phys.* **2**, 505–532 (1998). [arXiv:hep-th/9803131](https://arxiv.org/abs/hep-th/9803131)
13. E. Witten, Anti-de Sitter space and holography. *Adv. Theor. Math. Phys.* **2**, 253–291 (1998). [arXiv:hep-th/9802150](https://arxiv.org/abs/hep-th/9802150)
14. S.S. Gubser, I.R. Klebanov, A.M. Polyakov, Gauge theory correlators from noncritical string theory. *Phys. Lett. B* **428**, 105–114 (1998). [arXiv:hep-th/9802109](https://arxiv.org/abs/hep-th/9802109)
15. D. Kubiznak, R.B. Mann, Black hole chemistry. *Can. J. Phys.* **93**(9), 999–1002 (2015). [arXiv:1404.2126](https://arxiv.org/abs/1404.2126) [gr-qc]
16. S.J. Avis, C.J. Isham, D. Storey, Quantum field theory in anti-de Sitter spacetime. *Phys. Rev. D* **18**, 3565 (1978)
17. V.J. Rocha, Evaporation of large black holes in AdS: coupling to the Evaporon. *JHEP* **08**, 075 (2008). [arXiv:0804.0055](https://arxiv.org/abs/0804.0055) [hep-th]
18. A. Almheiri, D. Marolf, J. Polchinski, D. Stanford, J. Sully, An apology for firewalls. *JHEP* **09**, 018 (2013). [arXiv:1304.6483](https://arxiv.org/abs/1304.6483) [hep-th]
19. M. Van Raamsdonk, Evaporating firewalls. *JHEP* **11**, 038 (2014). [arXiv:1307.1796](https://arxiv.org/abs/1307.1796) [hep-th]
20. D.N. Page, Finite upper bound for the hawking decay time of an arbitrarily large black hole in anti-de Sitter spacetime. *Phys. Rev. D* **97**, 024004 (2018). [arXiv:1507.02682](https://arxiv.org/abs/1507.02682) [hep-th]
21. H. Xu, M.H. Yung, Black hole evaporation in Lovelock gravity with diverse dimensions. *Phys. Lett. B* **794**, 77 (2019). [arXiv:1904.06503](https://arxiv.org/abs/1904.06503) [gr-qc]
22. H. Xu, M.H. Yung, Black hole evaporation in conformal (Weyl) gravity. *Phys. Lett. B* **793**, 97 (2019). [arXiv:1811.07309](https://arxiv.org/abs/1811.07309) [gr-qc]
23. H. Xu, Y.C. Ong, Black hole evaporation in Hořava–Lifshitz gravity. *Eur. Phys. J. C* **80**(7), 679 (2020). [arXiv:2005.02086](https://arxiv.org/abs/2005.02086) [gr-qc]
24. M.S. Hou, H. Xu, Y.C. Ong, Hawking evaporation of black holes in massive gravity. *Eur. Phys. J. C* **80**(11), 1090 (2020). [arXiv:2008.10049](https://arxiv.org/abs/2008.10049) [hep-th]
25. D.G. Boulware, S. Deser, String generated gravity models. *Phys. Rev. Lett.* **55**, 2656 (1985)
26. C.G. Callan Jr., R.C. Myers, M.J. Perry, Black holes in string theory. *Nucl. Phys. B* **311**, 673–698 (1989)
27. D. Lovelock, The Einstein tensor and its generalizations. *J. Math. Phys.* **12**, 498 (1971)
28. C. Garraffo, G. Giribet, The Lovelock black holes. *Mod. Phys. Lett. A* **23**, 1801–1818 (2008). [arXiv:0805.3575](https://arxiv.org/abs/0805.3575) [gr-qc]
29. T. Clifton, P.G. Ferreira, A. Padilla, C. Skordis, Modified gravity and cosmology. *Phys. Rep.* **513**, 1–189 (2012). [arXiv:1106.2476](https://arxiv.org/abs/1106.2476) [astro-ph.CO]
30. R.G. Cai, Gauss–Bonnet black holes in AdS spaces. *Phys. Rev. D* **65**, 084014 (2002). [arXiv:hep-th/0109133](https://arxiv.org/abs/hep-th/0109133)

31. M. Cvetič, S. Nojiri, S.D. Odintsov, Black hole thermodynamics and negative entropy in de Sitter and anti-de Sitter Einstein–Gauss–Bonnet gravity. *Nucl. Phys. B* **628**, 295–330 (2002). [arXiv:hep-th/0112045](#)
32. T. Torii, H. Maeda, Spacetime structure of static solutions in Gauss–Bonnet gravity: neutral case. *Phys. Rev. D* **71**, 124002 (2005). [arXiv:hep-th/0504127](#)
33. R.A. Konoplya, A. Zhidenko, Long life of Gauss–Bonnet corrected black holes. *Phys. Rev. D* **82**, 084003 (2010). [arXiv:1004.3772](#) [hep-th]
34. W. Xu, H. Xu, L. Zhao, Gauss–Bonnet coupling constant as a free thermodynamical variable and the associated criticality. *Eur. Phys. J. C* **74**, 2970 (2014). [arXiv:1311.3053](#) [gr-qc]
35. H. Xu, W. Xu, L. Zhao, Extended phase space thermodynamics for third order Lovelock black holes in diverse dimensions. *Eur. Phys. J. C* **74**(9), 3074 (2014). [arXiv:1405.4143](#) [gr-qc]
36. H. Xu, Z.M. Xu, Maxwell’s equal area law for Lovelock thermodynamics. *Int. J. Mod. Phys. D* **26**(04), 1750037 (2016). [arXiv:1510.06557](#) [gr-qc]
37. Y. Sun, H. Xu, L. Zhao, Thermodynamics and holographic entanglement entropy for spherical black holes in 5D Gauss–Bonnet gravity. *JHEP* **1609**, 060 (2016). [arXiv:1606.06531](#) [gr-qc]
38. R. Araneda, R. Aros, O. Miskovic, R. Olea, Magnetic mass in 4D AdS gravity. *Phys. Rev. D* **93**(8), 084022 (2016). [arXiv:1602.07975](#) [hep-th]
39. R.A. Konoplya, A. Zhidenko, Eikonal instability of Gauss–Bonnet (anti-)de Sitter black holes. *Phys. Rev. D* **95**(10), 104005 (2017). [arXiv:1701.01652](#) [hep-th]
40. R.A. Konoplya, A. Zhidenko, Quasinormal modes of Gauss–Bonnet–AdS black holes: towards holographic description of finite coupling. *JHEP* **09**, 139 (2017). [arXiv:1705.07732](#) [hep-th]
41. D. Glavan, C. Lin, Einstein–Gauss–Bonnet gravity in four-dimensional spacetime. *Phys. Rev. Lett.* **124**(8), 081301 (2020). [arXiv:1905.03601](#)
42. R.G. Cai, L.M. Cao, N. Ohta, Black holes in gravity with conformal anomaly and logarithmic term in black hole entropy. *JHEP* **04**, 082 (2010). [arXiv:0911.4379](#) [hep-th]
43. R.G. Cai, Thermodynamics of conformal anomaly corrected black holes in AdS space. *Phys. Lett. B* **733**, 183–189 (2014). [arXiv:1405.1246](#) [hep-th]
44. G. Cognola, R. Myrzakulov, L. Sebastiani, S. Zerbini, Einstein gravity with Gauss–Bonnet entropic corrections. *Phys. Rev. D* **88**(2), 024006 (2013). [arXiv:1304.1878](#) [gr-qc]
45. M. Guo, P.C. Li, Innermost stable circular orbit and shadow of the 4D Einstein–Gauss–Bonnet black hole. *Eur. Phys. J. C* **80**(6), 588 (2020). [arXiv:2003.02523](#) [gr-qc]
46. P.G.S. Fernandes, Charged black holes in AdS spaces in 4D Einstein–Gauss–Bonnet gravity. *Phys. Lett. B* **805**, 135468 (2020). [arXiv:2003.05491](#) [gr-qc]
47. G. Anastasiou, O. Miskovic, R. Olea, I. Papadimitriou, Counterterms, Kounterterms, and the variational problem in AdS gravity. *JHEP* **08**, 061 (2020). [arXiv:2003.06425](#) [hep-th]
48. R. Kumar, S.G. Ghosh, Rotating black holes in 4D Einstein–Gauss–Bonnet gravity and its shadow. *JCAP* **07**(07), 053 (2020). [arXiv:2003.08927](#) [gr-qc]
49. C.Y. Zhang, P.C. Li, M. Guo, Greybody factor and power spectra of the Hawking radiation in the 4D Einstein–Gauss–Bonnet de Sitter gravity. *Eur. Phys. J. C* **80**(9), 874 (2020). [arXiv:2003.13068](#) [hep-th]
50. R.A. Konoplya, A.F. Zinhailo, Grey-body factors and Hawking radiation of black holes in 4D Einstein–Gauss–Bonnet gravity. *Phys. Lett. B* **810**, 135793 (2020). [arXiv:2004.02248](#) [gr-qc]
51. C.Y. Zhang, S.J. Zhang, P.C. Li, M. Guo, Superradiance and stability of the regularized 4D charged Einstein–Gauss–Bonnet black hole. *JHEP* **08**, 105 (2020). [arXiv:2004.03141](#) [gr-qc]
52. S. Mahapatra, A note on the total action of 4D Gauss–Bonnet theory. *Eur. Phys. J. C* **80**(10), 992 (2020). [arXiv:2004.09214](#) [gr-qc]
53. J. Arrechea, A. Delhom, A. Jiménez-Cano, Inconsistencies in four-dimensional Einstein–Gauss–Bonnet gravity. *Chin. Phys. C* **45**(1), 013107013107 (2021). [arXiv:2004.12998](#) [gr-qc]
54. K. Yang, B.M. Gu, S.W. Wei, Y.X. Liu, Born–Infeld black holes in 4D Einstein–Gauss–Bonnet gravity. *Eur. Phys. J. C* **80**(7), 662 (2020). [arXiv:2004.14468](#) [gr-qc]
55. A. Kumar, R. Kumar, Bardeen black holes in the novel 4D Einstein–Gauss–Bonnet gravity. [arXiv:2003.13104](#) [gr-qc]
56. M. Gürses, T.C. Şişman, B. Tekin, Is there a novel Einstein–Gauss–Bonnet theory in four dimensions? *Eur. Phys. J. C* **80**(7), 647 (2020). [arXiv:2004.03390](#) [gr-qc]
57. B.E. Panah, K. Jafarzade, S.H. Hendi, Charged 4D Einstein–Gauss–Bonnet–AdS black holes: shadow, energy emission, deflection angle and heat engine. *Nucl. Phys. B* **961**, 115269 (2020). [arXiv:2004.04058](#) [hep-th]
58. J. Arrechea, A. Delhom, A. Jiménez-Cano, Comment on “Einstein–Gauss–Bonnet gravity in four-dimensional spacetime. *Phys. Rev. Lett.* **125**(14), 149002 (2020). [arXiv:2009.10715](#) [gr-qc]
59. M. Gürses, T.C. Şişman, B. Tekin, Comment on “Einstein–Gauss–Bonnet gravity in 4-dimensional space-time”. *Phys. Rev. Lett.* **125**(14), 149001 (2020). [arXiv:2009.13508](#) [gr-qc]
60. F.W. Shu, Vacua in novel 4D Einstein–Gauss–Bonnet gravity: pathology and instability? *Phys. Lett. B* **811**, 135907 (2020). [arXiv:2004.09339](#) [gr-qc]
61. S.W. Hawking, D.N. Page, Thermodynamics of black holes in anti-de Sitter space. *Commun. Math. Phys.* **87**, 577 (1983)
62. Y.C. Ong, Y. Yao, Charged particle production rate from cosmic censorship in dilaton black hole spacetimes. *JHEP* **10**, 129 (2019). [arXiv:1907.07490](#) [gr-qc]
63. D. Birmingham, Topological black holes in Anti-de Sitter space. *Class. Quantum Gravity* **16**, 1197–1205 (1999). [arXiv:hep-th/9808032](#)
64. P. Landsberg, A. De Vos, The Stefan–Boltzmann constant in n-dimensional space. *J. Phys. A Math. Gen.* **22**, 1073 (1989)
65. T.R. Cardoso, A.S. de Castro, The blackbody radiation in D-dimensional universes. *Rev. Bras. Ens. Fis.* **27**, 559 (2005). [arXiv:quant-ph/0510002](#)
66. P. Chen, Y.C. Ong, D.H. Yeom, Black hole remnants and the information loss paradox. *Phys. Rep.* **603**, 1–45 (2015). [arXiv:1412.8366](#) [gr-qc]
67. R. Emparan, C.P. Herzog, The large D limit of Einstein’s equations. *Rev. Mod. Phys.* **92**(4), 045005 (2020). [arXiv:2003.11394](#) [hep-th]
68. F. Holdt-Sørensen, D.A. McGady, N. Wintergerst, Black hole evaporation and semiclassicality at large D. *Phys. Rev. D* **102**(2), 026016 (2020). [arXiv:1908.08083](#) [hep-th]
69. T. Torii, H. Maeda, Spacetime structure of static solutions in Gauss–Bonnet gravity: charged case. *Phys. Rev. D* **72**, 064007 (2005). [arXiv:hep-th/0504141](#)
70. G. Penington, Entanglement wedge reconstruction and the information paradox. *JHEP* **09**, 002 (2020). [arXiv:1905.08255](#) [hep-th]
71. A. Almheiri, R. Mahajan, J.E. Santos, Entanglement islands in higher dimensions. *Sci. Post Phys.* **9**(1), 001 (2020). [arXiv:1911.09666](#) [hep-th]
72. A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian, A. Tajdini, The entropy of Hawking radiation. [arXiv:2006.06872](#) [hep-th]