The superconducting dome for holographic doped Mott insulator with hyperscaling violation

Wenhe Cai^{1,2,a}, Sang-Jin Sin^{1,b}

¹ Department of Physics, Hanyang University, Seoul 04763, Korea

² Department of Physics, Shanghai University, Shanghai 200444, China

Received: 12 February 2021 / Accepted: 4 June 2021 / Published online: 30 June 2021

© The Author(s) 2021

Abstract We reconsider the holographic model featuring a superconducting dome on the temperature-doping phase diagram with a modified view on the role of the two charges. The first type charge with density ρ_A make the Mott insulator, and the second one with ρ_B is the extra charge by doping, so that the complex scalar describing the cooper pair condensation couples only with the second charge. We point out that the key role in creating the dome is played by the three point interaction $-c\chi^2 F_{\mu\nu} G^{\mu\nu}$. The T_c increases with their coupling. We also consider the effect of the quantum critical point hidden under the dome using the geometry of hyperscaling violation. Our results show that the dome size and optimal temperature increase with z whatever is θ , while we get bigger θ for larger (smaller) dome depending on $z > 2$ ($z < 2$). We also point out that the condensate increases for bigger value of θ but for smaller value of z .

1 Introduction

Holographic duality, also known as the gauge/gravity duality [1–3], has brought many insights in understanding strongly interacting electron systems, especially for condensed matter physics. For example, the way to calculate transports and spectral feature of strongly correlated system has been suggested and new mechanism of superconductivity has been suggested. See Refs. [4, 5] and references therein.

One important problem is to understand the phase diagram of the high T_c superconductivity: various high- T_c compounds all fit into a universal phase diagram where normal, superconducting, anti-ferromagnetic and pseudogap phases compete [6] and coexist. Recently, a holographic model was suggested which qualitatively realizes the phase diagram in the temperature-doping plane [7]. A simplified model was

proposed in [8] where the non-abelian gauge field is replaced by a higher power of scalar field that breaks the translation symmetry. Here metal insulator transition is still realized although anti-ferromagnetic origin of the insulating phase is undermined. In this model the metallic phase was characterized by the DC conductivity decreasing with temperature, and the pseudo-insulator characterized by the DC conductivity increasing with temperature (see the phase diagram in Fig. 1b).

In the previous papers [7, 8], the mobile carriers density is ρ_A , and $\rho_B = x\rho_A$ represents the impurities density. In this paper, we reinterpret the model such that it is more consistent with the doped Mott insulator. At zero doping, the lattice is half filled so that it has a Mott gap. The free electrons can not move around due to the strong Coulomb repulsion. We will make sure the ρ_A electric charges do not move by assuming that, in the conductivity calculation, the ρ_A charge does not respond to the external field while ρ_B charge responds. Then we dop the system with doping ratio x . For each dopant atom we can assume there are $1 + 1$ valence electrons: one contribute to the non-moving half filled state and the other contribute to itinerant electron whose number is say, $N_1 = xN$. These are the movable electrons. Therefore the density of the non-movable charge density is always (before and after doping,) $N/V = \rho_A$, and the movable charge density is $\rho_B = N_1/V = xN/V = x\rho_A$. We assume that as illustrated in Fig. 1a, ρ_A only contributes to the electrons in Mott insulator, so that it does not contribute to conductor or superconductor. On the other hand, the doped charges of density ρ_B contribute to the density of state near the Fermi surface, so that when $x = \rho_B/\rho_A$ is large enough, charge from the impurities begins to make the superconductor. With this setup, we studied the roles of the couplings on the phase diagram and evaluated parameteric dependence of the critical temperature at the optimal doping. It turns out that what makes the superconducting dome is the three points coupling between the

^a e-mail: whcai@shu.edu.cn (corresponding author)

^b e-mail: sjsin@hanyang.ac.kr

Fig. 1 **a** The density of the states for doped Mott insulator. The degrees of freedom in the red region are provided by non-moving charge with density ρ_A , while those of the blue region is provided by the doped charges ρ_N . **b** Phase diagram in the temperature-doping plane. The line connecting $(x_0, 0)$, $(0, T_0)$ denotes the phase boundary of the pseudo-

insulating phase. (x_1, x_2) is the interval of the superconducting dome. (x_0, T_0) denotes the optimal T_c . **c** The metal pseudo-insulator transition happens when the peak-position T_p is at 0. Here we used $\alpha = 1$ and $x = 3$

Fig. 2 Role of c . We plot the superconducting phase with different $c = 0, 0.01, 0.2, 0.5, 1, 1.5, 2$ $a = b = 0$. No dome without c . Larger dome with increasing c . Notice that $c \rightarrow 0$ is a singular limit

density of cooper pair and two kind of charges, namely

$$\mathcal{L}_{int} = -c\chi^2 F_{\mu\nu} G^{\mu\nu}, \quad (1)$$

where χ is the amplitude of the complex scalar describing the cooper pair condensation, and F and G are the field strengths of the two gauge fields created by the two kinds of charges. Increasing the coupling c makes the dome higher and bigger, therefore increasing this coupling is the key to increases the T_c . Also we find that by increasing the coupling b of the interaction term $\mathcal{L}_{int} = b\chi^2 G_{\mu\nu} G^{\mu\nu}$, one can make the width of the dome smaller to resemble the cuprate case.

Since the superconducting dome is believed to cover a quantum critical points (QCP), it is very interesting to explore the effects of dynamical exponents of the QCP. The hyperscaling violating geometry is precisely the geometry that realizes the symmetry of a general class of QCP. We generalize the model of [8] to a holographic model with hyperscaling violating geometry using the solution [9], where a black hole solution is derived with a dynamic exponent z and a hyperscaling violating exponent θ . We investigate the exponent

dependence of the dome size as well as that of the superconducting condensate. We found that the dome size and optimal temperature increase with z whatever is θ , while we get larger dome for bigger θ if $z > 2$ and vice versa. We also point out that the condensate increases for bigger value of θ but for smaller value of z .

The paper is organized as follows. In Sect. 2, we first review the doped holographic superconductors with broken translational symmetry. Then, we investigate impact of the coupling on the phase boundary, on the endpoints of the superconducting dome and on the optimal T_c . We single out the key parameter to create the dome: the three points coupling between the density of cooper pair and two different charge carriers. In Sect. 3, in order to explore the effects of QCP, we extend Baggioli and Goykhman's model to the doped holographic superconductor with hyperscaling violation. In Sect. 4, we focus on the superconducting condensate for $\theta = 0$ and $\theta = 1$. The Sect. 5 is the summary and discussion. In the Appendix A, we show that two methods to obtain the superconducting dome are equivalent. In the Appendix B, we figure out the role of parameters a, b, α in the superconducting dome.

2 Coupling dependence of the critical temperature

We first briefly review the doped holographic superconductors with broken translational symmetry. Baggioli and Goykhman introduced a doped holographic model with a momentum dissipation [8], and find the superconducting dome. There are two U(1) gauge fields A_μ and B_μ , two neutral scalar $\phi^I = \alpha x^I$, ($I = x, y$), the complex scalar field $\psi = \chi e^{i\theta}$. $A_\mu = (A_t(u), 0, 0, 0)$ is the bulk dual of the density of the charge carrier. $B_\mu = (B_t(u), 0, 0, 0)$ is the bulk dual of the density of impurity. The neutral and massless scalars are responsible for the breaking of translational sym-

Fig. 3 Role of c : **a** The dome-shaped superconducting phase with the coefficient c appearing in the expansion (6). **b, c** The left endpoint x_1 and the right endpoint x_2 of the superconducting region. We

fix $a = 10, b = 4/3, \alpha = 0$ and choose different c . The dashed line represents $(1, 0)$, and the solid line represents $(0, 1)$

metry. The complex scalar field represents the order parameter for superconducting phase transition. $x = \rho_B/\rho_A$ is called as dope parameter. u denotes holographic coordinate which is dual to the renormalization scale. The action is written as

$$S = \frac{1}{16\pi} \int d^4x \sqrt{-g} \left(R + \frac{6}{L^2} + \mathcal{L}_c + \mathcal{L}_s \right) \quad (2)$$

$$\begin{aligned} \mathcal{L}_c = & -\frac{Z_A(\chi)}{4} A_{\mu\nu} A^{\mu\nu} - \frac{Z_B(\chi)}{4} B_{\mu\nu} B^{\mu\nu} \\ & - \frac{Z_{AB}(\chi)}{2} A_{\mu\nu} B^{\mu\nu} \\ & - \frac{1}{2} (\partial_\mu \chi)^2 - H(\chi) (\partial_\mu \theta - q_A A_\mu - q_B B_\mu)^2 - V_{int}(\chi) \end{aligned} \quad (3)$$

$$\mathcal{L}_n = -2m^2 V(X). \quad (4)$$

$$\mathcal{L}_n = -2m^2 V(X). \quad (5)$$

We consider two interesting choices for (q_A, q_B) : $(1, 0)$ or $(0, 1)$. However, the density ρ_A only contributes to the Mott insulator, so $(0, 1)$ is the better choice for the couplings between the gauge fields and the complex scalar. Here

$$\begin{aligned} H(\chi) &= \frac{n\chi^2}{2}, \quad Z_A(\chi) = 1 - \frac{a\chi^2}{2}, \\ Z_B(\chi) &= 1 - \frac{b\chi^2}{2}, \quad Z_{AB}(\chi) = \frac{c\chi^2}{2}. \end{aligned} \quad (6)$$

$$V_{int}(\chi) = \frac{M^2 \chi^2}{2}. \quad (7)$$

In this paper, we consider the non-linear Lagrangian

$$V(X) = \frac{X}{2m^2} + \left(\frac{X}{2m^2} \right)^5, \quad X = \frac{1}{2} g^{\mu\nu} \partial_\mu \phi^I \partial_\nu \phi^I. \quad (8)$$

The corresponding equations of motion are as follows:

$$\begin{aligned} R_{\mu\nu} - \frac{Z_A}{2} A_{\nu\sigma} A_\mu^\sigma - \frac{Z_B}{2} B_{\nu\sigma} B_\mu^\sigma - \frac{Z_{AB}}{2} (A_{\nu\sigma} B_\mu^\sigma + A_{\mu\sigma} B_\nu^\sigma) \\ - \frac{1}{2} \partial_\mu \chi \partial_\nu \chi - H(\chi) (\partial_\mu \theta - q_A A_\mu - q_B B_\mu) (\partial_\nu \theta - q_A A_\nu - q_B B_\nu) \\ - \frac{1}{2} g_{\mu\nu} L = 0, \end{aligned}$$

$$\begin{aligned} \nabla_\nu (Z_A A^{\nu\mu} + Z_{AB} B^{\nu\mu}) + 2q_A H(\nabla^\mu \theta - q_A A^\mu - q_B B^\mu) &= 0, \\ \nabla_\nu (Z_B B^{\nu\mu} + Z_{AB} A^{\nu\mu}) + 2q_B H(\nabla^\mu \theta - q_A A^\mu - q_B B^\mu) &= 0, \\ \nabla_\mu \nabla^\mu \chi - \frac{1}{4} \partial_\chi Z_A A^2 - \frac{1}{4} \partial_\chi Z_B B^2 - \frac{1}{2} \partial_\chi Z_{AB} A \cdot B \\ - \partial_\chi H(\partial_\mu \theta - q_A A_\mu - q_B B_\mu)^2 - \partial_\chi V_{int} &= 0. \end{aligned}$$

We obtain the solutions by solving the background equations of motion,

$$ds^2 = \frac{L^2}{u^2} \left(-f(u) e^{-\tau(u)} dt^2 + dx^2 + dy^2 + \frac{du^2}{f(u)} \right), \quad (9)$$

$$f(u) = \frac{(u - u_h) \left(2u^2 (\alpha^2 L^2 u_h^2 - 2) + u^3 u_h^3 (\rho_A^2 + \rho_B^2) - 4uu_h - 4u_h^2 \right)}{4u_h^3}, \quad (10)$$

$$A_t(u) = \rho_A(u_h - u), \quad B_t(u) = \rho_B(u_h - u). \quad (11)$$

We emphasize that we take $(q_A, q_B) = (0, 1)$ in this paper. But for the comparison, we also calculate for the $(1, 0)$ case. There are three phases on the temperature and doping plane: superconducting, metallic and pseudo-insulating (see the phase diagram in [6]). As a generalization and comparison to the results in [8], it is interesting to discuss how much impact of the parameters a, b, c, α on $x_0, T_0, x_1, x_2, x_O, T_O$.

The DC conductivity can be calculated analytically following [10, 11] and the result is given by:

$$\sigma_{DC} = 1 + \frac{\rho_B^2 u_h^2}{2\alpha^2 \left(1 + 5 \frac{u_h^8 \alpha^8}{(2m^2)^4} \right)}. \quad (12)$$

Here u_h is the location of the horizon. Notice that we regards the ρ_A is not moving degree of freedom and does not contribute the the conductivity. There is other way to introduce insulator behavior by the coupling X with the field strength [12, 13]. Here, metal insulator transition is not our main concern. Following [14–16], we obtain the DC conductivity. The pseudo-insulating phase by the change from metallic ($d\sigma_{DC}/dT < 0$) to pseudo-insulating ($d\sigma_{DC}/dT > 0$)

is shown in Fig. 1c. We vary the ratio x and evaluate the movement of the peak-position. Our results imply that T_p decreases as x increases.

Now, we discuss the instability of a normal phase of the bulk system to determine whether a boundary system has a superconducting phase. The effective mass is read off where BF bound is violated. We first derive instability analytically in the normal state at zero temperature. Two endpoints x_1 and x_2 can be determined by the violated AdS_2 BF bound, namely $m_{eff}^2 L_2^2 < -1/4$.

We furthermore study the superconducting dome in a finite-temperature normal phase background. We begin with the lineared equation of motion for χ . The scalar χ represents the order parameter for superconducting phase transition (see more details in [7, 17]). Then we look for the finite temperature black brane solution to determine the superconducting dome, which satisfies the two boundary conditions. It is regular at the horizon, so it demands the expansion as follow

$$\begin{aligned} f(u) &= f'(u_h)(u - u_h) + \mathcal{O}(u - u_h)^2, \\ \chi(u) &= \chi(u_h) + \chi'(u_h)(u - u_h), \\ \tau(u) &= \tau(u_h) + \tau'(u_h)(u - u_h), \\ A_t &= \rho_A(u_h - u), \quad B_t = \rho_B(u_h - u). \end{aligned}$$

We solve the lineared equation for $\delta\chi$, which is considered as a probe on the AdS-RN background

$$\begin{aligned} \delta\chi''(u) + \left(\frac{f'(u)}{f(u)} - \frac{2}{u} \right) \delta\chi'(u) \\ + \frac{f(u) \left(u^4(a + x(bx + 2c)) - 2M^2 \right) + 2nu^2(u - u_h)^2(q_A + q_Bx)^2}{2u^2 f(u)^2} \delta\chi(u) = 0. \end{aligned} \quad (13)$$

There are five independent parameters u_h , $\chi(u_h)$, $\tau(u_h)$, ρ_A , ρ_B . This scaling symmetry can be used to set $\tau(u_h) = 0$, and we also set $\rho_A = 1$, $\rho_B = x$. Actually, we are left with two independent parameters x and u_h , namely doping and temperature. By instituting into equations of motion for $\delta\chi$, we solve background equations of motion numerically out to large value of r . Based on this initial condition. Then, we obtain the source free solution from horizon to the boundary. Afterwards, we keep the doping parameter x fixed, and find the maximal T_c (see Appendix A for details).

We focus on the three points coupling between the density of cooper pair and two kind of charges: $-c\chi^2 F_{\mu\nu} G^{\mu\nu}$, because the coupling c is the key parameter for the superconducting dome. We first discuss the role of c in the simplified model

$$Z_A(\chi) = 1, \quad Z_B(\chi) = 1, \quad Z_{AB}(\chi) = \frac{c\chi^2}{2}. \quad (14)$$

Our results are presented in Fig. 2. If c is negative, there is no instability solution. In this paper, we do not need to consider

Fig. 4 Phase diagram on the temperature-doping plane with different θ and fixed $z = 1$. In this figure, solid lines are for $(q_A, q_B) = (1, 0)$, and dashed lines are for $(q_A, q_B) = (0, 1)$. Black and red $\theta = 0, 0.1$, respectively. The critical temperature decreases as θ increases

Fig. 5 Phase diagram on the temperature-doping plane with different z and fixed $\theta = 0$. In this figure, solid lines are for $(q_A, q_B) = (1, 0)$, and dashed lines are for $(q_A, q_B) = (0, 1)$. Black and red represent $z = 1, 1.2$, respectively. The critical temperature increases as θ increases

the negative c . The superconducting dome occurs if $c \neq 0$ and the dome expands as c increases. Our result shows that the superconducting expands as c increase. If the cross term vanishes ($c = 0$, $Z_{AB} = 0$), there is no superconducting instability. A dome exists even at $a = b = 0$ if $c \neq 0$. Our result shows that c alone is almost enough to create the dome. Therefore, the coupling c is crucial to high T_c .

Then, we discuss the case of $a \neq 0$, $b \neq 0$. We find that the behavior of endpoints is the same for $(1, 0)$ or $(0, 1)$. Especially, x_1 moves left and x_2 moves right when c increase to make the dome bigger (see Fig. 3).

The optimal temperature T_O with $(q_A, q_B) = (1, 0)$ is always smaller than the case of $(0, 1)$. This result is not surprising because the coupling $Z_B = 1 - \frac{b\chi^2}{2}$ is stronger than the coupling $Z_A = 1 - \frac{a\chi^2}{2}$ in our choice. By the same reason, the superconducting dome of $(0, 1)$ theory is bigger than the one with $(1, 0)$ one. The roles of other parameters are given in the Appendix B. Perhaps the most interesting result is the fact that the coupling described by c alone can generate the superconducting dome and increasing its size, and the roles of other coupling Z_A Z_B are secondary ones for producing the dome.

Fig. 6 Phase diagram on the temperature-doping plane with different θ and fixed $z = 2.4$. In this figure, solid lines are for $(q_A, q_B) = (1, 0)$, and dashed lines are for $(q_A, q_B) = (0, 1)$. Black and red represent $\theta = 0.2, 0.4$, respectively. The critical temperature increases as z increases

3 The doped superconductor with hyperscaling violation

In the previous section, we studied the doped holographic superconductor with broken translational symmetry. In order to explore the effects of the QCP, it would be natural to extend the present analysis to the case of hyperscaling violation. We begin with the black hole solutions in Einstein–Maxwell–axion–dilaton theory with a dynamic exponent z and a hyperscaling violation exponent θ [9], where a $U(1)$ gauge field is considered as an auxiliary gauge field, leading to a Lifshitz-like vacuum. The neutral and massless scalars ϕ^I : $\phi^x = \alpha x$, $\phi^y = \alpha y$ generate momentum relaxation. We generalize this model by introducing a perturbative charge scalar $\psi = \chi e^{i\phi}$ in four-dimensional bulk spacetime. Now there are three gauge fields. A_1 is the auxiliary gauge field to support the Lifshitz gravity as usual, while A_2 and A_3 are the physical gauge fields which provide the finite chemical potentials. A_2 is the bulk dual of the density of the charge carrier ρ_A which describe the Mott insulator while A_3 is the bulk dual of the density of doped charge ρ_B which can move and provides the density of state at the Fermi surface. We describe the non-moving nature as the absence of the coupling of such charge with the electric field. Therefore our model is $q_B = 1, q_A = 0$. To compare with previous treatment, we also considered results coming from the $q_A = 1, q_B = 0$.

The new doping parameter is defined as

$$y = \frac{x}{1+x}, \quad x = \rho_B / \rho_A. \quad (15)$$

Here $0 < y < 1$. Since x is large in the superconducting dome, we consider y instead of x in the following calculation. The action is written as

$$S = \frac{1}{16\pi} \int d^4x \sqrt{-g} \left[R + \frac{6}{L^2} - \frac{1}{4} (e^{\lambda_1 \phi}) F_1^2 - \frac{1}{4} \left(e^{\lambda_2 \phi} + \frac{a \chi^2}{2} \right) F_2^2 \right]$$

$$\begin{aligned} & - \frac{1}{4} \left(e^{\lambda_3 \phi} + \frac{b \chi^2}{2} \right) F_3^2 - \frac{1}{2} \frac{c \chi^2}{2} F_2 \cdot F_3 \\ & - \frac{1}{2} (\partial \chi_i)^2 + H(\chi) (\partial_\mu \theta - q_A A_2 - q_B A_3)^2 \\ & - V_{int}(\chi) - \frac{1}{2} (\partial \phi)^2 - \frac{1}{2} e^{\eta \phi} (\partial \phi^I)^2 + V e^{\gamma \phi} \end{aligned} \quad (16)$$

Here we define the following coupling and potential

$$H(\chi) = \frac{n \chi^2}{2}, \quad V_{int}(\chi) = \frac{M \chi^2}{2}.$$

Following [9], we have

$$\begin{aligned} q_1 &= \sqrt{2} \sqrt{-2 + z + z^2 + \theta - z\theta} \\ Z_1 &= e^{\lambda_1 \phi} = u^{-(\theta-4)}, \quad Z_2 = e^{\lambda_2 \phi} = u^{(\theta-2z+2)}, \\ Z_3 &= e^{\lambda_3 \phi} = u^{(\theta-2z+2)} \\ Y &= e^{\eta \phi} = \frac{1}{Z_2}, \quad \phi = \sqrt{(2-\theta)(2z-2-\theta)} \ln \frac{1}{u}, \\ V &= (z-\theta+1)(z+2-\theta)u^{-\theta} \end{aligned}$$

The metric ansatz is

$$\begin{aligned} ds^2 &= \frac{1}{u^{2-\theta}} \left(-\frac{f(u)}{u^{2(z-1)}} e^{-\tau(u)} dt^2 + dx^2 + dy^2 + \frac{du^2}{f(u)} \right), \\ A_1 &= a_1(u)dt, \quad A_2 = a_2(u)dt, \\ A_3 &= a_3(u)dt, \\ \chi &= \chi(u), \quad \theta \equiv 0. \end{aligned} \quad (17)$$

where u is the radial bulk coordinate. We obtain the solution as follows,

$$\begin{aligned} f(u) &= 1 - \frac{m}{u^{\theta-z-2}} - \frac{\alpha^2}{(\theta-2)(z-2)u^{\theta-2z}} \\ &+ \frac{(\rho_A^2 + \rho_B^2)(\theta-2)}{2(\theta-2)u^{2\theta-2z-2}} \\ m &= -\frac{1}{u_h^{2+z-\theta}} \left(-1 + \frac{\alpha^2}{(-2+z)(-2+\theta)u_h^{-2z+\theta} - \frac{(\rho_A^2 + \rho_B^2)(-z+\theta)}{2(-2+\theta)u_h^{-2-2z+2\theta}}} \right) \\ a_1(u) &= \frac{q_1}{2+z-\theta} \left(\frac{1}{u^{2+z-\theta}} - \frac{1}{u_h^{2+z-\theta}} \right) \\ a_2(u) &= \mu_A - \rho_A u^{z-\theta}, \\ a_3(u) &= \mu_B - \rho_B u^{z-\theta}, \end{aligned} \quad (18)$$

When $z = 1, \theta = 0$, $f(u)$ return to the form in [8]. Following [7], we fix $a = 10, b = 4/3, c = 14/3, n = 1, \alpha = 1, M^2 = -5/4$. The superconducting dome survives the translational symmetry breaking [8]. Our calculation shows

Table 1 Optimal temperature T_O and SC dome with different z and θ : “-” denotes negative T_O . “•” denotes the monotonously decreasing T_c . “o” denotes the monotonously increasing T_c . “=” denotes that there is no instability solution. Here we fix $(1, 0)$

$z \setminus \theta$	1	11/10	12/10	13/10	14/10	15/10	16/10	17/10	19/10	21/10	22/10	24/10	26/10	28/10	3
- 1/10	0.051	0.066	0.078	0.088	0.097	=	=	-	-	-	-	-	-	-	-
0	0.038	0.056	0.071	0.083	0.093	0.102	-	0.116	0.127	0.136	0.140	0.146	0.152	0.157	0.161
1/10	0.021	0.044	0.061	0.076	0.093	0.098	-	0.114	0.126	0.136	0.140	0.147	0.153	0.158	0.162
2/10	-	0.028	0.050	0.067	0.081	0.093	-	0.112	0.126	0.136	0.141	0.148	0.154	0.159	0.163
3/10	-	0.008	0.035	0.057	0.074	0.088	-	0.109	0.126	0.137	0.141	0.149	0.156	0.161	0.165
4/10	-	-	-	0.044	0.065	0.081	0.095	0.093	o	o	o	0.151	0.158	0.163	0.167
5/10	-	-	-	0.028	0.053	0.073	0.090	o	o	o	o	0.154	0.161	0.166	0.170
6/10	-	-	-	0.008	0.039	0.063	0.083	o	o	o	o	o	0.165	0.170	0.173
7/10	-	-	-	-	0.022	0.051	o	o	o	o	o	o	o	0.174	0.178
8/10	-	-	-	-	-	0.037	o	o	o	o	o	o	o	o	0.184
9/10	-	-	-	-	-	0.023	0.053	o	o	o	o	o	o	o	o
1	•	•	0.028	-	-	•	0.043	o	o	o	o	o	o	o	o

Fig. 7 Relation between the optimal temperature T_O and z, θ . The maximum critical temperature defines the optima doping. Left: $(q_A, q_B) = (1, 0)$. Right: $(q_A, q_B) = (0, 1)$

the result still holds even for hyperscaling violating background. When $z = 1, \theta = 0$, our result could return to the one in [8]. The phase diagrams with different z and θ is presented in Figs. 4, 5 and 6.

The numerical optimal temperatures T_O with different z and θ are presented in Table 1.

The relation between the optimal temperature T_O and dynamical z with different θ is shown in Fig. 7. It shows that T_O increase as z increase. The result is in agreement with the numerical calculation without the mass of the probed scalar field in [18]. T_O decrease as θ increases when $z < 2$. However, we have the opposite behavior when $z > 2$. Figures 4 and 6 also indicate that for $z < z_c$ the optimal T_c decreases as θ increases, while for $z > z_c$, T_O increases as θ increases. The result is not in agreement with the previous work [18–20]. The difference is expected because our model is quite different from their model. There is only one gauge field in their model, so superconducting dome can not be generated.

Fig. 8 The cooper pair condensate $\langle O \rangle$ with fixed $\alpha = 1, x = 2$. The blue line is for $z = 1, \theta = 0$. The orange line is for $z = 3/2, \theta = 0$. The green line is for $z = 3/2, \theta = 1$. The condensate increases for bigger value of θ , but it is suppressed by z

4 Condensate for hyperscaling violation

We start with AdS black hole with dynamic exponent and hyperscaling violation exponent. The Hawking temperature is determined by

$$T = \frac{f'(r_0)}{4\pi}. \quad (20)$$

Therefore we define temperature as $T = r_0$. Consequently, we rewrite the equations of motion for A_t , B_t , χ in terms of the scaled fields: $A_t \rightarrow \frac{A_t}{r_0}$, $B_t \rightarrow \frac{B_t}{r_0}$, $\chi \rightarrow \frac{\chi}{r_0}$, $\mu \rightarrow \frac{\mu}{r_0}$. According to $\mu \rightarrow \frac{\mu}{r_0} = \frac{\mu}{T}$, fixing μ by changing T is equivalent to fixing T by changing μ . In order to simplify the following computations, we fix $r_0 = 1$. At the horizon, we demand the following expansion

$$\begin{aligned} A_t &= \alpha_0(r-1) + \alpha_1(r-1)^2 + \alpha_2(r-1)^3, \\ B_t &= \beta_0(r-1) + \beta_1(r-1)^2 + \beta_2(r-1)^3, \\ \chi &= \gamma_0 + \gamma_1(r-1) + \gamma_2(r-1)^2. \end{aligned}$$

Meanwhile we can impose the boundary condition $A_t = B_t = 0$, $A'_t = \alpha_0$, $B'_t = \beta_0$, $\chi = \gamma_0$. The horizon regularity keeps conditions for equations of motion

$$\begin{aligned} Z_A (2 A''_t + \tau' A'_t) + Z_{AB} (2 B''_t + \tau' B'_t) \\ + 2 \chi' (\dot{Z}_A A'_t + \dot{Z}_{AB} B'_t) - 4 q_A H \frac{q_A A_t + q_B B_t}{u^2 f} = 0, \end{aligned} \quad (21)$$

$$\begin{aligned} Z_B (2 B''_t + \tau' B'_t) + Z_{AB} (2 A''_t + \tau' A'_t) \\ + 2 \chi' (\dot{Z}_B B'_t + \dot{Z}_{AB} A'_t) - 4 q_B H \frac{q_A A_t + q_B B_t}{u^2 f} = 0, \end{aligned} \quad (22)$$

$$\begin{aligned} \chi'' + \left(\frac{f'}{f} - \frac{2}{u} - \frac{\tau'}{2} \right) \chi' - \frac{L^2}{u^2 f} \dot{V}_{int} \\ + \frac{e^\tau u^2}{2 f} \left(\dot{Z}_A \bar{A}_t'^2 + \dot{Z}_B \bar{B}_t'^2 + 2 \dot{Z}_{AB} A'_t B'_t \right) \\ + \frac{e^\tau \dot{H}}{f^2} (q_A A_t + q_B B_t)^2 = 0. \end{aligned} \quad (23)$$

After solving the conditions for α_0 , β_0 , γ_0 , α_1 , β_1 , γ_1 , α_2 , β_2 , γ_2 , we are left with three independent parameters α_0 , β_0 , γ_0 . By integrating out to infinity, these solutions of (21–23) for given α_0 , β_0 , γ_0 behave as

$$\begin{aligned} \chi &= \frac{\chi_1}{r} + \frac{\chi_2}{r^2}, \\ \chi_1 &= r \chi(r), \chi_2 = -r^2 (r \chi(r))', \\ A_t &= \mu_A - \frac{\rho}{r}, B_t = \mu_B - \frac{\rho}{r}. \end{aligned}$$

We set $\chi_1 = 0$ and $\mu_A = 1$ to determine a curve. Along this curve, we can calculate χ_2 and μ_B at infinity [21]. Somewhere along this curve χ_2 could be zero. To determine critical point, μ_c is defined by $\chi_1 = \chi_2 = 0$. Based on the approach in [22, 23], we explore the superconducting condensate with the dynamic exponent z and the hyperscaling violation exponent θ . Here we vary $1/\mu$ instead of varying temperature, and study the superconducting condensate with $\theta = 0$ for bosonic case and $\theta = 1$ for fermionic case. The cooper pair condensate $\langle O \rangle$ presented in Fig. 8 suggests that the couplings Z_A and Z_B are positive for all radial position in the case of broken translational symmetry and hyperscaling violation. The condensate becomes easier for bigger value of θ , but it is suppressed by z . The behavior is in agreement with the result in [18, 19]. However, the superconducting dome expands and the optimal temperature increases when z increases in Sect. 3. The results are not contradictory, since $\langle O \rangle$ is not only a function of T_c [24] but also the function of z and θ [18, 25, 26]. As z increases, the Fermi surface of cuprates becomes a Fermi arc, which means that it cannot be in a closed shape [27, 28]. So the number of superconducting electrons decrease. Consequently superconducting condensate decreases.

5 Conclusion and discussion

In this paper, we have studied the holographic theory which has the ability to realize the doped high-temperature superconductors. Based on the previous work on pseudo-insulator and superconducting phases on the doping-temperature plane, especially instability conditions at zero temperature and at finite temperature, we further evaluate the impact of coefficients a , b , c , α on the corner region of pseudo-insulator and the superconducting dome with broken translational symmetry. Our results show the three points coupling c is the key parameter. Afterwards, we extend our analysis to the case with a dynamic exponent and a hyperscaling violation exponent. Besides, we also calculate the superconducting condensate to verify the region of the couplings Z_A , Z_B .

We find which parameter has the most influence on the boundary of the pseudo-insulating phase (x_0 , T_0), two endpoints (x_1 , x_2) of the superconducting dome and the optimal temperature (x_O , T_O). As noted in Figs. 10 and 11, we notice that the left endpoint x_1 is more sensitive to a , but the right endpoint x_2 is more sensitive to b . Furthermore, our result indicates that the superconducting dome is expanding as the couplings Z_A , Z_B , Z_{AB} increase. The result is an advance in the research on high-temperature superconductivity.

In our calculation, q_B can be chosen to be zero, so ρ_B can be considered as the density of spin doping. One of the main progress of this paper is to show that the superconducting dome-shaped region with (0, 1) is much bigger than the one with (1, 0), and the three point interaction $-c \chi^2 F_{\mu\nu} G^{\mu\nu}$

alone is almost enough to create the dome. Our numerical calculation also shows the superconducting dome survives when $\theta \neq 0$. Let us combine our results with [7,8]. The superconducting phase with hyperscaling violation is in qualitative agreement with the case of $\theta = 0$. The depth of the superconducting dome increase as the dynamical exponent z increase. The depth first decrease then increase as the hyperscaling violation exponent θ increase. So the other important progress of this paper is that the superconducting dome-shaped region built in [7] can be expanded by hyperscaling violation exponent θ if z is large enough.

However, there are still some issues concerning the instability of our holographic setup in our calculations. First, gauge field B_μ is used to mimic the charge change due to doping. The rising part of the superconducting dome is attributed to the increasing of charge. So what about the descend part of the superconducting dome? Unfortunately, it seems hard to find out the reason of the descend part, since the Lagrangian indicates there is a symmetry between A_μ and B_μ . Second, our numerical results about the superconducting dome shown in Fig. 4 imply that $y = \frac{1+x}{x}$ can approach to 0.85, namely $\rho_B \gg \rho_A$. It suggests doping charge is dominant in this case. We leave these issues to a future study.

There are much more generalized form of hyperscaling violating solutions for black holes with multiple fields in [29]. It would be interesting to study the consequence of the such solution along the line we studied here.

Acknowledgements We would like to thank Xian-Hui Ge, Blaise Goutraux and Li Li for valuable comments and discussions. Wenhe Cai is also supported by NSFC China (Grant no. 11805117 and no. 11875184). This work is supported by Mid-career Researcher Program through the National Research Foundation of Korea Grant no. NRF-2016R1A2B3007687.

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: There are no external data associated with the manuscript.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.
Funded by SCOAP³.

Fig. 9 The dome-shaped region of the superconducting instability. We use two methods to obtain the critical temperature T_c as a function of doping parameter x . Here we fix the parameters as $M^2 = -5/4$, $m = 1$, $a = -10$, $b = -4/3$, $c = 14/3$, $n = 1$, $\rho_A = 1$, $\alpha = 0$, $\rho_B = x$ and $(0, 1)$

Appendix A: Methods to obtain the superconducting dome

In this appendix, we verify two methods to obtain the superconducting dome are equivalent. There are two methods to determine the superconducting dome. One is to keep the doping parameter x fixed. Then we obtain the maximal temperature indicated by $\bullet$. The other method is to keep the horizon u_h fixed. Then we obtain the maximal x indicated by $\blacklozenge$ and minimal x indicated by $\triangle$. There is one-to-one match between the critical temperature T and doping x .

Figure 9 shows that these two methods are equivalent. The endpoints of the superconducting region are $x_1 \approx 1.17$ and $x_2 \approx 7$. The result indicates that temperature gradually approaches to zero when x increases. If Z_A, Z_B, Z_{AB} are independent of χ , the dome vanish and the critical temperature always increases. In order to simplify the calculations, we choose the former method in this paper.

Appendix B: Role of a, b, α in the superconducting dome

In this appendix, we will show the role of a, b, α in Figs. 10, 11, 12 and 13. Most of the calculations are unchanged as we have calculated in Sect. 2. First, we figure out the role of a . The left endpoint x_1 moves right and the right endpoint x_2 moves left when a increases. The optimal temperature T_O increases as a decreases. The superconducting dome expands when a decreases.

Fig. 10 Left side: the dome-shaped superconducting phase with the coefficient a appearing in the expansion (6). Dome size increases for large Z_A . Right side: the left endpoint x_1 and the right endpoint x_2

of the superconducting region. We fix $b = 4/3$, $c = 14/3$, $\alpha = 0$ and choose different a . The dashed line represents $(1, 0)$, and the solid line represents $(0, 1)$. We find $x_1(a)$ is increasing function

Fig. 11 **a** The role of b . The dome-shaped superconducting phase with the coefficient b . Dome size increases for large Z_B . **b, c** The left endpoint x_1 and the right endpoint x_2 . We fix $a = 10$, $c = 14/3$, $\alpha = 0$. The dashed line represents $(1, 0)$, and the solid line represents $(0, 1)$

Fig. 12 The boundary of the pseudo-insulating phase x_0 and T_0 with the translational symmetry broken by the neutral scalars

Fig. 13 The roles of α . **a** Changing α doesn't make much difference, but interchanging q_A and q_B makes a big difference. Two curves for $\alpha = 0$ and $\alpha = 0.5$ cross at $\star$. **b, c** The left endpoint x_1 and the right endpoint

x_2 of the superconducting region. We fix $a = 10$, $b = 4/3$, $c = 14/3$ and choose different α . The dashed line represents $(1, 0)$, and the solid line represents $(0, 1)$

Then, we figure out the role of b . x_1 moves right and x_2 moves left to make the dome smaller when b increases. The dependence on b is similar to that on a , because a and b are symmetric in our model. As a result, we could conclude that the superconducting dome is expanding as the couplings $Z_A(\chi) = 1 - \frac{a\chi^2}{2}$, $Z_B(\chi) = 1 - \frac{b\chi^2}{2}$, $Z_{AB}(\chi) = \frac{c\chi^2}{2}$ increase. However, Fig. 13 shows that changing α doesn't make much difference.

Then, we deal with the superconducting dome with the coefficient α . We plot the dependence of the boundary of the pseudo-insulating phase (x_0, T_0) with the translational symmetry broken by the neutral scalars. As illustrated in Fig. 12, the pseudo-insulating phase occurs only if $\alpha \geq \alpha_c$ ($\alpha_c \approx 0.567358$).

Notice that the superconducting phase with $(q_A, q_B) = (1, 0)$ was already studied in [8], and our result in this case is consistent with theirs as one can see in Fig. 13.

References

1. J.M. Maldacena, The large N limit of superconformal field theories and supergravity. *Int. J. Theor. Phys.* **38**, 1113 (1999)
2. S.S. Gubser, I.R. Klebanov, A.M. Polyakov, Gauge theory correlators from noncritical string theory. *Phys. Lett. B* **428**, 105 (1998). [arXiv:hep-th/9802109](https://arxiv.org/abs/hep-th/9802109)
3. E. Witten, Anti-de Sitter space and holography. *Adv. Theor. Math. Phys.* **2**, 253 (1998). [arXiv:hep-th/9802150](https://arxiv.org/abs/hep-th/9802150)
4. S.A. Hartnoll, A. Lucas, S. Sachdev, Holographic quantum matter. [arXiv:1612.07324](https://arxiv.org/abs/1612.07324)
5. J. Zaanen, Y. Liu, Y.-W. Sun, K. Schalm, *Holographic Duality in Condensed Matter Physics* (Cambridge University Press, Cambridge, 2015)
6. A.L. Patrick, N. Naoto, W. Xiao-Gang, Doping a Mott insulator: physics of high-temperature superconductivity. *Rev. Mod. Phys.* **78**, 17 (2006). [arXiv:cond-mat/0410445](https://arxiv.org/abs/cond-mat/0410445)
7. E. Kiritsis, L. Li, Holographic competition of phases and superconductivity. *JHEP* **1601**, 147 (2016). [https://doi.org/10.1007/JHEP01\(2016\)147](https://doi.org/10.1007/JHEP01(2016)147). [arXiv:1510.00020](https://arxiv.org/abs/1510.00020) [cond-mat.str-el]
8. M. Baggioli, M. Goykhman, Under the dome: doped holographic superconductors with broken translational symmetry. *JHEP* **1601**, 011 (2016). [https://doi.org/10.1007/JHEP01\(2016\)011](https://doi.org/10.1007/JHEP01(2016)011). [arXiv:1510.06363](https://arxiv.org/abs/1510.06363) [hep-th]
9. X.H. Ge, Y. Tian, S.Y. Wu, S.F. Wu, Hyperscaling violating black hole solutions and Magneto-thermoelectric DC conductivities in holography. *Phys. Rev. D* **96**(4), 046015 (2017). <https://doi.org/10.1103/PhysRevD.96.046015>. <https://doi.org/10.1103/PhysRevD.97.089901>. [arXiv:1606.05959](https://arxiv.org/abs/1606.05959) [hep-th] (Erratum: [*Phys. Rev. D* **97**(8), 089901 (2018)])
10. M. Baggioli, O. Pujolas, Electron-phonon interactions, metal-insulator transitions, and holographic massive gravity. *Phys. Rev. Lett.* **114**(25), 251602 (2015). <https://doi.org/10.1103/PhysRevLett.114.251602>. [arXiv:1411.1003](https://arxiv.org/abs/1411.1003) [hep-th]
11. Y. Seo, G. Song, P. Kim, S. Sachdev, S.J. Sin, Holography of the Dirac fluid in graphene with two currents. *Phys. Rev. Lett.* **118**(3), 036601 (2017). <https://doi.org/10.1103/PhysRevLett.118.036601>. [arXiv:1609.03582](https://arxiv.org/abs/1609.03582) [hep-th]
12. M. Baggioli, O. Pujolas, On holographic disorder-driven metal-insulator transitions. *JHEP* **01**, 040 (2017). [https://doi.org/10.1007/JHEP01\(2017\)040](https://doi.org/10.1007/JHEP01(2017)040). [arXiv:1601.07897](https://arxiv.org/abs/1601.07897) [hep-th]
13. Y.S. An, T. Ji, L. Li, Magnetotransport and complexity of holographic metal-insulator transitions. [arXiv:2007.13918](https://arxiv.org/abs/2007.13918) [hep-th]
14. M. Blake, D. Tong, Universal resistivity from holographic massive gravity. *Phys. Rev. D* **88**(10), 106004 (2013). <https://doi.org/10.1103/PhysRevD.88.106004>. [arXiv:1308.4970](https://arxiv.org/abs/1308.4970) [hep-th]
15. A. Donos, J.P. Gauntlett, Thermoelectric DC conductivities from black hole horizons. *JHEP* **1411**, 081 (2014). [https://doi.org/10.1007/JHEP11\(2014\)081](https://doi.org/10.1007/JHEP11(2014)081). [arXiv:1406.4742](https://arxiv.org/abs/1406.4742) [hep-th]
16. S. Grozdanov, A. Lucas, S. Sachdev, K. Schalm, Absence of disorder-driven metal-insulator transitions in simple holographic models. *Phys. Rev. Lett.* **115**(22), 221601 (2015). <https://doi.org/10.1103/PhysRevLett.115.221601>. [arXiv:1507.00003](https://arxiv.org/abs/1507.00003) [hep-th]
17. F. Denef, S.A. Hartnoll, Landscape of superconducting membranes. *Phys. Rev. D* **79**, 126008 (2009). <https://doi.org/10.1103/PhysRevD.79.126008>. [arXiv:0901.1160](https://arxiv.org/abs/0901.1160) [hep-th]
18. S.J. Zhang, Q. Pan, E. Abdalla, Holographic superconductor in hyperscaling violation geometry with Maxwell-Dilaton coupling. [arXiv:1511.01841](https://arxiv.org/abs/1511.01841) [hep-th]
19. Q. Pan, S.J. Zhang, Revisiting holographic superconductors with hyperscaling violation. *Eur. Phys. J. C* **76**(3), 126 (2016). <https://doi.org/10.1140/epjc/s10052-016-3980-5>. [arXiv:1510.09199](https://arxiv.org/abs/1510.09199) [hep-th]
20. Z. Fan, Holographic superconductors with hyperscaling violation. *JHEP* **1309**, 048 (2013). [https://doi.org/10.1007/JHEP09\(2013\)048](https://doi.org/10.1007/JHEP09(2013)048). [arXiv:1305.2000](https://arxiv.org/abs/1305.2000) [hep-th]
21. M. Baggioli, M. Goykhman, Phases of holographic superconductors with broken translational symmetry. *JHEP* **1507**, 035 (2015). [https://doi.org/10.1007/JHEP07\(2015\)035](https://doi.org/10.1007/JHEP07(2015)035). [arXiv:1504.05561](https://arxiv.org/abs/1504.05561) [hep-th]
22. S.A. Hartnoll, C.P. Herzog, G.T. Horowitz, Building a holographic superconductor. *Phys. Rev. Lett.* **101**, 031601 (2008). <https://doi.org/10.1103/PhysRevLett.101.031601>. [arXiv:0803.3295](https://arxiv.org/abs/0803.3295) [hep-th]
23. S.A. Hartnoll, C.P. Herzog, G.T. Horowitz, Holographic superconductors. *JHEP* **0812**, 015 (2008). <https://doi.org/10.1088/1126-6708/2008/12/015>. [arXiv:0810.1563](https://arxiv.org/abs/0810.1563) [hep-th]
24. G. Siopsis, J. Therrien, Analytic calculation of properties of holographic superconductors. *JHEP* **05**, 013 (2010). [https://doi.org/10.1007/JHEP05\(2010\)013](https://doi.org/10.1007/JHEP05(2010)013). [arXiv:1003.4275](https://arxiv.org/abs/1003.4275) [hep-th]
25. J.W. Lu, Y.B. Wu, P. Qian, Y.Y. Zhao, X. Zhang, Lifshitz scaling effects on holographic superconductors. *Nucl. Phys. B* **887**, 112–135 (2014). <https://doi.org/10.1016/j.nuclphysb.2014.08.001>. [arXiv:1311.2699](https://arxiv.org/abs/1311.2699) [hep-th]
26. C.J. Luo, X.M. Kuang, F.W. Shu, Lifshitz holographic superconductor in Hořava-Lifshitz gravity. *Phys. Lett. B* **759**, 184–190 (2016). <https://doi.org/10.1016/j.physletb.2016.05.076>. [arXiv:1605.03260](https://arxiv.org/abs/1605.03260) [hep-th]
27. L.Q. Fang, X.H. Ge, X.M. Kuang, Holographic fermions in charged Lifshitz theory. *Phys. Rev. D* **86**, 105037 (2012). <https://doi.org/10.1103/PhysRevD.86.105037>. [arXiv:1201.3832](https://arxiv.org/abs/1201.3832) [hep-th]
28. J. Alsop, E. Papantonopoulos, G. Siopsis, K. Yeter, Duality between zeroes and poles in holographic systems with massless fermions and a dipole coupling. *Phys. Rev. D* **90**(12), 126013 (2014). <https://doi.org/10.1103/PhysRevD.90.126013>. [arXiv:1404.4010](https://arxiv.org/abs/1404.4010) [hep-th]
29. L. Li, Hyperscaling violating solutions in generalised EMD theory. *Phys. Lett. B* **767**, 278–284 (2017). <https://doi.org/10.1016/j.physletb.2017.02.004>. [arXiv:1608.03247](https://arxiv.org/abs/1608.03247) [hep-th]