



Erratum to: Sensitivity of multi-PMT optical modules in Antarctic ice to supernova neutrinos of MeV energy

C. J. Lozano Mariscal^{1,a}, L. Classen¹, M. A. Unland Elorrieta¹, A. Kappes¹

¹ Institut für Kernphysik, Westfälische Wilhelms-Universität Münster, Wilhelm-Klemm-Straße 9, 48149 Münster, Germany

Received: 2 May 2022 / Accepted: 11 July 2022
© The Author(s) 2022

Erratum to: Eur. Phys. J. C (2021) 81:1058

<https://doi.org/10.1140/epjc/s10052-021-09809-y>

We found an error in the code for calculating the CCSN detection range that led to double counting of signal events and thus to ranges that were too large by a factor of about 1.3. This results in the following corrections (old value → new value):

- The values for the detection ranges decrease as shown in the corrected Table 4. Figures 8, 9, 10 change accordingly. Correspondingly, the numbers change in:
 - Abstract: We find that exploiting temporal coincidences between signals in different photocathode segments, a $27 M_{\odot}$ progenitor mass CCSN can be detected up to a distance of (341 kpc → 269 kpc) with a false detection rate of 0.01 year^{-1} with a detector consisting of 10,000 sensors.
 - Section 4.2: The trigger condition ($m \geq 7, N_{\nu} \geq 7$) can be used to send supernova alerts with very high confidence (about one false detection per century), and identify CCSN at a distance of (341 kpc → 269 kpc) with 50% probability.
 - Section 4.2: With a relaxed set of conditions of ($m \geq 7, N_{\nu} \geq 6$), SNe up to (370 kpc → 291 kpc) can be detected with less than one false CCSN detection per year.
 - Section 4.2: For example, for a number of detected events $N_{\nu} = 5$ a background origin can be excluded at 3.2σ , while at least a corresponding number of events will be detected in 50% of cases from a $27 M_{\odot}$ CCSNe at a distance of (407 kpc → 322 kpc).

- Section 4.2: If $N_{\nu} = 7$ events with $m \geq 7$ are detected we obtain a 4.9σ confidence that such signal was not produced by background with a 50% detection probability at (341 kpc → 269 kpc) distance.
- Conclusions: For a detector equipped with 10,000 sensors consisting of 24 3-inch photomultipliers, we find that CCSNe up to a distance of (341 kpc → 269 kpc) can be identified with 50% probability with 0.01 false SN detection per year.
- Conclusions: If the arrival time of CCSN neutrinos is known from an independent observation with $\delta t = 1 \text{ h}$, a $27 M_{\odot}$ CCSN at ([407, 341] kpc → [322, 269] kpc) can be detected in 50% of cases and with a $[3.2, 4.9] \sigma$ certainty that the signal was not produced by background.
- Change in the 5σ detection horizons, in case the arrival time of the burst is known exactly:
 - Section 4.2: The 5σ discovery horizon in this scenario reaches (400 kpc → 315 kpc) for a $27 M_{\odot}$ CCSN using $m \geq 7$, and (300 kpc → 234 kpc) for the $9.6 M_{\odot}$ model.
- To reach a detection of one CCSN about every decade doubling the number of modules, the necessary noise reduction changes from a factor ~ 70 to a factor ~ 140 :
 - Abstract: Increasing the number of sensors to 20,000 and reducing the optical background by a factor of ($\sim 70 \rightarrow \sim 140$) expands the range such that a CCSN detection rate of (0.1 → 0.08) per year is achieved, while keeping the false detection rate at 0.01 year^{-1} .
 - Section 4.2: In contrast, doubling the number of modules installed would allow the false SN detection rate to be kept below 0.01 year^{-1} while expecting (1 → 0.8) CCSN detection per decade if the radioactive noise within the glass vessel can be reduced by a factor of about (70 → 140).

The original article can be found online at <https://doi.org/10.1140/epjc/s10052-021-09809-y>.

^ae-mail: c.lozano@wwu.de (corresponding author)

Table 4 False CCSN detection rate and range of supernova detection (50% probability) for different values of m and N_ν (see trigger conditions in Section 4; $\Delta t_{\text{coin}} = 20$ ns, $\Delta T_{\text{SN}} = 10$ s)

Trigger		False CCSN rate (year ⁻¹)	Range (kpc) 27 M _⊙ (9.6 M _⊙)
m	N_ν		
≥ 5	≥ 104	0.7	135 (101)
	≥ 107	< 0.01	133 (99)
≥ 6	≥ 17	0.9	245 (182)
	≥ 20	< 0.01	225 (167)
≥ 7	≥ 6	0.4	291 (216)
	≥ 7	0.01	269 (200)
≥ 8	≥ 3	0.2	275 (204)
	≥ 4	< 0.01	235 (174)




Fig. 8 Probability for the detection of a CCSN of 27 M_⊙ progenitor mass as a function of distance using the trigger conditions presented in Table 4

- Conclusions: Increasing the number of installed modules to 20,000 and using pressure vessels with significantly reduced optical background could extend the range such that one CCSN (per decade → every ~ 12 years) can be observed.

The conclusion from this work remains unchanged despite the reduced detection range: exploiting coincidences between detected photons within a segmented photosensor will significantly increase the sensitivity of sparsely instrumented neutrino telescopes to distant CCSNe.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article




Fig. 9 Detection prospects for a CCSN whose time is known to within 1 h ($m \geq 7$, $\Delta t_{\text{coin}} = 20$ ns, $\Delta T_{\text{SN}} = 10$ s). Left axis: Probability in σ that the signal is not produced by background fluctuations. Right axis: Distance at which a 27 M_⊙ progenitor mass CCSN is detected with 10% (upper boundary), 50% (middle mark) and 90% probability (lower boundary), when at least N_ν detected events are required




Fig. 10 CCSN detection rate for hypothetical detectors with 10,000 (upper) and 20,000 (lower) mDOMs as a function of the false SN detection rate and a reduction in radioactive noise compared to standard mDOMs. The CCSN detection rates have been calculated using the estimated CCSNe population from [16] based on actual observations and scaled to the star formation rate

are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP³. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.