



# Electromagnetic properties of the $\Sigma_c(2800)^+$ and $\Lambda_c(2940)^+$ states via light-cone QCD

Ulaş Ozdem<sup>a</sup>

Health Services Vocational School of Higher Education, Istanbul Aydın University, Sefakoy-Kucukcekmece, 34295 Istanbul, Türkiye

Received: 11 September 2023 / Accepted: 10 November 2023 / Published online: 24 November 2023  
© The Author(s) 2023

**Abstract** The  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states are among the most interesting and intriguing particles whose internal structures have not yet been elucidated. Inspired by this, the magnetic dipole moments of the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states with quantum numbers  $J^P = \frac{1}{2}^-$  and  $J^P = \frac{3}{2}^-$ , respectively, are analyzed in the framework of QCD light-cone sum rules, assuming that they have a molecule composed of a nucleon and a  $D^{(*)}$  meson. The magnetic dipole moments are obtained as  $\mu_{\Sigma_c^+} = 0.26 \pm 0.05 \mu_N$  and  $\mu_{\Lambda_c^+} = -0.31 \pm 0.04 \mu_N$ . The magnetic dipole moment is the leading-order response of a bound system to a weak external magnetic field. It therefore offers an excellent platform to explore the inner organization of hadrons governed by the quark-gluon dynamics of QCD. Comparison of the findings of this analysis with future experimental results on the magnetic dipole moments of the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states may shed light on the nature and internal organization of these states. The electric quadrupole and magnetic octupole moments of the  $\Lambda_c(2940)^+$  states have also been calculated, and these values are determined to be  $\mathcal{Q}_{\Lambda_c^+} = (0.65 \pm 0.25) \times 10^{-3} \text{ fm}^2$  and  $\mathcal{O}_{\Lambda_c^+} = -(0.38 \pm 0.10) \times 10^{-3} \text{ fm}^3$ , respectively. The values of the electric quadrupole and magnetic octupole moments show a non-spherical charge distribution. We hope that our estimates of the electromagnetic properties of the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states, together with the results of other theoretical studies of the spectroscopic parameters of these states, will be useful for their search in future experiments and will help us to define the exact internal structures of these states.

## 1 Introduction

In the past few years, many excited baryon states have been discovered experimentally. Elucidating and understanding the internal structure of excited hadronic states is an important problem in the field of nonperturbative QCD. The discovery of a large number of new heavy baryons at experimental facilities requires their theoretical identification and classification [1–4]. Theoretical investigations include the analysis of baryons containing a single heavy quark, which provides an excellent laboratory to study the dynamics of a light diquark in the environment of a heavy quark, allowing the predictions of different theoretical approaches to be tested and to improve the understanding of the non-perturbative nature of QCD. Significant experimental progress has been made in the field of singly heavy baryons in recent years [5–34]. More experimental research is needed because the evidence for the existence of some of these states is weak and the quantum numbers are not well determined. Therefore, their study is an active field in both experimental and theoretical research.

In 2005, the Belle Collaboration tentatively identified the  $\Sigma_c(2800)^{++}$ ,  $\Sigma_c(2800)^+$  and  $\Sigma_c(2800)^0$  as isospin triplet states in the  $\Lambda_c^+\pi^+$ ,  $\Lambda_c^+\pi^0$  and  $\Lambda_c^+\pi^-$  mass spectrum, respectively [15]. The measured mass differences and decay widths are listed as follows,

$$M_{\Sigma_c(2800)^{++}} - M_{\Lambda_c^+} = 514.5_{-3.1}^{+3.4+2.8}_{-4.9} \text{ MeV},$$
$$\Gamma_{\Sigma_c(2800)^{++}} = 75_{-13}^{+18+12}_{-11} \text{ MeV}, \quad (1)$$

$$M_{\Sigma_c(2800)^+} - M_{\Lambda_c^+} = 505.4_{-4.6}^{+5.8+12.4}_{-2.0} \text{ MeV},$$
$$\Gamma_{\Sigma_c(2800)^+} = 62_{-23}^{+37+52}_{-38} \text{ MeV}, \quad (2)$$

$$M_{\Sigma_c(2800)^0} - M_{\Lambda_c^+} = 515.4_{-3.1}^{+3.2+2.1}_{-6.0} \text{ MeV},$$
$$\Gamma_{\Sigma_c(2800)^0} = 61_{-13}^{+18+22}_{-13} \text{ MeV}. \quad (3)$$

<sup>a</sup>e-mail: [ulasozdem@aydin.edu.tr](mailto:ulasozdem@aydin.edu.tr) (corresponding author)

The  $\Sigma_c(2800)^0$  state is probably confirmed by the BaBar Collaboration [35] and; the mass and decay width are determined to be  $M_{\Sigma_c(2800)^0} = 2846 \pm 8 \pm 10$  MeV,  $\Gamma_{\Sigma_c(2800)^0} = 86^{+33}_{-22} \pm 7$  MeV. However, the measured width of this state is consistent with the Belle data, the mass value is about 50 MeV larger and somewhat inconsistent with the previous measurement. If the particle observed by the two experimental collaborations is indeed in the same state, then the discrepancy in the measured masses must be resolved. The quantum numbers of these states also remain to be determined.

In 2007, the BaBar Collaboration searched for charmed baryons in the  $D^0 p$  invariant mass spectrum and found the  $\Lambda_c(2940)^+$  [10]. The Belle Collaboration then confirmed the  $\Lambda_c(2940)^+$  in decay mode  $\Lambda_c(2940) \rightarrow \Sigma_c(2455)\pi$  [11]. In 2017, the LHCb collaboration measured the amplitude analysis of the  $\Lambda_b^0 \rightarrow D^0 p \pi^-$  process to determine the spin-parity of the  $\Lambda(2940)^+$  state, and the most likely spin-parity assignment for this state is  $J^P = \frac{3}{2}^-$  [12]. The masses and decay widths measured for this state by the three collaborations are given below,

$$M_{\Lambda_c(2940)^+} = 2939.8 \pm 1.3 \pm 1.0 \text{ MeV},$$

$$\Gamma_{\Lambda_c(2940)} = 17.5 \pm 5.0 \pm 5.9 \text{ MeV (BaBar)}, \quad (4)$$

$$M_{\Lambda_c(2940)^+} = 2938.0 \pm 1.3^{+2.0}_{-4.0} \text{ MeV},$$

$$\Gamma_{\Lambda_c(2940)} = 13^{+8+27}_{-5-7} \text{ MeV (Belle)}, \quad (5)$$

$$M_{\Lambda_c(2940)^+} = 2944.8^{+3.5}_{-2.5} \pm 0.4^{+0.1}_{-4.6} \text{ MeV},$$

$$\Gamma_{\Lambda_c(2940)} = 27.7^{+8.2}_{-6.0} \pm 0.9^{+5.2}_{-10.4} \text{ MeV (LHCb)}. \quad (6)$$

Following these experimental observations, understanding the internal structures and quantum numbers of the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states has been an important topic of research among theorists. To elucidate their underlying structure, they were studied in standard baryon [36–44] and molecular pentaquark states [45–63]. The  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states are still not yet fully understood, despite the considerable efforts that have been devoted to their study. All of these works, done with alternative structures however with predictions that are within error of the experimental observations, indicate the need for more for further study of the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states. Therefore, it is time for further efforts in the study of these states to have a better understanding of their properties.

If one examines the studies in the literature listed above, it can be seen that almost all the calculations are aimed at calculating the spectroscopic parameters of these states, and it is easy to see that the spectroscopic parameters alone are not sufficient to elucidate the controversial nature of these states. Hence, it is obvious that further analyses such as the electromagnetic form factors, the weak decays, and so on are needed to shed light on the internal structure of these states. The  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states are close to

the  $ND$  and  $ND^*$  thresholds, inspiring us that they might be the  $ND$  and  $ND^*$  molecular states. Then it is worthwhile to study the  $ND$  and  $ND^*$  interactions with various methods to further understand the nature of the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states. Based on this assumption, we consider the states  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  to be  $ND$  and  $ND^*$  bound states:  $\Sigma_c^+ = (|D^0 p\rangle + |D^+ n\rangle)/\sqrt{2}$ , and  $\Lambda_c^+ = (|D^{*0} p\rangle - |D^{*+} n\rangle)/\sqrt{2}$ ; and we explore the nature of these states in the framework of QCD light-cone sum rules. It is worth noting that, as mentioned above, the spin-parity quantum numbers of these states have not been experimentally determined. However, when considering the molecular state, studies in the literature have shown that the  $J^P = \frac{1}{2}^-$  assignment for the  $\Sigma_c(2800)^+$  state and the  $J^P = \frac{3}{2}^-$  assignment for the  $\Lambda_c(2940)^+$  state are more consistent with the experimental data. Therefore, in this study we assume that the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states have spin-parity quantum numbers  $J^P = \frac{1}{2}^-$  and  $J^P = \frac{3}{2}^-$ , respectively.

In the present paper, we proceed as follows. We derive QCD light-cone sum rules for molecular states in Sect. 2, with similar procedures as our previous studies on baryons [64–66] and molecular states [67–72]. In Sect. 3, we illustrate our numerical results and discussions, and the last section is devoted to the summary and concluding remarks. The analytical expressions obtained for the magnetic dipole moment results of the  $\Sigma_c^+$  state, electromagnetic multipole moments of the  $\Lambda_c(2940)^+$  state, and the photon DAs are presented in Appendix A B and C respectively.

## 2 QCD light-cone sum rules for the electromagnetic properties of the $\Sigma_c(2800)^+$ and $\Lambda_c(2940)^+$ states

As is well known, the QCD light-cone sum rules have been successfully applied to the calculation of masses, decay constants, form factors, magnetic dipole moments, etc. of standard hadrons, and are a powerful technique for the study of exotic hadron properties. The correlation function is evaluated both in terms of hadrons (the hadronic side) and in terms of quark-gluon degrees of freedom (the QCD side) according to the QCD light cone sum rules technique. Then, by equating these two different descriptions of the correlation function by quark-hadron duality, the physical quantities, i.e. the magnetic dipole moments, are calculated [73–75].

### 2.1 Magnetic dipole moments of the $\Sigma_c(2800)^+$ state

The first step to analyze magnetic dipole moments with the QCD light-cone sum rules is to write the following correlation function,

$$\Pi(p, q) = i \int d^4x e^{ip \cdot x} \langle 0 | T \{ J^{\Sigma_c^+}(x) \bar{J}^{\Sigma_c^+}(0) \} | 0 \rangle_\gamma, \quad (7)$$

where  $T$  is the time ordered product, sub-indices  $\gamma$  is the external electromagnetic field.  $J(x)$  is the interpolating current of the  $\Sigma_c^+$  state and is required to continue the analysis. This interpolating current is written as follows by the quantum numbers  $I(J^P) = 1(1/2^-)$ :

$$\begin{aligned}
 J^{\Sigma_c^+}(x) &= \frac{1}{\sqrt{2}} \left\{ |D^0 p\rangle + |D^+ n\rangle \right\} \\
 &= \frac{1}{\sqrt{2}} \left\{ [\bar{u}^d(x) i\gamma_5 c^d(x)] [\varepsilon^{abc} u^{aT}(x) C \gamma_\mu d^b(x) \right. \\
 &\quad \times \gamma^\mu \gamma_5 u^c(x)] - [\bar{d}^d(x) i\gamma_5 c^d(x)] [\varepsilon^{abc} u^{aT}(x) \\
 &\quad \times C \gamma_\mu d^b(x) \gamma^\mu \gamma_5 d^c(x)] \left. \right\}, \tag{8}
 \end{aligned}$$

where  $a, b, c$  and  $d$  are color indices and the  $C$  is the charge conjugation operator.

Let us start by obtaining the correlation function from the hadronic side. We insert a full set of intermediate state  $\Sigma_c$  with the same quantum numbers as the interpolating currents into the correlation function to get the hadronic representation of the correlation function. This gives us the following expression

$$\begin{aligned}
 \Pi^{Had}(p, q) &= \frac{\langle 0 | J^{\Sigma_c^+}(x) | \Sigma_c^+(p, s) \rangle}{[p^2 - m_{\Sigma_c^+}^2]} \\
 &\quad \times \langle \Sigma_c^+(p, s) | \Sigma_c^+(p + q, s) \rangle_\gamma \\
 &\quad \times \frac{\langle \Sigma_c^+(p + q, s) | \bar{J}^{\Sigma_c^+}(0) | 0 \rangle}{[(p + q)^2 - m_{\Sigma_c^+}^2]} + \dots \tag{9}
 \end{aligned}$$

The matrix elements in Eq. (9) can be written in terms of hadronic parameters and Lorentz invariant form factors in the following way:

$$\langle 0 | J^{\Sigma_c^+}(x) | \Sigma_c^+(p, s) \rangle = \lambda_{\Sigma_c^+} \gamma_5 u(p, s), \tag{10}$$

$$\langle \Sigma_c^+(p + q, s) | \bar{J}^{\Sigma_c^+}(0) | 0 \rangle = \lambda_{\Sigma_c^+} \gamma_5 \bar{u}(p + q, s), \tag{11}$$

$$\begin{aligned}
 \langle \Sigma_c^+(p, s) | \Sigma_c^+(p + q, s) \rangle_\gamma &= \varepsilon^\mu \bar{u}(p, s) \left[ [F_1(q^2) \right. \\
 &\quad \left. + F_2(q^2)] \gamma_\mu + F_2(q^2) \right. \\
 &\quad \left. \times \frac{(2p + q)_\mu}{2m_{\Sigma_c^+}} \right] u(p + q, s), \tag{12}
 \end{aligned}$$

where the  $u(p, s)$ ,  $u(p + q, s)$  and  $\lambda_{\Sigma_c^+}$  are the spinors and residue of the  $\Sigma_c^+$  state, respectively.

Substituting Eqs. (10)–(12) in Eq. (9) and doing some calculations, we get the following result for the hadronic side,

$$\Pi^{Had}(p, q) = \lambda_{\Sigma_c^+}^2 \gamma_5 \frac{(\not{p} + m_{\Sigma_c^+})}{[p^2 - m_{\Sigma_c^+}^2]} \varepsilon^\mu \left[ [F_1(q^2) + F_2(q^2)] \gamma_\mu \right.$$

$$\left. + F_2(q^2) \frac{(2p + q)_\mu}{2m_{\Sigma_c^+}} \right] \gamma_5 \frac{(\not{p} + \not{q} + m_{\Sigma_c^+})}{[(p + q)^2 - m_{\Sigma_c^+}^2]}. \tag{13}$$

To obtain the above expression, summing over the spins of  $\Sigma_c^+$  state,  $\sum_s u(p, s) \bar{u}(p, s) = \not{p} + m_{\Sigma_c^+}$  and  $\sum_s u(p + q, s) \bar{u}(p + q, s) = (\not{p} + \not{q}) + m_{\Sigma_c^+}$  have also been performed.

The magnetic dipole moments of hadrons are related to their magnetic form factors; more precisely, the magnetic dipole moments are equal to the magnetic form factor at zero momentum squared. The magnetic form factors ( $F_M(q^2)$ ), which are more directly accessible in experiments, are defined by the form factors  $F_1(q^2)$  and  $F_2(q^2)$

$$F_M(q^2) = F_1(q^2) + F_2(q^2). \tag{14}$$

Since we are dealing with a real photon, we can define the magnetic form factor  $F_M(q^2 = 0)$  in terms of the magnetic dipole moment  $\mu_{\Sigma_c^+}$ :

$$\mu_{\Sigma_c^+} = \frac{e}{2m_{\Sigma_c^+}} F_M(q^2 = 0). \tag{15}$$

Now we are ready to start the second step, which is the calculation of the correlation function using the QCD parameters. By explicitly using the interpolating currents in the correlation function, the second representation of the correlation function, the QCD side, is obtained. Then, all of the quark fields are contracted according to Wick’s theorem, and the desired results are obtained. As a result of the procedures described above, we obtain the following:

$$\begin{aligned}
 \Pi_{\Sigma_c^+}^{QCD}(p, q) &= -\frac{i}{2} \varepsilon^{abc} \varepsilon^{a'b'c'} \int d^4x e^{ip \cdot x} \langle 0 | \\
 &\quad \times \left\{ \text{Tr} \left[ \gamma_5 S_c^{dd'}(x) \gamma_5 S_u^{d'd}(-x) \right] \text{Tr} \right. \\
 &\quad \times \left[ \gamma_\alpha S_d^{bb'}(x) \gamma_\beta \tilde{S}_u^{aa'}(x) \right] (\gamma^\alpha \gamma_5 S_c^{cc'}(x) \gamma_5 \gamma^\beta) \\
 &\quad - \text{Tr} \left[ \gamma_5 S_c^{dd'}(x) \gamma_5 S_u^{d'd}(-x) \right] \text{Tr} \left[ \gamma^\alpha \gamma_5 S_u^{ca'}(x) \right. \\
 &\quad \times \gamma_\beta \tilde{S}_d^{bb'}(x) \gamma_\alpha S_u^{ac'}(x) \gamma_5 \gamma^\beta \left. \right] \\
 &\quad + \text{Tr} \left[ \gamma_5 S_c^{dd'}(x) \gamma_5 S_d^{d'd}(-x) \right] \text{Tr} \left[ \gamma_\alpha S_d^{bb'}(x) \right. \\
 &\quad \times \gamma_\beta \tilde{S}_u^{aa'}(x) \left. \right] (\gamma^\alpha \gamma_5 S_d^{cc'}(x) \gamma_5 \gamma^\beta) \\
 &\quad - \text{Tr} \left[ \gamma_5 S_c^{dd'}(x) \gamma_5 S_d^{d'd}(-x) \right] \text{Tr} \left[ \gamma^\alpha \gamma_5 S_d^{cb'}(x) \right. \\
 &\quad \times \gamma_\beta \tilde{S}_u^{aa'}(x) \gamma_\alpha S_d^{bc'}(x) \gamma_5 \gamma^\beta \left. \right\} | 0 \rangle_\gamma, \tag{16}
 \end{aligned}$$

where  $\tilde{S}_{c(q)}^{ij}(x) = C S_{c(q)}^{ijT}(x) C$  and,  $S_c(x)$  and  $S_q(x)$  are the

full propagators of the heavy and light quarks, which can be written as [76,77]

$$S_q(x) = S_q^{free} - \frac{\langle \bar{q}q \rangle}{12} \left( 1 - i \frac{m_q \not{x}}{4} \right) - \frac{\langle \bar{q}q \rangle}{192} m_0^2 x^2 \times \left( 1 - i \frac{m_q \not{x}}{6} \right) - \frac{ig_s}{32\pi^2 x^2} G^{\mu\nu}(x) \left[ k\sigma_{\mu\nu} + \sigma_{\mu\nu} k \right], \tag{17}$$

$$S_c(x) = S_c^{free} - \frac{g_s m_c}{16\pi^2} \int_0^1 dv G^{\mu\nu}(vx) \left[ (\sigma_{\mu\nu} \not{x} + \not{x} \sigma_{\mu\nu}) \times \frac{K_1(m_c \sqrt{-x^2})}{\sqrt{-x^2}} + 2\sigma_{\mu\nu} K_0(m_c \sqrt{-x^2}) \right], \tag{18}$$

and

$$S_q^{free} = \frac{1}{2\pi x^2} \left( i \frac{\not{x}}{x^2} - \frac{m_q}{2} \right), \tag{19}$$

$$S_c^{free} = \frac{m_c^2}{4\pi^2} \left[ \frac{K_1(m_c \sqrt{-x^2})}{\sqrt{-x^2}} + i \frac{\not{x} K_2(m_c \sqrt{-x^2})}{(\sqrt{-x^2})^2} \right], \tag{20}$$

where  $\langle \bar{q}q \rangle$  is light-quark condensate,  $G^{\mu\nu}$  is the gluon field strength tensor,  $v$  is line variable and  $K_i$ 's are modified Bessel functions of the second kind, respectively.

The correlation functions in Eq. (16) receive both perturbative contributions, that is, when a photon interacts with light/heavy quarks at short-distance and non-perturbative contributions, that is when a photon interacts with light quarks at large-distance. In the case of perturbative contributions, one of the light propagators or one of the heavy propagators of the quarks interacting perturbatively with the photon is replaced by the following

$$S^{free}(x) \rightarrow \int d^4y S^{free}(x-z) A(z) S^{free}(z), \tag{21}$$

and the rest of the propagators in Eq. (16) are considered full quark propagators, which include perturbative and non-perturbative contributions. The overall perturbative contribution is obtained by replacing the perturbatively interacting quark propagator with the photon as described above, and by replacing the surviving propagators with their free parts. Here we use  $A_\mu(z) = \frac{i}{2} z^\nu (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) e^{iq \cdot z}$ .

In the case of non-perturbative contributions, one of the light quark propagators in Eq. (17), which describes the photon emission at large distances, is replaced with

$$S_{\alpha\beta}^{ab}(x) \rightarrow -\frac{1}{4} [\bar{q}^a(x) \Gamma_i q^b(x)] (\Gamma_i)_{\alpha\beta}, \tag{22}$$

and in Eq. (16) the remaining four quark propagators are all considered to be full quark propagators. Here  $\Gamma_i$  represents the full set of Dirac matrices. Once Eq. (22) is inserted into

Eq. (16), matrix elements of type  $\langle \gamma(q) | \bar{q}(x) \Gamma_i q(0) | 0 \rangle$  and  $\langle \gamma(q) | \bar{q}(x) \Gamma_i G_{\alpha\beta} q(0) | 0 \rangle$  appear, which represent the non-perturbative contributions. We require these matrix elements, which are parameterized in terms of photon wave functions with certain twists, to calculate the non-perturbative contributions (see Ref. [78] for the explicit expressions of photon distribution amplitudes (DAs)). The QCD side of the correlation function can be acquired in terms of the quark-gluon parameters and the DAs of the photon with the help of Eqs. (16)–(22) and after Fourier transforms the x-space calculations to momentum space.

Finally, the Lorentz structure  $\not{x} \not{q}$  is chosen from both the hadronic and the QCD sides, and the coefficients of this structure are matched with each other. Then double Borel transformations and a continuum subtraction are performed. These are used to eliminate the effects of the continuum and higher states. This is followed by the required QCD light-cone sum rules for the magnetic dipole moments:

$$\mu_{\Sigma_c^+} = \frac{e \frac{m_{\Sigma_c^+}^2}{M^2}}{\lambda_{\Sigma_c^+}^2 m_{\Sigma_c^+}} \Delta_1^{QCD}(M^2, s_0). \tag{23}$$

The corresponding results for the  $\Delta_1^{QCD}(M^2, s_0)$  function can be found in Appendix. As you can see in the equation above, there are two extra parameters: the continuum threshold  $s_0$  and the Borel parameters  $M^2$ . In the numerical analysis section, we will explain how the working intervals of these extra parameters are determined.

### 2.2 Magnetic dipole, electric quadrupole and magnetic octupole moments of the $\Lambda_c^+$ state

The required correlation function for the electromagnetic multipole moments of the  $\Lambda_c^+$  state is given by

$$\Pi_{\mu\nu}(p, q) = i \int d^4x e^{ip \cdot x} \langle 0 | T \left\{ J_\mu^{\Lambda_c^+}(x) \bar{J}_\nu^{\Lambda_c^+}(0) \right\} | 0 \rangle_\gamma. \tag{24}$$

The interpolating current used for  $\Lambda_c^+$  state with isospin and spin-parity  $I(J^P) = 0(3/2^-)$  is as follows:

$$J_\mu^{\Lambda_c^+}(x) = \frac{1}{\sqrt{2}} \left\{ | D^{*0} p \rangle - | D^{*+} n \rangle \right\} = \frac{1}{\sqrt{2}} \left\{ [\bar{u}^d(x) \gamma_\mu c^d(x)] [\varepsilon^{abc} u^{aT}(x) C \gamma_\alpha d^b(x) \gamma^\alpha \times \gamma_5 u^c(x)] + [\bar{d}^d(x) \gamma_\mu c^d(x)] [\varepsilon^{abc} u^{aT}(x) C \gamma_\alpha \times d^b(x) \gamma^\alpha \gamma_5 d^c(x)] \right\}. \tag{25}$$

The correlation function obtained in terms of hadronic parameters is written as

$$\begin{aligned} \Pi_{\mu\nu}^{Had}(p, q) = & \frac{\langle 0 | J_{\mu}^{\Lambda_c^+}(x) | \Lambda_c^+(p, s) \rangle}{[p^2 - m_{\Lambda_c^+}^2]} \\ & \times \langle \Lambda_c^+(p, s) | \Lambda_c^+(p + q, s) \rangle_{\gamma} \\ & \times \frac{\langle \Lambda_c^+(p + q, s) | \bar{J}_{\nu}^{\Lambda_c^+}(0) | 0 \rangle}{[(p + q)^2 - m_{\Lambda_c^+}^2]} + \dots \end{aligned} \quad (26)$$

The matrix elements  $\langle 0 | J_{\mu}^{\Lambda_c^+}(x) | \Lambda_c^+(p, s) \rangle$ ,  $\langle \Lambda_c^+(p + q, s) | \bar{J}_{\nu}^{\Lambda_c^+}(0) | 0 \rangle$  and  $\langle \Lambda_c^+(p) | \Lambda_c^+(p + q) \rangle_{\gamma}$  are expressed as follows,

$$\langle 0 | J_{\mu}^{\Lambda_c^+}(x) | \Lambda_c^+(p, s) \rangle = \lambda_{\Lambda_c^+} u_{\mu}(p, s), \quad (27)$$

$$\langle \Lambda_c^+(p + q, s) | \bar{J}_{\nu}^{\Lambda_c^+}(0) | 0 \rangle = \lambda_{\Lambda_c^+} \bar{u}_{\nu}(p + q, s), \quad (28)$$

$$\begin{aligned} \langle \Lambda_c^+(p, s) | \Lambda_c^+(p + q, s) \rangle_{\gamma} = & -e \bar{u}_{\mu}(p, s) \left[ F_1(q^2) g_{\mu\nu} \not{\epsilon} \right. \\ & - \frac{1}{2m_{\Lambda_c^+}} \left[ F_2(q^2) g_{\mu\nu} \right. \\ & + F_4(q^2) \frac{q_{\mu} q_{\nu}}{(2m_{\Lambda_c^+})^2} \left. \right] \not{\epsilon} \not{q} \\ & \left. + F_3(q^2) \frac{1}{(2m_{\Lambda_c^+})^2} q_{\mu} q_{\nu} \not{\epsilon} \right] \\ & \times u_{\nu}(p + q, s), \end{aligned} \quad (29)$$

where  $F_i$ 's are the Lorentz invariant form factors and; the  $u_{\mu}(p, s)$ ,  $u_{\nu}(p + q, s)$  and  $\lambda_{\Lambda_c^+}$  are the spinors and residue of the  $\Lambda_c^+$  state, respectively.

In principle, we can use Eqs. (26)–(29) to get the final expression of the hadronic side of the correlation function, but we run into two problems: not all Lorentz structures are independent, and the correlation function also receives contributions from spin-1/2 particles, which must be eliminated for the calculations to be reliable. The matrix element of the current  $J_{\mu}$  between spin-1/2 pentaquarks and vacuum is nonzero and is determined as

$$\begin{aligned} \langle 0 | J_{\mu}(0) | B(p, s = 1/2) \rangle \\ = (A p_{\mu} + B \gamma_{\mu}) u(p, s = 1/2). \end{aligned} \quad (30)$$

As is seen the unwanted spin-1/2 contributions are proportional to  $\gamma_{\mu}$  and  $p_{\mu}$ . By multiplying both sides with  $\gamma^{\mu}$  and using the condition  $\gamma^{\mu} J_{\mu} = 0$  one can determine the constant A in terms of B. To obtain only independent structures and to eliminate the spin-1/2 contaminations in the correlation function, we apply the ordering for Dirac matrices as  $\gamma_{\mu} \not{\epsilon} \not{q} \gamma_{\nu}$  and eliminate terms with  $\gamma_{\mu}$  at the beginning,  $\gamma_{\nu}$  at the end and those proportional to  $p_{\mu}$  and  $p_{\nu}$  [79]. Applying

the above-mentioned manipulations, the correlation function becomes the following

$$\begin{aligned} \Pi_{\mu\nu}^{Had}(p, q) = & \frac{\lambda_{\Lambda_c^+}^2}{[(p + q)^2 - m_{\Lambda_c^+}^2][p^2 - m_{\Lambda_c^+}^2]} \\ & \times \left[ g_{\mu\nu} \not{\epsilon} \not{q} F_1(q^2) - m_{\Lambda_c^+} g_{\mu\nu} \not{\epsilon} \not{q} F_2(q^2) \right. \\ & - \frac{F_3(q^2)}{4m_{\Lambda_c^+}^2} q_{\mu} q_{\nu} \not{\epsilon} \not{q} - \frac{F_4(q^2)}{4m_{\Lambda_c^+}^3} (\epsilon \cdot p) q_{\mu} q_{\nu} \not{\epsilon} \not{q} \\ & \left. + \dots \right]. \end{aligned} \quad (31)$$

Here, summation over spins of  $\Lambda_c$  state is applied as:

$$\begin{aligned} \sum_s u_{\mu}(p, s) \bar{u}_{\nu}(p, s) = & -(\not{p} + m_{\Lambda_c^+}) \left[ g_{\mu\nu} - \frac{1}{3} \gamma_{\mu} \gamma_{\nu} \right. \\ & \left. - \frac{2 p_{\mu} p_{\nu}}{3 m_{\Lambda_c^+}^2} + \frac{p_{\mu} \gamma_{\nu} - p_{\nu} \gamma_{\mu}}{3 m_{\Lambda_c^+}} \right]. \end{aligned} \quad (32)$$

The final expression obtained for the hadronic representation of the correlation function together with the chosen Lorentz structures is written as follows:

$$\begin{aligned} \Pi_{\mu\nu}^{Had}(p, q) = & \Pi_1^{Had} g_{\mu\nu} \not{\epsilon} \not{q} + \Pi_2^{Had} g_{\mu\nu} \not{\epsilon} \not{q} \\ & + \Pi_3^{Had} q_{\mu} q_{\nu} \not{\epsilon} \not{q} \\ & + \Pi_4^{Had} (\epsilon \cdot p) q_{\mu} q_{\nu} \not{\epsilon} \not{q} + \dots, \end{aligned} \quad (33)$$

where  $\Pi_1^{Had}$ ,  $\Pi_2^{Had}$ ,  $\Pi_3^{Had}$  and  $\Pi_4^{Had}$  are functions of the form factors  $F_1(q^2)$ ,  $F_2(q^2)$ ,  $F_3(q^2)$  and  $F_4(q^2)$ , respectively; and dots denote other independent structures.

The form factors of the magnetic dipole,  $G_M(q^2)$ , electric quadrupole,  $G_Q(q^2)$ , and magnetic octupole,  $G_O(q^2)$  are expressed concerning the form factors  $F_i(q^2)$  in the following manner [80–83]:

$$\begin{aligned} G_M(q^2) = & \left[ F_1(q^2) + F_2(q^2) \right] \left( 1 + \frac{4}{5} \tau \right) - \frac{2}{5} \left[ F_3(q^2) \right] \\ & + \left[ F_4(q^2) \right] \tau (1 + \tau), \\ G_Q(q^2) = & \left[ F_1(q^2) - \tau F_2(q^2) \right] - \frac{1}{2} \left[ F_3(q^2) - \tau F_4(q^2) \right] \\ & \times (1 + \tau), \\ G_O(q^2) = & \left[ F_1(q^2) + F_2(q^2) \right] - \frac{1}{2} \left[ F_3(q^2) + F_4(q^2) \right] \\ & \times (1 + \tau), \end{aligned} \quad (34)$$

where  $\tau = -\frac{q^2}{4m_{\Lambda_c^+}^2}$ . In the static limit, the electromagnetic multipole form factors are given by the form factors  $F_i(q^2 = 0)$  as

$$\begin{aligned}
 G_M(q^2 = 0) &= F_1(q^2 = 0) + F_2(q^2 = 0), \\
 G_Q(q^2 = 0) &= F_1(q^2 = 0) - \frac{1}{2}F_3(q^2 = 0), \\
 G_O(q^2 = 0) &= F_1(q^2 = 0) + F_2(q^2 = 0) - \frac{1}{2}[F_3(q^2 = 0) \\
 &\quad + F_4(q^2 = 0)]. \tag{35}
 \end{aligned}$$

The magnetic dipole ( $\mu_{\Lambda_c^+}$ ), electric quadrupole ( $Q_{\Lambda_c^+}$ ), and magnetic octupole ( $O_{\Lambda_c^+}$ ) moments are written as follows:

$$\begin{aligned}
 \mu_{\Lambda_c^+} &= \frac{e}{2m_{\Lambda_c^+}} G_M(q^2 = 0), \\
 Q_{\Lambda_c^+} &= \frac{e}{m_{\Lambda_c^+}^2} G_Q(q^2 = 0), \\
 O_{\Lambda_c^+} &= \frac{e}{2m_{\Lambda_c^+}^3} G_O(q^2 = 0). \tag{36}
 \end{aligned}$$

The hadronic side of the correlation function is complete when the procedures described above have been applied to it. The next step is to determine the QCD side of the correlation function in terms of the QCD parameters, such as the quark-gluon parameters and the photon DAs. The following result is obtained by repeating the steps of the previous subsection:

$$\begin{aligned}
 \Pi_{\mu\nu}^{QCD}(p, q) &= -\frac{i}{2}\varepsilon^{abc}\varepsilon^{a'b'c'} \int d^4x e^{ip\cdot x} \langle 0 | \\
 &\quad \times \left\{ \text{Tr} \left[ \gamma_\mu S_c^{dd'}(x) \gamma_\nu S_u^{d'd}(-x) \right] \text{Tr} \right. \\
 &\quad \times \left[ \gamma_\alpha S_d^{bb'}(x) \gamma_\beta \tilde{S}_u^{aa'}(x) \right] (\gamma^\alpha \gamma_5 S_u^{cc'}(x) \gamma_5 \gamma^\beta) \\
 &\quad - \text{Tr} \left[ \gamma_\mu S_c^{dd'}(x) \gamma_\nu S_u^{d'd}(-x) \right] \text{Tr} \left[ \gamma^\alpha \gamma_5 S_u^{ca'}(x) \right. \\
 &\quad \times \left. \gamma_\beta \tilde{S}_d^{bb'}(x) \gamma_\alpha S_u^{ac'}(x) \gamma_5 \gamma^\beta \right] \\
 &\quad + \text{Tr} \left[ \gamma_\mu S_c^{dd'}(x) \gamma_\nu S_d^{d'd}(-x) \right] \text{Tr} \left[ \gamma_\alpha S_d^{bb'}(x) \right. \\
 &\quad \times \left. \gamma_\beta \tilde{S}_u^{aa'}(x) \right] (\gamma^\alpha \gamma_5 S_d^{cc'}(x) \gamma_5 \gamma^\beta) \\
 &\quad - \text{Tr} \left[ \gamma_\mu S_c^{dd'}(x) \gamma_\nu S_d^{d'd}(-x) \right] \text{Tr} \left[ \gamma^\alpha \gamma_5 S_d^{cb'}(x) \right. \\
 &\quad \times \left. \gamma_\beta \tilde{S}_u^{aa'}(x) \gamma_\alpha S_d^{bc'}(x) \gamma_5 \gamma^\beta \right] \left. \right\} | 0 \rangle_\gamma. \tag{37}
 \end{aligned}$$

The final expression obtained for the QCD representation of the correlation function together with the chosen Lorentz structures is written as follows

$$\begin{aligned}
 \Pi_{\mu\nu}^{QCD}(p, q) &= \Pi_1^{QCD} g_{\mu\nu} \not{p} \not{q} \\
 &\quad + \Pi_2^{QCD} g_{\mu\nu} \not{q} + \Pi_3^{QCD} q_\mu q_\nu \not{q} \\
 &\quad + \Pi_4^{QCD} (\varepsilon.p) q_\mu q_\nu \not{p} \not{q} + \dots, \tag{38}
 \end{aligned}$$

where  $\Pi_1^{QCD}$ ,  $\Pi_2^{QCD}$ ,  $\Pi_3^{QCD}$  and  $\Pi_4^{QCD}$  are functions of the QCD degrees of freedom and photon DAs parameters.

Both hadronic and quark-gluon parameters have been used to obtain the correlation function. To analyze the magnetic dipole and higher multipole moments, the QCD

and hadronic representations of the correlation function are matched through the quark-hadron duality ansatz. By matching the coefficients of the structures  $g_{\mu\nu} \not{p} \not{q}$ ,  $g_{\mu\nu} \not{q}$ ,  $q_\mu q_\nu \not{q}$  and  $(\varepsilon.p) q_\mu q_\nu \not{p} \not{q}$ , respectively for the  $F_1$ ,  $F_2$ ,  $F_3$  and  $F_4$  we obtain QCD light-cone sum rules for these four invariant form factors. The explicit forms of the expressions obtained for form factors  $F_1$ ,  $F_2$ ,  $F_3$ , and  $F_4$  are given in Appendix B.

Analytical results are obtained for  $\Sigma_c^+$  and  $\Lambda_c^+$  states. Performing numerical calculations for these states would be the next step.

### 3 Numerical illustrations and discussion of the results

The numerical analysis for the magnetic dipole moments of the  $\Sigma_c^+$  and  $\Lambda_c^+$  states is presented in this section. The QCD parameters that are used in our calculations are as follows:  $m_u = m_d = 0$ ,  $m_c = 1.27 \pm 0.02$  GeV [84],  $m_{\Sigma_c^+} = 2792_{-5}^{+14}$  MeV,  $m_{\Lambda_c^+} = 2939.6_{-1.5}^{+1.3}$  MeV [84],  $\lambda_{\Sigma_c^+} = (7.98_{-1.56}^{+1.98}) \times 10^{-4}$  GeV<sup>6</sup>,  $\lambda_{\Lambda_c^+} = (8.87_{-1.59}^{+1.84}) \times 10^{-4}$  GeV<sup>6</sup> [52],  $\langle \bar{q}q \rangle = (-0.24 \pm 0.01)^3$  GeV<sup>3</sup> [85],  $m_0^2 = 0.8 \pm 0.1$  GeV<sup>2</sup>,  $\langle g_s^2 G^2 \rangle = 0.88$  GeV<sup>4</sup> [86],  $\chi = -2.85 \pm 0.5$  GeV<sup>-2</sup> [87] and  $f_{3\gamma} = -0.0039$  GeV<sup>2</sup> [78].

One of the important input parameters in the numerical analysis of the calculation of the magnetic dipole and higher multipole moments by the QCD light-cone sum rule method are the photon DAs. The photon DAs and the parameters that are used in these DAs are listed in Appendix C.

In addition to these numerical parameters, which we have already mentioned in the previous section, the sum rules also depend on the auxiliary parameters: Borel mass squared parameter  $M^2$  and continuum threshold  $s_0$ . Physical measurable such as magnetic dipole and higher multipole moments should be varied as slightly as possible by these extra parameters. Therefore, we are looking for working intervals for these additional parameters such that the results for the magnetic dipole moments are almost independent of these parameters. The standard prescription of the technique used, the Operator Product Expansion (OPE) convergence, and the Pole Contribution (PC) dominance are taken into account in the determination of the working ranges for the parameters  $M^2$  and  $s_0$ . For the characterization of the above restrictions, it is convenient to use the equations below:

$$\text{PC} = \frac{\Delta(M^2, s_0)}{\Delta(M^2, \infty)} \geq 30\%, \tag{39}$$

and

$$\text{OPE} = \frac{\Delta \text{DimN}(M^2, s_0)}{\Delta(M^2, s_0)} \leq 5\%, \tag{40}$$

where  $\Delta \text{DimN}(M^2, s_0) = \Delta \text{Dim}(10+11+12)(M^2, s_0)$ .

Based on these restrictions, the working intervals for the auxiliary parameters are as follows:

$$3.0 \text{ GeV}^2 \leq M^2 \leq 5.0 \text{ GeV}^2,$$

$$11.2 \text{ GeV}^2 \leq s_0 \leq 12.6 \text{ GeV}^2,$$

for the  $\Sigma_c^+$  state, and

$$3.0 \text{ GeV}^2 \leq M^2 \leq 5.0 \text{ GeV}^2,$$

$$12.3 \text{ GeV}^2 \leq s_0 \leq 13.7 \text{ GeV}^2,$$

for the  $\Lambda_c^+$  state.

Our numerical calculations show that the magnetic dipole moments of these states PC vary within the interval  $38\% \leq PC \leq 58\%$  and  $33\% \leq PC \leq 57\%$  for the  $\Sigma_c^+$  and  $\Lambda_c^+$  states, respectively, corresponding to the upper and lower bounds of the Borel mass parameter, by considering these working intervals for the helping parameters. Analyzing the convergence of the OPE, it can be seen that the contribution of the higher twist and higher dimensional terms in the OPE are 3.62 and 3.87% for the  $\Sigma_c^+$  and  $\Lambda_c^+$  states, respectively, of the total and the series shows good convergence. It follows that the constraints imposed by the dominance of PC and the convergence of OPE are satisfied by the working regions determined for  $M^2$  and  $s_0$ . After the determination of the working intervals of  $M^2$  and  $s_0$ , we now study the dependence of the magnetic dipole moments on  $M^2$  for different fixed values of  $s_0$ . From Fig. 1 we can see that the magnetic dipole moments show good stability with respect to the variation of  $M^2$  in its working interval.


To give numerical values for the magnetic dipole moments of the  $\Sigma_c^+$  and  $\Lambda_c^+$  states, we have determined all the necessary parameters. Following our extensive numerical calculations, the magnetic dipole moments, both in their natural unit ( $\frac{e}{2m_B}$ ) and in the nuclear magneton ( $\mu_N$ ), are given as follows:

$$\mu_{\Sigma_c^+} = 0.78 \pm 0.24 \frac{e}{2m_{\Sigma_c^+}} = 0.26 \pm 0.05 \mu_N, \quad (41)$$

$$\mu_{\Lambda_c^+} = -0.97 \pm 0.13 \frac{e}{2m_{\Lambda_c^+}} = -0.31 \pm 0.04 \mu_N. \quad (42)$$

It should be emphasized here that the numerical computations consider uncertainties in the input parameters, ambiguities entering the photon DAs, as well as uncertainties due to variations of the auxiliary parameters  $M^2$  and  $s_0$ .

The magnitude of magnetic dipole moments can provide information about their experimental measurability, and the results obtained for magnetic dipole moments show that they are experimentally accessible. The facilities such as LHCb, Belle II, BESIII and so on may be able to measure magnetic dipole moments of  $\Sigma_c^+$  and  $\Lambda_c^+$  states with the increased



**Fig. 1** The dependencies of magnetic dipole moments of the  $\Sigma_c^+$  and  $\Lambda_c^+$  states; on  $M^2$  at three different values of  $s_0$

luminosity in future runs and our predictions. A comparison of the estimates obtained in this study with the results obtained by different approaches will be a test of the consistency of our results. It is highly recommended to study the magnetic dipole moments of  $\Sigma_c^+$  and  $\Lambda_c^+$  states with Lattice QCD. In Refs. [88,89], the authors have calculated the magnetic dipole moment and the mass of the  $\Lambda(1405)$  baryon by assuming it as a  $\bar{K}N$  molecular configuration via lattice QCD. The magnetic dipole moments of the  $\Sigma_c^+$  and  $\Lambda_c^+$  states can be calculated similarly with the help of lattice QCD. Compared with the predicted magnetic dipole moments of lattice QCD, the results of the present study can provide further information for the experimental search for the  $\Sigma_c^+$  and  $\Lambda_c^+$  states, and the experimental measurements for these magnetic dipole moments could be an important test for the  $DN^{(*)}$  molecular configuration of these states. Our under-

standing of their properties and nature will be enhanced by these efforts.

The electric quadrupole ( $\mathcal{Q}_{\Lambda_c^+}$ ) and the magnetic octupole ( $\mathcal{O}_{\Lambda_c^+}$ ) of the  $\Lambda_c^+$  state are also calculated. The predicted results are as follows

$$\mathcal{Q}_{\Lambda_c^+} = (0.65 \pm 0.25) \times 10^{-3} \text{ fm}^2 \quad (43)$$

$$\mathcal{O}_{\Lambda_c^+} = -(0.38 \pm 0.10) \times 10^{-3} \text{ fm}^3. \quad (44)$$

These results are small but non-zero, implying that the  $\Lambda_c^+$  state charge distribution is non-spherical. Furthermore, the electric quadrupole moment has a positive sign, corresponding to the prolate charge distribution.

#### 4 Summary and concluding remarks

Inspired by the controversial and intriguing nature of  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states, the magnetic dipole moments of these states have been calculated in the framework of the QCD light-cone sum rules, using the photon DAs and the assumption that the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  are hadronic molecular states. The obtained result may be useful in determining the exact nature of these states. To understand the inner structure as well as the geometric shape of the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states, the confirmation of our predictions with different theoretical models and future experiments can be very helpful. The  $\Lambda_c(2940)^+$  state's electric quadrupole and magnetic octupole moments are also extracted. These values indicate that the charge distribution of the  $\Lambda_c(2940)^+$  state is non-spherical. We hope that our estimates of the electromagnetic properties of the  $\Sigma_c(2800)^+$  and  $\Lambda_c(2940)^+$  states, together with the results of other theoretical studies of the spectroscopic parameters of these states, will be useful for their search in future experiments and will help us to define the exact internal structures of these states.

**Data Availability Statement** This manuscript has no associated data or the data will not be deposited. [Authors' comment: All the numerical and mathematical data have been included in the paper and we have no other data regarding this paper.]

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Funded by SCOAP<sup>3</sup>. SCOAP<sup>3</sup> supports the goals of the International Year of Basic Sciences for Sustainable Development.

#### Appendix A Explicit forms of the $\Delta_1^{QCD}(M^2, s_0)$ function

In the present appendix, we present explicit expressions of the analytical expressions obtained for the magnetic dipole moment of the  $\Sigma_c^+$  state as follows:

$$\begin{aligned} \Delta_1^{QCD}(M^2, s_0) = & -\frac{P_1 P_2^2 (e_d + e_u)}{110592 m_c \pi^4} \left[ (m_c^4 I[-2, 0] \right. \\ & - I[0, 0]) I_3[h_\gamma] + (-m_c^4 I[-2, 0] + I[0, 0]) \mathbb{A}[u_0] \\ & + \chi \left( 4m_c^6 I[-3, 1] + m_c^4 (-m_0^2 I[-2, 0] + 2I[-2, 1]) \right. \\ & \left. \left. + m_0^2 I[0, 0] - 2I[0, 1] \right) \varphi_\gamma[u_0] \right] \\ & - \frac{P_1 P_2}{3538944 m_c^2 \pi^6} \left[ 288(e_d + e_u) \left( m_c^8 (I[-4, 2] \right. \right. \\ & - 2I[-3, 1]) + m_c^6 (m_0^2 (2I[-3, 1] + I[-2, 0]) \\ & - 4(I[-3, 2] + I[-2, 1])) + m_c^4 (3I[-2, 2] \\ & + m_0^2 (I[-2, 1] - I[-1, 0]) - 2I[-1, 1]) + m_0^2 I[0, 1]) \\ & + (e_d + e_u) \left( -39m_c^4 (m_c^4 I[-4, 2] - I[-2, 2]) I_2[S] \right. \\ & - 12m_c^4 (7m_c^4 I[-4, 2] - 13m_c^2 I[-3, 2] + 6I[-2, 2]) \\ & \times I_3[h_\gamma] + 6(14m_c^8 I[-4, 2] \\ & - 13m_c^6 I[-3, 2] - I[0, 2]) \mathbb{A}[u_0] \\ & - 8\chi (3m_c^{10} I[-5, 3] + 5m_c^8 I[-4, 3] \\ & - 2m_c^4 I[-2, 3] + 6I[0, 3]) \varphi_\gamma[u_0] \left. \right) \\ & - f_{3\gamma} \pi^2 \left( 52(e_d + 2e_u) (m_c^4 I[-2, 1] \right. \\ & + I[0, 1]) I_1[\mathcal{V}] - 96(e_d - 2e_u) \\ & \times (4m_c^6 I[-3, 1] - m_0^2 m_c^4 I[-2, 0] \\ & \left. \left. + m_0^2 I[0, 0] - 4I[0, 1]) \psi^a[u_0] \right) \right] \\ & + \frac{P_2^2}{18432 m_c \pi^4} \left[ 18(e_d + e_u) \right. \\ & \times \left( 4m_c^8 (I[-4, 2] - 2I[-3, 1]) \right. \\ & \left. \left. + 8m_c^6 (m_0^2 (2I[-3, 1] + I[-2, 0]) - 2(I[-3, 2] \right. \right. \end{aligned}$$



$$\begin{aligned}
 &+ I[-2, 1]) - m_c^4(m_0^4 I[-2, 0] \\
 &- 12I[-2, 2] - 8m_0^2 I[-2, 1] \\
 &- I[-1, 0]) + 8I[-1, 1]) + m_0^4 I[0, 0] \\
 &+ 8m_0^2 I[0, 1]) + (e_d + e_u) \left( 9m_c^4(m_c^4 I[-4, 2] \right. \\
 &- m_c^2(m_0^2 I[-3, 1] + 2I[-3, 2]) - m_0^2 I[-2, 1] \\
 &+ I[-2, 2]) \times I_3[h_\gamma] - 9(2m_c^8 I[-4, 2] \\
 &- 2m_c^6(m_0^2 I[-3, 1] + I[-3, 2]) - m_0^2 m_c^4 I[-2, 1] \\
 &+ m_0^2 I[0, 1]) \mathbb{A}[u_0] \\
 &+ 9(m_c^8 I[-4, 2] - m_0^2 m_c^6 I[-3, 1] - m_c^4 I[-2, 2] \\
 &+ m_0^2 I[0, 1]) I_2[\mathcal{S}] + \chi \left( m_c^8(4m_c^2 I[-5, 3] - 9m_0^2 I[-4, 2] \right. \\
 &+ 6I[-4, 3]) + 9m_0^2 m_c^6 I[-3, 2] - 2m_c^4 I[-2, 3] \\
 &+ 8I[0, 3]) \varphi_\gamma[u_0] \Big) \\
 &+ 6f_{3\gamma} \pi^2 \left( (e_d + 2e_u)(m_c^4 I[-2, 1] \right. \\
 &+ I[0, 1]) I_1[\mathcal{V}] - 4(e_d - 2e_u)(2m_c^6 I[-3, 1] \\
 &- m_0^2 m_c^4 I[-2, 0] + m_0^2 I[0, 0] - 2I[0, 1]) \psi^a[u_0] \Big) \\
 &+ \frac{P_2}{491520m_c^2 \pi^6} \left[ 40(e_d + e_u) \left( m_c^{12} (3I[-6, 4] \right. \right. \\
 &+ 4I[-5, 3]) - 3m_c^{10} (m_0^2(4I[-5, 3] - 3I[-4, 2]) \\
 &+ 4(I[-5, 4] - I[-4, 3])) \\
 &- 3m_c^8 (-5I[-4, 4] + 6m_0^2(I[-4, 3] + I[-3, 2]) \\
 &- 4I[-3, 3]) \\
 &+ m_c^6 (-6I[-3, 4] + m_0^2(-6I[-3, 3] + 9I[-2, 2]) \\
 &+ 4I[-2, 3]) + 32m_c^2 I[0, 3] - 36m_0^2 I[0, 3]) \Big) \\
 &+ (e_u + e_d) \left( -5m_c^8 \left( (m_c^4 I[-6, 4] - 3I[-4, 4]) I_2[\mathcal{S}] \right. \right. \\
 &+ (m_c^4 I[-6, 4] \\
 &- 3m_c^2 I[-5, 4] + 3I[-4, 4]) I_6[h_\gamma] \Big) \\
 &+ 5m_c^6 (-2I_2[\mathcal{S}] + I_3[h_\gamma]) I[-3, 4] \\
 &+ m_c^6 (2m_c^6 I[-6, 4] - 3m_c^4 I[-5, 4] + I[-3, 4]) \mathbb{A}[u_0] \\
 &+ 2\chi \left( m_c^6(m_c^6 I[-6, 5] + 3m_c^4 I[-5, 5] \right. \\
 &+ 3m_c^2 I[-4, 5] + I[-3, 5]) + 8I[0, 5]) \varphi_\gamma[u_0] \Big)
 \end{aligned}$$

$$\begin{aligned}
 &+ f_{3\gamma} \pi^2 \left( - (e_d + 2e_u) \left( 4m_c^8 I[-4, 3] \right. \right. \\
 &+ m_c^6 (-3m_0^2 I[-3, 2] \\
 &+ 4I[-3, 3]) + 3m_0^2 I[0, 2] + 8I[0, 3]) I_1[\mathcal{V}] \\
 &+ 8(e_d - 2e_u) \left( 2m_c^{10} I[-5, 3] + 3m_0^2 m_c^8 I[-4, 2] \right. \\
 &\left. \left. - 2m_c^6 I[-3, 3] - 3m_0^2 I[0, 2] \right) \psi^a[u_0] \right) \Big), \tag{A1}
 \end{aligned}$$

where  $P_1 = \langle g_s^2 G^2 \rangle$  and  $P_2 = \langle \bar{q}q \rangle$  are gluon and light-quark condensates, respectively. It should also be noted that in the  $\Delta_1^{QCD}(M^2, s_0)$  function we have, for simplicity, presented only those terms that make significant contributions to the numerical values of the magnetic dipole moment and have ignored many higher dimensional terms, although they are included in the numerical calculations. The functions  $I[n, m]$ ,  $I_1[\mathcal{F}]$ ,  $I_2[\mathcal{F}]$  and  $I_3[\mathcal{F}]$  are defined as:

$$\begin{aligned}
 I[n, m] &= \int_{m_c^2}^{s_0} ds \int_{m_c^2}^s dl e^{-s/M^2} l^n (s-l)^m, \\
 I_1[\mathcal{F}] &= \int D\alpha_i \int_0^1 dv \mathcal{F}(\alpha_{\bar{q}}, \alpha_q, \alpha_g) \delta'(\alpha_{\bar{q}} + v\alpha_g - u_0), \\
 I_2[\mathcal{F}] &= \int D\alpha_i \int_0^1 dv \mathcal{F}(\alpha_{\bar{q}}, \alpha_q, \alpha_g) \delta(\alpha_{\bar{q}} + v\alpha_g - u_0), \\
 I_3[\mathcal{F}] &= \int_0^1 du \mathcal{F}(u), \tag{A2}
 \end{aligned}$$

where  $\mathcal{F}$  represents the corresponding photon DAs and The measure  $\mathcal{D}\alpha_i$  is defined as

$$\int \mathcal{D}\alpha_i = \int_0^1 d\alpha_{\bar{q}} \int_0^1 d\alpha_q \int_0^1 d\alpha_g \delta'(1 - \alpha_{\bar{q}} - \alpha_q - \alpha_g).$$

### Appendix B Explicit forms of the sum rules for $F_i$ form factors

In the present appendix, we present explicit expressions of the analytical expressions obtained for the magnetic dipole, electric quadrupole, and magnetic octupole moments of the  $\Lambda_c^+$  state. These multipole moments are written as follows:

$$\begin{aligned}
 \mu_{\Lambda_c^+} &= \frac{e}{2m_{\Lambda_c^+}} F_1 + F_2, \\
 Q_{\Lambda_c^+} &= \frac{e}{m_{\Lambda_c^+}^2} F_1 - \frac{1}{2} F_3, \\
 O_{\Lambda_c^+} &= \frac{e}{2m_{\Lambda_c^+}^3} F_1 + F_2 - \frac{1}{2} [F_3 + F_4], \tag{B1}
 \end{aligned}$$

where

$$\begin{aligned}
 F_1 = & -\frac{e^{\frac{m_c^2}{M^2}}}{\lambda_{\Lambda_c^+}^2} \\
 & \times \left\{ -\frac{P_1 P_2^2 (e_d + e_u)}{13824 m_c^2 \pi^4} \left[ I[0, 0] \right. \right. \\
 & - 2m_c^4 I[2, 0] + m_c^6 I[3, 0] \\
 & + \frac{P_1 P_2 (e_d + e_u)}{589824 m_c \pi^6} \left[ (m_c^6 I[-3, 1] + 2m_c^4 I[-2, 1] \right. \\
 & + m_c^2 I[-1, 1] + 4I[0, 1]) I_1[S] \\
 & - (m_c^6 I[-3, 1] + 2m_c^4 I[-2, 1] \\
 & + m_c^2 I[-1, 1] + 4I[0, 1]) I_1[\tilde{S}] \\
 & - 20(4m_c^6 I[-3, 1] + m_0^2 I[0, 0] \\
 & + m_c^4(4I[-2, 1] - m_0^2 I[2, 0])) \left. \right] \\
 & + \frac{P_2^2 (e_d + e_u)}{2048 m_c^2 \pi^4} \left[ 4m_c^{10} I[-5, 2] \right. \\
 & - 8m_0^2 m_c^8 I[-4, 1] - m_0^4 I[0, 0] \\
 & + 8m_0^2 I[0, 1] + m_c^6(-4I[-3, 2] \\
 & + m_0^4 I[3, 0]) \left. \right] \\
 & - \frac{P_1 (e_d + e_u)}{1769472 m_c^2 \pi^8} \left[ 6m_c^{12} I[-6, 3] \right. \\
 & + 8m_c^{10} I[-5, 3] + 3m_c^8 I[-4, 3] + 6m_c^6 I[-3, 3] \\
 & + 5m_c^4 I[-2, 3] + 28I[0, 3] \left. \right] \\
 & - \frac{P_2 (e_d + e_u)}{12288 m_c \pi^6} \\
 & \times \left[ m_c^4(4m_c^6 I[-5, 3] + m_c^4(-9m_0^2 I[-4, 2] \right. \\
 & + 12I[-4, 3]) + 6m_c^2(3m_0^2 I[-3, 2] + 2I[-3, 3]) \\
 & - 9m_0^2 I[-2, 2] + 4I[-2, 3]) + 32I[0, 3] \left. \right] \\
 & - \frac{(e_d + e_u)}{1638400 m_c^2 \pi^8} \left[ m_c^{16} I[-8, 5] \right. \\
 & - 10m_c^{12} I[-6, 5] - 20m_c^{10} I[-5, 5] \\
 & - 15m_c^8 I[-4, 5] - 4m_c^6 I[-3, 5] - 48I[0, 5] \left. \right] \left. \right\}, \quad (B2) \\
 F_2 = & \frac{m_{\Lambda_c^+} e^{\frac{m_c^2}{M^2}}}{\lambda_{\Lambda_c^+}^2}
 \end{aligned}$$

$$\begin{aligned}
 & \times \left\{ \frac{P_1 P_2 (e_d + e_u)}{36864 m_c^2 \pi^6} \right. \\
 & \times \left[ 2m_c^8 (I[-4, 2] - I[-3, 1]) \right. \\
 & + m_c^4(4I[-2, 2] - m_0^2 I[-1, 0] - 2I[-1, 1]) \\
 & + m_0^2 I[0, 1] \\
 & + m_c^6(-6I[-3, 2] - 4I[-2, 1] + m_0^2(I[-3, 1] \\
 & + I[2, 0])) \left. \right] \\
 & + \frac{P_2^2 (e_d + e_u)}{1024 m_c^2 \pi^4} \\
 & \times \left[ 4m_c^8 (I[-4, 2] - 2I[-3, 1]) \right. \\
 & + m_0^4 I[0, 0] + 8m_0^2 I[0, 1] \\
 & - m_c^4(-12I[-2, 2] - 8m_0^2(I[-2, 1] \\
 & - I[-1, 0]) + 8I[-1, 1] + m_0^4 I[2, 0]) \\
 & + 8m_c^6(-2(I[-3, 2] + I[-2, 1]) \\
 & + m_0^2(2I[-3, 1] + I[2, 0])) \left. \right] \\
 & + \frac{P_2 (e_d + e_u)}{12288 m_c^2 \pi^6} \left[ 2m_c^{12} (I[-6, 4] + 2I[-5, 3]) \right. \\
 & - 3m_c^{10} (3I[-5, 4] + m_0^2(5I[-5, 3] \\
 & - 3I[-4, 2]) - 4I[-4, 3]) \\
 & - 6m_c^8 (m_0^2(4I[-4, 3] + 3I[-3, 2]) - 2(I[-4, 4] \\
 & + I[-3, 3])) + m_c^6(-5I[-3, 4] \\
 & - 9m_0^2(I[-3, 3] - I[-2, 2]) \\
 & + 4I[-2, 3]) - 48m_0^2 I[0, 3] + 32m_c^2 I[0, 3] \left. \right] \left. \right\}, \quad (B3)
 \end{aligned}$$

$$\begin{aligned}
 F_3 = & \frac{4m_{\Lambda_c^+} e^{\frac{m_c^2}{M^2}}}{\lambda_{\Lambda_c^+}^2} \left\{ -\frac{P_1 P_2 (e_d + e_u)}{55296 m_c^2 \pi^6} \right. \\
 & \times \left[ m_c^{10} (I[-5, 1] + I[4, 0]) - 3m_c^8 (I[-4, 1] + I[3, 0]) \right. \\
 & + 3m_c^6(-3I[-3, 1] + I[2, 0]) \\
 & - m_0^2 I[3, 0] - m_c^4(5I[-2, 1] \\
 & + I[-1, 0] - 3m_0^2 I[2, 0]) - 16I[0, 1] \left. \right] \\
 & - \frac{P_2 (e_d + e_u)}{12288 m_c^2 \pi^6}
 \end{aligned}$$

$$\begin{aligned} & \times \left[ m_c^6 \left( 3m_c^8 (I[-7, 3] - I[-6, 2]) \right. \right. \\ & + m_c^6 \left( -6m_0^2 (I[-6, 2] + I[-5, 1]) + 4(I[-6, 3] \right. \\ & + 3I[-5, 2]) \left. \right) \\ & - 6m_c^4 \left( I[-5, 3] + 3m_0^2 (-2I[-5, 2] + I[-4, 1]) \right. \\ & + 3I[-4, 2] \left. \right) - 6m_c^2 \left( 3m_0^2 (3I[-4, 2] + I[-3, 1]) \right. \\ & + 2(I[-4, 3] - I[-3, 2]) \left. \right) - 5I[-3, 3] \\ & + 6m_0^2 (4I[-3, 2] - I[-2, 1]) - 3I[-2, 2] \left. \right) \\ & \left. - 48m_0^2 m_c^2 I[0, 1] - 16I[0, 3] \right\}, \end{aligned} \tag{B4}$$

and,

$$\begin{aligned} F_4 = & \frac{4m_{\Lambda_c^+}^3 e^{\frac{m_{\Lambda_c^+}^2}{M^2}}}{\lambda_{\Lambda_c^+}^2} \\ & \times \left\{ \frac{P_1 P_2 (e_d + e_u)}{27648\pi^6} \left[ -I[2, 0] + 3m_c^2 I[3, 0] \right. \right. \\ & \left. \left. - 3m_c^4 I[4, 0] + m_c^6 I[5, 0] \right] \right. \\ & + \frac{P_2 (e_d + e_u)}{2048m_c^2 \pi^6} \left[ m_c^6 \left( -m_c^8 I[-7, 2] \right. \right. \\ & + 2m_c^6 (m_0^2 I[-6, 1] + 2I[-6, 2]) \\ & + 6m_c^4 (m_0^2 I[-5, 1] - I[-5, 2]) + m_c^2 (6m_0^2 \\ & \times I[-4, 1] + 4I[-4, 2]) + 2m_0^2 I[-3, 1] \\ & \left. \left. - I[-3, 2] \right) + 16m_0^2 I[0, 1] \right] \left. \right\}. \end{aligned} \tag{B5}$$

Definitions of  $P_1$ ,  $P_2$ ,  $I[n, m]$ ,  $I_1[\mathcal{F}]$ ,  $I_2[\mathcal{F}]$ , and  $I_3[\mathcal{F}]$  in Eqs. (B2)–(B5) are given at end of Appendix A

### Appendix C Distribution amplitudes of the photon

In this appendix, the matrix elements  $\langle \gamma(q) | \bar{q}(x) \Gamma_i q(0) | 0 \rangle$  and  $\langle \gamma(q) | \bar{q}(x) \Gamma_i G_{\mu\nu} q(0) | 0 \rangle$  associated with the photon DAs are presented as follows [78]:

$$\begin{aligned} \langle \gamma(q) | \bar{q}(x) \gamma_\mu q(0) | 0 \rangle &= e_q f_{3\gamma} \left( \varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \\ & \times \int_0^1 du e^{i\bar{u}qx} \psi^v(u) \\ \langle \gamma(q) | \bar{q}(x) \gamma_\mu \gamma_5 q(0) | 0 \rangle &= -\frac{1}{4} e_q f_{3\gamma} \varepsilon_{\mu\nu\alpha\beta} \varepsilon^\nu q^\alpha x^\beta \end{aligned}$$

$$\begin{aligned} & \times \int_0^1 du e^{i\bar{u}qx} \psi^a(u) \\ \langle \gamma(q) | \bar{q}(x) \sigma_{\mu\nu} q(0) | 0 \rangle &= -ie_q \langle \bar{q}q \rangle (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \\ & \times \int_0^1 du e^{i\bar{u}qx} \left( \chi \varphi_\gamma(u) + \frac{x^2}{16} \mathbb{A}(u) \right) \\ & - \frac{i}{2(qx)} e_q \bar{q}q \left[ x_\nu \left( \varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \right. \\ & \left. - x_\mu \left( \varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \right] \int_0^1 du e^{i\bar{u}qx} h_\gamma(u) \\ \langle \gamma(q) | \bar{q}(x) g_s G_{\mu\nu}(vx) q(0) | 0 \rangle &= -ie_q \langle \bar{q}q \rangle (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \\ & \times \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{S}(\alpha_i) \\ \langle \gamma(q) | \bar{q}(x) g_s \tilde{G}_{\mu\nu}(vx) i\gamma_5 q(0) | 0 \rangle &= -ie_q \langle \bar{q}q \rangle (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \\ & \times \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \tilde{\mathcal{S}}(\alpha_i) \\ \langle \gamma(q) | \bar{q}(x) g_s \tilde{G}_{\mu\nu}(vx) \gamma_\alpha \gamma_5 q(0) | 0 \rangle &= e_q f_{3\gamma} q_\alpha (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \\ & \times \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{A}(\alpha_i) \\ \langle \gamma(q) | \bar{q}(x) g_s G_{\mu\nu}(vx) i\gamma_\alpha q(0) | 0 \rangle &= e_q f_{3\gamma} q_\alpha (\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \\ & \times \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{V}(\alpha_i) \\ \langle \gamma(q) | \bar{q}(x) \sigma_{\alpha\beta} g_s G_{\mu\nu}(vx) q(0) | 0 \rangle &= e_q \langle \bar{q}q \rangle \left\{ \left[ \left( \varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \right. \right. \\ & \times \left( g_{\alpha\nu} - \frac{1}{qx} (q_\alpha x_\nu + q_\nu x_\alpha) \right) q_\beta \\ & - \left( \varepsilon_\mu - q_\mu \frac{\varepsilon x}{qx} \right) \left( g_{\beta\nu} - \frac{1}{qx} (q_\beta x_\nu + q_\nu x_\beta) \right) q_\alpha \\ & - \left( \varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \left( g_{\alpha\mu} - \frac{1}{qx} (q_\alpha x_\mu + q_\mu x_\alpha) \right) q_\beta \\ & \left. \left. + \left( \varepsilon_\nu - q_\nu \frac{\varepsilon x}{qx} \right) \left( g_{\beta\mu} - \frac{1}{qx} (q_\beta x_\mu + q_\mu x_\beta) \right) q_\alpha \right] \right. \\ & \times \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_1(\alpha_i) \\ & + \left[ \left( \varepsilon_\alpha - q_\alpha \frac{\varepsilon x}{qx} \right) \left( g_{\mu\beta} - \frac{1}{qx} (q_\mu x_\beta + q_\beta x_\mu) \right) q_\nu \right. \\ & - \left( \varepsilon_\alpha - q_\alpha \frac{\varepsilon x}{qx} \right) \left( g_{\nu\beta} - \frac{1}{qx} (q_\nu x_\beta + q_\beta x_\nu) \right) q_\mu \\ & - \left( \varepsilon_\beta - q_\beta \frac{\varepsilon x}{qx} \right) \left( g_{\mu\alpha} - \frac{1}{qx} (q_\mu x_\alpha + q_\alpha x_\mu) \right) q_\nu \\ & \left. \left. + \left( \varepsilon_\beta - q_\beta \frac{\varepsilon x}{qx} \right) \left( g_{\nu\alpha} - \frac{1}{qx} (q_\nu x_\alpha + q_\alpha x_\nu) \right) q_\mu \right] \right\} \end{aligned}$$

$$\begin{aligned}
 & \times \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_2(\alpha_i) && + (\zeta_1 - \zeta_1^+)(1 - \alpha_g)(1 - 2\alpha_g) \\
 & + \frac{1}{qx} (q_\mu x_\nu - q_\nu x_\mu)(\varepsilon_\alpha q_\beta - \varepsilon_\beta q_\alpha) && + \zeta_2 [3(\alpha_{\bar{q}} - \alpha_q)^2 - \alpha_g(1 - \alpha_g)]. \tag{C1} \\
 & \times \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_3(\alpha_i) \\
 & + \frac{1}{qx} (q_\alpha x_\beta - q_\beta x_\alpha)(\varepsilon_\mu q_\nu - \varepsilon_\nu q_\mu) \\
 & \times \left. \int \mathcal{D}\alpha_i e^{i(\alpha_{\bar{q}} + v\alpha_g)qx} \mathcal{T}_4(\alpha_i) \right\},
 \end{aligned}$$

where  $\varphi_\gamma(u)$  is the DA of leading twist-2,  $\psi^v(u)$ ,  $\psi^a(u)$ ,  $\mathcal{A}(\alpha_i)$  and  $\mathcal{V}(\alpha_i)$ , are the twist-3 amplitudes, and  $h_\gamma(u)$ ,  $\mathbb{A}(u)$ ,  $\mathcal{S}(\alpha_i)$ ,  $\tilde{\mathcal{S}}(\alpha_i)$ ,  $\mathcal{T}_1(\alpha_i)$ ,  $\mathcal{T}_2(\alpha_i)$ ,  $\mathcal{T}_3(\alpha_i)$  and  $\mathcal{T}_4(\alpha_i)$  are the twist-4 photon DAs.

The expressions of the DAs that are entered into the matrix elements above are as follows:

$$\begin{aligned}
 \varphi_\gamma(u) &= 6u\bar{u} \left( 1 + \varphi_2(\mu) C_2^{\frac{3}{2}}(u - \bar{u}) \right), \\
 \psi^v(u) &= 3 \left( 3(2u - 1)^2 - 1 \right) \\
 & \quad + \frac{3}{64} \left( 15w_\gamma^V - 5w_\gamma^A \right) \\
 & \quad \times \left( 3 - 30(2u - 1)^2 + 35(2u - 1)^4 \right), \\
 \psi^a(u) &= \left( 1 - (2u - 1)^2 \right) \left( 5(2u - 1)^2 - 1 \right) \frac{5}{2} \left( 1 \right. \\
 & \quad \left. + \frac{9}{16} w_\gamma^V - \frac{3}{16} w_\gamma^A \right), \\
 h_\gamma(u) &= -10 \left( 1 + 2\kappa^+ \right) C_2^{\frac{1}{2}}(u - \bar{u}), \\
 \mathbb{A}(u) &= 40u^2\bar{u}^2 \left( 3\kappa - \kappa^+ + 1 \right) \\
 & \quad + 8(\zeta_2^+ - 3\zeta_2) [u\bar{u}(2 + 13u\bar{u}) \\
 & \quad + 2u^3(10 - 15u + 6u^2) \ln(u) \\
 & \quad + 2\bar{u}^3(10 - 15\bar{u} + 6\bar{u}^2) \ln(\bar{u})], \\
 \mathcal{A}(\alpha_i) &= 360\alpha_q\alpha_{\bar{q}}\alpha_g^2 \left( 1 + w_\gamma^A \frac{1}{2}(7\alpha_g - 3) \right), \\
 \mathcal{V}(\alpha_i) &= 540w_\gamma^V (\alpha_q - \alpha_{\bar{q}})\alpha_q\alpha_{\bar{q}}\alpha_g^2, \\
 \mathcal{T}_1(\alpha_i) &= -120(3\zeta_2 + \zeta_2^+)(\alpha_{\bar{q}} - \alpha_q)\alpha_{\bar{q}}\alpha_q\alpha_g, \\
 \mathcal{T}_2(\alpha_i) &= 30\alpha_g^2(\alpha_{\bar{q}} - \alpha_q) \left( (\kappa - \kappa^+) + (\zeta_1 - \zeta_1^+)(1 - 2\alpha_g) \right. \\
 & \quad \left. + \zeta_2(3 - 4\alpha_g) \right), \\
 \mathcal{T}_3(\alpha_i) &= -120(3\zeta_2 - \zeta_2^+)(\alpha_{\bar{q}} - \alpha_q)\alpha_{\bar{q}}\alpha_q\alpha_g, \\
 \mathcal{T}_4(\alpha_i) &= 30\alpha_g^2(\alpha_{\bar{q}} - \alpha_q) \left( (\kappa + \kappa^+) + (\zeta_1 + \zeta_1^+)(1 - 2\alpha_g) \right. \\
 & \quad \left. + \zeta_2(3 - 4\alpha_g) \right), \\
 \mathcal{S}(\alpha_i) &= 30\alpha_g^2 \{ (\kappa + \kappa^+)(1 - \alpha_g) \\
 & \quad + (\zeta_1 + \zeta_1^+)(1 - \alpha_g)(1 - 2\alpha_g) \\
 & \quad + \zeta_2 [3(\alpha_{\bar{q}} - \alpha_q)^2 - \alpha_g(1 - \alpha_g)] \}, \\
 \tilde{\mathcal{S}}(\alpha_i) &= -30\alpha_g^2 \{ (\kappa - \kappa^+)(1 - \alpha_g)
 \end{aligned}$$

The numerical values of the parameters used in the DAs are  $\varphi_2(1 \text{ GeV}) = 0$ ,  $w_\gamma^V = 3.8 \pm 1.8$ ,  $w_\gamma^A = -2.1 \pm 1.0$ ,  $\kappa = 0.2$ ,  $\kappa^+ = 0$ ,  $\zeta_1 = 0.4$ , and  $\zeta_2 = 0.3$ .

References

1. F.-K. Guo, C. Hanhart, U.-G. Meißner, Q. Wang, Q. Zhao, B.-S. Zou, Hadronic molecules. *Rev. Mod. Phys.* **90**(1), 015004 (2018). <https://doi.org/10.1103/RevModPhys.90.015004> [arXiv:1705.00141](https://arxiv.org/abs/1705.00141). [Erratum: *Rev. Mod. Phys.* 94, 029901 (2022)]
2. S.L. Olsen, T. Skwarnicki, D. Zieminska, Nonstandard heavy mesons and baryons: experimental evidence. *Rev. Mod. Phys.* **90**(1), 015003 (2018). <https://doi.org/10.1103/RevModPhys.90.015003>. [arXiv:1708.04012](https://arxiv.org/abs/1708.04012)
3. N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C.-P. Shen, C.E. Thomas, A. Vairo, C.-Z. Yuan, The XYZ states: experimental and theoretical status and perspectives. *Phys. Rep.* **873**, 1–154 (2020). <https://doi.org/10.1016/j.physrep.2020.05.001>. [arXiv:1907.07583](https://arxiv.org/abs/1907.07583)
4. H.-X. Chen, W. Chen, X. Liu, Y.-R. Liu, S.-L. Zhu, An updated review of the new hadron states. *Rep. Prog. Phys.* **86**(2), 026201 (2023). <https://doi.org/10.1088/1361-6633/aca3b6>. [arXiv:2204.02649](https://arxiv.org/abs/2204.02649)
5. H. Albrecht et al., Observation of a new charmed baryon. *Phys. Lett. B* **317**, 227–232 (1993). [https://doi.org/10.1016/0370-2693\(93\)91598-H](https://doi.org/10.1016/0370-2693(93)91598-H)
6. K.W. Edwards et al., Observation of excited baryon states decaying to Lambda(c)+ pi+ pi-. *Phys. Rev. Lett.* **74**, 3331–3335 (1995). <https://doi.org/10.1103/PhysRevLett.74.3331>
7. H. Albrecht et al., Evidence for Lambda(c)+ (2593) production. *Phys. Lett. B* **402**, 207–212 (1997). [https://doi.org/10.1016/S0370-2693\(97\)00503-0](https://doi.org/10.1016/S0370-2693(97)00503-0)
8. P.L. Frabetti et al., An observation of an excited state of the Lambda(c)+ baryon. *Phys. Rev. Lett.* **72**, 961–964 (1994). <https://doi.org/10.1103/PhysRevLett.72.961>
9. M. Artuso et al., Observation of new states decaying into Lambda+(c) pi- pi+. *Phys. Rev. Lett.* **86**, 4479–4482 (2001). <https://doi.org/10.1103/PhysRevLett.86.4479>. [arXiv:hep-ex/0010080](https://arxiv.org/abs/hep-ex/0010080)
10. B. Aubert et al., Observation of a charmed baryon decaying to D0p at a mass near 2.94-GeV/c\*\*2. *Phys. Rev. Lett.* **98**, 012001 (2007). <https://doi.org/10.1103/PhysRevLett.98.012001>. [arXiv:hep-ex/0603052](https://arxiv.org/abs/hep-ex/0603052)
11. K. Abe et al., Experimental constraints on the possible J\*\*P quantum numbers of the Lambda(c)(2880)+. *Phys. Rev. Lett.* **98**, 262001 (2007). <https://doi.org/10.1103/PhysRevLett.98.262001>. [arXiv:hep-ex/0608043](https://arxiv.org/abs/hep-ex/0608043)
12. R. Aaij et al., Study of the D0p amplitude in Lambda\_b0 -> D0p pi- decays. *JHEP* **05**, 030 (2017). [https://doi.org/10.1007/JHEP05\(2017\)030](https://doi.org/10.1007/JHEP05(2017)030). [arXiv:1701.07873](https://arxiv.org/abs/1701.07873)
13. V.V. Ammosov et al., Observation of the production of charmed Sigma(c)\*++ baryons in neutrino interactions at the SKAT bubble chamber. *JETP Lett.* **58**, 247–251 (1993)
14. G. Brandenburg et al., Observation of two excited charmed baryons decaying into Lambda(c)+ pi+- . *Phys. Rev. Lett.* **78**, 2304–2308 (1997). <https://doi.org/10.1103/PhysRevLett.78.2304>
15. R. Mizuk et al., Observation of an is triplet of excited charmed baryons decaying to Lambda+(c) pi. *Phys. Rev. Lett.* **94**, 122002 (2005). <https://doi.org/10.1103/PhysRevLett.94.122002>. [arXiv:hep-ex/0412069](https://arxiv.org/abs/hep-ex/0412069)

16. P. Avery et al., Observation of a narrow state decaying into  $\Xi(c)^+ \pi^-$ . Phys. Rev. Lett. **75**, 4364–4368 (1995). <https://doi.org/10.1103/PhysRevLett.75.4364>. arXiv:hep-ex/9508010
17. L. Gibbons et al., Observation of an excited charmed baryon decaying into  $\Xi(c)^0 \pi^+$ . Phys. Rev. Lett. **77**, 810–813 (1996). <https://doi.org/10.1103/PhysRevLett.77.810>
18. P.L. Frabetti et al., Observation of a narrow state decaying into  $\Xi(c)^0 \pi^+$ . Phys. Lett. B **426**, 403–410 (1998). [https://doi.org/10.1016/S0370-2693\(98\)00348-7](https://doi.org/10.1016/S0370-2693(98)00348-7)
19. S.E. Csorna et al., Evidence of new states decaying into  $\Xi$ -prime(c)  $\pi$ . Phys. Rev. Lett. **86**, 4243–4246 (2001). <https://doi.org/10.1103/PhysRevLett.86.4243>. arXiv:hep-ex/0012020
20. J.P. Alexander et al., Evidence of new states decaying into  $\Xi(c)^* \pi$ . Phys. Rev. Lett. **83**, 3390–3393 (1999). <https://doi.org/10.1103/PhysRevLett.83.3390>. arXiv:hep-ex/9906013
21. R. Aaij et al., Observation of new  $\Xi_c^0$  baryons decaying to  $\Lambda_c^+ K^-$ . Phys. Rev. Lett. **124**(22), 222001 (2020). <https://doi.org/10.1103/PhysRevLett.124.222001>. arXiv:2003.13649
22. B. Aubert et al., A study of anti-B  $\rightarrow$   $\Xi(c)$  anti-Lambda(c) and anti-B  $\rightarrow$  anti-Lambda+(c) anti-Lambda-(c) anti-K decays at BABAR. Phys. Rev. D **77**, 031101 (2008). <https://doi.org/10.1103/PhysRevD.77.031101>. arXiv:0710.5775
23. R. Chistov et al., Observation of new states decaying into Lambda(c)+ K- pi+ and Lambda(c)+ K0(S) pi-. Phys. Rev. Lett. **97**, 162001 (2006). <https://doi.org/10.1103/PhysRevLett.97.162001>
24. B. Aubert et al., A study of excited charm-strange baryons with evidence for new baryons  $\Xi(c)(3055)^+$  and  $\Xi(c)(3123)^+$ . Phys. Rev. D **77**, 012002 (2008). <https://doi.org/10.1103/PhysRevD.77.012002>. arXiv:0710.5763
25. B. Aubert et al., Observation of an excited charm baryon Omega(C)\* decaying to Omega(C)0 gamma. Phys. Rev. Lett. **97**, 232001 (2006). <https://doi.org/10.1103/PhysRevLett.97.232001>. arXiv:hep-ex/0608055
26. R. Aaij et al., Observation of five new narrow  $\Omega_c^0$  states decaying to  $\Xi_c^+ K^-$ . Phys. Rev. Lett. **118**(18), 182001 (2017). <https://doi.org/10.1103/PhysRevLett.118.182001>. arXiv:1703.04639
27. R. Aaij et al., Observation of excited  $\Lambda_b^0$  baryons. Phys. Rev. Lett. **109**, 172003 (2012). <https://doi.org/10.1103/PhysRevLett.109.172003>. arXiv:1205.3452
28. R. Aaij et al., Observation of a new baryon state in the  $\Lambda_b^0 \pi^+ \pi^-$  mass spectrum. JHEP **06**, 136 (2020). [https://doi.org/10.1007/JHEP06\(2020\)136](https://doi.org/10.1007/JHEP06(2020)136). arXiv:2002.05112
29. R. Aaij et al., Observation of new resonances in the  $\Lambda_b^0 \pi^+ \pi^-$  system. Phys. Rev. Lett. **123**(15), 152001 (2019). <https://doi.org/10.1103/PhysRevLett.123.152001>. arXiv:1907.13598
30. R. Aaij et al., Observation of two resonances in the  $\Lambda_b^0 \pi^\pm$  systems and precise measurement of  $\Sigma_b^\pm$  and  $\Sigma_b^{*\pm}$  properties. Phys. Rev. Lett. **122**(1), 012001 (2019). <https://doi.org/10.1103/PhysRevLett.122.012001>. arXiv:1809.07752
31. A.M. Sirunyan et al., Observation of a new excited beauty strange baryon decaying to  $\Xi_b^- \pi^+ \pi^-$ . Phys. Rev. Lett. **126**(25), 252003 (2021). <https://doi.org/10.1103/PhysRevLett.126.252003>. arXiv:2102.04524
32. R. Aaij et al., Observation of a new  $\Xi_b^-$  resonance. Phys. Rev. Lett. **121**(7), 072002 (2018). <https://doi.org/10.1103/PhysRevLett.121.072002>. arXiv:1805.09418
33. R. Aaij et al., Observation of a new  $\Xi_b^0$  state. Phys. Rev. D **103**(1), 012004 (2021). <https://doi.org/10.1103/PhysRevD.103.012004>. arXiv:2010.14485
34. R. Aaij et al., First observation of excited  $\Omega_b^-$  states. Phys. Rev. Lett. **124**(8), 082002 (2020). <https://doi.org/10.1103/PhysRevLett.124.082002>. arXiv:2001.00851
35. B. Aubert et al., Measurements of B(anti-B0  $\rightarrow$  Lambda(c)+ anti-p) and B(B-  $\rightarrow$  Lambda(c)+ anti-p pi-) and studies of Lambda(c)+ pi- resonances. Phys. Rev. D **78**, 112003 (2008). <https://doi.org/10.1103/PhysRevD.78.112003>. arXiv:0807.4974
36. K.-L. Wang, Y.-X. Yao, X.-H. Zhong, Q. Zhao, Strong and radiative decays of the low-lying S- and P-wave singly heavy baryons. Phys. Rev. D **96**(11), 116016 (2017). <https://doi.org/10.1103/PhysRevD.96.116016>. arXiv:1709.04268
37. H.-M. Yang, H.-X. Chen, P-wave charmed baryons of the SU(3) flavor  $6_F$ . Phys. Rev. D **104**(3), 034037 (2021). <https://doi.org/10.1103/PhysRevD.104.034037>. arXiv:2106.15488
38. D. Jia, W.-N. Liu, A. Hosaka, Regge behaviors in orbitally excited spectroscopy of charmed and bottom baryons. Phys. Rev. D **101**(3), 034016 (2020). <https://doi.org/10.1103/PhysRevD.101.034016>. arXiv:1907.04958
39. Y.-H. Zhou, W.-J. Wang, L.-Y. Xiao, X.-H. Zhong, Strong decays of the low-lying 1P- and 1D-wave  $\Sigma_c$  baryons. Phys. Rev. D **108**(1), 014019 (2023). <https://doi.org/10.1103/PhysRevD.108.014019>. arXiv:2303.13774
40. W.-J. Wang, L.-Y. Xiao, X.-H. Zhong, Strong decays of the low-lying  $\rho$ -mode 1P-wave singly heavy baryons. Phys. Rev. D **106**(7), 074020 (2022). <https://doi.org/10.1103/PhysRevD.106.074020>. arXiv:2208.10116
41. J.-J. Guo, P. Yang, A. Zhang, Strong decays of observed  $\Lambda_c$  baryons in the  $^3P_0$  model. Phys. Rev. D **100**(1), 014001 (2019). <https://doi.org/10.1103/PhysRevD.100.014001>. arXiv:1902.07488
42. Q.-F. Lü, L.-Y. Xiao, Z.-Y. Wang, X.-H. Zhong, Strong decay of  $\Lambda_c(2940)$  as a 2P state in the  $\Lambda_c$  family. Eur. Phys. J. C **78**(7), 599 (2018). <https://doi.org/10.1140/epjc/s10052-018-6083-7>. arXiv:1806.01076
43. S.-Q. Luo, B. Chen, Z.-W. Liu, X. Liu, Resolving the low mass puzzle of  $\Lambda_c(2940)^+$ . Eur. Phys. J. C **80**(4), 301 (2020). <https://doi.org/10.1140/epjc/s10052-020-7874-1>. arXiv:1910.14545
44. K. Gong, H.-Y. Jing, A. Zhang, Possible assignments of highly excited  $\Lambda_c(2860)^+$ ,  $\Lambda_c(2880)^+$  and  $\Lambda_c(2940)^+$ . Eur. Phys. J. C **81**(5), 467 (2021). <https://doi.org/10.1140/epjc/s10052-021-09255-w>
45. S. Sakai, F.-K. Guo, B. Kubis, Extraction of ND scattering lengths from the  $\Lambda_b \rightarrow \pi^- p D^0$  decay and properties of the  $\Sigma_c(2800)^+$ . Phys. Lett. B **808**, 135623 (2020). <https://doi.org/10.1016/j.physletb.2020.135623>. arXiv:2004.09824
46. L. Zhao, H. Huang, J. Ping, ND and NB systems in quark delocalization color screening model. Eur. Phys. J. A **53**(2), 28 (2017). <https://doi.org/10.1140/epja/i2017-12219-4>. arXiv:1612.00350
47. Z.-Y. Wang, J.-J. Qi, X.-H. Guo, K.-W. Wei, Study of molecular ND bound states in the Bethe–Salpeter equation approach. Phys. Rev. D **97**(9), 094025 (2018). <https://doi.org/10.1103/PhysRevD.97.094025>
48. D. Zhang, D. Yang, X.-F. Wang, K. Nakayama, Possible S-wave  $ND^{(*)}$  and  $N\bar{B}^{(*)}$  bound states in a chiral quark model. (2019). arXiv:1903.01207
49. M.-J. Yan, F.-Z. Peng, M. Pavon Valderrama, Molecular charmed baryons and pentaquarks from light-meson exchange saturation. (2023). arXiv:2304.14855
50. J.-R. Zhang, S-wave  $D^{(*)}N$  molecular states:  $\Sigma_c(2800)$  and  $\Lambda_c(2940)^{+?}$ . Phys. Rev. D **89**(9), 096006 (2014). <https://doi.org/10.1103/PhysRevD.89.096006>. arXiv:1212.5325
51. B. Wang, L. Meng, S.-L. Zhu,  $D^{(*)}N$  interaction and the structure of  $\Sigma_c(2800)$  and  $\Lambda_c(2940)$  in chiral effective field theory. Phys. Rev. D **101**(9), 094035 (2020). <https://doi.org/10.1103/PhysRevD.101.094035>. arXiv:2003.05688
52. Q. Xin, X.-S. Yang, Z.-G. Wang, The singly-charmed pentaquark molecular states via the QCD sum rules. (2023). arXiv:2307.08926
53. X.-G. He, X.-Q. Li, X. Liu, X.-Q. Zeng, Lambda+(c)(2940): a possible molecular state? Eur. Phys. J. C **51**, 883–889 (2007). <https://doi.org/10.1140/epjc/s10052-007-0347-y>. arXiv:hep-ph/0606015

54. Y. Yan, X. Hu, Y. Wu, H. Huang, J. Ping, Y. Yang, Pentaquark interpretation of  $\Lambda_c$  states in the quark model. *Eur. Phys. J. C* **83**(6), 524 (2023). <https://doi.org/10.1140/epjc/s10052-023-11709-2>. arXiv:2211.12129
55. Z.-L. Zhang, Z.-W. Liu, S.-Q. Luo, F.-L. Wang, B. Wang, H. Xu,  $\Lambda_c(2910)$  and  $\Lambda_c(2940)$  as conventional baryons dressed with the  $D^*N$  channel. *Phys. Rev. D* **107**(3), 034036 (2023). <https://doi.org/10.1103/PhysRevD.107.034036>. arXiv:2210.17188
56. Y. Huang, J. He, J.-J. Xie, L.-S. Geng, Production of the  $\Lambda_c(2940)$  by kaon-induced reactions on a proton target. *Phys. Rev. D* **99**(1), 014045 (2019). <https://doi.org/10.1103/PhysRevD.99.014045>. arXiv:1610.06994
57. X.-Y. Wang, A. Guskov, X.-R. Chen,  $\Lambda_c^*(2940)^+$  photoproduction off the neutron. *Phys. Rev. D* **92**(9), 094032 (2015). <https://doi.org/10.1103/PhysRevD.92.094032>. arXiv:1509.02602
58. Y. Dong, A. Faessler, T. Gutsche, S. Kumano, V.E. Lyubovitskij, Strong three-body decays of  $\Lambda_c(2940)^+$ . *Phys. Rev. D* **83**, 094005 (2011). <https://doi.org/10.1103/PhysRevD.83.094005>. arXiv:1103.4762
59. J. He, Y.-T. Ye, Z.-F. Sun, X. Liu, The observed charmed hadron  $\Lambda_c(2940)^+$  and the  $D^*N$  interaction. *Phys. Rev. D* **82**, 114029 (2010). <https://doi.org/10.1103/PhysRevD.82.114029>. arXiv:1008.1500
60. Y. Dong, A. Faessler, T. Gutsche, S. Kumano, V.E. Lyubovitskij, Radiative decay of  $\Lambda_c(2940)^+$  in a hadronic molecule picture. *Phys. Rev. D* **82**, 034035 (2010). <https://doi.org/10.1103/PhysRevD.82.034035>. arXiv:1006.4018
61. Y. Dong, A. Faessler, T. Gutsche, V.E. Lyubovitskij, Strong two-body decays of the  $\Lambda_c(2940)^+$  in a hadronic molecule picture. *Phys. Rev. D* **81**, 014006 (2010). <https://doi.org/10.1103/PhysRevD.81.014006>. arXiv:0910.1204
62. P.G. Ortega, D.R. Entem, F. Fernandez, Quark model description of the  $\Lambda_c(2940)^+$  as a molecular  $D^*N$  state and the possible existence of the  $\Lambda_b(6248)$ . *Phys. Lett. B* **718**, 1381–1384 (2013). <https://doi.org/10.1016/j.physletb.2012.12.025>. arXiv:1210.2633
63. Y. Dong, A. Faessler, T. Gutsche, V.E. Lyubovitskij, Role of the hadron molecule  $\Lambda_c(2940)$  in the  $p\bar{p} \rightarrow pD^0\bar{\Lambda}_c(2286)$  annihilation reaction. *Phys. Rev. D* **90**(9), 094001 (2014). <https://doi.org/10.1103/PhysRevD.90.094001>. arXiv:1407.3949
64. U. Özdem, Magnetic moments of doubly heavy baryons in light-cone QCD. *J. Phys. G* **46**(3), 035003 (2019). <https://doi.org/10.1088/1361-6471/aaaffc>. arXiv:1804.10921
65. U. Özdem, Magnetic dipole moments of the spin- $\frac{3}{2}$  doubly heavy baryons. *Eur. Phys. J. A* **56**(2), 34 (2020). <https://doi.org/10.1140/epja/s10050-020-00049-4>. arXiv:1906.08353
66. U. Özdem, Magnetic dipole moments of bottom-charm baryons in light-cone QCD. *Eur. Phys. J. C* **83**(10), 887 (2023). <https://doi.org/10.1140/epjc/s10052-023-12055-z>. arXiv:2305.10063
67. U. Özdem, Magnetic dipole moments of the hidden-charm pentaquark states:  $P_c(4440)$ ,  $P_c(4457)$  and  $P_{cs}(4459)$ . *Eur. Phys. J. C* **81**(4), 277 (2021). <https://doi.org/10.1140/epjc/s10052-021-09070-3>. arXiv:2102.01996
68. U. Özdem, Electromagnetic properties of  $\bar{D}^{(*)}\Xi'_c, \bar{D}^{(*)}\Lambda_c, \bar{D}_s^{(*)}\Lambda_c$  and  $\bar{D}_s^{(*)}\Xi_c$  pentaquarks. *Phys. Lett. B* **846**, 13827 (2023). <https://doi.org/10.1016/j.physletb.2023.138267>. arXiv:2303.10649
69. U. Özdem, Investigation of magnetic moment of  $P_{cs}(4338)$  and  $P_{cs}(4459)$  pentaquark states. *Phys. Lett. B* **836**, 137635 (2023). <https://doi.org/10.1016/j.physletb.2022.137635>. arXiv:2208.07684
70. U. Özdem, Magnetic moment of the  $\Xi_b(6227)$  as a molecular pentaquark state. *Eur. Phys. J. Plus* **137**(1), 103 (2022). <https://doi.org/10.1140/epjp/s13360-022-02339-w>. arXiv:2109.09313
71. U. Özdem, Magnetic moments of the doubly charged axial-vector  $T_{cc++}$  states. *Phys. Rev. D* **105**(5), 054019 (2022). <https://doi.org/10.1103/PhysRevD.105.054019>. arXiv:2112.10402
72. U. Özdem, Electromagnetic properties of doubly heavy pentaquark states. *Eur. Phys. J. Plus* **137**, 936 (2022). <https://doi.org/10.1140/epjp/s13360-022-03125-4>. arXiv:2201.00979
73. V.L. Chernyak, I.R. Zhitnitsky, B meson exclusive decays into baryons. *Nucl. Phys. B* **345**, 137–172 (1990). [https://doi.org/10.1016/0550-3213\(90\)90612-H](https://doi.org/10.1016/0550-3213(90)90612-H)
74. V.M. Braun, I.E. Filyanov, QCD sum rules in exclusive kinematics and pion wave function. *Z. Phys. C* **44**, 157 (1989). <https://doi.org/10.1007/BF01548594>
75. I.I. Balitsky, V.M. Braun, A.V. Kolesnichenko, Radiative decay  $\sigma^+ \rightarrow p \gamma$  in quantum chromodynamics. *Nucl. Phys. B* **312**, 509–550 (1989). [https://doi.org/10.1016/0550-3213\(89\)90570-1](https://doi.org/10.1016/0550-3213(89)90570-1)
76. K.-C. Yang, W.Y.P. Hwang, E.M. Henley, L.S. Kisslinger, QCD sum rules and neutron proton mass difference. *Phys. Rev. D* **47**, 3001–3012 (1993). <https://doi.org/10.1103/PhysRevD.47.3001>
77. V.M. Belyaev, B.Y. Blok, Charmed baryons in quantum chromodynamics. *Z. Phys. C* **30**, 151 (1986). <https://doi.org/10.1007/BF01560689>
78. P. Ball, V.M. Braun, N. Kivel, Photon distribution amplitudes in QCD. *Nucl. Phys. B* **649**, 263–296 (2003). [https://doi.org/10.1016/S0550-3213\(02\)01017-9](https://doi.org/10.1016/S0550-3213(02)01017-9). arXiv:hep-ph/0207307
79. V.M. Belyaev, B.L. Ioffe, Determination of the baryon mass and baryon resonances from the quantum-chromodynamics sum rule. Strange baryons. *Sov. Phys. JETP* **57**, 716–721 (1983)
80. H.J. Weber, H. Arenhovel, Isobar configurations in nuclei. *Phys. Rep.* **36**, 277–348 (1978). [https://doi.org/10.1016/0370-1573\(78\)90187-4](https://doi.org/10.1016/0370-1573(78)90187-4)
81. S. Nozawa, D.B. Leinweber, Electromagnetic form-factors of spin 3/2 baryons. *Phys. Rev. D* **42**, 3567–3571 (1990). <https://doi.org/10.1103/PhysRevD.42.3567>
82. V. Pascalutsa, M. Vanderhaeghen, S.N. Yang, Electromagnetic excitation of the  $\Delta(1232)$ -resonance. *Phys. Rep.* **437**, 125–232 (2007). <https://doi.org/10.1016/j.physrep.2006.09.006>. arXiv:hep-ph/0609004
83. G. Ramalho, M.T. Pena, F. Gross, Electric quadrupole and magnetic octupole moments of the  $\Delta$ . *Phys. Lett. B* **678**, 355–358 (2009). <https://doi.org/10.1016/j.physletb.2009.06.052>. arXiv:0902.4212
84. R.L. Workman et al., *Rev. Part. Phys. PTEP* **2022**, 083C01 (2022). <https://doi.org/10.1093/ptep/ptac097>
85. B.L. Ioffe, QCD at low energies. *Prog. Part. Nucl. Phys.* **56**, 232–277 (2006). <https://doi.org/10.1016/j.pnpnp.2005.05.001>. arXiv:hep-ph/0502148
86. R.D. Matheus, S. Narison, M. Nielsen, J.M. Richard, Can the  $X(3872)$  be a  $1^{++}$  four-quark state? *Phys. Rev. D* **75**, 014005 (2007). <https://doi.org/10.1103/PhysRevD.75.014005>. arXiv:hep-ph/0608297
87. J. Rohrwild, Determination of the magnetic susceptibility of the quark condensate using radiative heavy meson decays. *JHEP* **09**, 073 (2007). <https://doi.org/10.1088/1126-6708/2007/09/073>. arXiv:0708.1405
88. J.M.M. Hall, W. Kamleh, D.B. Leinweber, B.J. Menadue, B.J. Owen, A.W. Thomas, R.D. Young, Lattice QCD evidence that the  $\Lambda(1405)$  resonance is an antikaon-nucleon molecule. *Phys. Rev. Lett.* **114**(13), 132002 (2015). <https://doi.org/10.1103/PhysRevLett.114.132002>. arXiv:1411.3402
89. J.M.M. Hall, W. Kamleh, D.B. Leinweber, B.J. Menadue, B.J. Owen, A.W. Thomas, Light-quark contributions to the magnetic form factor of the  $\Lambda(1405)$ . *Phys. Rev. D* **95**(5), 054510 (2017). <https://doi.org/10.1103/PhysRevD.95.054510>. arXiv:1612.07477