

Theory of the $n = 2$ levels in muonic helium-3 ions

Beatrice Franke^{1,2,a}, Julian J. Krauth^{1,3,b}, Aldo Antognini^{4,5}, Marc Diepold¹, Franz Kottmann⁴, and Randolph Pohl^{3,1,c}

¹ Max-Planck-Institut für Quantenoptik, 85748 Garching, Germany

² TRIUMF, 4004 Wesbrook Mall, Vancouver, BC V6T 2A3, Canada

³ Johannes Gutenberg-Universität Mainz, QUANTUM, Institut für Physik & Exzellenzcluster PRISMA, 55099 Mainz, Germany

⁴ Institute for Particle Physics and Astrophysics, ETH Zurich, 8093 Zurich, Switzerland

⁵ Paul Scherrer Institute, 5232 Villigen, Switzerland

Received 30 April 2017 / Received in final form 13 July 2017

Published online 21 December 2017

© The Author(s) 2017. This article is published with open access at Springerlink.com

Abstract. The present knowledge of Lamb shift, fine-, and hyperfine structure of the 2S and 2P states in muonic helium-3 ions is reviewed in anticipation of the results of a first measurement of several 2S → 2P transition frequencies in the muonic helium-3 ion, $\mu^3\text{He}^+$. This ion is the bound state of a single negative muon μ^- and a bare helium-3 nucleus (helion), $^3\text{He}^{++}$.

A term-by-term comparison of all available sources, including new, updated, and so far unpublished calculations, reveals reliable values and uncertainties of the QED and nuclear structure-dependent contributions to the Lamb shift and the hyperfine splitting. These values are essential for the determination of the helion rms charge radius and the nuclear structure effects to the hyperfine splitting in $\mu^3\text{He}^+$. With this review we continue our series of theory summaries in light muonic atoms [see A. Antognini et al., *Ann. Phys.* **331**, 127 (2013); J.J. Krauth et al., *Ann. Phys.* **366**, 168 (2016); and M. Diepold et al. [arXiv:1606.05231](https://arxiv.org/abs/1606.05231) (2016)].

1 Introduction

Laser spectroscopy of light muonic atoms and ions, where a single negative muon orbits a bare nucleus, holds the promise for a vastly improved determination of nuclear parameters, compared to the more traditional methods of elastic electron scattering and precision laser spectroscopy of regular electronic atoms.

The CREMA collaboration has so far determined the charge radii of the proton and the deuteron, by measuring several transitions in muonic hydrogen (μp) [1–3] and muonic deuterium (μd) [4,5]. Interestingly, both values differ by as much as six standard deviations from the respective CODATA-2014 values [6], which contain data from laser spectroscopy in atomic hydrogen/deuterium and electron scattering. This discrepancy has been coined “proton radius puzzle” [7–9]. However, the discrepancy exists for the deuteron, too. Interestingly, for the proton and the deuteron, the muonic isotope shift is compatible with the electronic one from the 1S–2S transition in H and D [10,11]. The respective radii are

$$\begin{aligned} r_p(\mu\text{p}) &= 0.84087(26)^{\text{exp}}(29)^{\text{th}} \\ &= 0.84087(39) \text{ fm} \end{aligned} \quad [1,2] \quad (1)$$

^a e-mail: bfranke@triumf.ca

^b e-mail: jkrauth@uni-mainz.de

^c e-mail: pohl@uni-mainz.de

$$r_p(\text{CODATA}'14) = 0.87510(610) \text{ fm} \quad [6] \quad (2)$$

$$\begin{aligned} r_d(\mu\text{d}) &= 2.12562(13)^{\text{exp}}(77)^{\text{th}} \\ &= 2.12562(78) \text{ fm} \end{aligned} \quad [4] \quad (3)$$

$$r_d(\text{CODATA}'14) = 2.14130(250) \text{ fm.} \quad [6] \quad (4)$$

Very recently, the CREMA collaboration has measured a total of five transitions in muonic helium-3 and -4 ions [12], which have been analyzed now.

These measurements will help to improve our understanding of nuclear model theories [13,14] and shed more light on the proton radius puzzle. Several ideas exist to solve the puzzle [15], some within the standard model [16,17] and others proposing muon specific forces beyond the standard model [18–21]. These ideas lead to predictions which can be tested with precise charge radius determinations in muonic helium ions.

The measurement of the charge radius in both, helium-3 and helium-4 ions will in addition help understand the discrepancy between several measurements of the helium isotope shift in electronic helium [22–26] which yield the difference of the squared charge radii (see Fig. 1).

Several other experiments are on the way to contribute to the puzzle in the future [15] by precision spectroscopy measurements in electronic hydrogen [28–30] and He^+ [31,32], as well as by electron scattering at very low Q^2




Fig. 1. Difference of squared helion-to-alpha particle charge radii as obtained from laser spectroscopy of transitions in regular, electronic helium-3 and helium-4 atoms [22–24] when combined with accurate theory (*[26], **[25]). A 4σ discrepancy persists. Also shown are the individual theory uncertainties which enter $r_h^2 - r_\alpha^2$ ($\mu^4\text{He}^+$: [27], $\mu^3\text{He}^+$: this work), as well as the expected uncertainty from our laser spectroscopy of the Lamb shift in muonic helium ions. Note that the combination of the two theoretical uncertainties should contain correlations which will partly cancel in the total uncertainty.

[33,34] and muon-scattering [35]. The He^+ spectroscopy, in combination with our measurement in muonic helium ions, will be able to determine the Rydberg constant independently from hydrogen and deuterium. This is particularly interesting as the proton charge radius and the Rydberg constant are highly correlated which means that a change in the Rydberg constant could also resolve the puzzle [29].

The determination of the helion charge radius from muonic helium spectroscopy requires accurate knowledge of the corresponding theory. Similar to muonic hydrogen [3], deuterium [5], and helium-4 ions [27], we feel therefore obliged to summarize the current knowledge on the state of theory contributions to the Lamb shift, fine-, and hyperfine structure in muonic helium-3 ions.

The accuracy to be expected from the experiment will be on the order of 20 GHz, which corresponds to ~ 0.08 meV.¹ In order to exploit the experimental precision, theory should, ideally, be accurate to a level of

$$\sigma_{\text{theory}} \sim \mathcal{O}(0.01 \text{ meV}). \quad (5)$$

This would result in a nearly hundred-fold better accuracy in the helion rms charge radius r_h compared to the value from electron scattering of

$$r_h = 1.973(14) \text{ fm}, \quad (6)$$

deduced by Sick [36].

A more precise value has been given by Angeli et al. [37], which should be discarded. Their value is based on a charge radius extraction from $\mu^4\text{He}^+$ by Carboni et al. [38] and on the isotope shift measurement from Shiner et al. [22]. The Carboni measurement has however shown to be wrong [39], and the more recent measurement of the electronic isotope shift by van Rooij et al. [23] disagrees by 4σ from the Shiner one [22], see Figure 1.

We anticipate here that the total uncertainty in the theoretical calculation of the Lamb shift transition amounts to 0.52 meV (corresponding to a relative uncertainty of $\sim 0.03\%$), neglecting the charge radius contribution to be

extracted from the $\mu^3\text{He}^+$ measurement. This value is completely dominated by the two-photon exchange contributions which are difficult to calculate but have seen wonderful progress in recent years [14,40,41]. The total uncertainty of the pure QED contributions (without the two-photon exchange) amounts to 0.04 meV and is thus in the desired order of magnitude. Note that while the theory uncertainty from the two-photon exchange in r_p is of similar size as the experimental uncertainty (Eq. (1)), already for μd the theory uncertainty is vastly dominant (Eq. (3)). Experiments with muonic atoms are thus a sensitive tool to determine the two-photon exchange contributions.

2 Overview

The $n = 2$ energy levels of the muonic helium-3 ion are sketched in Figure 2. The helion has nuclear spin $I = 1/2$, just as the proton. Hence the level scheme is very similar to the one of muonic hydrogen. However, the helion magnetic moment $g = -2.127\,625\,308(25)$ [6] (here given in units of the nuclear magneton) is negative, which swaps the ordering of the hyperfine levels.

A note on the sign convention of the Lamb shift contributions used in this article: The 2S level is shifted below the 2P levels due to the Lamb shift. This means that, fundamentally, the 2S Lamb shift should be given a *negative* sign.

However, following long-established conventions we assign the *measured* $2S_{1/2} \rightarrow 2P_{1/2}$ energy difference a *positive* sign, i.e. $E(2P) - E(2S) > 0$. This is in accord with almost all publications we review here and we will mention explicitly when we have inverted the sign with respect to the original publications where the authors calculated level shifts.

Moreover, we obey the traditional definition of the Lamb shift as the terms beyond the Dirac equation and the leading order recoil corrections, i.e. excluding effects of the hyperfine structure. In particular, this means that the mixing of the hyperfine levels (Sect. 5) does *not* influence the Lamb shift.

The Lamb shift is dependent on the rms charge radius of the nucleus and is treated in Section 3. We split the Lamb

¹ 1 meV \cong 241.799 GHz.




Fig. 2. The 2S and 2P energy levels in the muonic helium-3 ion. The inset on the right displays the shift Δ of the 2P levels due to the mixing of levels with same quantum number F , as described in Section 5. The figure is not to scale.

shift contributions into *nuclear structure-independent* contributions and *nuclear structure-dependent* ones. The latter are composed out of one-photon exchange diagrams which represent the finite size effect and two-photon exchange diagrams which contain the polarizability contributions.

In Section 4, we treat the 2S hyperfine structure, which depends on the Zemach radius. It also has two-photon exchange contributions. However, these have not been calculated yet and can only be estimated with a large uncertainty.

In Section 5, we compile the 2P level structure which includes fine- and hyperfine splitting, and the mixing of the hyperfine levels [42].

For the theory compilation presented here, we use the calculations from many sources mentioned in the following. The names of the authors of the respective groups are ordered alphabetically.

The first source is E. Borie who was one of the first to publish detailed calculations of many terms involved in the Lamb shift of muonic atoms. Her most recent calculations for μp , μd , $\mu^4\text{He}^+$, and $\mu^3\text{He}^+$ are all found in her reference [43]. Several updated versions of this paper are available on the arXiv. In this work we always refer to [44] which is version-7, the most recent one at the time of this writing.

The second source is the group of Elekina, Faustov, Krutov, and Martynenko et al. (termed “Martynenko group” in here for simplicity). The calculations we use in here are found in Krutov et al. [45] for the Lamb shift, in Martynenko et al. [46,47] and Faustov et al. [48] for the 2S hyperfine structure, and Elekina et al. [49] for the 2P fine- and hyperfine structure.

Jentschura and Wundt calculated some Lamb shift contributions in their references [50,51]. They are referred to as “Jentschura” for simplicity.

The group of Ivanov, Karshenboim, Korzinin, and Sheyuto is referred to “Karshenboim group” for simplicity. Their calculations are found in Korzinin et al. [52] and in Karshenboim et al. [53] for Lamb shift and fine structure contributions.

The group of Bacca, Barnea, Hernandez, Ji, and Nevo Dinur, situated at TRIUMF and Hebrew University, has performed ab initio calculations on two-photon exchange contributions of the Lamb shift. Their calculations are found in Nevo Dinur et al. [14] and Hernandez et al. [40]. For simplicity we refer to them as “TRIUMF-Hebrew group”.

A recent calculation of the two-photon exchange using scattering data and dispersion relations has been performed by Carlson, Gorchtein, and Vanderhaeghen [41].

Item numbers # in our tables follow the nomenclature in references [3,5]. In the tables, we usually identify the “source” of all values entering “our choice” by the first letter of the (group of) authors given in adjacent columns (e.g. “B” for Borie). We denote as average “avg.” in the tables the center of the band covered by all values v_i under consideration, with an uncertainty of half the spread, i.e.

$$\text{avg.} = \frac{1}{2} [\text{MAX}(v_i) + \text{MIN}(v_i)] \pm \frac{1}{2} [\text{MAX}(v_i) - \text{MIN}(v_i)]. \quad (7)$$

If individual uncertainties are provided by the authors we add these in quadrature. We would like to point out that uncertainties due to uncalculated higher order terms are often not indicated explicitly by the authors. In the case some number is given, we include it in our sum. But in general our method cannot account for uncertainty estimates of uncalculated higher order terms.

Throughout the paper, Z denotes the nuclear charge with $Z = 2$ for the helion and alpha particle, α is the fine structure constant, $m_r = 199 m_e$ is the reduced mass of the muon-nucleon system. “VP” is short for “vacuum polarization”, “SE” is “self-energy”, “RC” is “recoil correction”. “Perturbation theory” is abbreviated as “PT”, and SOPT and TOPT denote 2nd and 3rd order perturbation theory, respectively.




Fig. 3. Item #1, the leading order 1-loop electron vacuum polarization (eVP), also called Uehling term.

3 Lamb shift in muonic helium-3

3.1 Nuclear structure-independent contributions

Nuclear structure-independent contributions have been calculated by Borie, Martynenko group, Karshenboim group, and Jentschura. The contributions are listed in Table 1, labeled with #*i*. The leading contribution is the one-loop electron vacuum polarization (eVP) of order $\alpha(Z\alpha)^2$, the so-called Uehling term (see Fig. 3). It accounts for 99.5% of the radius-independent part of the Lamb shift, so it is very important that this contribution is well understood. There are two different approaches to calculate this term.

Borie [44] (p. 4, Tab.) and the Karshenboim group [52] (Tab. I) use relativistic Dirac wavefunctions to calculate a relativistic Uehling term (item #3). A relativistic recoil correction (item #19) has to be added to allow comparison to nonrelativistic calculations (see below). Borie provides the value of this correction explicitly in [44] Table 6, whereas the Karshenboim group only gives the total value which includes the correction, thus corresponding to (#3 + #19).

Nonrelativistic calculations of the Uehling term (item #1) exist from the Martynenko group [45] (No. 1, Tab. 1) and Jentschura [51], which are in very good agreement. Additionally, a relativistic correction (item #2) has to be applied. This relativistic correction already accounts for relativistic recoil effects (item #19). Item #2 has been calculated by the Martynenko group [45] (No. 7+10, Tab. 1), Borie [44] (Tab. 1), Jentschura [50,51] (Eq. 17), and Karshenboim et al. [53], which agree well within all four groups, however do not have to be included in Borie's and Korzinin et al.'s value because their relativistic Dirac wavefunction approach already accounts for relativistic recoil effects.

Both approaches agree well within the required uncertainty. As *our choice* for the Uehling term with relativistic correction (#1 + #2) or (#3 + #19) we take the average

$$\Delta E(\text{Uehling} + \text{rel. corr.}) = 1642.3962 \pm 0.0018 \text{ meV.} \quad (8)$$

Item #4, the second largest contribution in this section, is the two-loop eVP of order $\alpha^2(Z\alpha)^2$, the so-called Källén-Sabry term [54] (see Fig. 4). It has been calculated by Borie [44] (p. 4, Tab.) and the Martynenko group [45] (No. 2, Tab. 1) which agree within 0.0037 meV. As *our choice* we take the average.




Fig. 4. Item #4, the two-loop eVP (Källén-Sabry) contribution. This is Figures 1b–1d from the Martynenko group [45].




Fig. 5. Item #5, the one-loop eVP in 2-Coulomb lines.

Item #5 is the one-loop eVP in two Coulomb lines of order $\alpha^2(Z\alpha)^2$ (see Fig. 5). It has been calculated by Borie [44] (Tab. 6), the Martynenko group [45] (No. 9, Tab. 1), and Jentschura [50] (Eq. 13) of whom the latter two obtain the same result, which differs from Borie by 0.0033 meV. As *our choice* we adopt the average.

The Karshenboim group [52] (Tab. I) has calculated the sum of item #4 and #5, the two-loop eVP (Källén-Sabry) and one-loop eVP in two Coulomb lines (Figs. 4 and 5). Good agreement between all groups is observed.

Item #6+7 is the third order eVP of order $\alpha^3(Z\alpha)^2$. It has been calculated by the Martynenko group [45] (No. 4 + 11 + 12, Tab. 1) and the Karshenboim group [52] (Tab. I). Borie [44] (p. 4) adopts the value from Karshenboim et al., Martynenko et al. and Karshenboim et al. differ by 0.004 meV, which is in agreement considering the uncertainty of 0.003 meV given by the Martynenko group. As *our choice* we adopt the average and obtain an uncertainty of 0.0036 meV via Gaussian propagation of uncertainty.

Item #29 is the second order eVP of order $\alpha^2(Z\alpha)^4$. It has been calculated by the Martynenko group [45] (No. 8 + 13, Tab. 1) and the Karshenboim group [52] (Tab. VIII). Their values did agree in the case of μd , however for $\mu^3\text{He}^+$ they differ by 0.004 meV. This difference is twice as large as the value from Martynenko et al. but this contribution is small, so the uncertainty is not at all dominating. We reflect the difference by adopting the average as *our choice*.

Items #9, #10, and #9a are the terms of the Light-by-light (LbL) scattering contribution (see Fig. 6). The sum of the LbL terms is calculated by the Karshenboim group [52] (Tab. I). Borie [44] also lists the value from Karshenboim et al. Item #9 is the *Wichmann-Kroll* term, or “1:3” LbL, which is of order $\alpha(Z\alpha)^4$. This item has also been calculated by Borie [44] (p. 4) and the Martynenko group [45] (No. 5, Tab. 1) who obtain the same result. Item #10 is the *virtual Delbrück* or “2:2” LbL, which is of order $\alpha^2(Z\alpha)^3$.




Fig. 6. The three contributions to Light-by-light scattering: (a) Wichmann-Kroll or “1:3” term, item #9, (b) Virtual Delbrück or “2:2” term, item #10, and (c) inverted Wichmann-Kroll or “3:1” term, item #9a[†].




Fig. 8. Item #11, muon self-energy corrections to the electron vacuum polarization $\alpha^2(Z\alpha)^4$. This figure is Figure 2 from Jentschura [55]. It corresponds to Figure 6(a) from Karshenboim [52].




Fig. 7. Item #20, the muon-self energy (a) and the muon vacuum polarization (b), $\alpha(Z\alpha)^4$.

Item #9a is the *inverted Wichmann-Kroll* term, or “3:1” LbL, which is of order $\alpha^3(Z\alpha)^2$. The sum of the latter two is also given by the Martynenko group [45] (No. 6, Tab. 1). As *our choice* we use the one from Karshenboim et al., who are the first and only group to calculate all three LbL contributions. The groups are in agreement when taking into account the uncertainty of 0.0006 meV given by Karshenboim et al.

Item #20 is the contribution from muon self-energy (μ SE) and muon vacuum polarization (μ VP) of order $\alpha(Z\alpha)^4$ (see Fig. 7). This item constitutes the third largest term in this section.² This item has been calculated by Borie [44] (Tabs. 2 and 6) and the Martynenko group [45] (No. 24, Tab. 1). They differ by 0.001 meV. As *our choice* we adopt the average.

Items #11, #12, #30, #13, and #31 are all corrections to VP or μ SE and of order $\alpha^2(Z\alpha)^4$.

Item #11 is the μ SE correction to eVP (see Fig. 8). It has been calculated by all four groups. Martynenko et al. calculate this term (Eq. 99) in [45], however in their table (No. 28) they use the more exact calculation from Jentschura. Jentschura [50] (Eq. 29), and the Karshenboim group [52] (Tab. VIII a) are in excellent agreement. Borie [44] (Tab. 16) differs significantly because she only calculates a part of this contribution in her App. C. This value does not enter her sum and thus is also not considered in here. On p. 12 of [44] she states that this value should be considered as an uncertainty. As *our choice* we adopt the number from Jentschura and Karshenboim et al.

² In ordinary hydrogen-like atoms this term is the leading order Lamb shift contribution: The leptons in the loop are the same as the orbiting lepton. This term can thus be rescaled from well-known results in hydrogen.




Fig. 9. Item #12, *eVP loop in SE* are radiative corrections with VP effects. This is Figure 11(b) from a publication by the Martynenko group [45] which is the same as Figure 4 in Pachucki [56]. It is Karshenboim’s Figure 6(d) in reference [52].




Fig. 10. Item #30, hadronic VP in SE contribution, corresponds to Figure 6(e) in Karshenboim et al.’s [52].

Item #12 is the eVP in μ SE (see Fig. 9). This item has been calculated by the Martynenko group [45] (No. 27, Tab. 1) and the Karshenboim group [52] (Tab. VIII d), which are in perfect agreement. On p. 10 of [44] Borie mentions that she included the “fourth order electron loops” in “muon Lamb shift, higher order” term, which is our item #21. As we include item #21 from Borie, we will not on top include item #12.

Item #30 is the hadronic vacuum polarization (hVP) in μ SE (see Fig. 10). This item has only been calculated by the Karshenboim group [52] (Tab. VIII e) which we adopt as *our choice*.

Item #13 is the mixed eVP + μ VP (see Fig. 11). The calculations from Borie [44] (p. 4) and the Martynenko group [45] (No. 3, Tab. 1) roughly agree, whereas the value from the Karshenboim group [52] (Tab. VIII b) is 0.002 meV larger. As *our choice* we take the average.




Fig. 11. Item #13, the mixed eVP- μ VP contribution.




Fig. 12. Item #31, the mixed eVP- and hadronic VP contribution, comes from the Uehling correction to the hadronic VP correction. See Figure 6(c) in Karshenboim et al.'s [52].




Fig. 13. Item #32, muon VP in SE contribution, is automatically included in a rescaled electronic ${}^3\text{He}^+$ QED value of higher order SE contributions (see text).

Item #31 is the mixed eVP + hVP (see Fig. 12) which has only been calculated by the Karshenboim group [52] (Tab. VIII c). We adopt their value as *our choice*.

Item #32, the muon VP in SE correction shown in Figure 13 is not included as a separate item in our Table 1. It should already be automatically included in the QED contribution which has been rescaled from the QED of electronic ${}^3\text{He}^+$ by a simple mass replacement $m_e \rightarrow m_\mu$ [57]. This is the case only for QED contributions where the particle in the loop is the same as the bound particle - like in this case, a muon VP correction in a muonic atom. The size of this item #32 can be estimated from the relationship found by Borie [58], that the ratio of hadronic to muonic VP is 0.66. With the Karshenboim group's value of item #30 [52] one would obtain a value for item #32 of $-0.0004/0.66 \text{ meV} = -0.0006 \text{ meV}$. This contribution is contained in our item #21, together with the dominating item #12 (see also p. 10 of Ref. [44]).

Item #21 is a higher-order correction to μSE and μVP of order $\alpha^2(Z\alpha)^4$ and $\alpha^2(Z\alpha)^6$. This item has only been calculated by Borie [44] (Tabs. 2 and 6). On p. 10 she

points out that this contribution includes the “fourth order electron loops”, which is our item #12. It also contains our item #32. We adopt her value as *our choice*.

Item #14 is the hadronic VP of order $\alpha(Z\alpha)^4$. It has been calculated by Borie [44] (Tab. 6) and the Martynenko group [45] (No. 29, Tab. 1). Borie assigns a 5% uncertainty to their value. However, in her reference [44] there are two different values of item #14, the first on p. 5 (0.219 meV) and the second in Table 6 on p. 16 (0.221 meV). Regarding the given uncertainty this difference is not of interest. In our Table 1, we report the larger value which is further from that of the Martynenko group in order to conservatively reflect the scatter. Martynenko et al. did not assign an uncertainty to their value. However, for μd [59] they estimated an uncertainty of 5%. As *our choice* we take the average of their values and adopt the uncertainty of 5% (0.011 meV).

Item #17 is the Barker-Glover correction [60]. It is a recoil correction of order $(Z\alpha)^4 m_r^3/M^2$ and includes the nuclear Darwin-Foldy term that arises due to the Zitterbewegung of the nucleus. As already discussed in App. A of [5], we follow the atomic physics convention [61], which is also adopted by CODATA in their report from 2010 [62] and 2014 [6]. This convention implies that item #17 is considered as a recoil correction to the energy levels and not as a part of the rms charge radius. This term has been calculated by Borie [44] (Tab. 6), the Martynenko group [45] (No. 21, Tab. 1), and Jentschura [51] and [50] (Eq. A.3). As *our choice* we use the number given by Borie and Jentschura as they give one more digit.

Item #18 is the term called “recoil, finite size” by Borie. It is of order $(Z\alpha)^5 \langle r \rangle_{(2)}/M$ and is linear in the *first* Zemach moment. It has first been calculated by Friar [63] (see Eq. F5 in App. F) for hydrogen and has later been given by Borie [44] for μd , $\mu^4\text{He}^+$, and $\mu^3\text{He}^+$. We discard item #18 because it is considered to be included in the elastic TPE [64,65]. It has also been discarded in μp [3], μd [5], and $\mu^4\text{He}^+$ [27]. For the muonic helium-3 ion, item #18 in [44] (Tab. 6) amounts to 0.4040 meV, which is five times larger than the experimental uncertainty of about 0.08 meV (see Eq. 5), so it is important that the treatment of this contribution is well understood.

Item #22 and #23 are relativistic recoil corrections of order $(Z\alpha)^5$ and $(Z\alpha)^6$, respectively. Item #22 has been calculated by Borie [44] (Tab. 6), the Martynenko group [45] (No. 22, Tab. 1), and Jentschura [50] (Eq. 32). They agree perfectly. Item #23 has only been calculated by the Martynenko group [45] (No. 23, Tab. 1) whose value we adopt as *our choice*.

Item #24 are higher order radiative recoil corrections of order $\alpha(Z\alpha)^5$ and $(Z^2\alpha)(Z\alpha)^4$. This item has been calculated by Borie [44] (Tab. 6) and the Martynenko group [45] (No. 25, Tab. 1). Their values differ by 0.015 meV. As *our choice* we adopt the average.

Item #28 is the radiative (only eVP) recoil of order $\alpha(Z\alpha)^5$. It consists of three terms which have been calculated by Jentschura and Wundt [50] (Eq. 46). We adopt their value as *our choice*. Note that a second value (0.0072 meV) is found in [51]. However, this value is just one of the three terms, namely the seagull term, and is already included in #28 (see [50], Eq. 46).

The total sum of the QED contributions without explicit nuclear structure dependence is summarized in Table 1 and amounts to

$$\Delta E_{r\text{-indep.}}^{\text{LS}} = 1644.3466 \pm 0.0146 \text{ meV.} \quad (9)$$

Note that Borie, on p.15 in reference [44] attributes an uncertainty of 0.6 meV to her total sum. The origin of this number remains unclear [66]. Its order of magnitude is neither congruent with the other uncertainties given in reference [44] nor with other uncertainties collected in our summary. Thus it will not be taken into account.

3.2 Nuclear structure contributions

Terms that depend on the nuclear structure are separated into one-photon exchange (OPE) contributions and two-photon exchange (TPE) contributions.

The OPE terms (also called *radius-dependent contributions*) represent the finite size effect which is by far the largest part of the nuclear structure contributions and are discussed in Section 3.2.1. They are parameterizable with a coefficient times the rms charge radius squared. These contributions are QED interactions with nuclear form factor insertions.

The TPE terms can be written as a sum of elastic and inelastic terms, where the latter describe the polarizability of the nucleus. These involve contributions from strong interaction and therefore are much more complicated to evaluate, which explains why the dominant uncertainty originates from the TPE part. The TPE contributions are discussed in more detail in Section 3.2.2.

The main nuclear structure corrections to the nS states have been given up to order $(Z\alpha)^6$ by Friar [63] (see Eq. (43a) therein)

$$\begin{aligned} \Delta E_{\text{fin.size}} &= \frac{2\pi Z\alpha}{3} |\Psi(0)|^2 \\ &\times \left(\langle r^2 \rangle - \frac{Z\alpha m_r}{2} \langle r^3 \rangle_{(2)} + (Z\alpha)^2 (F_{\text{REL}} + m_r^2 F_{\text{NREL}}) \right), \end{aligned} \quad (10)$$

where $\Psi(0)$ is the muon wave function at the origin, $\langle r^2 \rangle$ is the second moment of the charge distribution of the nucleus, i.e. the square of the rms charge radius, r_E^2 . $\langle r^3 \rangle_{(2)}$ is the Friar moment,³ and F_{REL} and F_{NREL} contain various moments of the nuclear charge distribution (see Eq. (43b) and (43c) in Ref. [63]). Analytic expressions for some simple model charge distributions are listed in App. E of reference [63].

As the Schrödinger wavefunction at the origin $\Psi(0)$ is nonzero only for S states, it is in leading order only the S states which are affected by the finite size. However, using the Dirac wavefunction a nonzero contribution appears for the $2P_{1/2}$ level [68]. This contribution affects the values

³ $\langle r^3 \rangle_{(2)}$ has been called “third Zemach moment” in [63]. To avoid confusion with the Zemach radius r_Z in the 2S hyperfine structure we adopt the term “Friar moment”, as recently suggested by Karshenboim et al. [67].




Fig. 14. Item #r1, the leading nuclear finite size correction stems from a one-photon interaction with a helion form factor insertion, indicated by the thick dot.

for the Lamb shift and the fine structure and is taken into account in the section below.

The Friar moment $\langle r^3 \rangle_{(2)}$ has not been included in μd [5] because of a cancellation [69–71] with a part of the inelastic nuclear polarizability contributions. The TRIUMF-Hebrew group pointed out [14,40], that in the case of $\mu^3\text{He}^+$ however, a smaller uncertainty might be achieved treating each term separately. This discussion is not finished yet and we will therefore continue with the more conservative treatment as before. See Section 3.2.2.

3.2.1 One-photon exchange contributions (finite size effect)

Finite size contributions have been calculated by Borie ([44] Tab. 14), the Martynenko group ([45] Tab. 1), and the Karshenboim group ([53] Tab. III). All of these contributions are listed in Table 2, labeled with # ri .

Most of the terms, given in Table 2, can be parameterized as $c \cdot r_E^2$ with coefficients c in units of meV fm^{-2} . Borie and Karshenboim et al. have provided the contributions in this parameterization, whereas Martynenko et al. provide the total value in units of energy. However, the value of their coefficients can be obtained by dividing their numbers by r_E^2 . The value they used for the charge radius r_E is 1.9660 fm^4 [73]. In this way the numbers from Martynenko et al. can be compared with the ones from the other groups.

Item #r1, the leading term of equation (10), is the one-photon exchange with a helion form factor (FF) insertion (see Fig. 14). Item #r1 is of order $(Z\alpha)^4 m_r^3$ and accounts for 99% of the OPE contributions. Borie ([44] Tab. 14, b_a), the Martynenko group ([45] No. 14), and the Karshenboim group ([53] Tab. III, $\Delta_{FNS}^{(0)}$) obtain the same result which we adopt as *our choice*. This contribution is much larger than the following terms, but its absolute precision is worse, which we indicate by introducing an uncertainty. For that we take the value from Borie which is given with one more digit than the values of the other authors and attribute an uncertainty of 0.0005 meV, which may arise from rounding.

Item #r2 and #r2' are the radiative correction of order $\alpha(Z\alpha)^5$. The equation used for the calculation of item #r2 is given in equation (10) of [74]. It has been calculated by Borie [44] (Tab. 14, b_b) and the Martynenko group [45] (No. 26, only Eq. (92)). Note that the value from the Martynenko group was published with a wrong sign.

⁴ This value has been introduced by Borie [44] as an average of several previous measurements [23,24,72].

Table 1. All known **nuclear structure-independent** contributions to the Lamb shift in $\mu^3\text{He}^+$. Values are in meV. Item numbers “#” in the 1st column follow the nomenclature of references [3,5], which in turn follow the supplement of reference [1]. Items “#” with a dagger † were labeled “New” in reference [3], but we introduced numbers in reference [5] for definiteness. For Borie [44] we refer to the most recent arXiv version-7 which contains several corrections to the published paper [43] (available online 6 Dec. 2011). For the Martynenko group, numbers #1 to #29 refer to rows in Table I of reference [45]. Numbers in parentheses refer to equations in the respective paper.

#	Contribution	Borie (B) [44]	Martynenko group (M) Krutov et al. [45]	Jentschura (J) Jentschura, Wundt [50] Jentschura [51]	Karshenboim group (K) Karshenboim et al. [53] Korzinin et al. [52]	Our choice Value	Source Fig.
1	NR one-loop electron VP (eVP)		1641.8862 #1	1641.885 [51]		1642.3962 ± 0.0018	avg 3
2	Rel. corr. (Breit-Pauli)	(0.50934) ^a Tab. 1	0.5093 #7+#10	0.509344 [50](17), [51]	(0.509340) [53] Tab. IV		
3	Rel. one-loop eVP	1642.412 Tab. p. 4				11.4089 ± 0.0019	avg. 4
19	Rel. RC to eVP, $\alpha(Z\alpha)^4$	-0.0140 Tab. 1+6				1.6757 ± 0.0017	avg. 5
	Sum of the above	1642.3980 3+19	1642.3955 1+2	1642.3943 1+2	1642.3954 [52] Tab. I	1642.3962 ± 0.0018	avg 3
4	Two-loop eVP (Källén-Sabry)	11.4107 Tab. p. 4	11.4070 #2				
5	One-loop eVP in 2-Coulomb lines $\alpha^2(Z\alpha)^2$	1.674 Tab. 6	1.6773 #9	1.677290 [50](13)			
	Sum of 4 and 5	13.0847 4+5	13.0843 4+5		13.0843 [52] Tab. I	(13.0846) ^b	
6+7	Third order VP	0.073(3) p. 4	0.0689 #4+#12+#11		0.073(3) [52] Tab. I	0.0710 ± 0.0036	avg.
29	Second-order eVP contribution $\alpha^2(Z\alpha)^4 m$		0.0018 #8+#13		0.00558 [52] Tab. VIII “eVP2”	0.0037 ± 0.0019	avg
9	Light-by-light “1,3”: Wichmann-Kroll	-0.01969 p. 4	-0.0197 #5				6a
10	Virtual Delbrück, “2,2” LbL		} 0.0064 #6				6b
9a†	“3,1” LbL						6c
	Sum: Total light-by-light scatt.		-0.0134(6) p. 5+Tab. 6		-0.0134(6) [52] Tab. I	-0.0134 ± 0.0006	K
20	μ SE and μ VP	-10.827368 Tab. 2+6	-10.8286 #24			-10.8280 ± 0.0006	avg.
11	Muon SE corr. to eVP $\alpha^2(Z\alpha)^4$	(-0.1277) ^c Tab. 16	-0.0627 #28	-0.06269 [50](29)	-0.06269 [52] Tab. VIII (a)	-0.06269	J, K
12	eVP loop in self-energy $\alpha^2(Z\alpha)^4$	incl. in 21	-0.0299 #27		-0.02992 [52] Tab. VIII (d)	incl. in 21	B
30	Hadronic VP loop in self-energy $\alpha^2(Z\alpha)^4 m$				-0.00040(4) [52] Tab. VIII (e)	-0.00040 ± 0.00004	K 10
13	Mixed eVP + μ VP	0.00200 p. 4	0.0022 #3		0.00383 [52] Tab. VIII (b)	0.0029 ± 0.0009	avg 11
31	Mixed eVP + hadronic VP	-0.033749 Tab. 2+6			0.0024(2) [52] Tab. VIII (c)	0.0024 ± 0.0002	K 12
21	Higher-order corr. to μ SE and μ VP	-0.031749 13+21	-0.0277 12+13			-0.033749	B
	Sum of 12, 30, 13, 31, and 21				-0.0241(2) 12+30+13+31	-0.0288	sum
14	Hadronic VP	0.221(11) Tab. 6	0.2170 #29			0.219 ± 0.011	avg.
17	Recoil corr. $(Z\alpha)^4 m_r^3/M^2$ (Barker-Glover)	0.12654 Tab. 6	0.1265 #21	0.12654 [50](A.3) [51](15)		0.12654	B, J
18	Recoil, finite size	(0.4040(10)) ^d					
22	Rel. RC $(Z\alpha)^5$	-0.55811 p. 9+Tab. 6	-0.5581 #22	-0.558107 [50](32)		-0.558107	J
23	Rel. RC $(Z\alpha)^6$		0.0051 #23			0.0051	M
24	Higher order radiative recoil corr.	-0.08102 p. 9+Tab. 6	-0.0656 #25			-0.0733 ± 0.0077	avg.
28†	Rad. (only eVP) RC $\alpha(Z\alpha)^5$			0.004941		0.004941	J
	Sum		1644.3431			1644.3466 ± 0.0146	

^a Does not contribute to the sum in Borie’s approach.

^b Sum of our choice of item #4 and #5, written down for comparison with the Karshenboim group.

^c In App. C of [44], incomplete. Does not contribute to the sum in Borie’s approach, see text.

^d Is not included, because it is a part of the TPE, see text.

^e Including item #18 and #r3† yields 1644.9169 meV, which is Borie’s value from reference [44] page 15. On that page she attributes an uncertainty of 0.6 meV to that value. This number is far too large to be correct, so we ignore it.

Very recently the Martynenko group updated their calculation of higher-order finite size corrections [75] using more realistic, measured nuclear form factors. The results contain a coefficient (in our work termed item #r2) which agrees with the old value, and an additional, previously unknown term which cannot be parametrized with r_h^2 and therefore is given as a constant. This constant is found in our Table 2 as item #r2'. In reference [75] the values are given for the 1S state but can easily be transferred to the 2S state via the $1/n^3$ scaling. For the 2S state this results in

$$\begin{aligned} & 1/8 \times (-0.6109) \text{ meV} \\ &= 1/8 \times (-0.1946 r_h^2 - 0.1412) \text{ meV} \quad (11) \\ &= -0.0243 \text{ meV/fm}^2 r_h^2 - 0.0177 \text{ meV}. \end{aligned}$$

Borie and Martynenko get the same result for item #r2, which we adopt as *our choice*. Additionally we adopt the constant term from Martynenko as item #r2'.

Item #r3 and #r3' are the finite size corrections of order $(Z\alpha)^6$. They have first been calculated in reference [63]. Item #r3 and #r3' consider third-order perturbation theory in the finite size potential correction and relativistic corrections of the Schrödinger wave functions. There are also corrections in the TPE of the same order $(Z\alpha)^6$, but these are of different origin. Borie [44] (Tab. 14, b_c and Tab. 6) and the Martynenko group [45] (Eq. (91)) follow the procedure in reference [63] and then separate their terms into a part with an explicit r_E^2 dependence (item #r3) and another one which is usually evaluated with an exponential charge distribution, since a model independent calculation of this term is prohibitively difficult [44]. Differences in sorting the single terms have already been noticed in the μd case [5], where we mentioned that e.g. the term $\langle r^2 \rangle \langle \ln(\mu r) \rangle$ in F_{REL} of equation 10 is attributed to #r3 and #r3' by Martynenko et al. and Borie, respectively. The difference in this case amounts to 0.007 meV for #r3'. Note that in equation (91) from the Martynenko group [45], the charge radius has to be inserted in units of GeV^{-1} , with $r_E = 1.966 \text{ fm} \cong 9.963 \text{ GeV}^{-1}$.

Item #r4 is the one-loop eVP correction (*Uehling*) of order $\alpha(Z\alpha)^4$. It has been calculated by all three groups, Borie [44] (Tab. 14, b_d), Martynenko et al. [45] (No. 16, Eq. (69)), and Karshenboim et al. [53] (Tab. III, $\Delta E_{FNS}^{(2)}$). On p. 31 of [44], Borie notes that she included the correction arising from the Källén-Sabry potential in her b_d . This means that her value already contains item #r6, which is the two-loop eVP correction of order $\alpha^2(Z\alpha)^4$. Item #r6 has been given explicitly only by the Martynenko group [45] (No. 18, Eq. 73). The sum of Martynenko et al.'s #r4 and #r6 differs by 0.016 meV/fm^2 from Borie's result. Using a charge radius of 1.9660 fm this corresponds to roughly 0.06 meV and, hence, causes the largest uncertainty in the radius-dependent OPE part. The origin of this difference is not clear [66,76]. A clarification of this difference is desired but does not limit the extraction of the charge radius. As *our choice* we take the average of the sum (#r4+#r6) of these two groups. The resulting average does also reflect the value for #r4 provided by Karshenboim et al. [53].

Item #r5 is the one-loop eVP correction (*Uehling*) in second order perturbation theory (SOPT) of order $\alpha(Z\alpha)^4$. It has been calculated by all three groups, Borie [44] (Tab. 14, b_e), the Martynenko group [45] (No. 17, Eq. 70), and the Karshenboim group [53] (Tab. III, $\Delta E_{FNS}^{(1)}$). On p. 31 of [44], Borie notes that she included the two-loop corrections to ϵ_{VP2} in her b_e . This means that her value already contains item #r7, which is the two-loop eVP in SOPT of order $\alpha^2(Z\alpha)^4$. Item #r7 has only been given explicitly by the Martynenko group [45] (No. 19). The sum of Martynenko et al.'s #r5+#r7 differs by 0.003 meV from Borie's result. As *our choice* we take the average of the sum (#r5+#r7) of these two groups. Again here, *our choice* reflects the value for #r5 provided by Karshenboim et al. [53], too.

Item #r8 is the finite size correction to the $2P_{1/2}$ level of order $(Z\alpha)^6$. It has only been calculated by Borie [44] (Tab. 14, $b(2p_{1/2})$). This correction is the smallest in this section and is the only term which affects the $2P_{1/2}$ level. In consequence, the effect on the Lamb shift is inverse, i.e. if the $2P$ level is lifted "upwards", the Lamb shift gets larger. Thus, in contrast to Borie, we include this correction with a positive sign. At the same time this term decreases the fine structure ($2P_{3/2} - 2P_{1/2}$ energy difference) and is hence listed in Table 4 as item #f10 with a negative sign.

The total sum of the QED contributions with an explicit dependence of r_E^2 is summarized in Table 2 and amounts to

$$\begin{aligned} & \Delta E_{\text{r-dep.}}^{\text{LS}}(r_E^2) \\ &= -103.5184(98) \text{ meV fm}^{-2} r_E^2 \\ &+ 0.1354(33) \text{ meV}. \quad (12) \end{aligned}$$

3.2.2 Two-photon exchange contributions to the Lamb shift

Historically, the two-photon exchange (TPE) contribution to the Lamb shift (LS) in muonic atoms has been considered the sum of the two parts displayed in Figures 15a,b and 15c,d, respectively:

$$\Delta E_{\text{TPE}}^{\text{LS}} = \Delta E_{\text{Friar}}^{\text{LS}} + \Delta E_{\text{inelastic}}^{\text{LS}} \quad (13)$$

with the elastic "Friar moment" contribution $\Delta E_{\text{Friar}}^{\text{LS}}$ ⁵ and the inelastic part $\Delta E_{\text{inelastic}}^{\text{LS}}$, frequently termed "polarizability".

The elastic part, $\Delta E_{\text{Friar}}^{\text{LS}}$ is shown in Figures 15a and 15b. It is sensitive to the shape of the nuclear charge distribution, beyond the leading $\langle r^2 \rangle$ dependence discussed in Section 3.2.1. This part is traditionally parameterized as being proportional to the third power of the rms charge radius and it already appeared in equation (10) as the second term proportional to $\langle r^3 \rangle_{(2)}$. The coefficient depends on the assumed radial charge distribution.

⁵ formerly known as "third Zemach moment", see footnote³ on p. 7 for disambiguation.

Table 2. Coefficients of the **nuclear structure-dependent** one-photon exchange (OPE) contributions to the Lamb shift of $\mu^3\text{He}^+$. The values from the Martynenko group shown here are the published ones divided by $(1.9660\text{ fm})^2$, which is the radius they used. The numbers $\#i$ from the Martynenko group refer to rows in Table 1 of reference [45] and numbers in parenthesis to equations therein. KS: Källén-Sabry, VP: vacuum polarization, SOPT: second-order perturbation theory. Values are in meV/fm^2 , except for $\#r2'$ and $\#r3'$.

#	Contribution	Borie (B)		Martynenko group (M)		Karshenboim group (K)		Our choice		Source
		Borie [44]	Tab. 14	Krutov et al. [45]	Faustov et al. [75]	Karshenboim et al. [53]	Value	Value		
r1	Leading fin. size corr., $(Z\alpha)^4$	-102.520	b_a	-102.52	#14, (61)	-102.52	$\Delta E_{FNS}^{(0)}$	-102.520	± 0.0010	B,M,K
r2	Radiative corr., $\alpha(Z\alpha)^5$	-0.0243 ^a	b_b	-0.0243 ^b	#26, (92)	-0.0243		-0.0243		B,M
r3	Finite size corr. order $(Z\alpha)^6$	-0.1275	b_c	-0.1301	#26, (91)	-0.1288		-0.1288	± 0.0013	avg.
r4	Uehling corr. (+KS), $\alpha(Z\alpha)^4$			-0.3310	#16, (69)	-0.323	$\Delta E_{FNS}^{(2)}$			
r6	Two-loop VP corr., $\alpha^2(Z\alpha)^4$	-0.3176	b_d	-0.0026	#18, (73)				± 0.0080	avg.
sum	r4+r6			-0.3336				-0.3256		
r5	One-loop VP in SOPT, $\alpha(Z\alpha)^4$			-0.5196	#17, (70)	-0.520	$\Delta E_{FNS}^{(1)}$			
r7	Two-loop VP in SOPT, $\alpha^2(Z\alpha)^4$			-0.0063	#19 ^c					
sum	r5+r7		b_e	-0.5259		-0.5238		-0.5238	± 0.0021	avg.
r8	Corr. to the $2P_{1/2}$ level	0.00409 ^d	$b(2p_{1/2})$			0.00409		0.00409		B
	Sum of coefficients	-103.507(5) ^e		-103.5339		-103.37	ΔE_{FNS}	-103.5184	$\pm 0.0098^f$	
r2'	Rad. corr. $\alpha(Z\alpha)^5$ [meV] ^g			0.0177 meV	[75]			0.0177 meV		M
r3'	Remaining order $(Z\alpha)^6$ [meV] ^h	0.121 meV	Tab. 6	0.11445 meV	(91)			0.1177 meV	± 0.0033	avg.
Sum		-103.507 $r_h^2 + 0.121$ meV		-103.5339 $r_h^2 + 0.1322$ meV		-103.37 r_h^2		-103.5184(98) $r_h^2 + 0.1354(33)$ meV		

^a Borie uses equation (10) of [74] to calculate this term. For further explanations, see text.

^b The value in equation 92 of [45] was published with a wrong sign.

^c This term is represented by Figure 9(a,b,c,d) from the Martynenko group [45]. This figure includes equation (76) therein.

^d The sign is explained in the text.

^e The summed coefficient is given in reference [44] on p. 15, where Borie indicates the uncertainty of 0.005 meV.

^f This uncertainty is the one obtained from averaging the above values (0.0084 meV) and the one given by Borie in her sum of (0.005 meV) added in quadrature.

^g Belongs to $\#r2$. Not parametrizable with r_h^2 .

^h Belongs to $\#r3$. Depends on the charge distribution in a non-trivial way, see text.




Fig. 15. (a)+(b) Elastic $\Delta E_{\text{Friar}}^{\text{LS}}$, and (c)+(d) inelastic $\Delta E_{\text{inelastic}}^{\text{LS}}$ two-photon exchange (TPE) contribution. The thick dots in (a) indicate helion form factor insertions. The blob in (c) and (d) represents all possible excitations of the nucleus.

The inelastic part, $\Delta E_{\text{inelastic}}^{\text{LS}}$ is shown in Figures 15c and 15d. It stems from virtual excitations of the nucleus. The inelastic contributions are notoriously the least well-known theory contributions and limit the extraction of the charge radius from laser spectroscopy of the Lamb shift.

Equation (13) is valid for the nuclear contributions as well as for the nucleon contributions. This means that elastic and inelastic parts have to be evaluated for both, respectively.

The nuclear parts of $\Delta E_{\text{TPE}}^{\text{LS}}$ are then given as $\delta E_{\text{Friar}}^A$ and $\delta E_{\text{inelastic}}^A$ for a nucleus with A nucleons, and the nucleon parts as $\delta E_{\text{Friar}}^N$ and $\delta E_{\text{inelastic}}^N$.

With that, the total (nuclear and nucleon) TPE is given as⁶

$$\Delta E_{\text{TPE}}^{\text{LS}} = \delta E_{\text{Friar}}^A + \delta E_{\text{Friar}}^N + \delta E_{\text{inelastic}}^A + \delta E_{\text{inelastic}}^N. \quad (14)$$

We refer here to two calculations of the TPE contributions. The first stems from the TRIUMF-Hebrew group, who perform ab initio calculations using two different nuclear potentials. They have published two papers on the TPE in muonic helium-3 ions: Detailed calculations are given in Nevo Dinur et al. [14], and updated results are found in Hernandez et al. [40]. The second calculation has been performed by Carlson et al. [41], who obtain the TPE from inelastic structure functions via dispersion relations.

The two calculations are very different, so that comparisons of any but the total value may be inexact [41]. An attempt to compare the different approaches is given in Table II of reference [41]. Here, we want to refer to this table only and later compare the total values as suggested. Note that we proceed differently to our previous compilation for μd [5] (Tab. 3), where we listed and compared 16

⁶ Compared to the notation of the TRIUMF-Hebrew group [14], the terms in equation (14) correspond to δZ_{em}^A , δZ_{em}^N , $\delta \rho_{\text{pol}}^A$, and $\delta \rho_{\text{pol}}^N$, respectively.

individual terms (labeled #p1...16) which together yield the sum of the four terms of equation (14).

The nuclear Friar moment contribution is calculated by the TRIUMF-Hebrew group to be $\delta E_{\text{Friar}}^A = 10.49(24)$ meV [14,40]. Previous values have been given by Borie [44] (10.258(305) meV) and Krutov et al., [45] (10.50(10) meV)⁷ using a Gaussian charge distribution and assuming an rms radius of 1.966(10) fm. These uncertainties do not include the (rather large) dependence of the calculation on the charge distribution [36,45]. This type of uncertainty is gauged within the ab-initio calculation of [14] by using two different state-of-the-art nuclear potentials. We therefore use the more recent value provided by the TRIUMF-Hebrew group. Their value also agrees with a value of 10.87(27) meV which is obtained in [14] from the third Zemach moment $\langle r^3 \rangle_{(2)} = 28.15(70) \text{ fm}^3$ that was extracted from electrons scattering off ^3He by Sick [36].

The nuclear polarizability contribution from the TRIUMF-Hebrew group is $\delta E_{\text{inelastic}}^A = 4.16(17)$ meV [14, 40]. The first calculation of the nuclear polarizability contribution in $\mu^3\text{He}^+$ has been published in 1961 [77]. The recent value from the TRIUMF-Hebrew group replaces a former one of 4.9 meV from Rinker [78] which has been used for more than 40 years now.

As mentioned before, the total TPE contribution has a nuclear part and a nucleon part. The nucleon Friar moment contribution from the TRIUMF-Hebrew group amounts to $\delta E_{\text{Friar}}^N = 0.52(3)$ meV. They obtain this value using $\delta E_{\text{Friar}}^N(\mu p) = 0.0247(13)$ meV from μp and scale it according to equation (17) in reference [14]. This procedure has also been done in [5] for μd .⁸ $\delta E_{\text{Friar}}^N(\mu p)$ is a sum of the elastic term (0.0295(13) meV) and the non-pole term (−0.0048 meV) which have been obtained by Carlson et al. in reference [79].

The nucleon polarizability contribution from the TRIUMF-Hebrew group amounts to $\delta E_{\text{inelastic}}^N = 0.28(12)$ meV. It is obtained using the proton polarizability contribution from μp and scaling it with the number of protons and neutrons,⁹ as well as with the wavefunction overlap, according to equation (19) of reference [14]. Furthermore it is corrected for estimated medium effects and possible nucleon-nucleon interferences. The proton polarizability contribution used here amounts to 0.0093(11) meV and is the sum of an inelastic term (0.00135 meV [81]) and the proton subtraction term $\delta_{\text{subtraction}}^p = -0.0042(10)$ meV which has been calculated for muonic hydrogen in reference [82].

⁷ Sum of 10.28(10) meV and 0.2214(22) meV, which correspond to line 15 and 20 from Table 1 in reference [45], respectively.

⁸ In equation (12) of reference [5], we used a scaling of the nucleon TPE contribution by the reduced mass ratio to the third power, which is only correct for $\delta E_{\text{inelastic}}^N$. $\delta E_{\text{Friar}}^N$ should be scaled with the fourth power [14,71]. This is due to an additional m_r scaling factor compared to the proton polarizability term. This mistake has no consequences for μd yet, as the nuclear uncertainty is much larger, but the correct scaling is relevant for $\mu^3\text{He}^+$ and $\mu^4\text{He}^+$.

⁹ Assuming isospin symmetry, the value of the neutron polarizability contribution used in [14] is the same as the one of the proton, but an additional uncertainty of 20% is added, motivated by studies of the nucleon polarizabilities [80].

Summing up all nuclear and nucleon contributions evaluated by the TRIUMF-Hebrew group [14,40] yields a total value of the $\Delta E_{\text{TPE}}^{\text{LS}}$ of [14,40]

$$\begin{aligned} \Delta E_{\text{TPE}}^{\text{LS}}(\text{nuclear potentials}) & \\ &= \delta E_{\text{Friar}}^A + \delta E_{\text{Friar}}^N + \delta E_{\text{inelastic}}^A + \delta E_{\text{inelastic}}^N \\ &= 15.46(39) \text{ meV}.^{10} \end{aligned} \quad (15)$$

Recently, Carlson et al. [41] have also calculated the TPE in $\mu^3\text{He}^+$. Their result of

$$\Delta E_{\text{TPE}}^{\text{LS}}(\text{dispersion relations}) = 15.14(49) \text{ meV} \quad (16)$$

is in agreement with the one from the TRIUMF-Hebrew group. As *our choice* we take the average of equations (15) and (16) and remain with

$$\Delta E_{\text{TPE}}^{\text{LS}} = 15.30(52) \text{ meV}. \quad (17)$$

As conservative uncertainty we use the larger one (from Eq. (16)) and add in quadrature half the spread. A weighted average of the two values (Eqs. (15) and (16)) which would reduce the total uncertainty is not adequate as certain contributions are effectively fixed by the same data [83].

3.3 Total Lamb shift in $\mu^3\text{He}^+$

Collecting the radius-independent (mostly) QED contributions listed in Table 1 and summarized in equation (9), the radius-dependent contributions listed in Table 2 and summarized in equation (12), and the complete TPE contribution $\Delta E_{\text{TPE}}^{\text{LS}}$ from equation (17), we obtain for the $2S \rightarrow 2P$ energy difference in $\mu^3\text{He}^+$

$$\begin{aligned} \Delta E(2S_{1/2} \rightarrow 2P_{1/2}) &= 1644.3466(146) \text{ meV} \\ &+ 0.1354(33) \text{ meV} - 103.5184(98)r_h^2 \text{ meV/fm}^2 \\ &+ 15.3000(5200) \text{ meV} \\ &= 1659.78(52) \text{ meV} - 103.518(10)r_h^2 \text{ meV/fm}^2, \end{aligned} \quad (18)$$

where in the last step we have rounded the values to reasonable accuracies.

One should note that the uncertainty of 0.52 meV from the nuclear structure corrections $\Delta E_{\text{TPE}}^{\text{LS}}$, equation (17), is about 30 times larger than the combined uncertainty of all radius-independent terms summarized in Table 1, and 13 times larger than the uncertainty in the coefficient of the r_h^2 -dependent term (which amounts to 0.038 meV for $r_h = 1.966 \text{ fm}$). A further improvement of the two-photon exchange contributions in light muonic atoms is therefore strongly desirable.

4 2S hyperfine splitting

The 2S hyperfine splitting (HFS) in muonic helium-3 ions has been calculated by Borie [44] and Martynenko [47].

¹⁰ As explained in the introduction, we use a different sign convention, which explains the minus sign in references [14,40].

(There is also the more recent paper [46] from Martynenko et al., but it is less detailed and reproduces all numbers from [47], with one exception to be discussed for #h27.) The values are summarized in Table 3 and labeled with #hi.

We also adapted the ordering according to increasing order/complexity of the terms and grouped them thematically as: Fermi energy with anomalous magnetic moment and relativistic corrections discussed in Section 4.1, vacuum polarization and vertex corrections in Section 4.2, nuclear structure contributions and corrections listed in Section 4.3, and the weak interaction contribution in Section 4.4.

4.1 Fermi energy with muon anomalous magnetic moment and Breit corrections

4.1.1 h1 and h4 Fermi energy and muon AMM correction

Item #h1 is the Fermi energy ΔE_{Fermi} which defines the main splitting of the $2S$ hyperfine levels. Borie and the Martynenko group have both calculated the Fermi energy, however, their values disagree by 0.055 meV (see Tab. 3). For the calculation Borie uses equation (13) in her reference [44]. Martynenko uses equation (6) in his reference [47]. The Fermi energy is calculated using fundamental constants only. Thus we repeated the calculation for both equations, the one from Borie and the one from Martynenko which resulted to be the same: Both equations yield the same result, as they should, which is

$$\Delta E_{\text{Fermi}} = \frac{8(\alpha^4 Z^3)m_r^3}{3n^3 m_\mu m_p} \mu_h = -171.3508 \text{ meV}, \quad (19)$$

where m_μ is the muon mass, m_p is the proton mass, m_r is the reduced mass, and μ_h is the helion magnetic moment to nuclear magneton ratio of $\mu_h = -2.127\,625\,308(25)$ [6]. We use the value in equation (19) as *our choice*. This value agrees neither with Borie's value (-171.3964 meV) nor with the one from the Martynenko group (-171.341 meV).

The value for the Fermi energy corrected for the muon anomalous magnetic moment (AMM) a_μ is then also updated to

$$\Delta E_{\text{Fermi,AMM}} = \Delta E_{\text{Fermi}} \cdot (1 + a_\mu) = -171.5506 \text{ meV} \quad (20)$$

with a correction of -0.1998 meV .

All further corrections from Borie given as coefficients ϵ , are applied to this value analogous to

$$\Delta E_{\text{Fermi,AMM}} \cdot (1 + \epsilon). \quad (21)$$

Note, that in Table 3, for the contributions given by Borie, we use her coefficients but apply them to our value of the Fermi Energy given in equation (20). The value for the Fermi Energy in equation (20) is evaluated to a precision of 0.0001 meV. If the number of significant digits from Borie's coefficients is too small to yield this precision we attribute a corresponding uncertainty. For example item #h28* has the coefficient $\epsilon_{2\gamma} = 0.0013$; here the coefficient is only precise up to a level of 0.00005, which we include

as uncertainty. This uncertainty is propagated upon multiplication with the Fermi energy (Eq. (20)) and then yields 0.0086 meV.

4.1.2 h2 Relativistic Breit correction

Item #h2 is the relativistic Breit correction of order $(Z\alpha)^6$. It is given congruently by both authors as $\Delta E_{F,\text{rel}}^B = -0.0775 \text{ meV}$ and $\Delta E_{F,\text{rel}}^M = -0.078 \text{ meV}$, respectively. We take the number from Borie as *our choice*, which is given with one more digit and attribute an uncertainty of 0.0001 meV due to the precision in her coefficient.

4.2 Vacuum polarization and vertex corrections

4.2.1 h8 and h9: Electron vacuum polarization in a one-photon one-loop interaction (h8) and in a one-photon two-loop interaction (h9)

The Feynman diagrams corresponding to #h8 and #h9 are analogous to those shown in Figures 3 and 4, respectively, and constitute the analogs to the Uehling- and Källén-Sabry contributions in the Lamb shift. #h8 is of order $\alpha(Z\alpha)^4$, #h9 is of order $\alpha^2(Z\alpha)^4$.

Borie calculates the main electron VP contribution (“by modification of the magnetic interaction between muon and nucleus”), which is a one-photon one-loop interaction. It amounts to a correction $\epsilon_{VP1} = 0.00315$, which results in an energy shift of -0.5405 meV (#h8). She also gives $\epsilon_{VP1} = 2.511 \cdot 10^{-5}$ for one-photon two-loop interactions, resulting in -0.0043 meV (#h9). These terms are evaluated on p.21 of her document [44], using her equation (16).

Martynenko calculates these contributions to be -0.540 meV and -0.004 meV , respectively. These values are found in the table in reference [47].

Martynenko mentions that his value for our item #h9 consists of his equations (15,16). The numerical result from equation (15) corresponds to two separate loops (see our Fig. 4a) and is given as -0.002 meV , whereas equation (16) describes the two nested two-loop processes where an additional photon is exchanged within the electron VP loop (see our Figs. 4b and 4c). One can conclude that its numerical value is also -0.002 meV .

Both authors give congruent results within their precisions, as *our choice* we write down the numbers by Borie which are given with one more digit. We attribute an uncertainty to item #h8 due to the precision in Borie’s coefficient.

4.2.2 h5 and h7: Electron vacuum polarization in SOPT in one loop (h5) and two loops (h7)

Items #h5 and #h7 are the SOPT contributions to items #h8 and #h9, respectively.

Borie’s value for our item #h5 is given by the coefficient $\epsilon_{VP2} = 0.00506$ and her value for our item #h7 by $\epsilon_{VP2} = 3.928 \cdot 10^{-5}$. This results in energy shifts of $-0.8680(9) \text{ meV}$ and -0.0067 meV , respectively (those values are for point nuclei; the finite size correction is

taken into account in our #h25 and #h26). The uncertainty in item #h5 originates from the precision of ϵ_{VP2} .

The corresponding values from Martynenko are -0.869 meV (#h5) and -0.010 meV (#h7).

Due to slight differences between the two authors, as *our choice* we take the average of items #h5 and #h7, respectively. The uncertainty of item #h5 is the above uncertainty and half the spread between both authors added in quadrature.

4.2.3 h13 and h14: Vertex correction ($\hat{=}$ self energy happening at the muon-photon vertex)

Item #h13 is the muon self-energy contribution of order $\alpha(Z\alpha)^5$ (it is the analogue to a part of item #20 in the Lamb shift, see Fig. 7a). It has only been calculated by Borie as

$$\epsilon_{\text{vertex}} = \alpha(Z\alpha) \left(\ln 2 - \frac{5}{2} \right) = -0.9622 \cdot 10^{-4} \cdot Z. \quad (22)$$

Its numerical value is thus 0.0330 meV, however this includes a muon VP contribution of -0.0069 meV (#h12, see Sect. 4.2.4). For our item #h13, we use the value from Borie as *our choice*. We therefore should not include #h12, which is discussed later.

Borie also cites a higher order correction of Brodsky and Erickson [84] which results in a correction of $-0.211 \cdot 10^{-4} \hat{=} -0.0036 \text{ meV}$ (#h14). Very probably the sign of the energy shift is not correct because the coefficient is negative, but the Fermi energy of helium-3 also has a negative sign, thus the energy shift should be positive. (The analogous contributions in muonic hydrogen and deuterium are negative, which is a further hint to a wrong sign since the helium-3 Fermi energy is negative, contrary to hydrogen and deuterium.)

4.2.4 h12: Muon VP and muon VP SOPT

Item #h12 is the one-loop muon vacuum polarization. Borie on p.19 (below the equation of ϵ_{vertex}) of reference [44] gives the coefficient as $0.3994 \cdot 10^{-4} \cdot Z$. In combination with the Fermi energy this yields -0.0069 meV . Martynenko obtains a value of -0.007 meV which is congruent to Borie’s value. However, Borie’s value of this contribution is already included in our item #h13, which has been discussed in the previous section. Hence, we do not include it separately in ‘our choice’.

4.2.5 h18 Hadronic vacuum polarization

Item #h18 is the hadronic vacuum polarization. Borie gives this contribution as $\epsilon_{hVP} = 0.2666 \cdot 10^{-4} \cdot Z$, which amounts to -0.0091 meV on p. 19 of her paper. This contribution is analogous to our Figure 3, but with a hadronic loop in the photon line. Since Martynenko does not provide a value for hadronic VP in muonic helium-3 ions, we use Borie’s value as ‘our choice’.

4.3 Nuclear structure and finite size corrections

Analogously to Section 3.2, we categorize the nuclear structure contributions to the 2S HFS as one-photon exchange (OPE) and two-photon exchange (TPE) processes, respectively. We list first the by far dominant contribution to nuclear structure: the Zemach term, which is an elastic TPE process. The following subsections describe the known elastic TPE corrections in the 2S HFS. So far, to our knowledge there are yet no calculations with respect to the *inelastic* TPE contribution to the 2S HFS. Thus we only give a simplified estimate with a large uncertainty. Later the section is concluded with the one-photon exchange (OPE) corrections to nuclear structure in the 2S HFS.

4.3.1 h20 Zemach term and h23, h23b*, h28* nuclear recoil

Item #h20 is the elastic TPE and the main finite size correction to the 2S HFS. This correction arises due to the extension of the magnetization density (Bohr-Weisskopf effect) and is also called the Zemach term [85]. The Zemach term is usually parameterized as [86]

$$\Delta E_{\text{Zemach}}^{\text{HFS}} = -\Delta E_{\text{Fermi,AMM}} 2(Z\alpha)m_r r_Z \quad (23)$$

with m_r being the reduced mass and r_Z the Zemach radius of the nucleus [36]

$$r_Z = -\frac{4}{\pi} \int_0^\infty [G_E(q)G_M(q) - 1] \frac{dq}{q^2}. \quad (24)$$

Here, $G_E(q)$ and $G_M(q)$ are the electric and magnetic form factors of the nucleus, respectively.

The corresponding coefficient to the Fermi energy in equation (23) is given by Borie on p. 23 of [44] as

$$\epsilon_{\text{Zem}} = -2(Z\alpha)m_r r_Z = -0.01506 \text{ fm}^{-1} r_Z. \quad (25)$$

With our Fermi energy from equation (20), item #h20 is

$$\Delta E_{\text{Zemach}}^{\text{HFS}} = 2.5836 r_Z \text{ meV/fm} = 6.5312(413) \text{ meV}, \quad (26)$$

where, in the second step, we inserted the most recent Zemach radius from Sick [36] ($r_Z = 2.528(16) \text{ fm}$).

Note that Borie's published value of $\Delta E_{\text{Zemach}}^{\text{HFS}}$ differs from the one given here, because she uses a different Zemach radius of $r_Z = 2.562 \text{ fm}$, assuming a Gaussian charge distribution.

Martynenko, in his reference [47], gives a value of $\Delta E_{\text{str}}^{\text{HFS}} = 6.047 \text{ meV}$. This value contains a recoil contribution and is thus not directly comparable with our item #h20. However, this value has been updated [73] and is now available as two separate values of $\Delta E_{\text{str}}^{\text{HFS}} = 6.4435 \text{ meV} = (6.4085 + 0.0350_{\text{recoil}}) \text{ meV}$. The first can be compared to equation (26). The second is the recoil correction and listed in our table as item #h23. Martynenko notes [47] that changing from a Gaussian to a dipole parameterization results in a change of the final number of 2%.




Fig. 16. (a) Item #h15, μ SE contribution to the elastic two-photon exchange; (b) item #h16 the vertex correction to the elastic two-photon exchange, which results in two terms (the vertex correction can take place either at one or the other photon); and (c) item #h17, spanning photon contribution to the elastic two-photon exchange, also referred to as jellyfish diagram.

Regarding our item #h20, we do not consider the respective value from Martynenko because it is model-dependent and therefore carries a large uncertainty. This uncertainty can be avoided using the model-independent Zemach radius from Sick and the coefficient given by Borie as stated above.

A new contribution which hasn't been calculated for μp and μd is our item #h23b*. It is an additional recoil contribution which amounts to 0.038 meV . It has only been calculated by Martynenko and we adopt his value as *our choice*. In order to account for the precision given by Martynenko, we write $0.0380(5) \text{ meV}$.

Another contribution which has not been calculated for μp and μd is item #h28*. It is a two-photon recoil correction, calculated by Borie in 1980 [87], who followed the procedure of Grotch and Yennie [88]. This contribution is not listed in Borie's recent reference [44], but should be included [89]. It is given by $\epsilon_{2\gamma} = 0.0013$ and therefore results in $-0.2230(86) \text{ meV}$, using our Fermi energy from equation (20). The attributed uncertainty originates from the number of significant digits in $\epsilon_{2\gamma}$ (the value of the coefficient is considered to be accurate only to ± 0.00005). Regarding the contributions given by Martynenko, no overlap is found, which is why we list this item separately.

4.3.2 h24 electron VP contribution to two-photon exchange

Item #h24, the electron VP contribution to the 2S HFS elastic two-photon exchange in muonic helium-3 ions is only calculated by Martynenko [47]. The corresponding Feynman diagrams are shown in Figure 4 of his helium 2S HFS paper [47]. These are analogous to our Figure 15, but with a VP loop in one of the exchange photons. A numerical value of the contribution is given in his equation (38) of 0.095 meV and thus enters *our choice*, where we write $0.0950(5) \text{ meV}$ and therefore account for the precision given by Martynenko.

4.3.3 h15, h16, h17 radiative corrections to the elastic two-photon exchange

Items #h15, #h16, and #h17 are radiative corrections to the elastic two-photon exchange in the 2S hyperfine structure and represented in Figure 16. They are partially given

in Martynenko's reference [47], but have been updated [76] and result to be -0.0101 meV (#h15), 0.0333 meV (#h16), and 0.0074 meV (#h17). These numbers include recoil corrections and are based on equations (24)–(27) from the Martynenko group [48] and use a dipole parameterization of the helion form factor, as well as $r_h = 1.966$ fm. For the moment, we will adapt these preliminary numbers including recoil considerations into *our choice*.

4.3.4 h22 inelastic two-photon exchange in the hyperfine structure

In contrast to the Lamb shift, no calculations are available for the inelastic two-photon exchange (polarizability contribution) in the 2S HFS. We give an estimate of this value by calculating the ratio between the polarizability contribution and the Zemach term in the 1S ground state of (electronic) ${}^3\text{He}^+$ and assume the ratio to be similar for the 2S state in $\mu^3\text{He}^+$.

The 1S Zemach term for electronic ${}^3\text{He}^+$ is found by using equation (23), but with the muon mass replaced by the electron mass and $n = 1$. Using the Zemach radius r_Z from Friar and Payne [90] a value of 1717 kHz is obtained. In order to obtain the total sum (polarizability + Zemach) of 1442 kHz [90], a polarizability term of order -300 kHz is missing. The ratio is then roughly $-1/6$. The Zemach term for muonic helium-3 ions (our item h20), obtained above, yields $\Delta E_{Zem} \approx 6.5$ meV. The estimate for the polarizability contribution consequently follows with $\Delta E_{pol}^{HFS} \approx -1.0 \pm 1.0$ meV, which includes a conservative 100% uncertainty.

4.3.5 h25 and h26 finite size correction to electron VP

Borie gives the electron VP contributions #h8 and #h5 (eVP processes in OPE, see Section 4.2) which are based on a point nucleus. Additionally, she provides modified contributions which include the finite size effect on electron VP. These are $\epsilon'_{VP1} = 0.00295$ and $\epsilon'_{VP2} = 0.00486$, respectively. The difference between those values and #h8 and #h5 constitute finite size corrections. Multiplied with the Fermi energy (including the AMM), these yield $0.0343(9)$ meV each and we attribute them to #h25 and #h26, analogous to the previous CREMA summaries. The uncertainty originates from the precision in Borie's coefficients. Note that these are OPE processes.

4.3.6 h27 and h27b nuclear structure correction in leading order and SOPT

This correction is only given by Martynenko. The two terms are found in Figure 5a,b of reference [47], for leading and second order, respectively. This correction is also an OPE process. Care has to be taken here because this contribution is given as 0.272 meV in [47], but as 0.245 meV in a 2010 follow up paper [46] (however, this is the only term that changed between [47] and [46]). As compared to muonic deuterium, Martynenko only gives the sum (h27 + h27b) and not the single contributions. In [47] the formulas he uses to calculate h27 and h27b

are explicitly given as

$$\Delta E_{1\gamma, str}^{HFS} = -\frac{4}{3}(Z\alpha)^2 m_r^2 r_M^2 \cdot E_{\text{Fermi}} \cdot \frac{1 - n^2}{4n^2} \quad (27)$$

$$\begin{aligned} \Delta E_{str, SOPT}^{HFS}(2S) \\ = \frac{4}{3}(Z\alpha)^2 m_1^2 r_E^2 \cdot E_{\text{Fermi}}(2S) \cdot (\ln(Z\alpha) - \ln 2), [-5pt] \end{aligned} \quad (28)$$

where m_r is the reduced mass of the muon, m_1 is the muon mass, and r_E and r_M are the charge and magnetic radii, respectively. Martynenko states to use $r_E \approx r_M = 1.844 \pm 0.045$ fm which is known to be outdated.

However, inserting Martynenko's Fermi energy, the radius he used, and fundamental constants into equations (27) and (28) yields a sum of 0.2251 ± 0.0001 meV which is neither congruent with [47] nor [46].

Using Sick's 2014 values [36] for the charge and magnetic radii yields 0.2577 ± 0.0001 meV.

In the course of some private communications with Martynenko, he provided us his most current value of 0.2421 meV for the sum of h27 + h27b, and we use this preliminarily as *our choice*.

4.4 h19 weak interaction

The contribution of the weak interaction to the 2S HFS of helium-3 is only given by Borie. She cites Eides [91] and provides $\epsilon_{\text{weak}} = 1.5 \cdot 10^{-5} \hat{=} -0.0026$ meV, which we adopt as *our choice*.

4.5 Total 2S HFS contribution

In total, the 2S HFS contributions are given by

$$\begin{aligned} \Delta E^{HFS}(2S_{1/2}^{F=1} - 2S_{1/2}^{F=0}) &= -172.7457(89) \text{ meV} \\ &\quad + 2.5836 \text{ meV/fm } r_Z + \Delta E_{pol}^{HFS} \\ &= -166.2145(423) - 1.0(1.0) \text{ meV} \\ &= -167.2(1.0) \text{ meV.} \end{aligned} \quad (29)$$

Here, in the first line, we separated out the Zemach contribution and the estimate of the polarizability contribution. In the second line, the Zemach radius $r_Z = 2.528(16)$ fm [36] is inserted and the estimated value of ΔE_{pol}^{HFS} is shown. The polarizability is the dominant source of uncertainty in the hyperfine structure and prevents a precise determination of the helion Zemach radius from the measured transitions in the muonic helium-3 ion [92]. A calculation of the polarizability contribution is therefore highly desirable. Until then a precise measurement of the 1S or 2S HFS in muonic helium-3 ions can be used to experimentally determine a value of the polarizability contribution ΔE_{pol}^{HFS} . In essence, the measurement of the 2S HFS by the CREMA collaboration can be used to give the total TPE contribution to the HFS, $\Delta E_{TPE}^{HFS} = 2.5836$ meV/fm $r_Z + \Delta E_{pol}^{HFS}$ with an expected uncertainty of 0.1 meV.

Table 3. All contributions to the **2S hyperfine splitting (HFS)**. The item numbers *hi* in the first column follow the entries in Table 3 of reference [3]. However, the terms are now sorted by increasing complexity, analogous to their order in the text. For Martynenko, numbers #1 to #13 refer to rows in Table I of his reference [47], whereas numbers in parentheses refer to equations therein. Borie [44] gives the values as coefficients ϵ to be multiplied with the sum of (h1+h4) of ‘our choice’ values. We list the resulting values in meV. AMM: anomalous magnetic moment, PT: perturbation theory, VP: vacuum polarization, SOPT: second order perturbation theory, TOPT: third order perturbation theory. All values are in meV. Values in brackets do not contribute to the total sum.

	Contribution	Borie (B)		Martynenko group (M)		Our choice	
		Borie [44]	Value	Martynenko [47]	Value	Source	
h1	Fermi splitting, $(Z\alpha)^4$		-171.341	#1, (6)	-171.3508 ^a		
h4	μ AMM corr., $\alpha(Z\alpha)^4$	(-0.1999)	-0.200	#2, (7)	-0.1998		
sum	(h1+h4)	-171.5963	(-171.541)				
h2	Breit corr., $(Z\alpha)^6$	-0.0775 ± 0.0001	-0.078	#3, (8)	-0.0775 ± 0.0001	B	
h8	One-loop eVP in OPE, $\alpha(Z\alpha)^4$ (ϵ VP1)	-0.5404 ± 0.0009	-0.540	#4, (12)	-0.5404 ± 0.0009	B	
h9	Two-loop eVP in OPE, $\alpha^2(Z\alpha)^4$ (ϵ VP1)	-0.0043	-0.004	#5, (15,16)	-0.0043	B	
h5	One-loop eVP in OPE, SOPT, $\alpha(Z\alpha)^4$ (ϵ VP2)	-0.8680 ± 0.0009	-0.869	#7, (24)	-0.8685 ± 0.0010	avg.	
h7	Two-loop eVP in OPE, SOPT, $\alpha^2(Z\alpha)^4$ (ϵ VP2)	-0.0067	-0.010	#8, (29,30)	-0.0084 ± 0.0017	avg.	
h13	Vertex, $\alpha(Z\alpha)^5$	0.0330			0.0330	B	
h14	Higher order corr. of (h13), part with $\ln(\alpha)$	0.0036 ^b			0.0036	B	
h12	one-loop μ VP in 1γ int., α^6	(-0.0069) incl. in h13	-0.007	#6, (12)	incl. in h13	B	
h18	Hadronic VP, α^6	-0.0091			-0.0091	B	
h20	Fin. size (Zemach) corr. to ΔE_{Fermi} , $(Z\alpha)^5$	6.5312 ^c (=2.5836 r_Z /fm)	6.4085 (± 0.1) ^d	priv.comm.	2.5836 r_Z /fm	B	
h23	Recoil of order $(Z\alpha)(m_1/m_2)\ln(m_1/m_2)E_F$		0.0350	priv.comm.	0.0350	M	
h23b*	Recoil of order $(Z\alpha)^2(m_1/m_2)E_F$		0.038	#13, (48)	0.0380 ± 0.0005	M	
h28*	Two-photon recoil	-0.2230 ± 0.0086		[87]	-0.2230 ± 0.0086	B	
h24	eVP in two-photon-exchange, α^6		0.095	#10, (38)	0.0950 ± 0.0005	M	
h15	muon self energy contribution in TPE, w/recoil		-0.0101	priv.comm.	-0.0101	M	
h16	vertex correction contribution in TPE, w/recoil		0.0333	priv.comm.	0.0333	M	
h17	jelly fish correction contribution in TPE, w/recoil		0.0074	priv.comm.	0.0074	M	
h22a	Helion polarizability, $(Z\alpha)^5$						
h22b	Helion internal polarizability, $(Z\alpha)^5$						
sum	(h22a+h22b)				(-1.0 ± 1.0) ^e		
h25	eVP corr. to fin. size in OPE (sim. to ϵ VP2)	0.0343 ± 0.0009			$0.0343^f \pm 0.0009$	B	
h26	eVP corr. to fin. size in OPE (sim. to ϵ VP1)	0.0343 ± 0.0009			0.0343 ± 0.0009	B	
h27+h27b	Nucl. struct. corr. in SOPT, $\alpha(Z\alpha)^5$		0.2421	priv.comm.	0.2421	M	
h19	Weak interact. contr.	-0.0026 ± 0.0001			-0.0026 ± 0.0001	B	
Sum		-166.6988 ^g	-165.1998 ^h		-172.7457 ± 0.0089 2.5836 r_Z/fm -1.0 ± 1.0		

^a calculated in this work and given in equation (19).

^b The sign from Borie is wrong and has been corrected here, see Section 4.2.3.

^c Calculated by combining Borie’s coefficient with Sick’s r_Z .

^d This uncertainty reflects the change in this contribution when moving from dipole parameterization to a Gaussian one.

^e Is a preliminary estimate, see text. It is therefore listed separately in the sum below.

^f Difference of two terms in Borie [44], see also Section 4.3.5.

^g Borie’s sum given in this table differs from her published one of -166.3745 meV [44]. This is because we used an updated value of the Fermi energy (see Sect. 4.1.1), a different value for the Zemach radius r_Z (see Sect. 4.3.1), and included item #h28* which has not been considered in reference [44].

^h Martynenko’s sum given in this table is different from the (superseded) published one of -166.615 meV [47] because several terms have been changed and added upon private communication.

5 2P levels

5.1 2P fine structure

Fine structure (FS) contributions have been calculated by Borie [44] (Tab. 7), the Martynenko group [49] (Tab. 1), and the Karshenboim group [53] (Tab. 4) and [52] (Tab. 9). All of these contributions are listed in Tab. 4 and labeled with #fi.

The leading fine structure contribution of order $(Z\alpha)^4$ has been calculated by Borie using the Dirac wavefunctions (same as in Lamb shift). Her result (our item #f1) has to be corrected by a recoil term (item #f2) in order to be compared with the result from the Martynenko group. They use a nonrelativistic approach (our item #f3) and then add relativistic corrections (our item #f4a+b). Their total results differ by 0.02 meV. We take the average as *our choice* and remain with an uncertainty of 0.01 meV. This is by far the dominant uncertainty in the 2P fine structure.

Item #f5a and #f5b are the one-loop eVP of order $\alpha(Z\alpha)^4$ in leading order and SOPT. Item #f13* is the one-loop eVP contribution of order $\alpha^2(Z\alpha)^4$ in SOPT. All three items are given individually by the Martynenko group [49] in lines 5, 7, and 9 of their Tab. 1. In Tab. 7 of [44], Borie's term "Uehling(VP)" presumably contains all these three items. Karshenboim et al. [53] (Tab. 4) also calculate the sum of these items. All agree within 0.0009 meV and we take the average as *our choice* which coincides with Borie's value.

Item #f6a and #f6b are the two-loop eVP (*Källén-Sabry*) contribution of order $\alpha^2(Z\alpha)^4$ in leading order and SOPT. These terms have been calculated by Martynenko et al. [49] (Tab. 1, line 10+11 and 12+13, respectively). Borie [44] and the Karshenboim group [52] (Tab. IX) only calculated our item #f6b. We therefore adopt the value provided by the Martynenko group for item #f6a and the Karshenboim group's value of #f6b as they included some higher order terms as well.

Items #f7a, #f7d, and #f7e are of order $\alpha^2(Z\alpha)^4$ and have been calculated with high accuracy by the Karshenboim group [52] (Tab. IX). They correspond to the same Feynman diagrams as the Lamb shift items #11, #12, and #30, shown in Figures 8, 9, and 10, respectively. We adopt the values from the Karshenboim group as *our choice*.

Item #f11* is a contribution of order $\alpha(Z\alpha)^6$ which has been calculated by Martynenko et al. [49] (Tab. 1, line 8). Item #f12* is the one-loop μVP of order $\alpha(Z\alpha)^4$ which has been calculated by the Martynenko group as well [49] (Tab. 1, line 6). We adopt both of these values as *our choice*.

The sum of items #f8 and #f9 is the muon anomalous magnetic moment (AMM) contribution of order $(Z\alpha)^4$. These items are labeled by Borie [44] as "second order" and "higher orders", respectively. Martynenko et al. [49] (Tab. 1, line 2) provide the sum of these. Both groups agree very well. As *our choice* we adopt the average.

Item #f10 is the finite size correction to the $2P_{1/2}$ level of order $(Z\alpha)^6$ which has only been calculated by Borie [44]. It is the same correction which appears in the radius dependent part of the Lamb shift as #r8, with opposite sign and evaluated with a helium charge radius of

1.966(10) fm [44]. We adopt Borie's value as *our choice* and add the uncertainty which we obtain from the given charge radius.

The total sum of the FS contributions is summarized in Table 4 and amounts to

$$\Delta E_{\text{FS}} = 144.7993 \text{ meV} \pm 0.0101 \text{ meV}. \quad (30)$$

It will enter the calculation of the 2P hyperfine structure in the following section. Note, that the uncertainty originates only from differences in the treatment of Dirac term (sum of items #f1 to #f4).

5.2 2P hyperfine structure

The 2P hyperfine splitting is described by the Breit Hamiltonian. Off-diagonal terms appear in the matrix representation of this Hamiltonian in the basis of $2P_{1/2}^{F=1}$, $2P_{1/2}^{F=0}$, $2P_{3/2}^{F=2}$, and $2P_{3/2}^{F=1}$. These terms lead to a mixing of energy levels with same quantum number F (see Fig. 2). This has first been calculated by Brodsky and Parsons [42] for hydrogen and later has also been evaluated for muonic hydrogen by Pachucki [56]. In previous publications [3,5], we also discussed the mixing of hyperfine states.

The traditional way [42,56] is to calculate the FS (without perturbations from the HFS F state mixing) and then include the so obtained FS in the evaluation of the Breit matrix. The centroids of the diagonal elements are now the virtual levels $2P_{1/2}$ and $2P_{3/2}$. *Afterwards* the mixing is included (via diagonalization) which means that the actual centroid is not at the position of the virtual levels anymore.

The 2P hyperfine structure has been calculated by Borie [44] (Tab. 9) and Martynenko et al. [49] (Tab. 2). We also calculated the splittings following Pachucki [56], who did the evaluation for μp . The values which are listed in our Table 5 are not the published values, but the values which result when including our FS value from Section 5.1.

Borie in her Table 9 lists the energies of the four 2P hyperfine levels relative to the $2P_{1/2}$ fine structure state where she already included the F state mixing. We reproduced her results using equations given in her Table 9 and then inserted our ΔE_{FS} from our equation (30). The result is listed in the second column of Table 5. Borie mentions, she used the shielded helion magnetic moment, whereas the (unshielded) magnetic moment should be used. The change, however, appears only on the seventh digit and is therefore negligible.

In their Table 2, Martynenko et al. provide the total splittings of the hyperfine structure levels, and at the end of their Section 3, they list the term $\Delta = 0.173$ meV originating from the mentioned F state mixing. In order to include this term, the numbers in their Table 2 first have to be divided according to the weight given by the number of m_F states. Δ has then to be added to the two $F = 1$ states. Furthermore, for the $2P_{3/2}$ states, we add our ΔE_{FS} . The result is listed in the third column of our Table 5.

Additionally, following Pachucki [56], we repeat his calculations in μp for $\mu^3\text{He}^+$. The off-diagonal elements are

Table 4. Contributions to the **2P fine structure**. Items # with an asterisk * denote new contributions in this compilation. The items #f7a, #f7d, and #f7e originate from the same graphs as the Lamb shift items #11, #12, and #30, respectively. VP: vacuum polarization, AMM: anomalous magnetic moment, KS: Källén-Sabry. All values are in meV.

#	Contribution	Borie (B) Borie [44] Tab. 7	Martynenko group (M) Elekina et al. [49] Tab. 1	Karshenboim group (K) Karshenboim et al. [53] Korzinin et al. [52]	Our choice
f1	Dirac	144.4157			
f2	Recoil	-0.1898			
f3	Contrib. of order $(Z\alpha)^4$	144.18648	1.1		
f4a	Contrib. of order $(Z\alpha)^6$	0.01994	1.3		
f4b	Contrib. of order $(Z\alpha)^6 m^2/M$	-0.00060	1.4		
sum	(f1+f2) or (f3+f4)	144.2259	144.20582		144.2159 \pm 0.0100 avg.
f5a	eVP corr. (Uehling), $\alpha(Z\alpha)^4$		0.12925		
f5b	eVP corr. SOPT, $\alpha(Z\alpha)^4$		0.14056		
f13*	eVP corr. SOPT, $\alpha^2(Z\alpha)^4$		0.00028		
sum	f5+f13*	0.2696	0.27009	0.26920 [53] Tab. IV	0.2696 \pm 0.0004 avg.
f6a	two-loop eVP corr. (KS), $\alpha^2(Z\alpha)^4$		0.00098		0.0010 M
f6b	two-loop eVP in SOPT, $\alpha^2(Z\alpha)^4$	0.0021	0.00234	0.00242 [52] Tab. IX “eVP2”	0.0024 K
f7a	$\alpha^2(Z\alpha)^4 m$, like #11			0.000606 [52] Tab. IX (a)	0.0006 K
f7d	$\alpha^2(Z\alpha)^4 m$, like #12			0.00164 [52] Tab. IX (d)	0.0016 K
f7e	$\alpha^2(Z\alpha)^4 m$, like #30*			0.000019(2) [52] Tab. IX (e)	0.0000 K
f11*	$\alpha(Z\alpha)^6$		-0.00055		-0.0006 M
f12*	one-loop μ VP, $\alpha(Z\alpha)^4$		0.00001		0.0000 M
f8	AMM (second order)	0.3232			
f9	AMM (higher orders)	0.0012			
sum	Total AMM (f8+f9)	0.3244	0.32446		0.3244 avg.
f10	Finite size, $(Z\alpha)^{6a}$	-0.0158			-0.0158 \pm 0.0002 B
Sum		144.8062	144.80315		144.7993 \pm 0.0101

^aThis is item #r8, evaluated for a helion radius of 1.966(10) fm [44], see text. The uncertainty is propagated from the charge radius, but is negligible.

Table 5. 2P levels from fine- and hyperfine splitting. All values are in meV relative to the $2P_{1/2}$ level. The columns labeled with Borie and Martynenko include their HFS calculations, but our value of the fine structure ($2P_{3/2}-2P_{1/2}$ energy splitting) $\Delta E_{\text{FS}} = 144.7993(101)$ meV from equation (30). The column ‘following [56]’ is calculated in this work following the treatment of Pachucki for $\mu^3\text{He}^+$, also including our value of the fine structure. Uncertainties arise from differences between the published values and from the uncertainty in the fine structure value ΔE_{FS} .

	Borie [44]	Martynenko [49]	Following [56]	Our choice
$2P_{1/2}^{\text{F}=1}$	-14.7877	-14.8080	-14.7990	-14.7979(102)
$2P_{1/2}^{\text{F}=0}$	43.8458	43.9049	43.8797	43.8754(296)
$2P_{3/2}^{\text{F}=2}$	135.7580	135.7552	135.7527	135.7554(27)(101) _{FS}
$2P_{3/2}^{\text{F}=1}$	160.0410	160.0459	160.0494	160.0452(42)(101) _{FS}

given by equation (85) of [56]

$$\begin{aligned} & \langle 2P_{1/2}^{\text{F}=1} | V | 2P_{3/2}^{\text{F}=1} \rangle \\ &= \frac{1}{3} (Z\alpha)^4 \frac{m_r^3}{m_\mu m_h} (1 + \kappa) \left(1 + \frac{m_\mu}{m_h} \frac{1 + 2\kappa}{1 + \kappa} \right) \left(-\frac{\sqrt{2}}{48} \right), \end{aligned} \quad (31)$$

where we included the correct Z scaling. m_r is the reduced mass of the muonic helium-3 ion, m_μ (m_h) is the mass of the muon (helion), and $\kappa = -4.18415^{11}$ is the helion anomalous magnetic moment. The diagonal terms are given by equation (86) therein

$$\begin{aligned} & E_{\text{HFS}}(2P_{1/2}) \\ &= \frac{1}{3} (Z\alpha)^4 \frac{m_r^3}{m_\mu m_h} (1 + \kappa) \left(\frac{1}{3} + \frac{a_\mu}{6} + \frac{1}{12} \frac{m_\mu}{m_h} \frac{1 + 2\kappa}{1 + \kappa} \right) \end{aligned} \quad (32)$$

$$\begin{aligned} & E_{\text{HFS}}(2P_{3/2}) \\ &= \frac{1}{3} (Z\alpha)^4 \frac{m_r^3}{m_\mu m_h} (1 + \kappa) \left(\frac{2}{15} - \frac{a_\mu}{30} + \frac{1}{12} \frac{m_\mu}{m_h} \frac{1 + 2\kappa}{1 + \kappa} \right) \end{aligned} \quad (33)$$

with the anomalous magnetic moment of the muon $a_\mu = 1.165\,920\,89(63) \times 10^{-3}$ [6].

Furthermore, Pachucki adds corrections due to vacuum polarization in his equations (89) and (90). With correct Z scaling these are

$$\delta E_{\text{HFS}}(2P_{1/2}) = \frac{1}{3} (Z\alpha)^4 \frac{m_r^3}{m_\mu m_h} (1 + \kappa) \cdot 0.00022 \quad (34)$$

¹¹ The helion anomalous magnetic moment is obtained using the respective equation on p.17 of Borie’s reference [44], where this magnitude is denoted as κ_2 .

$$\delta E_{\text{HFS}}(2P_{3/2}) = \frac{1}{3} (Z\alpha)^4 \frac{m_r^3}{m_\mu m_h} (1 + \kappa) \cdot 0.00008. \quad (35)$$

They have to be added to equations (32) and (33), respectively. Diagonalizing the matrix given in equation (91) of reference [56] with entries determined by the above equations yields the values given as *our choice* in Table 5. The diagonalization yields an F mixing of $\Delta = 0.1724$ meV. In the same manner as for the sections above, *our choice* in Table 5 takes into account the spread of values from the different authors and additionally the uncertainty of our value of the fine structure which we obtained in Section 5.1. It is astonishing that the splitting of the $2P_{1/2}$ states differs by as much as 0.04 meV between Borie and Martynenko. These states do not overlap with the nucleus, so it should be possible to determine them to much better precision. A precise calculation of these splittings is therefore highly welcome.

6 Summary

We have compiled all available contributions necessary to extract a charge radius of the helion from the Lamb shift measurement in muonic helium-3 ions, performed by the CREMA collaboration.

The total of the Lamb shift contributions are summarized in equation (18).

The nuclear structure-independent contributions of the Lamb shift, given in Table 1, show good agreement within the four (groups of) authors. The uncertainty is dominated by the hadronic VP (#14) and higher order radiative recoil corrections (#24). The total uncertainty in Table 1, however, is in the order of 0.01 meV and therefore sufficiently good (see also Eq. (5)).

The nuclear structure-dependent part of the Lamb shift completely dominates the theoretical uncertainties. The one-photon exchange (finite size) contributions, where the coefficients are given in Table 2, have an uncertainty which corresponds to 0.04 meV, which already is above

the “ideal” precision, mentioned in the introduction. This uncertainty is dominated by a disagreement in the terms #r4 and #r6. The much larger uncertainty, however, stems from the two-photon exchange contributions (TPE), given in equation (17). Recently, two groups have published new calculations on the TPE with a precision of about 3% (~ 0.5 meV). In terms of the helion charge radius this uncertainty corresponds to about

$$\sigma_{\text{theory}}(r_h) \approx \pm 0.0013 \text{ fm.} \quad (36)$$

The expected experimental uncertainty will be about an order of magnitude smaller. Thus, improving the theoretical uncertainty directly improves the extraction of the charge radius.

Isotope shift measurements generally benefit from cancellations of theory contributions that limit the absolute charge radii [11,24]. For the present case of the muonic helium isotope shift it will be useful to exploit possible correlations between the nuclear and nucleon structure contributions, which dominate the total uncertainty of the muonic radii. The correlations could lead to a reduction of the uncertainty of the muonic isotope shift determination and shed light on the 4σ discrepancy in the electronic isotope shift measurements, see Figure 1. A further investigation of these correlations is therefore desired.

The total of the 2S HFS contributions are given in Table 3 and summarized in equation (29). The uncertainty in the 2S HFS is completely dominated by the polarizability contribution, where no calculation exists. We have given a very rough estimate. The second largest uncertainty in the 2S HFS originates from the Zemach radius term (Bohr-Weisskopf effect). The upcoming results of the CREMA experiment will be able to extract a value for the TPE in the 2S hyperfine splitting (sum of polarizability and Zemach radius contribution) from measured data. In this case the uncertainty will be limited by the experimental uncertainty.

For the 2P levels, we collect all fine structure terms from the various authors (Tab. 4) which are then used to calculate the hyperfine structure by means of the Breit matrix. The results are compared with two other groups (Tab. 5). Here, the largest uncertainty originates from the leading order contributions (#f1 to #f4) in the fine structure (which is still sufficiently good) and from differing published values of the $2P_{3/2}$ splitting. A clarification of this difference would be very welcome.

Note added in proof: After this manuscript was accepted for publication, a paper by Karshenboim et al. [93] about the Lamb shift theory in muonic helium and tritium was published. They discuss the 2S-2P Lamb shift and the 2P fine- and hyperfine structure. The 2S hyperfine structure is not treated therein. The comparison of their values with ours has to be done carefully because Karshenboim et al. treat the mixing of the hyperfine levels (Brodsky Parsons contribution) differently. In their work the mixing is added as a perturbation to the fine structure. The traditional way, however, is to use the unperturbed fine structure and add the mixing as a perturbation to the

hyperfine levels, which is what we do. Comparing the values one therefore has to subtract/add the Brodsky Parsons term printed in bold italic in [93]. Furthermore Karshenboim et al. neglect some known higher order terms and increase the uncertainty due to estimates of non-listed higher order contributions. The comparison with the values in Ref. [93] yields the following (the numbers shown here are *adapted* to the traditional treatment of the Brodsky Parsons contribution): For the radius-independent QED Lamb shift without TPE, Karshenboim et al. obtain a value of $1644.35(2)$ meV which is in very good agreement with ours (Eq. (9)). In order to compare the radius-dependent (finite size) part we use a helion charge radius of 1.966 fm [44]. The value of Karshenboim et al. is then $-399.69(23)^{\text{theo}}$ meV which differs by $0.33(23)$ meV (1.4σ) from our value of $-400.02(4)^{\text{theo}}$ meV. This difference is the largest between our values and the ones from Karshenboim et al. For the 2P fine structure, Karshenboim et al. obtain a value of $144.800(5)$ meV $- 0.004r_h^2/fm^2$ meV which differs by 0.0142 meV (1.3σ) from ours. Regarding the $2P_{1/2}$ hyperfine structure, the value from Karshenboim et al. of $-58.7150(7)$ meV differs by 0.0417 meV (1.3σ) and has by far the smaller uncertainty. In our case the uncertainty arises from the huge difference between Borie and Martynenko. The $2P_{3/2}$ splitting of $-24.2925(7)$ meV agrees very well with our value. However, all these differences are considerably smaller than the uncertainty of the two-photon contribution which we assumed to be 0.52 meV while Karshenboim et al. increase it to 0.86 meV. The final result for the charge radius will therefore not be changed significantly.

We are grateful to E. Borie and A.P. Martynenko for insightful comments and for providing us with previously unpublished results. We thank M. Gorchtein and N. Nevo Dinur for helpful discussions about the two-photon exchange in muonic helium-3 ions and the treatment of the Friar moment contribution. We acknowledge valuable contributions in general from S. Bacca, N. Barnea, M. Birse, E. Borie, C.E. Carlson, M. Eides, J.L. Friar, M. Gorchtein, F. Hagelstein, C. Ji, S. Karshenboim, A.P. Martynenko, J. McGovern, N. Nevo Dinur, K. Pachucki, and M. Vanderhaeghen and are thankful for their valuable remarks and insightful discussions. We proactively thank a future generation of motivated theorists for all future critical compilations of theory terms in light muonic atoms/ions.

The authors acknowledge support from the European Research Council (ERC) through StG. #279765 and CoG. #725039, and the Swiss National Science Foundation SNF, Projects 200021L_138175 and 200021_165854.

Open access funding provided by Max Planck Society.

Author contribution statement

B.F. and J.J.K. set up the tables and wrote the manuscript. Both contributed equally to the paper. The paper was written under the supervision of and includes many comments and suggestions from A.A., F.K., and R.P., whereas M.D. participated in the discussion. All authors discussed the paper and participated in the review.

Open Access This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- R. Pohl, A. Antognini, F. Nez et al., *Nature* **466**, 213 (2010)
- A. Antognini, F. Nez, K. Schuhmann et al., *Science* **339**, 417 (2013)
- A. Antognini, F. Kottmann, F. Biraben et al., *Ann. Phys.* **331**, 127 (2013)
- R. Pohl, F. Nez, L.M.P. Fernandes et al., *Science* **353**, 669 (2016)
- J.J. Krauth, M. Diepold, B. Franke et al., *Ann. Phys.* **366**, 168 (2016)
- P.J. Mohr, D.B. Newell, B.N Taylor, *Rev. Mod. Phys.* **88**, 035009 (2016)
- R. Pohl, R. Gilman, G.A. Miller et al., *Ann. Rev. Nucl. Part. Sci.* **63**, 175 (2013)
- C.E. Carlson, *Prog. Part. Nucl. Phys.* **82**, 59 (2015)
- R.J. Hill, *EPJ Web Conf.* **137**, 01023 (2017)
- C.G. Parthey, A. Matveev, J. Alnis et al., *Phys. Rev. Lett.* **104**, 233001 (2010)
- U.D. Jentschura, A. Matveev, C.G. Parthey et al., *Phys. Rev. A* **83**, 042505 (2011)
- A. Antognini, F. Nez, F.D. Amaro et al., *Can. J. Phys.* **89**, 47 (2010)
- R. Machleidt, D. Entem, *Phys. Rep.* **503**, 1 (2011)
- N. Nevo Dinur, C. Ji, S. Bacca et al., *Phys. Lett. B* **755**, 380 (2016)
- A. Antognini, K. Schuhmann, F.D. Amaro et al., *EPJ Web Conf.* **113**, 01006 (2016)
- G.A. Miller, *Phys. Lett. B* **718**, 1078 (2013)
- U.D. Jentschura, *Phys. Rev. A* **92**, 012123 (2015)
- D. Tucker-Smith, I. Yavin, *Phys. Rev. D* **83**, 101702 (2011)
- B. Batell, D. McKeen, M. Pospelov, *Phys. Rev. Lett.* **107**, 011803 (2011)
- S.G. Karshenboim, D. McKeen, M. Pospelov, *Phys. Rev. D* **90**, 073004 (2014)
- C.E. Carlson, M. Freid, *Phys. Rev. D* **92**, 095024 (2015)
- D. Shiner, R. Dixson, V. Vedantham, *Phys. Rev. Lett.* **74**, 3553 (1995)
- R. van Rooij, J.S. Borbely, J. Simonet et al., *Science* **333**, 196 (2011)
- P. Cancio Pastor, L. Consolino, G. Giusfredi et al., *Phys. Rev. Lett.* **108**, 143001 (2012)
- V. Patkóš, V.A. Yerokhin, K. Pachucki, *Phys. Rev. A* **94**, 052508 (2016)
- V. Patkóš, V.A. Yerokhin, K. Pachucki, *Phys. Rev. A* **95**, 012508 (2017)
- M. Diepold, J.J. Krauth, B. Franke et al. [[arXiv:1606.05231](https://arxiv.org/abs/1606.05231)] (2016)
- A.C. Vutha, N. Bezginov, I. Ferchichi et al., *Bull. Am. Phys. Soc.* **57**, Q1.138 (2012)
- A. Beyer, J. Alnis, K. Khabarova et al., *Ann. D: Phys. (Berlin)* **525**, 671 (2013)
- E. Peters, D.C. Yost, A. Matveev et al., *Ann. D: Phys. (Berlin)* **525**, L29 (2013)
- M. Herrmann, M. Haas, U. Jentschura et al., *Phys. Rev. A* **79**, 052505 (2009)
- D.Z. Kandula, C. Gohle, T.J. Pinkert et al., *Phys. Rev. Lett.* **105**, 063001 (2010)
- M. Mihovilovic, H. Merkel, *AIP Conf. Proc.* **1563**, 187 (2013)
- A. Gasparian, *EPJ Web Conf.* **73**, 07006 (2014)
- R. Gilman, *AIP Conf. Proc.* **1563**, 167 (2013)
- I. Sick, *Phys. Rev. C* **90**, 064002 (2014)
- I. Angeli, K. Marinova, *At. Data Nucl. Data Tables* **99**, 69 (2013)
- G. Carboni, G. Gorini, E. Iacopini et al., *Phys. Lett. B* **73**, 229 (1978)
- P. Hauser, H.P. von Arb, A. Bianchetti et al., *Phys. Rev. A* **46**, 2363 (1992)
- O.J. Hernandez, N. Nevo Dinur, C. Ji et al., *Hyp. Interact.* **237**, 158 (2016)
- C.E. Carlson, M. Gorchtein, M. Vanderhaeghen, *Phys. Rev. A* **95**, 012506 (2017)
- S.J. Brodsky, R.G. Parsons, *Phys. Rev.* **163**, 134 (1967)
- E. Borie, *Ann. Phys.* **327**, 733 (2012)
- E. Borie, Lamb shift in light muonic atoms – revisited, [arXiv:1103.1772-v7](https://arxiv.org/abs/1103.1772-v7) [physic.atom-ph] (2014)
- A. Krutov, A. Martynenko, G. Martynenko et al., *J. Exp. Theor. Phys.* **120**, 73 (2015)
- A.P. Martynenko, E.N. Elekina, *Phys. At. Nucl.* **73**, 2074 (2010)
- A.P. Martynenko, *J. Exp. Theor. Phys.* **106**, 690 (2008)
- R. Faustov, A. Martynenko, G. Martynenko et al., *Phys. Lett. B* **733**, 354 (2014)
- E.N. Elekina, A.P. Martynenko, *Phys. At. Nucl.* **73**, 1828 (2010)
- U.D. Jentschura, B.J. Wundt, *Eur. Phys. J. D* **65**, 357 (2011)
- U.D. Jentschura, *Phys. Rev. A* **84**, 012505 (2011)
- E.Y. Korzinin, V.G. Ivanov, S.G. Karshenboim, *Phys. Rev. D* **88**, 125019 (2013)
- S.G. Karshenboim, V.G. Ivanov, E.Y. Korzinin, *Phys. Rev. A* **85**, 032509 (2012)
- G. Källén, A. Sabry, *Dan. Mat. Fys. Medd.* **29**, 1 (1955)
- U.D. Jentschura *Ann. Phys.* **326**, 500 (2011)
- K. Pachucki, *Phys. Rev. A* **53**, 2092 (1996)
- S.G. Karshenboim, Private communication (2015)
- E. Borie, *Z. Phys. A* **302**, 187 (1981)
- A.A. Krutov, A.P. Martynenko, *Phys. Rev. A* **84**, 052514 (2011)
- W. Barker, F. Glover, *Phys. Rev.* **99**, 317 (1955)
- U.D. Jentschura, *Eur. Phys. J. D* **61**, 7 (2011)
- P.J. Mohr, B.N. Taylor, D.B. Newell, *Rev. Mod. Phys.* **84**, 1527 (2012)
- J.L. Friar, *Ann. Phys.* **122**, 151 (1979)
- K. Pachucki, Private communication (2015)
- V.A. Yerokhin, Nuclear recoil in the Lamb shift of hydrogen-like atoms, in *ECT* Workshop on the Proton Radius Puzzle* (2016)
- E. Borie, Private communication (2017)
- S.G. Karshenboim, E.Y. Korzinin, V.A. Shelyuto et al., *Phys. Rev. D* **91**, 073003 (2015)
- V.G. Ivanov, S.G. Karshenboim, in *Lamb Shift in Light Hydrogen-Like Atoms* (Springer, Berlin, Heidelberg, 2001), pp. 637–650
- J. Friar, G. Payne, *Phys. Rev. A* **56**, 5173 (1997)
- K. Pachucki, *Phys. Rev. Lett.* **106**, 193007 (2011)
- J.L. Friar, *Phys. Rev. C* **88**, 034003 (2013)

72. I. Sick, in *Precise Radii of Light Nuclei from Electron Scattering* (Springer, Berlin, Heidelberg, (2008), pp. 57–77
73. A.P. Martynenko, Private communication (2016)
74. M.I. Eides, H. Grotch, Phys. Rev. A **56**, R2507 (1997)
75. R.N. Faustov, A.P. Martynenko, F.A. Martynenko et al., [[arXiv:1706.01060](https://arxiv.org/abs/1706.01060)] [hep-ph]
76. A.P. Martynenko, Private communication (2017)
77. C. Joachain, Nucl. Phys. **25**, 317 (1961)
78. G.A. Rinker, Phys. Rev. A **14**, 18 (1976)
79. C.E. Carlson, M. Vanderhaeghen, Phys. Rev. A **84**, 020102(R) (2011)
80. L.S. Myers, J.R.M. Annand, J. Brudvik et al., Phys. Rev. Lett. **113**, 262506 (2014)
81. C.E. Carlson, M. Gorchtein, M. Vanderhaeghen, Phys. Rev. A **89**, 022504 (2014)
82. M.C. Birse, J.A. McGovern, Eur. Phys. J. A **48**, 120 (2012)
83. M. Gorchtein, Private communication (2016)
84. S.J. Brodsky, G.W. Erickson, Phys. Rev. **148**, 26 (1966)
85. A.C. Zemach, Phys. Rev. **104**, 1771 (1956)
86. J.L. Friar, I. Sick, Phys. Lett. B **579**, 285 (2004)
87. E. Borie, Z. Phys. A **297**, 17 (1980)
88. H. Grotch, D.R. Yennie, Rev. Mod. Phys. **41**, 350 (1969)
89. E. Borie, Private communication (2015)
90. J.L. Friar, G.L. Payne, Phys. Rev. C **72**, 014002 (2005)
91. M.I. Eides, Phys. Rev. A **85**, 034503 (2012)
92. CREMA Collaboration, to be published
93. S.G. Karshenboim, E.Y. Korzinin, V.A. Shelyuto et al., Phys. Rev. A **96**, 022505 (2017)