



Correction

Correction: Method for indirect radiofluorination with [^{18}F]FDG by biorthogonal reaction

Gergana Simeonova^{1,2,a} , Boyan Todorov^{2,b}, and Valentina Lyubomirova²

¹ UMHAT “St. Marina”, Cl. in Nuclear Medicine, 1 Hristo Smirnenski Blvd., Varna, Bulgaria

² Department of Analytical Chemistry, Faculty of Chemistry and Pharmacy, 1 James Boucher Str., Sofia, Bulgaria

Published online 3 July 2023

© The Author(s), under exclusive licence to EDP Sciences, Springer-Verlag GmbH Germany, part of Springer Nature 2023

Correction: Eur. Phys. J. Spec. Top.
<https://doi.org/10.1140/epjs/s11734-023-00885-7>

In this article the captions of Figs. 6 and 7 show mistakes.

The correct captions are as follows:

Fig. 6 Chromatogram from reaction mixture TLC spots: 1. tetrazine as referent; 2. unreacted [^{18}F]FDG; 3. product [^{18}F]FDG-Tz; 4. catalyst; 5. unreacted tetrazine and radio TLC : Peak 1—unreacted [^{18}F]FDG with $\text{RF}_1 = 0.058$ Peak 2—product [^{18}F]FDG-Tz with $\text{RF}_2 = 0.269$

Fig. 7 HPLC chromatogram from reaction mixture with corresponding RT: [^{19}F]FDG—1.2 min; [^{19}F]FDG-Tz—3.5 min ; Tz—6.0 min ; Cat. —8.0 min

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1140/epjs/s11734-023-00885-7>.

^a e-mail: gerimm@abv.bg (corresponding author)

^b e-mail: ahbt@chem.uni-sofia.bg