

Research Article

An Order on Subsets of Cone Metric Spaces and Fixed Points of Set-Valued Contractions

M. Asadi,¹ H. Soleimani,¹ and S. M. Vaezpour^{2,3}

¹ Department of Mathematics, Science and Research Branch, Islamic Azad University (IAU),
14778 93855 Tehran, Iran

² Department of Mathematics, Amirkabir University of Technology,
15916 34311 Tehran, Iran

³ Department of Mathematics, Newcastle University, Newcastle, NSW 2308, Australia

Correspondence should be addressed to S. M. Vaezpour, vaez@aut.ac.ir

Received 16 April 2009; Revised 19 August 2009; Accepted 22 September 2009

Recommended by Marlene Frigon

In this paper at first we introduce a new order on the subsets of cone metric spaces then, using this definition, we simplify the proof of fixed point theorems for contractive set-valued maps, omit the assumption of normality, and obtain some generalization of results.

Copyright © 2009 M. Asadi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction and Preliminary

Cone metric spaces were introduced by Huang and Zhang [1]. They replaced the set of real numbers by an ordered Banach space and obtained some fixed point theorems for mapping satisfying different contractions [1]. The study of fixed point theorems in such spaces followed by some other mathematicians, see [2–8]. Recently Wardowski [9] was introduced the concept of set-valued contractions in cone metric spaces and established some end point and fixed point theorems for such contractions. In this paper at first we will introduce a new order on the subsets of cone metric spaces then, using this definition, we simplify the proof of fixed point theorems for contractive set-valued maps, omit the assumption of normality, and obtain some generalization of results.

Let E be a real Banach space. A nonempty convex closed subset $P \subset E$ is called a cone in E if it satisfies.

- (i) P is closed, nonempty, and $P \neq \{0\}$,
- (ii) $a, b \in \mathbb{R}$, $a, b \geq 0$, and $x, y \in P$ imply that $ax + by \in P$,
- (iii) $x \in P$ and $-x \in P$ imply that $x = 0$.

The space E can be partially ordered by the cone $P \subset E$; that is, $x \leq y$ if and only if $y - x \in P$. Also we write $x \ll y$ if $y - x \in P^\circ$, where P° denotes the interior of P .

A cone P is called normal if there exists a constant $K > 0$ such that $0 \leq x \leq y$ implies $\|x\| \leq K\|y\|$.

In the following we always suppose that E is a real Banach space, P is a cone in E , and \leq is the partial ordering with respect to P .

Definition 1.1 (see [1]). Let X be a nonempty set. Assume that the mapping $d : X \times X \rightarrow E$ satisfies

- (i) $0 \leq d(x, y)$ for all $x, y \in X$ and $d(x, y) = 0$ iff $x = y$
- (ii) $d(x, y) = d(y, x)$ for all $x, y \in X$
- (iii) $d(x, y) \leq d(x, z) + d(z, y)$ for all $x, y, z \in X$.

Then d is called a cone metric on X , and (X, d) is called a cone metric space.

In the following we have some necessary definitions.

- (1) Let (M, d) be a cone metric space. A set $A \subseteq M$ is called *closed* if for any sequence $\{x_n\} \subseteq A$ convergent to x , we have $x \in A$.
- (2) A set $A \subseteq M$ is called *sequentially compact* if for any sequence $\{x_n\} \subseteq A$, there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ is convergent to an element of A .
- (3) Denote $N(M)$ a collection of all nonempty subsets of M , $C(M)$ a collection of all nonempty closed subsets of M and $K(M)$ a collection of all nonempty sequentially compact subsets of M .
- (4) An element $x \in M$ is said to be an *endpoint* of a set-valued map $T : M \rightarrow N(M)$, if $Tx = \{x\}$. We denote a set of all endpoints of T by $\text{End}(T)$.
- (5) An element $x \in M$ is said to be a *fixed point* of a set-valued map $T : M \rightarrow N(M)$, if $x \in Tx$. Denote $\text{Fix}(T) = \{x \in M \mid x \in Tx\}$.
- (6) A map $f : M \rightarrow \mathbb{R}$ is called *lower semi-continuous*, if for any sequence $\{x_n\}$ in M and $x \in M$, such that $x_n \rightarrow x$ as $n \rightarrow \infty$, we have $f(x) \leq \liminf_{n \rightarrow \infty} f(x_n)$.
- (7) A map $f : M \rightarrow E$ is called *have lower semi-continuous property*, and denoted by *lsc property* if for any sequence $\{x_n\}$ in M and $x \in M$, such that $x_n \rightarrow x$ as $n \rightarrow \infty$, then there exists $N \in \mathbb{N}$ that $f(x) \leq f(x_n)$ for all $n \geq N$.
- (8) P called *minihedral* cone if $\sup\{x, y\}$ exists for all $x, y \in E$, and *strongly minihedral* if every subset of E which is bounded from above has a supremum [10]. Let (M, d) a cone metric space, cone P is strongly minihedral and hence, every subset of P has infimum, so for $A \in C(M)$, we define $d(x, A) = \inf_{y \in A} d(x, y)$.

Example 1.2. Let $E := \mathbb{R}^n$ with $P := \{(x_1, x_2, \dots, x_n) : x_i \geq 0 \text{ for all } i = 1, 2, \dots, n\}$. The cone P is normal, minihedral and strongly minihedral with $P^0 \neq \emptyset$.

Example 1.3. Let $D \subseteq \mathbb{R}^n$ be a compact set, $E := C(D)$, and $P := \{f \in E : f(x) \geq 0 \text{ for all } x \in D\}$. The cone P is normal and minihedral but is not strongly minihedral and $P^0 \neq \emptyset$.

Example 1.4. Let (X, S, μ) be a finite measure space, S countably generated, $E := L^p(X)$, $(1 < p < \infty)$, and $P := \{f \in E : f(x) \geq 0 \mu \text{ a.e. on } X\}$. The cone P is normal, minihedral and strongly minihedral with $P^0 = \emptyset$.

For more details about above examples, see [11].

Example 1.5. Let $E := C^2([0, 1], \mathbb{R}^+)$ with norm $\|f\| = \|f\|_\infty + \|f'\|_\infty$ and $P := \{f \in E : f \geq 0\}$ that is not normal cone by [12] and not minihedral by [10].

Example 1.6. Let $E := \mathbb{R}^2$ and $P := \{(x_1, 0) : x_1 \geq 0\}$. This P is strongly minihedral but not minihedral by [10].

Throughout, we will suppose that P is strongly minihedral cone in E with nonempty interior and \leq be a partial ordering with respect to P .

2. Main Results

Let (M, d) be a cone metric space and $T : M \rightarrow C(M)$. For $x, y \in M$, Let

$$\begin{aligned} D(x, Ty) &= \{d(x, z) : z \in Ty\}, \\ S(x, Ty) &= \{u \in D(x, Ty) : \|u\| = \inf\{\|v\| : v \in D(x, Ty)\}\}. \end{aligned} \quad (2.1)$$

At first we prove the closedness of $\text{Fix}(T)$ without the assumption of normality.

Lemma 2.1. *Let (M, d) be a complete cone metric space and $T : M \rightarrow C(M)$. If the function $f(x) = \inf_{y \in Tx} \|d(x, y)\|$ for $x \in M$ is lower semi-continuous, then $\text{Fix}(T)$ is closed.*

Proof. Let $x_n \in Tx_n$ and $x_n \rightarrow x$. We show that $x \in Tx$. Since

$$\begin{aligned} f(x) &\leq \liminf_{n \rightarrow \infty} f(x_n) = \liminf_{n \rightarrow \infty} \inf_{y \in Tx_n} \|d(x_n, y)\|, \\ &\leq \liminf_{n \rightarrow \infty} \|d(x_n, x_n)\| = 0, \end{aligned} \quad (2.2)$$

so $f(x) = 0$ which implies $d(y_n, x) \rightarrow 0$ for some $y_n \in Tx$. Let $c \in E$ with $c \gg 0$ then, there exists N such that for $n \geq N$, $d(y_n, x) \ll (1/2)c$. Now, for $n > m$, we have,

$$d(y_n, y_m) \leq d(y_n, x) + d(x, y_m) \ll \frac{1}{2}c + \frac{1}{2}c = c. \quad (2.3)$$

So $\{y_n\}$ is a Cauchy sequence in complete metric space, hence there exist $y^* \in M$ such that $y_n \rightarrow y^*$. Since Tx is closed, thus $y^* \in Tx$. Now by uniqueness of limit we conclude that $x = y^* \in Tx$. \square

Definition 2.2. Let A and B are subsets of E , we write $A \leq B$ if and only if there exist $x \in A$ such that for all $y \in B$, $x \leq y$. Also for $x \in E$, we write $x \leq B$ if and only if $\{x\} \leq B$ and similarly $A \leq x$ if and only if $A \leq \{x\}$.

Note that $aA + B := \{ax + y : x \in A, y \in B\}$, for every scalar $a \in \mathbb{R}^+$ and A, B subsets of E .

The following lemma is easily proved.

Lemma 2.3. Let $A, B, C \subseteq E$, $x, y \in E$, $a \in \mathbb{R}^+$, and $a \neq 0$.

- (1) If $A \leq B$, and $B \leq C$, then $A \leq C$,
- (2) $A \leq B \Leftrightarrow aA \leq aB$,
- (3) If $x \leq B$, then $ax \leq aB$,
- (4) If $A \leq y$, then $aA \leq ay$,
- (5) $x \leq y \Leftrightarrow \{x\} \leq \{y\}$,
- (6) If $A \leq B$, then $A \leq B + P$.

The order " \leq " is not antisymmetric, thus this order is not partially order.

Example 2.4. Let $E := \mathbb{R}$ and $P := \mathbb{R}^+$. Put $A := [1, 3]$ and $B := [1, 4]$ so $A \leq B$, $B \leq A$ but $A \neq B$.

Theorem 2.5. Let (M, d) be a complete cone metric space, $T : M \rightarrow C(M)$, a set-valued map and the function $f : M \rightarrow P$ defined by $f(x) = d(x, Tx)$, $x \in M$ with lsc property. If there exist real numbers $a, b, c, e \geq 0$ and $q > 1$ with $k := aq + b + ceq < 1$ such that for all $x \in M$ there exists $y \in Tx$:

$$\begin{aligned} d(x, y) &\leq qD(x, Tx), \\ D(y, Tx) &\leq ed(x, y), \\ D(y, Ty) &\leq ad(x, y) + bD(x, Tx) + cD(y, Tx), \end{aligned} \tag{2.4}$$

then $\text{Fix}(T) \neq \emptyset$.

Proof. Let $x \in M$, then there exists $y \in Tx$ such that

$$\begin{aligned} D(y, Ty) &\leq ad(x, y) + bD(x, Tx) + cD(y, Tx) \\ &\leq (aq + b + ceq)D(x, Tx) = kD(x, Tx). \end{aligned} \tag{2.5}$$

Let $x_0 \in M$, there exist $x_1 \in Tx_0$ such that $D(x_1, Tx_1) \leq kD(x_0, Tx_0)$ and $d(x_0, x_1) \leq qD(x_0, Tx_0)$. Continuing this process, we can iteratively choose a sequence $\{x_n\}$ in M such that $x_{n+1} \in Tx_n$, $D(x_n, Tx_n) \leq k^n D(x_0, Tx_0)$, and $d(x_n, x_{n+1}) \leq qD(x_n, Tx_n) \leq qk^n D(x_0, Tx_0)$. So, for $n > m$, we have,

$$\begin{aligned} \{d(x_n, x_m)\} &\leq \{d(x_n, x_{n-1}) + d(x_{n-1}, x_{n-2}) + \cdots + d(x_{m+1}, x_m)\} \\ &\leq q(k^{n-1} + k^{n-2} + \cdots + k^m)D(x_0, Tx_0) \\ &\leq qk^m(1 + k + k^2 + \cdots)D(x_0, Tx_0) \\ &\leq q\frac{k^m}{1-k}D(x_0, Tx_0). \end{aligned} \tag{2.6}$$

Therefore, for every $u_0 \in D(x_0, Tx_0)$, $d(x_n, x_m) \leq q(k^m/(1-k))u_0$. Let $c \in E$ and $c \gg 0$ be given. Choose $\delta > 0$ such that $c + N_\delta(0) \subseteq P$, where $N_\delta(0) = \{x \in E : \|x\| < \delta\}$. Also, choose a $N \in \mathbb{N}$ such that $q(k^m/(1-k))u_0 \in N_\delta(0)$, for all $m \geq N$. Then $q(k^m/(1-k))u_0 \ll c$, for all $m \geq N$. Thus $d(x_n, x_m) \leq q(k^m/(1-k))u_0 \ll c$ for all $n > m$. Namely, $\{x_n\}$ is Cauchy sequence in complete cone metric space, therefore $x_n \rightarrow x^*$ for some $x^* \in M$. Now we show that $x^* \in Tx^*$.

Let $u_n \in D(x_n, Tx_n)$ hence there exists $t_n \in Tx_n$ such that $0 \leq u_n = d(x_n, t_n) \leq k^n u_0$ for all $u_0 \in D(x_0, Tx_0)$. Now $k^n u_0 \rightarrow 0$ as $n \rightarrow \infty$ so for all $0 \ll c$ there exists $N \in \mathbb{N}$ such that $0 \leq u_n = d(x_n, t_n) \leq k^n u_0 \ll c$ for all $n \geq N$.

According to *lsc property* of f , for all $c \gg 0$ there exists $N \in \mathbb{N}$ such that for all $n \geq N$

$$f(x^*) \leq f(x_n) = \inf_{y \in Tx_n} d(x_n, y) \leq d(x_n, t_n) \ll c. \quad (2.7)$$

So $0 \leq f(x^*) \ll c$ for all $c \gg 0$. Namely, $f(x^*) = 0$ thus $d(y_n, x^*) \rightarrow 0$ for some $y_n \in Tx^*$, and by the closedness of Tx^* we have $x^* \in Tx^*$. \square

We notice that $d(x_n, x) \rightarrow 0$ implies that for all $c \gg 0$ there exists $N \in \mathbb{N}$ such that $d(x_n, x) \ll c$ for all $n \geq N$, but the inverse is not true.

Example 2.6. Let $M = E := C^2([0, 1], \mathbb{R}^+)$ with norm $\|f\| = \|f\|_\infty + \|f'\|_\infty$ and $P := \{f \in E : f \geq 0\}$ that is not normal cone by [12]. Consider $x_n := (1 - \sin nt)/(n + 2)$ and $y_n := (1 + \sin nt)/(n + 2)$ so $0 \leq x_n \leq x_n + y_n \rightarrow 0$ and $\|x_n\| = \|y_n\| = 1$, (see [10]) Define cone metric $d : M \times M \rightarrow E$ with $d(f, g) = f + g$, for $f \neq g$, $d(f, f) = 0$. Since $0 \leq x_n \ll c$, namely, $d(x_n, 0) \ll c$ but $d(x_n, 0) \not\rightarrow 0$. Indeed $x_n \rightarrow 0$ in (M, d) but $x_n \not\rightarrow 0$ in E . Even for $n > m$, $d(x_n, x_m) = x_n + x_m \ll c$ and $\|d(x_n, x_m)\| = \|x_n + x_m\| = 2$ in particular $d(x_n, x_{n+1}) \ll c$ but $d(x_n, x_{n+1}) \not\rightarrow 0$.

Example 2.7. Let $M = E := C^2([0, 1], \mathbb{R})$ with norm $\|f\| = \|f\|_\infty + \|f'\|_\infty$ and $P := \{f \in E : f \geq 0\}$ that is not normal cone. Define cone metric $d : M \times M \rightarrow E$ with $d(f, g) = f^2 + g^2$, for $f \neq g$, $d(f, f) = 0$ and set-valued mapping $T : M \rightarrow C(M)$ by $Tf = \{-f, 0, f\}$. In this space every Cauchy sequence converges to zero. The function $F(f) = d(f, Tf) = \inf_{g \in Tf} d(f, g) = \inf\{0, f^2, 2f^2\} = 0$ have *lsc property*. Also we have $D(f, Tf) = \{0, f^2, 2f^2\}$ and $D(f, Tg) = \{f^2, f^2 + g^2\}$. Now for $q > 1$, $e \geq 1$, $a, b, c \geq 0$, $k = aq + b + ceq < 1$ and for all $f \in M$ take $g := 0 \in Tf$. Therefore, it satisfies in all of the hypothesis of Theorem 2.5. So T has a fixed point $f \in Tf$. For sample take $a = b = c = 1/6$, $e = 1$, and $q = 2$.

Theorem 2.8. Let (M, d) be a complete cone metric space, $T : M \rightarrow K(M)$, a set-valued map, and a function $f : M \rightarrow P$ defined by $f(x) = d(x, Tx)$, $x \in M$ with *lsc property*. The following conditions hold:

- (i) if there exist real numbers $a, b, c, e \geq 0$ and $q > 1$ with $k := aq + b + ceq < 1$ such that for all $x \in M$, there exists $y \in Tx$:

$$\begin{aligned} d(x, y) &\leq qS(x, Tx), \\ S(y, Tx) &\leq ed(x, y), \\ S(y, Ty) &\leq ad(x, y) + bS(x, Tx) + cS(y, Tx), \end{aligned} \quad (2.8)$$

then $\text{Fix}(T) \neq \emptyset$,

(ii) if there exist real numbers $a, b, c, e \geq 0$ and $q > 1$ with $k := aq + b + ceq < 1$ such that for all $x \in M$ and $y \in Tx$:

$$\begin{aligned} d(x, y) &\leq qS(x, Tx), \\ S(y, Tx) &\leq ed(x, y), \\ S(y, Ty) &\leq ad(x, y) + bS(x, Tx) + cS(y, Tx), \end{aligned} \tag{2.9}$$

then $\text{Fix}(T) = \text{End}(T) \neq \emptyset$.

Proof. (i) It is obvious that $S(x, Tx) \subseteq D(x, Tx)$. It is enough to show that $S(x, Tx) \neq \emptyset$ for all $x \in M$. However $S(x, Tx) = \emptyset$ for some $x \in M$, it implies $d(x, y) \leq \emptyset$ for some $y \in Tx$, and this is a contradiction.

(ii) By (i), there exists $x^* \in M$ such that $x^* \in Tx^*$. Then for $y \in Tx^*$ and $0 \in S(x^*, Tx^*)$ we have $d(x^*, y) \leq (1/b)S(x^*, Tx^*)$. Therefore, $d(x^*, y) \leq (1/b)0 = 0$. This implies that $x^* = y \in Tx^*$. \square

Corollary 2.9. Let (M, d) be a complete cone metric space, $T : M \rightarrow C(M)$, a set-valued map, and the function $f : M \rightarrow P$ defined by $f(x) = d(x, Tx)$, for $x \in M$ with lsc property. If there exist real numbers $a, b \geq 0$ and $q > 1$ with $k := aq + b < 1$ such that for all $x \in M$ there exists $y \in Tx$ with

$$\begin{aligned} d(x, y) &\leq qD(x, Tx), \\ D(y, Ty) &\leq ad(x, y) + bD(x, Tx), \end{aligned} \tag{2.10}$$

then $\text{Fix}(T) \neq \emptyset$.

To have Theorems 3.1 and 3.2 in [9], as the corollaries of our theorems we need the following lemma and remarks.

Lemma 2.10. Let (M, d) be a cone metric space, P a normal cone with constant one and $T : M \rightarrow C(M)$, a set-valued map, then

$$\|d(x, Tx)\| = \left\| \inf_{y \in Tx} d(x, y) \right\| = \inf_{y \in Tx} \|d(x, y)\|. \tag{2.11}$$

Proof. Put $\alpha := \inf_{y \in Tx} \|d(x, y)\|$ and $\beta := \inf_{y \in Tx} d(x, y)$ we show that $\alpha = \|\beta\|$.

Let $y \in Tx$ then $\beta \leq d(x, y)$ and so $\|\beta\| \leq \|d(x, y)\|$, which implies $\|\beta\| \leq \alpha$.

For the inverse, let for all $0 \leq r \leq \alpha$. Then $r \leq \|d(x, y)\|$ for all $y \in Tx$.

Since $\beta := \inf_{y \in Tx} d(x, y)$, for every c that $c \gg 0$ there exists $y \in Tx$ such that $d(x, y) < \beta + c$, so $r \leq \|d(x, y)\| < \|\beta + c\| \leq \|\beta\| + \|c\|$, for all $c \gg 0$. Thus $r \leq \|\beta\|$. \square

Remark 2.11. By Proposition 1.7.59, page 117 in [11], if E is an ordered Banach space with positive cone P , then P is a normal cone if and only if there exists an equivalent norm $|\cdot|$ on E which is monotone. So by renorming the E we can suppose P is a normal cone with constant one.

Remark 2.12. Let (M, d) be a cone metric space, P a normal cone with constant one, $T : M \rightarrow C(M)$, a set-valued map, the function $f : M \rightarrow P$ defined by $f(x) = d(x, Tx)$, $x \in M$ with *lsc property*, and $g : E \rightarrow \mathbb{R}^+$ with $g(x) = \|x\|$. Then $g \circ f(x) = \inf_{y \in Tx} \|d(x, y)\|$, is lower semi-continuous.

Now the Theorems 3.1 and 3.2 in [9] is stated as the following corollaries without the assumption of normality, and by Lemma 2.10 and Remarks 2.11, 2.12 we have the same theorems.

Corollary 2.13 (see [9, Theorem 3.1]). *Let (M, d) be a complete cone metric space, $T : M \rightarrow C(M)$, a set-valued map and the function $f : M \rightarrow P$ defined by $f(x) = d(x, Tx)$, $x \in M$ with *lsc property*. If there exist real numbers $0 \leq \lambda < 1$, $\lambda < b \leq 1$ such that for all $x \in M$ there exists $y \in Tx$ one has $D(y, Ty) \leq \lambda d(x, y)$ and $bd(x, y) \leq D(x, Tx)$ then $\text{Fix}(T) \neq \emptyset$.*

Corollary 2.14 (see [9, Theorem 3.2]). *Let (M, d) be a complete cone metric space, $T : M \rightarrow K(M)$, a set-valued map and the function $f : M \rightarrow P$ defined by $f(x) = d(x, Tx)$, $x \in M$ with *lsc property*. The following hold:*

- (i) *if there exist real numbers $0 \leq \lambda < 1$, $\lambda < b \leq 1$ such that for all $x \in M$ there exists $y \in Tx$ one has $S(y, Ty) \leq \lambda d(x, y)$ and $bd(x, y) \leq S(x, Tx)$, then $\text{Fix}(T) \neq \emptyset$,*
- (ii) *if there exist real numbers $0 \leq \lambda < 1$, $\lambda < b \leq 1$ such that for all $x \in M$ and every $y \in Tx$ one has $S(y, Ty) \leq \lambda d(x, y)$ and $bd(x, y) \leq S(x, Tx)$, then $\text{Fix}(T) = \text{End}(T) \neq \emptyset$.*

Definition 2.15. For $A \subseteq M$, $T : M \rightarrow C(M)$ where T is a set-valued map we define

$$\overline{D}(A, TA) := \bigcup_{x \in A} D(x, Tx), \quad \underline{D}(A, TA) := \bigcap_{x \in A} D(x, Tx). \quad (2.12)$$

Note that $T^2x = T(Tx)$ for $x \in M$.

The following theorem is a reform of Theorem 2.5.

Theorem 2.16. *Let (M, d) be a complete cone metric space, $T : M \rightarrow C(M)$, a set-valued map, and the function $f : M \rightarrow P$ defined by $f(x) = d(x, Tx)$, $x \in M$ with *lsc property*. If there exists $0 \leq k < 1$ such that*

$$\overline{D}(Tx, T^2x) \leq k \underline{D}(M, TM). \quad (2.13)$$

for all $x \in M$. Then $\text{Fix}(T) \neq \emptyset$.

Proof. For every $x \in M$, then there exist $y \in Tx$ and $z \in Ty$ such that $d(y, z) \leq kd(x, t)$, for all $t \in Tx$. Let $x_n \in M$, there exist $x_{n+1} \in Tx_n$ and $x_{n+2} \in Tx_{n+1}$ such that $d(x_{n+1}, x_{n+2}) \leq kd(x_n, x_{n+1})$, since $x_{n+1} \in Tx_n$. Thus $d(x_n, x_{n+1}) \leq k^n d(x_0, x_1)$. The remaining is same as the proof of Theorem 2.5. \square

References

- [1] L.-G. Huang and X. Zhang, "Cone metric spaces and fixed point theorems of contractive mappings," *Journal of Mathematical Analysis and Applications*, vol. 332, no. 2, pp. 1468–1476, 2007.

- [2] M. Abbas and B. E. Rhoades, "Fixed and periodic point results in cone metric spaces," *Applied Mathematics Letters*, vol. 22, no. 4, pp. 511–515, 2009.
- [3] M. Arshad, A. Azam, and P. Vetro, "Some common fixed point results in cone metric spaces," *Fixed Point Theory and Applications*, vol. 2009, Article ID 493965, 11 pages, 2009.
- [4] D. Ilić and V. Rakočević, "Common fixed points for maps on cone metric space," *Journal of Mathematical Analysis and Applications*, vol. 341, no. 2, pp. 876–882, 2008.
- [5] D. Ilić and V. Rakočević, "Quasi-contraction on a cone metric space," *Applied Mathematics Letters*, vol. 22, no. 5, pp. 728–731, 2009.
- [6] S. Janković, Z. Kadelburg, S. Radenović, and B. E. Rhoades, "Assad-Kirk-type fixed point theorems for a pair of nonself mappings on cone metric spaces," *Fixed Point Theory and Applications*, vol. 2009, Article ID 761086, 16 pages, 2009.
- [7] G. Jungck, S. Radenović, S. Radojević, and V. Rakočević, "Common fixed point theorems for weakly compatible pairs on cone metric spaces," *Fixed Point Theory and Applications*, vol. 2009, Article ID 643840, 13 pages, 2009.
- [8] P. Raja and S. M. Vaezpour, "Some extensions of Banach's contraction principle in complete cone metric spaces," *Fixed Point Theory and Applications*, vol. 2008, Article ID 768294, 11 pages, 2008.
- [9] D. Wardowski, "Endpoints and fixed points of set-valued contractions in cone metric spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 1-2, pp. 512–516, 2009.
- [10] K. Deimling, *Nonlinear Functional Analysis*, Springer, Berlin, Germany, 1985.
- [11] Z. Denkowski, S. Migórski, and N. S. Papageorgiou, *An Introduction to Nonlinear Analysis: Applications*, Kluwer Academic Publishers, Boston, Mass, USA, 2003.
- [12] Sh. Rezapour and R. Hamlbarani, "Some notes on the paper "Cone metric spaces and fixed point theorems of contractive mappings"," *Journal of Mathematical Analysis and Applications*, vol. 345, no. 2, pp. 719–724, 2008.