

# FIRST- AND SECOND-ORDER DYNAMIC EQUATIONS WITH IMPULSE

F. M. ATICI AND D. C. BILES

*Received 3 December 2004 and in revised form 11 February 2005*

We present existence results for discontinuous first- and continuous second-order dynamic equations on a time scale subject to fixed-time impulses and nonlinear boundary conditions.

## 1. Introduction

We first briefly survey the recent results for existence of solutions to first-order problems with fixed-time impulses. Periodic boundary conditions using upper and lower solutions were considered in [19], using degree theory. A nonlinear alternative of Leray-Schauder type was used in [15] for initial conditions or periodic boundary conditions. The monotone iterative technique was employed in [14] for antiperiodic and nonlinear boundary conditions. Lower and upper solutions and periodic boundary conditions were studied in [20]. Semilinear damped initial value problems in a Banach space using fixed point theory were investigated in [6]. In [9], existence of solutions for the differential equation  $u'(t) = q(u(t))g(t, u(t))$  subject to a general boundary condition is proven, in which  $g$  is Carathéodory and  $q \in L^\infty$ , and existence of lower and upper solutions is assumed. Schauder's fixed point theorem was used there. This generalized an earlier result found in [18]. It appears that little has been done concerning dynamic equations with impulses on time scales (see [4, 5, 16] for earlier results). In Section 2, the present paper uses ideas from [9] to prove an existence result for discontinuous dynamic equations on a time scale subject to fixed-time impulses and nonlinear boundary conditions.

The study of boundary value problems for nonlinear second-order differential equations with impulses has appeared in many papers (see [10, 11, 13] and the references therein). In Section 3, we use ideas from [12, 16] to prove an existence result for second-order dynamic equations on a time scale subject to fixed-time impulses and nonlinear boundary conditions. Nonlinear boundary conditions cover, among others, the periodic and the Dirichlet conditions, and have been introduced for ordinary differential equations by Adje in [1]. Assuming the existence of a lower and an upper solution, we prove that the solution of the boundary value problem stays between them.

In [2], it was shown that the upper and lower solution method will not work for first-order dynamic equations involving  $\Delta$ -derivatives, unless restrictive assumptions are

made. Hence, in Section 2, we work with the  $\nabla$ -derivative. In Section 3, we can use the more conventional  $\Delta$ -derivative.

The monographs [17, 21] are good general references on impulsive differential equations—discussion of applications may be found in these books. Applications of the results given in this paper could involve those typically modelled on time scales which are subjected to sudden major influences, for example, an insect population sprayed with an insecticide or a financial market affected by a major terrorist attack.

For our purposes, we let  $\mathbb{T}$  be a time scale (a closed subset of  $\mathbb{R}$ ), let  $[a, b]$  be the closed and bounded interval in  $\mathbb{T}$ , that is,  $[a, b] := \{t \in \mathbb{T} : a \leq t \leq b\}$  and  $a, b \in \mathbb{T}$ . For the readers' convenience, we state a few basic definitions on a time scale  $\mathbb{T}$  [7, 8].

Obviously a time scale  $\mathbb{T}$  may or may not be connected. Therefore, we have the concept of *forward* and *backward jump operators* as follows: define  $\sigma, \rho : \mathbb{T} \rightarrow \mathbb{T}$  by

$$\sigma(t) = \inf\{s \in \mathbb{T} : s > t\}, \quad \rho(t) = \sup\{s \in \mathbb{T} : s < t\}. \quad (1.1)$$

If  $\sigma(t) = t$ ,  $\sigma(t) > t$ ,  $\rho(t) = t$ ,  $\rho(t) < t$ , then  $t \in \mathbb{T}$  is called *right dense* (rd), *right scattered*, *left dense*, *left scattered*, respectively. We also define the *graininess function*  $\mu : \mathbb{T} \rightarrow [0, \infty)$  as  $\mu(t) = \sigma(t) - t$ . The sets  $\mathbb{T}^\kappa$ ,  $\mathbb{T}_\kappa$  which are derived from  $\mathbb{T}$  are as follows: if  $\mathbb{T}$  has a left-scattered maximum  $t_1$ , then  $\mathbb{T}^\kappa = \mathbb{T} - \{t_1\}$ , otherwise  $\mathbb{T}^\kappa = \mathbb{T}$ . If  $\mathbb{T}$  has a right-scattered minimum  $t_2$ , then  $\mathbb{T}_\kappa = \mathbb{T} - \{t_2\}$ , otherwise  $\mathbb{T}_\kappa = \mathbb{T}$ . If  $f : \mathbb{T} \rightarrow \mathbb{R}$  is a function, we define the functions  $f^\sigma : \mathbb{T}^\kappa \rightarrow \mathbb{R}$  by  $f^\sigma(t) = f(\sigma(t))$  for all  $t \in \mathbb{T}^\kappa$ ,  $f^\rho : \mathbb{T}_\kappa \rightarrow \mathbb{R}$  by  $f^\rho(t) = f(\rho(t))$  for all  $t \in \mathbb{T}_\kappa$  and  $\sigma^0(t) = \rho^0(t) = t$ .

If  $f : \mathbb{T} \rightarrow \mathbb{R}$  is a function and  $t \in \mathbb{T}^\kappa$ , then the *delta derivative* of  $f$  at a point  $t$  is defined to be the number  $f^\Delta(t)$  (provided it exists) with the property that, for each  $\varepsilon > 0$ , there is a neighborhood of  $U_1$  of  $t$  such that

$$|[f(\sigma(t)) - f(s)] - f^\Delta(t)[\sigma(t) - s]| \leq \varepsilon |\sigma(t) - s|, \quad (1.2)$$

for all  $s \in U_1$ . If  $t \in \mathbb{T}_\kappa$ , then we define the *nabla derivative* of  $f$  at a point  $t$  to be the number  $f^\nabla(t)$  (provided it exists) with the property that, for each  $\varepsilon > 0$ , there is a neighborhood of  $U_2$  of  $t$  such that

$$|[f(\rho(t)) - f(s)] - f^\nabla(t)[\rho(t) - s]| \leq \varepsilon |\rho(t) - s|, \quad (1.3)$$

for all  $s \in U_2$ .

*Remark 1.1.* If  $\mathbb{T} = \mathbb{R}$ , then  $f^\Delta(t) = f^\nabla(t) = f'(t)$ , and if  $\mathbb{T} = \mathbb{Z}$ , then  $f^\Delta(t) = \Delta f(t) = f(t+1) - f(t)$  and  $f^\nabla(t) = \nabla f(t) = f(t) - f(t-1)$ .

A function  $F : \mathbb{T} \rightarrow \mathbb{R}$  is called a  $\Delta$ -antiderivative of  $f : \mathbb{T} \rightarrow \mathbb{R}$  provided  $F^\Delta(t) = f(t)$  holds for all  $t \in \mathbb{T}^k$ . Then the Cauchy  $\Delta$ -integral from  $a$  to  $t$  of  $f$  is defined by

$$\int_a^t f(s) \Delta s = F(t) - F(a) \quad \forall t \in \mathbb{T}. \quad (1.4)$$

A function  $\Phi : \mathbb{T} \rightarrow \mathbb{R}$  is called a  $\nabla$ -antiderivative of  $f : \mathbb{T} \rightarrow \mathbb{R}$  provided  $\Phi^\nabla(t) = f(t)$  for all  $t \in \mathbb{T}_k$ . We then define the Cauchy  $\nabla$ -integral from  $a$  to  $t$  of  $f$  by

$$\int_a^t f(s) \nabla s = \Phi(t) - \Phi(a) \quad \forall t \in \mathbb{T}. \tag{1.5}$$

Note that, in the case  $\mathbb{T} = \mathbb{R}$ , we have

$$\int_a^b f(t) \Delta t = \int_a^b f(t) \nabla t = \int_a^b f(t) dt, \tag{1.6}$$

and, in the case  $\mathbb{T} = \mathbb{Z}$ , we have

$$\int_a^b f(t) \Delta t = \sum_{k=a}^{b-1} f(k), \quad \int_a^b f(t) \nabla t = \sum_{k=a+1}^b f(k), \tag{1.7}$$

where  $a, b \in \mathbb{T}$  with  $a \leq b$ .

There are two types of impulse effects that are studied in the literature. The first is the “fixed-time impulse”: a set of times  $0 < t_1 < t_2 < \dots < t_n < T$  is specified, and the solution is required to satisfy

$$u(t_k^+) = I_k(u(t_k)) \tag{1.8}$$

for  $k = 1, 2, \dots, n$ , where the functions  $I_k$  provide the “impulse.” Also studied are “variable-time impulses,” in which a set of curves  $t = \tau_1(x), t = \tau_2(x), \dots, t = \tau_n(x)$  is given, and the solution satisfies  $u(t^+) = I_k(u(t))$  for  $t = \tau_k(u(t))$ ,  $k = 1, 2, \dots, n$ . Impulses of both types introduce discontinuities in the solution. As mentioned in [17] and other works in the reference list, applications involving impulse effects can be found in biology, medicine, physics, economics, pharmacokinetics, and engineering. In this paper, we consider fixed-time impulses. Without loss of generality, we investigate systems with a single impulse.

### 2. First order

Let  $0, t_1, T \in \mathbb{T}$  with  $0 < t_1 < T$  and  $t_1$  right dense. Let  $J = [0, T] \cap \mathbb{T}$ ,

$$u^\nabla(t) = g(t, u(t)), \quad t \in J \setminus \{t_1\}, \tag{2.1}$$

$$u(t_1^+) = I(u(t_1)), \tag{2.2}$$

$$B(u(0), u(T)) = 0. \tag{2.3}$$

Note that (2.3) covers as special cases many initial and boundary conditions found in the literature. Let  $J_1 = [0, t_1] \cap J$ ,  $J_2 = (t_1, T] \cap J$ . Define  $\int_a^b y(s) \nabla s = \int_{[a,b]} y(s) \nabla s$ , where the integrals in Section 2 are with respect to the Lebesgue  $\nabla$ -measure as defined by Atici and Guseinov in [3].

Let  $u_i$  be the restriction of  $u : J \rightarrow \mathbb{R}$  to  $J_i$ ,  $i = 1, 2$ , then

$$\begin{aligned} \mathcal{C}(J_1) &= \{u : J_1 \rightarrow \mathbb{R} : u \text{ is continuous on } J_1\}, \\ \mathcal{C}(J_2) &= \{u : J_2 \rightarrow \mathbb{R} : u \text{ is continuous on } J_2 \text{ and } u(t_1^+) \text{ exists}\}, \\ A &= \{u : J \rightarrow \mathbb{R} : u_1 \in \mathcal{C}(J_1) \text{ and } u_2 \in \mathcal{C}(J_2)\}. \end{aligned} \tag{2.4}$$

For  $u \in A$ , let  $\|u\| = \sup\{|u(t)| : t \in J\}$ .  $(A, \|\cdot\|)$  is a Banach space. For  $u, v \in A$ ,

$$[u, v] \equiv \{w \in A : u(t) \leq w(t) \leq v(t) \quad \forall t \in J\}. \tag{2.5}$$

*Definition 2.1.*  $u : \mathbb{T} \rightarrow \mathbb{R}$  is a solution of (2.1)–(2.3) if

- (i)  $u \in A$ ,
- (ii)

$$\begin{aligned} u(t) &= u(0) + \int_0^t g(s, u(s)) \nabla s, \quad t \in J_1, \\ u(t) &= I(u(t_1)) + \int_{t_1}^t g(s, u(s)) \nabla s, \quad t \in J_2, \end{aligned} \tag{2.6}$$

- (iii)  $B(u(0), u(T)) = 0$ .

We call  $\alpha : J \rightarrow \mathbb{R}$  a lower solution of (2.1)–(2.3) if

- (i)  $\alpha \in A$ ,
- (ii)  $\alpha(b) - \alpha(a) \leq \int_a^b g(s, \alpha(s)) \nabla s$  for  $a \leq b$  and  $a, b \in J_1$ , or  $a, b \in J_2$ ,
- (iii)  $\alpha(t_1^+) \leq I(\alpha(t_1))$ ,
- (iv)  $B(\alpha(0), \alpha(T)) \leq 0$ .

We call  $\beta : J \rightarrow \mathbb{R}$  an upper solution of (2.1)–(2.3) if it satisfies the same assumptions, but replace  $\leq$  with  $\geq$ .

Let  $p(t, x) = \max\{\alpha(t), \min\{x, \beta(t)\}\}$ . We have the following assumptions throughout this section:

- (1) for each  $x \in \mathbb{R}$ ,  $g(\cdot, x)$  is Lebesgue  $\nabla$ -measurable on  $J$ ,
- (2) for a.e.  $(\nabla) t \in J$ ,  $g(t, \cdot)$  is continuous,
- (3) there is an  $h : J \rightarrow [0, \infty)$ ,  $\int_0^T h(s) \nabla s < \infty$  such that  $|g(t, p(t, x))| \leq h(t)$  a.e.  $(\nabla)$  on  $J$ , for all  $x \in \mathbb{R}$ ,
- (4)  $I : \mathbb{R} \rightarrow \mathbb{R}$  is continuous and nondecreasing,
- (5)  $B : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous and for each  $x \in [\alpha(0), \beta(0)]$ ,  $B(x, \cdot)$  is nonincreasing.

*Note.* a.e.  $(\nabla)$  denotes the Lebesgue  $\nabla$ -measure.

**THEOREM 2.2.** *Assume that conditions (1)–(5) are satisfied and  $\alpha, \beta$  are lower and upper solutions of (2.1)–(2.3) with  $\alpha(t) \leq \beta(t)$  for all  $t \in J$ . Then, there exists a solution  $u$  to (2.1)–(2.3) such that  $u \in [\alpha, \beta]$ .*

*Proof.* Our proof follows that of Cabada and Liz [9]. Define an operator  $G : A \rightarrow A$  by

$$Gu(t) = p(0, \bar{u}(0)) + \int_0^t g(s, p(s, u(s))) \nabla s, \quad t \in J_1, \tag{2.7}$$

$$Gu(t) = I(p(t_1, u(t_1))) + \int_{t_1}^t g(s, p(s, u(s))) \nabla s, \quad t \in J_2, \tag{2.8}$$

where  $\bar{u}(0) \equiv u(0) - B(u(0), u(T))$ .

*Claim 2.3.* If  $u$  is a fixed point of the operator  $G$ , then  $u$  is a solution of (2.1)–(2.3) such that  $u \in [\alpha, \beta]$ .

*Proof of Claim 2.3.* We assume that  $u \in A$  satisfies

$$u(t) = p(0, \bar{u}(0)) + \int_0^t g(s, p(s, u(s))) \nabla s, \quad t \in J_1, \tag{2.9}$$

$$u(t) = I(p(t_1, u(t_1))) + \int_{t_1}^t g(s, p(s, u(s))) \nabla s, \quad t \in J_2. \tag{2.10}$$

*Subclaim 1.*  $u(t) \in [\alpha(t), \beta(t)]$ , for all  $t \in J$ .

Note that, by letting  $t = 0$  in the right-hand side of (2.9), we have  $\int_{\emptyset} g(s, p(s, u(s))) \nabla s = 0$  and hence  $u(0) = p(0, \bar{u}(0))$  which is in  $[\alpha(0), \beta(0)]$  by the definition of  $p$ . Suppose there exists a  $t'_1 \in (0, t_1] \cap J$  such that  $\alpha(t'_1) > u(t'_1)$ . Since  $\alpha(0) \leq u(0)$ , there exists a  $t'_2 \in [0, t'_1] \cap J$  such that  $\alpha(t'_2) \leq u(t'_2)$  and  $\alpha(t) > u(t)$  on  $(t'_2, t'_1] \cap J$ . Then,  $g(t, p(t, u(t))) = g(t, \alpha(t))$  for all  $t \in (t'_2, t'_1] \cap J$ . We then have, for any  $t \in (t'_2, t'_1] \cap J$ ,

$$\begin{aligned} \int_0^t g(s, p(s, u(s))) \nabla s &= u(t) - p(0, \bar{u}(0)) = u(t'_2) - p(0, \bar{u}(0)) + u(t) - u(t'_2), \\ \int_0^{t'_2} g(s, p(s, u(s))) \nabla s &+ u(t) - u(t'_2) \\ \implies u(t) - u(t'_2) &= \int_0^t g(s, p(s, u(s))) \nabla s - \int_0^{t'_2} g(s, p(s, u(s))) \nabla s \\ &= \int_{t'_2}^t g(s, p(s, u(s))) \nabla s = \int_0^{t'_2} g(s, \alpha(s)) \nabla s. \end{aligned} \tag{2.11}$$

From assumption (ii) of the definition of lower solution, we have

$$\alpha(t) - \alpha(t'_2) \leq \int_{t'_2}^t g(s, \alpha(s)) \nabla s. \tag{2.12}$$

We then have

$$u(t) - u(t'_2) = \int_{t'_2}^t g(s, \alpha(s)) \nabla s \geq \alpha(t) - \alpha(t'_2) \tag{2.13}$$

and recalling that  $\alpha(t'_2) \leq u(t'_2)$  and  $u(t) < \alpha(t)$ , this is a contradiction. Hence,  $\alpha \leq u$  on  $J_1$ . Similarly,  $u \leq \beta$  on  $J_1$ .

We then have

$$\alpha(t_1) \leq u(t_1) \leq \beta(t_1), \tag{2.14}$$

and using the fact that  $I$  is nondecreasing, we have

$$\alpha(t_1^+) \leq I(\alpha(t_1)) \leq I(u(t_1)) \leq I(\beta(t_1)) \leq \beta(t_1^+). \tag{2.15}$$

We also have  $I(u(t_1)) = I(p(t_1, u(t_1))) = u(t_1^+)$  and hence from (2.15) we conclude

$$\alpha(t_1^+) \leq u(t_1^+) \leq \beta(t_1^+). \tag{2.16}$$

We may now proceed as before to get  $\alpha \leq u \leq \beta$  on  $J_2$ , establishing Subclaim 1.

We may apply Subclaim 1 to (2.9) to verify that  $u$  satisfies the first equation in property (ii) of a solution to (2.1)–(2.3), and apply Subclaim 1 to (2.10) to verify that  $u$  satisfies the second equation in property (ii).

*Subclaim 2.*  $\bar{u}(0) \in [\alpha(0), \beta(0)]$ .

Suppose that  $\alpha(0) > \bar{u}(0) = u(0) - B(u(0), u(T))$ . Thus,  $u(0) = p(0, \bar{u}(0)) = \alpha(0)$  and hence  $B(u(0), u(T)) > 0$ . Using assumption (5), we have  $B(\alpha(0), \alpha(T)) \geq B(\alpha(0), u(T)) > 0$ , which contradicts  $\alpha$  being a lower solution of (2.1)–(2.3). We then have  $\alpha(0) \leq \bar{u}(0)$  and, similarly,  $\bar{u}(0) \leq \beta(0)$ , establishing Subclaim 2.

As a result of Subclaim 2, we have  $u(0) = p(0, \bar{u}(0)) = \bar{u}(0) = u(0) - B(u(0), u(T))$  and hence  $B(u(0), u(T)) = 0$ , establishing Claim 2.3. □

*Claim 2.4.*  $G : A \rightarrow A$  has a fixed point.

*Proof of Claim 2.4.* We will apply Schauder’s fixed point theorem.

Let  $K = \|\alpha\| + \|\beta\|$ . Define  $w : J \rightarrow \mathbb{R}$  by  $w(t) = K + \int_0^t h(s) \nabla s$ .

Let

$$S = \{u \in A : |u(0)| \leq K, |u(t_1^+)| \leq K, |u(b) - u(a)| \leq w(b) - w(a) \text{ on } 0 \leq a \leq b \leq t_1 \text{ or } t_1 < a \leq b \leq T, \text{ where } a, b \in J\}. \tag{2.17}$$

It can be shown that  $S$  is a convex and compact subset of  $(A, \|\cdot\|)$ .

*Subclaim 3.*  $G(S) \subseteq S$ .

Let  $u \in S$  and consider  $Gu$ . Let  $t = 0$  in (2.7) to obtain

$$|Gu(0)| = |p(0, \bar{u}(0))| \leq \max\{|\alpha(0)|, |\beta(0)|\} \leq K. \tag{2.18}$$

Note that  $\alpha(t_1) \leq p(t_1, u(t_1)) \leq \beta(t_1)$ , hence

$$\begin{aligned} \alpha(t_1^+) &\leq I(\alpha(t_1)) \leq I(p(t_1, u(t_1))) \leq I(\beta(t_1)) \leq \beta(t_1^+), \\ |I(p(t_1, u(t_1)))| &\leq \max\{|\alpha(t_1^+)|, |\beta(t_1^+)|\} \leq K. \end{aligned} \tag{2.19}$$

Let  $t \downarrow t_1$  in (2.8) to obtain

$$|Gu(t_1^+)| = |I(p(t_1, u(t_1)))| \leq K. \tag{2.20}$$

Let  $a, b \in J$  with  $0 \leq a \leq b \leq t_1$ ,

$$|Gu(b) - Gu(a)| = \left| \int_a^b g(s, p(s, u(s))) \nabla s \right| \leq \int_a^b h(s) \nabla s = w(b) - w(a). \tag{2.21}$$

Similar results hold for  $t_1 < a \leq b \leq T$ .

*Subclaim 4.*  $G : S \rightarrow S$  is continuous.

Let  $\{u_n\}_{n=1}^\infty \subseteq S$  which converges to  $u \in S$  in the space  $(A, \|\cdot\|)$ . Note that  $u_n \rightarrow u$  uniformly on compact subsets of  $J$ . Let  $n \in \mathbb{N}$  and  $t \in J_1$ , then

$$\begin{aligned} Gu(t) - Gu_n(t) &= Gu(t) - p(0, \bar{u}(0)) - [Gu_n(t) - p(0, \bar{u}_n(0))] + p(0, \bar{u}(0)) - p(0, \bar{u}_n(0)) \\ &= \int_0^t g(s, p(s, u(s))) \nabla s - \int_0^t g(s, p(s, u_n(s))) \nabla s + p(0, \bar{u}(0)) - p(0, \bar{u}_n(0)) \end{aligned} \tag{2.22}$$

and hence

$$\begin{aligned} |Gu(t) - Gu_n(t)| &\leq \int_0^{t_1} |g(s, p(s, u(s))) - g(s, p(s, u_n(s)))| \nabla s + p(0, \bar{u}(0)) - p(0, \bar{u}_n(0)). \end{aligned} \tag{2.23}$$

Now take  $\lim_{n \rightarrow \infty}$  and apply the Lebesgue dominated convergence theorem and the continuity of  $g$  in its second variable and of  $p$  to conclude

$$\lim_{n \rightarrow \infty} |Gu_n(t) - Gu(t)| = 0. \tag{2.24}$$

Note that (2.23) does not involve  $t$  in its right-hand side, so we can conclude that the convergence is uniform on  $J_1$ .

A similar argument shows that  $Gu_n \rightarrow G$  uniformly on compact subsets of  $J_2$ .

Hence, by Subclaims 3 and 4, Schauder’s fixed point theorem applies to  $G$ , finishing the proof of Claim 2.4. □

Claims 2.3 and 2.4 yield the desired result. □

*Example 2.5.* Let  $\mathbb{T} = [0, 1] \cup [2, 3]$ ,  $t_1 = 2$ ,  $g(t, x) = t^2 + x^2$ ,  $I(x) = x + 1$ ,  $u(0) = 0$ . (Note that  $I$  is not bounded, as required in [4].)

$\alpha(t) = 0$  is a lower solution .

To construct  $\beta$  on  $[0, 1]$ , we solve  $\beta' = 1 + \beta^2 (\geq t^2 + \beta^2)$ ,  $\beta(0) = 0$ . Then this implies that  $\beta(t) = \tan t$ . By considering boundary conditions, we have

$$\begin{aligned} \beta(2) &= \beta(0) + \int_0^2 \beta^\nabla(s) \nabla s = \beta(0) + \int_0^1 \beta'(s) ds + \beta^\nabla(2), \\ \beta(2) = \tan 1 + \frac{\beta(2) - \beta(1)}{2 - 1} &\implies \beta(2) = \beta(2) \implies \beta(2) \text{ is arbitrary, let } \beta(2) = 1, \\ \beta(2^+) &= I(\beta(2)) = 2. \end{aligned} \tag{2.25}$$

To construct  $\beta$  on  $[2, 3]$ , we solve  $\beta' = 9 + \beta^2 (\geq t^2 + \beta^2)$ ,  $\beta(2) = 2$ , then we have

$$\beta = 3 \tan \left( \tan^{-1} \left( \frac{2}{3} \right) + 3t - 6 \right). \tag{2.26}$$

Applying Theorem 2.2, we know there exists a solution  $u$  such that  $0 \leq u(t) \leq \beta(t)$  for  $t \in \mathbb{T}$ .

### 3. Second order

In this section, we are concerned with second-order dynamic equations with functional boundary conditions and impulse:

$$y^{\Delta\Delta}(t) = f(t, y^\sigma(t)), \quad t \in \mathbb{T}^{\kappa^2} \equiv [a, b] \setminus \{t_1\}, \tag{3.1}$$

$$L_1(y(a), y^\Delta(a), y(\sigma^2(b)), y^\Delta(\sigma(b))) = 0, \tag{3.2}$$

$$L_2(y(a), y(\sigma^2(b))) = 0, \tag{3.3}$$

$$y(t_1^+) - y(t_1^-) = r_1, \tag{3.4}$$

$$y^\Delta(t_1^+) - y^\Delta(t_1^-) = I(y(t_1), y^\Delta(t_1^-)), \tag{3.5}$$

where  $t_1 \in \mathbb{T}$  with  $a < t_1 < b$  and  $t_1$  right-dense,  $r_1 \in \mathbb{R}$ ,  $I$  is a real-valued function and  $J = [a, b]$ . We set  $y^\Delta(t_1) = y^\Delta(t_1^+)$  if  $t_1$  is left-scattered, and  $y^\Delta(t_1) = y^\Delta(t_1^-)$  if  $t_1$  is left-dense. We note that these impulses are different from those studied in [16]. (3.2) and (3.3) cover many conditions found in the literature such as separated and nonseparated boundary conditions, respectively,

$$\begin{aligned} L_1(x, y, z, w) &= x, & L_2(x, y) &= y, \\ L_1(x, y, z, w) &= y - z, & L_2(x, y) &= y - x, \end{aligned} \tag{3.6}$$

as in [7, Chapter 4].

Let  $J_1 = [a, t_1]$ ,  $J_2 = (t_1, b]$ . We define the following spaces of functions.

Let  $y_i$  be the restriction of  $y : J \rightarrow \mathbb{R}$  to  $J_i$ ,  $i = 1, 2$ , then

$$\begin{aligned} \mathcal{C}'(J_1) &= \{y : J_1 \rightarrow \mathbb{R} : y \text{ and } y^\Delta \text{ are continuous on } J_1\}, \\ \mathcal{C}'(J_2) &= \{y : J_2 \rightarrow \mathbb{R} : y \text{ and } y^\Delta \text{ are continuous on } J_2 \text{ and } y(t_1^+) \text{ and } y^\Delta(t_1^+) \text{ exist}\}, \\ A &= \{y : J \rightarrow \mathbb{R} : y_1 \in \mathcal{C}'(J_1) \text{ and } y_2 \in \mathcal{C}'(J_2)\}. \end{aligned} \tag{3.7}$$

For  $y \in A$ , let  $\|y\| = \sup\{|y(t)| : t \in J\}$ .  $(A, \|\cdot\|)$  is a Banach space. For  $x, y \in A$ ,

$$[x, y] \equiv \{z \in A : x(t) \leq z(t) \leq y(t) \quad \forall t \in J\}. \tag{3.8}$$

Now we introduce the concept of lower and upper solutions of problem (3.1)–(3.5) as follows.



*Definition 3.1.* The functions  $\alpha$  and  $\beta$  are, respectively, a lower and an upper solution of problem (3.1)–(3.5) if the following properties hold:

- (i)  $\alpha, \beta \in A$ ;
- (ii)

$$\begin{aligned} \alpha^{\Delta\Delta}(t) &\geq f(t, \alpha^\sigma(t)) \quad \text{on } t \in [a, b] \setminus \{t_1\}, \\ L_1(\alpha(a), \alpha^\Delta(a), \alpha(\sigma^2(b)), \alpha^\Delta(\sigma(b))) &\geq 0, \\ L_2(\alpha(a), \alpha(\sigma^2(b))) = 0, \quad L_2(\alpha(a), \cdot) &\text{ is injective,} \\ \alpha(t_1^+) - \alpha(t_1^-) &= r_1, \\ \alpha^\Delta(t_1^+) - \alpha^\Delta(t_1^-) &\geq I(\alpha(t_1), \alpha^\Delta(t_1^-)); \end{aligned} \tag{3.9}$$

- (iii)

$$\begin{aligned} \beta^{\Delta\Delta}(t) &\leq f(t, \beta^\sigma(t)) \quad \text{on } t \in [a, b] \setminus \{t_1\}, \\ L_1(\beta(a), \beta^\Delta(a), \beta(\sigma^2(b)), \beta^\Delta(\sigma(b))) &\leq 0, \\ L_2(\beta(a), \beta(\sigma^2(b))) = 0, \quad L_2(\beta(a), \cdot) &\text{ is injective,} \\ \beta(t_1^+) - \beta(t_1^-) &= r_1, \\ \beta^\Delta(t_1^+) - \beta^\Delta(t_1^-) &\leq I(\beta(t_1), \beta^\Delta(t_1^-)). \end{aligned} \tag{3.10}$$

We assume the following conditions are satisfied for the functions  $f$ ,  $L_1$  and  $L_2$ , and  $I$ .

- (F) The function  $f : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous.
- (L)  $L_1 \in C(\mathbb{R}^4, \mathbb{R})$  is nondecreasing in the second variable, nonincreasing in the fourth. Moreover,  $L_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$  is a continuous function and it is nonincreasing with respect to its first variable.
- (I)  $I$  is continuous and strictly increasing with respect to the first variable and non-increasing in the second variable.

We consider the following modified truncated problem:

$$y^{\Delta\Delta}(t) - y^\sigma(t) = f(t, p(\sigma(t), y^\sigma(t))) - p(\sigma(t), y^\sigma(t)), \quad t \in \mathbb{T}^{\kappa^2} \equiv [a, b] \setminus \{t_1\}, \tag{3.11}$$

$$y(a) = L_1^*(y(a), y^\Delta(a), y(\sigma^2(b)), y^\Delta(\sigma(b))), \tag{3.12}$$

$$y(\sigma^2(b)) = L_2^*(y(a), y(\sigma^2(b))), \tag{3.13}$$

$$r_1 = y(t_1^+) - y(t_1^-), \tag{3.14}$$

$$y^\Delta(t_1^+) - y^\Delta(t_1^-) = I(y(t_1), y^\Delta(t_1^-)), \tag{3.15}$$

where  $p(t, y) = \min\{\max\{\alpha(t), y\}, \beta(t)\}$ ,

$$\begin{aligned} L_1^*(x, y, z, w) &= p(a, x + L_1(x, y, z, w)) \quad \forall (x, y, z, w) \in \mathbb{R}^4, \\ L_2^*(x, y) &= p(\sigma^2(b), y - L_2(x, y)) \quad \forall (x, y) \in \mathbb{R}^2. \end{aligned} \tag{3.16}$$

**THEOREM 3.2.** Assume that conditions (F) and (L) are satisfied. If there exist a lower solution  $\alpha$  and an upper solution  $\beta$  of (3.1)–(3.5) such that  $\alpha \leq \beta$  on  $\mathbb{T}$ , then the BVP (3.11)–(3.15) has a solution.

*Proof.* It is not difficult to verify that the problem

$$\begin{aligned} y^{\Delta\Delta} - y^\sigma &= 0, \quad t \in [a, b], \\ y(a) &= y(\sigma^2(b)) = 0, \end{aligned} \tag{3.17}$$

has only the trivial solution.

By using [7, Theorem 4.67 and Corollary 4.74], we have that for every  $h \in \mathbb{C}_{rd}[a, b]$  and  $A, B \in \mathbb{R}$ , the problem

$$\begin{aligned} y^{\Delta\Delta} - y^\sigma &= h(t), \quad t \in [a, b], \\ y(a) &= A, \quad y(\sigma^2(b)) = B, \end{aligned} \tag{3.18}$$

has a solution  $y(t)$  if and only if the operator

$$Qy(t) = Ay_1(t) + By_2(t) + \int_a^{\sigma(b)} G(t, s)h(s)\Delta s \tag{3.19}$$

has a fixed point. Here  $y_1(t), y_2(t)$  are the solutions of the linear homogeneous equation  $y^{\Delta\Delta} - y^\sigma = 0, t \in [a, b]$  and satisfy the boundary conditions  $y_1(a) = 1, y_1(\sigma^2(b)) = 0$  and  $y_2(a) = 0, y_2(\sigma^2(b)) = 1$ .

$G$  is called the Green's function of the Dirichlet problem. One can verify that (see [7, page 169]) it is continuous in  $[a, \sigma^2(b)] \times [a, \sigma^2(b)]$  and  $G^\Delta(\cdot, s)$  is continuous at  $t \neq s = \sigma(s)$  and bounded in  $[a, \sigma^2(b)]$ .

Define

$$\begin{aligned} Qy(t) &= L_1^*(y(a), y^\Delta(a), y(\sigma^2(b)), y^\Delta(\sigma(b)))y_1(t) + L_2^*(y(a), y(\sigma^2(b)))y_2(t) \\ &\quad + \int_a^{\sigma(b)} G(t, s)\{f(s, p(\sigma(s), y^\sigma(s))) - p(\sigma(s), y(\sigma(s)))\}\Delta s + L(t, y(t)), \end{aligned} \tag{3.20}$$

where

$$L(t, y) = \begin{cases} \frac{y_2(t)}{W} [y_1(t_1)I(y(t_1), y^\Delta(t_1^-)) - r_1 y_1^\Delta(t_1)], & a \leq t \leq t_1, \\ \frac{y_1(t)}{W} [y_2(t_1)I(y(t_1), y^\Delta(t_1^-)) - r_1 y_2^\Delta(t_1)], & t_1 \leq t \leq \sigma^2(b), \end{cases} \tag{3.21}$$

where  $W = y_2(t_1)y_1^\Delta(t_1) - y_1(t_1)y_2^\Delta(t_1)$ .

One can easily observe that  $Q$  has a fixed point  $y$  if and only if  $y$  is a solution of (3.11)–(3.15).

Since  $L_1, L_2, p,$  and  $G$  are bounded and continuous, it can be shown that there exists  $R > 0$  such that the compact operator  $Q : S \rightarrow S$  where  $S = \{y \in A : \|y\| \leq R\}$ .

Since  $S$  is a closed, bounded, and convex set, in view of the Tychonoff-Schauder fixed point theorem, there is at least one fixed point of  $Q$ . □

**THEOREM 3.3.** *Assume that conditions (F) and (I) are satisfied. Let  $\alpha$  and  $\beta$  be a lower and upper solution, respectively, of the problem (3.1)–(3.5) such that  $\alpha \leq \beta$  on  $\mathbb{T}$ . Then every solution of the BVP (3.11)–(3.15) belongs to the sector  $[\alpha, \beta]$ .*

*Proof.* Let  $y$  be a solution of (3.11)–(3.15), by definition of  $L_1^*$  and  $L_2^*$ , we know that  $y(a) \in [\alpha(a), \beta(a)]$  and  $y(\sigma^2(b)) \in [\alpha(\sigma^2(b)), \beta(\sigma^2(b))]$ . We will prove that  $y \in [\alpha, \beta]$  for  $t \in (a, \sigma^2(b))$ .

Consider  $z(t) = y(t) - \beta(t)$ . By definition of  $\beta$ ,  $z$  is continuous on  $[a, \sigma^2(b)]$ . Suppose, to the contrary, there is a  $t^* \in (a, t_1) \cup (t_1, \sigma^2(b))$  such that  $(y - \beta)(t^*) = \max_{t \in \mathbb{T}} \{y(t) - \beta(t)\} > 0$ .

Suppose that  $t^*$  is left scattered. In this case, we have that

$$y^\Delta(t^*) \leq \beta^\Delta(t^*), \quad y^{\Delta\Delta}(\rho(t^*)) \leq \beta^{\Delta\Delta}(\rho(t^*)). \tag{3.22}$$

Consequently, by using condition  $F$ , we arrive at the following contradiction:

$$\begin{aligned} 0 &> y^{\Delta\Delta}(\rho(t^*)) - \beta^{\Delta\Delta}(\rho(t^*)) - (y(t^*) - \beta(t^*)) \\ &\geq f(\rho(t^*), \beta(t^*)) - f(\rho(t^*), \beta(t^*)) = 0. \end{aligned} \tag{3.23}$$

When  $t^*$  is left dense the contradiction holds in a similar way.

Now suppose  $t^* = t_1$ .

*Case 1.*  $t_1$  is left scattered.

Then we have  $z^\Delta(t_1^+) \leq 0$  and  $z^\Delta(t_1^+) = z^\Delta(t_1)$ . Consequently, by using condition (I), we arrive at the following contradiction:

$$\begin{aligned} 0 &= z^\Delta(t_1^+) - z^\Delta(t_1^-) \\ &= y^\Delta(t_1^+) - y^\Delta(t_1^-) - [\beta^\Delta(t_1^+) - \beta^\Delta(t_1^-)] \\ &\geq I(y(t_1), y^\Delta(t_1)) - I(\beta(t_1), \beta^\Delta(t_1)) \\ &> I(\beta(t_1), y^\Delta(t_1)) - I(\beta(t_1), \beta^\Delta(t_1)) \geq 0. \end{aligned} \tag{3.24}$$

*Case 2.*  $t_1$  is left dense.

For sufficiently small  $\epsilon > 0$ , we have

$$z^\Delta(s) \geq 0 \quad \text{for } s \in (t_1 - \epsilon, t_1), \quad z^\Delta(s^*) \leq 0 \quad \text{for } s^* \in (t_1, t_1 + \epsilon). \tag{3.25}$$

Then the contradiction holds in a similar way.

Analogously, the fact that  $\alpha(t) \leq y(t)$  for all  $t \in \mathbb{T}$  can be shown. □

**THEOREM 3.4.** *Assume that (L) holds. If  $y \in [\alpha, \beta]$  is a solution of (3.11)–(3.15), then  $y$  satisfies equalities (3.1)–(3.5).*

*Proof.* If  $y(\sigma^2(b)) - L_2(y(a), y(\sigma^2(b))) < \alpha(\sigma^2(b))$ , the definition of  $L_2^*$  gives us that  $y(\sigma^2(b)) = \alpha(\sigma^2(b))$ .

Now using (L), we obtain a contradiction:

$$\begin{aligned} \alpha(\sigma^2(b)) &> y(\sigma^2(b)) - L_2(y(a), y(\sigma^2(b))) \\ &\geq \alpha(\sigma^2(b)) - L_2(\alpha(a), \alpha(\sigma^2(b))) = \alpha(\sigma^2(b)). \end{aligned} \tag{3.26}$$

Analogously, we arrive at  $\alpha(\sigma^2(b)) \leq y(\sigma^2(b)) - L_2(y(a), y(\sigma^2(b))) \leq \beta(\sigma^2(b))$  and (3.13) implies that  $L_2(y(a), y(\sigma^2(b))) = 0$ .

To prove that  $L_1(y(a), y^\Delta(a), y(\sigma^2(b)), y^\Delta(\sigma(b))) = 0$ , it is enough using (3.12) to show

$$\alpha(a) \leq y(a) + L_1(y(a), y^\Delta(a), y(\sigma^2(b)), y^\Delta(\sigma(b))) \leq \beta(a). \quad (3.27)$$

If  $y(a) + L_1(y(a), y^\Delta(a), y(\sigma^2(b)), y^\Delta(\sigma(b))) < \alpha(a)$ , then  $y(a) = \alpha(a)$  implies that  $0 = L_2(y(a), y(\sigma^2(b))) = L_2(\alpha(a), \alpha(\sigma^2(b)))$ .

Since  $L_2$  is injective with respect to the second variable, we have  $y(\sigma^2(b)) = \alpha(\sigma^2(b))$ .

Using the definition of  $L_1$ , we obtain a contradiction:

$$\begin{aligned} \alpha(a) &> y(a) + L_1(y(a), y^\Delta(a), y(\sigma^2(b)), y^\Delta(\sigma(b))) \\ &\geq \alpha(a) + L_1(\alpha(a), \alpha^\Delta(a), \alpha(\sigma^2(b)), \alpha^\Delta(\sigma(b))) \geq \alpha(a). \end{aligned} \quad (3.28)$$

Here we used the fact that  $y \in [\alpha, \beta]$ , and  $\alpha(a) = y(a)$ ,  $\alpha(\sigma^2(b)) = y(\sigma^2(b))$ , consequently, it follows that  $y^\Delta(a) \geq \alpha^\Delta(a)$  and  $y^\Delta(\sigma(b)) \leq \alpha^\Delta(\sigma(b))$ . The other inequality holds similarly.  $\square$

*Example 3.5.* Let  $\mathbb{T}$  be any time scale and let the point  $1/2$  be a right-dense point in  $\mathbb{T} \cap [0, 1]$ . We define  $f$ ,  $L_1$ , and  $L_2$  in the following way:

$$\begin{aligned} f(t, y) &= y \sinh((y-1)^2), & L_1(x, y, z, w) &= 1-x, \\ L_2(x, y) &= -y, & I(x, y) &= x-1. \end{aligned} \quad (3.29)$$

Next we consider the following boundary value problem:

$$y^{\Delta\Delta}(t) = f(t, y^\sigma(t)), \quad t \in [0, 1]^{k^2} \setminus \left\{ \frac{1}{2} \right\}, \quad (3.30)$$

$$y(0) = 1, \quad (3.31)$$

$$y(1) = 0, \quad (3.32)$$

$$y\left(\frac{1^+}{2}\right) - y\left(\frac{1^-}{2}\right) = -1, \quad (3.33)$$

$$y^\Delta\left(\frac{1^+}{2}\right) - y^\Delta\left(\frac{1^-}{2}\right) = I\left(y\left(\frac{1}{2}\right), y^\Delta\left(\frac{1^-}{2}\right)\right). \quad (3.34)$$

One can easily verify that

$$\alpha(t) = \begin{cases} 0, & \text{if } t \in \left[0, \frac{1}{2}\right], \\ 2(t-1), & \text{if } t \in \left(\frac{1}{2}, 1\right], \end{cases} \quad (3.35)$$

is a lower solution and

$$\beta(t) = \begin{cases} 2t+1, & \text{if } t \in \left[0, \frac{1}{2}\right], \\ -2(t-1), & \text{if } t \in \left(\frac{1}{2}, 1\right], \end{cases} \quad (3.36)$$

is an upper solution of the problem (3.30)–(3.34).

Theorem 3.3 assures that there exists a solution  $y(t)$  of the problem (3.30)–(3.34) such that  $y \in [\alpha, \beta]$ . We note that

$$y(t) = \begin{cases} 1, & \text{if } t \in \left[0, \frac{1}{2}\right], \\ 0, & \text{if } t \in \left(\frac{1}{2}, 1\right], \end{cases} \quad (3.37)$$

is one such solution.

## References

- [1] A. Adje, *Sur et Sous-Solutions dans les Equations Differentielles Discontinues avec des Conditions aux Limites non Linéaires*, Ph.D. thesis, Université Catholique de Louvain-la-Neuve, Belgium, 1987.
- [2] F. M. Atici and D. C. Biles, *First order dynamic inclusions on time scales*, J. Math. Anal. Appl. **292** (2004), no. 1, 222–237.
- [3] F. M. Atici and G. Sh. Guseinov, *On Green's functions and positive solutions for boundary value problems on time scales*, J. Comput. Appl. Math. **141** (2002), no. 1-2, 75–99.
- [4] M. Benchohra, J. Henderson, S. K. Ntouyas, and A. Ouahab, *On first order impulsive dynamic equations on time scales*, J. Difference Equ. Appl. **10** (2004), no. 6, 541–548.
- [5] M. Benchohra, S. K. Ntouyas, and A. Ouahab, *Existence results for second order boundary value problem of impulsive dynamic equations on time scales*, J. Math. Anal. Appl. **296** (2004), no. 1, 65–73.
- [6] M. Benchohra, S. K. Ntouyas, and A. Ouahabi, *Existence results for impulsive semilinear damped differential equations*, Int. J. Appl. Math. **11** (2002), no. 1, 77–93.
- [7] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales, an Introduction with Applications*, Birkhäuser Boston, Massachusetts, 2001.
- [8] ———, *Advances in Dynamic Equations on Time Scales*, Birkhäuser Boston, Massachusetts, 2003.
- [9] A. Cabada and E. Liz, *Discontinuous impulsive differential equations with nonlinear boundary conditions*, Nonlinear Anal. **28** (1997), no. 9, 1491–1497.
- [10] ———, *Boundary value problems for higher order ordinary differential equations with impulses*, Nonlinear Anal. **32** (1998), no. 6, 775–786.
- [11] A. Cabada, E. Liz, and S. Lois, *Green's function and maximum principle for higher order ordinary differential equations with impulses*, Rocky Mountain J. Math. **30** (2000), no. 2, 435–446.
- [12] V. Doddaballapur, P. W. Eloe, and Y. Zhang, *Quadratic convergence of approximate solutions of two-point boundary value problems with impulse*, Proceedings of the Third Mississippi State Conference on Difference Equations and Computational Simulations (Mississippi, 1997), Electron. J. Differ. Equ. Conf., vol. 1, Southwest Texas State Univ., Texas, 1998, pp. 81–95.
- [13] L. H. Erbe and X. Liu, *Existence results for boundary value problems of second order impulsive differential equations*, J. Math. Anal. Appl. **149** (1990), no. 1, 56–69.
- [14] D. Franco and J. J. Nieto, *First-order impulsive ordinary differential equations with anti-periodic and nonlinear boundary conditions*, Nonlinear Anal. **42** (2000), no. 2, 163–173.
- [15] M. Frigon and D. O'Regan, *Existence results for first-order impulsive differential equations*, J. Math. Anal. Appl. **193** (1995), no. 1, 96–113.
- [16] J. Henderson, *Double solutions of impulsive dynamic boundary value problems on a time scale*, J. Difference Equ. Appl. **8** (2002), no. 4, 345–356.
- [17] V. Lakshmikantham, D. D. Bañov, and P. S. Simeonov, *Theory of Impulsive Differential Equations*, Series in Modern Applied Mathematics, vol. 6, World Scientific, New Jersey, 1989.

- [18] E. Liz, *Existence and approximation of solutions for impulsive first order problems with nonlinear boundary conditions*, *Nonlinear Anal.* **25** (1995), no. 11, 1191–1198.
- [19] C. Pierson-Gorez, *Impulsive differential equations of first order with periodic boundary conditions*, *Differential Equations Dynam. Systems* **1** (1993), no. 3, 185–196.
- [20] J. Qi and K. Wang, *Upper and lower solutions for impulsive differential equations with application to ODE*, *Northeast. Math. J.* **18** (2002), no. 3, 189–196.
- [21] A. M. Samoilenko and N. A. Perestyuk, *Impulsive Differential Equations*, *World Scientific Series on Nonlinear Science. Series A: Monographs and Treatises*, vol. 14, World Scientific, New Jersey, 1995.

F. M. Atici: Department of Mathematics, Western Kentucky University, Bowling Green, KY 42101, USA

*E-mail address:* ferhan.atici@wku.edu

D. C. Biles: Department of Mathematics, Western Kentucky University, Bowling Green, KY 42101, USA

*E-mail address:* daniel.biles@wku.edu