MEETING ABSTRACT

Open Access

Evaluation of hTERC (3q26) gain by fluorescence in situ hybridization (FISH) to predict the behavior of oral epithelial precursor lesions

Silvia Gabba¹, Elisa Giannini¹, Luisa Borghi¹, Giovanni Fellegara¹, Chiara Marcialis¹, Maria Grugni¹, Maria Pia Foschini², Elena Repetti¹, Tshering Dorji¹, Giorgio Vittadini^{3*}, Vittorio Grazioli¹

From EPMA-World Congress 2013
Brussels, Belgium. 20-21 September 2013

Oral Squamous Cell Carcinoma (OSCC) is the most common tumor of the oral cavity. Approximately 500.000 new cases of OSCC are diagnosed worldwide each year with the incidence seemingly increasing. Chemo- and radiotherapy are potential curative procedures although destructive surgery is often necessary if early diagnosis is not achieved. Such procedures impact heavily on human and social costs. Infection of oral mucosa by oncogenic HPV is strongly associated with OSCC among subjects with established risk factors such as tobacco and alcohol abuse. The development of OSCC is a multistep process in which accumulated genetic and epigenetic events lead to cell cycle deregulation and chromosomal abnormalities. Gain of chromosome 3 (which contains human telomerase RNA gene, hTERC, located on chromosome 3q26) appears to be an important selective advantage in the progression of cervical intraepithelial neoplasia to invasive cancer [1] as well as in the progression to invasive head and neck carcinoma [2]. We utilized automated interphase FISH to evaluate the gain of chromosome 3 (3q26) and 7 (CEP) in oral epithelial precursor lesions in order to determine whether aneuploidy is already present and thus whether it is possible to anticipate diagnosis and clinical interventions. Retrospective in blind analysis on 40 oral samples with long-term follow up (ranging from 5 to 10 years) and with initial histological diagnosis of mild and moderate dysplasia, will be run. Preliminary results (8 samples) showed that tumor cells were already present

Figure 1

only in the cases that progressed to cancer (4/8) [figure 1]; these preliminary findings confirm the usefulness of FISH as a reflex test in case of morphologically assessed pre-cancerous lesions, and highlight the potential value of this technique for early diagnosis and thus earlier treatment with less destructive therapeutic approaches.

Authors' details

¹Laboratorio Analisi, Centro Diagnostico Italiano, Milano, Italy. ²Anatomia patologica, Università degli Studi di Bologna, Italy. ³Bracco Imaging, Centro Ricerche Bracco, San Donato Milanese, Italy.

Published: 11 February 2014

References

1. Heselmeyer K, Schrock E, du Manoir S, et al: Gain of chromosome 3q defines the transition from severe dysplasia to invasive carcinoma of the uterine cervix. *Proc Natl Acad Sci* 1996, **93**:479-484.

* Correspondence: Giorgio.Vittadini@bracco.com

³Bracco Imaging, Centro Ricerche Bracco, San Donato Milanese, Italy
Full list of author information is available at the end of the article

2. Singh JP, Stoffel A, Gogineni S, et al: **Amplification of the 3q26.3 Locus is Associated with Progression to Invasive Cancer and is a Negative Prognostic Factor in Head and Neck Squamous Cell Carcinomas.** *Am J Pathol* 2002, **161**:365-371.

doi:10.1186/1878-5085-5-S1-A57

Cite this article as: Gabba et al.: Evaluation of hTERC (3q26) gain by fluorescence in situ hybridization (FISH) to predict the behavior of oral epithelial precursor lesions. *EPMA Journal* 2014 **5**(Suppl 1):A57.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit