MEETING ABSTRACT

Open Access

Personalized medicine in common mental disorders

Hans Jörgen Grabe

From EPMA-World Congress 2013
Brussels, Belgium. 20-21 September 2013

Scientific objectives

Mental disorders affect up to 40 % of all individuals living in Europe at least once a life time. Serious mental disorders with chronic or relapsing course like major depression bipolar disorder and schizophrenia are disabling and often disintegrating disorders. The early identification of subjects being at risk for serious mental disorders holds the promise of early intervention and better long-term outcomes. However, still reliable early indicators are largely missing to date. Like in dementia research major research initiatives need to be developed to address those urgent issues.

Technological approaches

New MRI imaging technique, genotyping, gene methylation analyses, metabolom analyses, validated questionnaires for the assessment of psychosocial risk factors.

Results interpretation

Advanced statistical modeling of brain imaging data, support vector machine approaches, systems integration of OMICS data; data mining approaches for risk marker identification.

Outlook and expert recommendations

The identification of peripheral risk indicators of severe mental diseases is probably limited to rare but highly penetrative genetic variants. Possibly gene-methylation patterns that emerge in the light of abuse and traumatization be constitute another source of valuable information. The non-invasive MRI imaging technologies might be very promising but still time consuming and expensive. Especially functional MRI studies are only feasible in specialist centers. The most promising approach

seems to create a set of specific markers throughout the different levels of information to integrate biological, psychosocial and structural information on each subject.

Published: 11 February 2014

References

1. Grabe HJ, Schwahn C, Appel K, Mahler J, Schulz A, Spitzer C, Barnow S, John U, Freyberger HJ, Rosskopf D, Völzke H: **Update on the 2005 paper: Moderation of mental and physical distress by polymorphisms in the 5-HT transporter gene by interacting with social stressors and chronic disease burden.** *Mol Psychiatry* 2011, **16**(4):354-6.
2. Appel K, Schwahn C, Mahler J, Schulz A, Spitzer C, Fenske K, Stender J, Barnow S, John U, Teumer A, Biffar R, Nauck M, Völzke H, Freyberger HJ, Grabe HJ: **Moderation of Adult Depression by a Polymorphism in the FKBP5 Gene and Childhood Physical Abuse in the General Population.** *Neuropsychopharmacology* 2011, **36**(10):1982-91.
3. Grabe HJ, Lange M, Wolff B, Völzke H, Lucht M, Freyberger HJ, John U, Cascorbi I: **Mental and physical distress is modulated by a polymorphism in the 5-HT transporter gene interacting with social stressors and chronic disease burden.** *Mol Psychiatry* 2005, **10**:220-4.
4. Stein JL, Medland SE, Vasquez AA, Hibar DP, Grabe HJ, Meyer-Lindenberg A, Morris DW, Müller-Myhsok B, Nichols TE, Ophoff RA, Paus T, Pausova Z, Penninx BW, Potkin SG, Sämann PG, Saykin AJ, Schumann G, Smoller JW, Wardlaw JM, Weale ME, Martin NG, Franke B, Wright MJ, Thompson PM, for the Enhancing Neuroimaging Genetics through Meta-Analysis (ENIGMA) Consortium: **Common genetic polymorphisms are associated with human hippocampal and intracranial volumes.** *Nat Genet* 2012, **44**(5):552-561.

doi:10.1186/1878-5085-5-S1-A94

Cite this article as: Grabe: Personalized medicine in common mental disorders. *EPMA Journal* 2014 **5**(Suppl 1):A94.

Correspondence: grabeh@uni-greifswald.de
Department of Psychiatry and Psychotherapy, Ellernholzstraße 1-2, 17475
Greifswald, Germany