

RESEARCH

Open Access



# Genomic insights into the evolution and mechanisms of carbapenem-resistant hypervirulent *Klebsiella pneumoniae* co-harboring $bla_{KPC}$ and $bla_{NDM}$ : implications for public health threat mitigation

Qian Wang<sup>1†</sup>, Yue Liu<sup>1†</sup>, Ran Chen<sup>1,2</sup>, Meng Zhang<sup>2</sup>, Zaifeng Si<sup>2</sup>, Yueling Wang<sup>1</sup>, Yan Jin<sup>1</sup>, Yuanyuan Bai<sup>1</sup>, Zhen Song<sup>1</sup>, Xinglun Lu<sup>1</sup>, Mingju Hao<sup>3\*</sup> and Yingying Hao<sup>1,2\*</sup>

## Abstract

**Background** Carbapenem-resistant hypervirulent *Klebsiella pneumoniae* (CR-hvKP) co-producing  $bla_{KPC}$  and  $bla_{NDM}$  poses a serious threat to public health. This study aimed to investigate the mechanisms underlying the resistance and virulence of CR-hvKP isolates collected from a Chinese hospital, with a focus on  $bla_{KPC}$  and  $bla_{NDM}$  dual-positive hvKP strains.

**Methods** Five CR-hvKP strains were isolated from a teaching hospital in China. Antimicrobial susceptibility and plasmid stability testing, plasmid conjugation, pulsed-field gel electrophoresis, and whole-genome sequencing (WGS) were performed to examine the mechanisms of resistance and virulence. The virulence of CR-hvKP was evaluated through serum-killing assay and *Galleria mellonella* lethality experiments. Phylogenetic analysis based on 16 highly homologous carbapenem-resistant *K. pneumoniae* (CRKP) producing KPC-2 isolates from the same hospital was conducted to elucidate the potential evolutionary pathway of CRKP co-producing NDM and KPC.

**Results** WGS revealed that five isolates individually carried three unique plasmids: an IncFIB/IncHI1B-type virulence plasmid, IncFII/IncR-type plasmid harboring KPC-2 and IncC-type plasmid harboring NDM-1. The conjugation test results indicated that the transference of KPC-2 harboring IncFII/IncR-type plasmid was unsuccessful on their own, but could be transferred by forming a hybrid plasmid with the IncC plasmid harboring NDM. Further genetic analysis confirmed that the pJNKPN26-KPC plasmid was entirely integrated into the IncC-type plasmid via the copy-in route, which was mediated by *TnA51* and IS26.

**Conclusion** KPC-NDM-CR-hvKP likely evolved from a KPC-2-CRKP ancestor and later acquired a highly transferable  $bla_{NDM-1}$  plasmid. ST11-KL64 CRKP exhibited enhanced plasticity. The identification of KPC-2-NDM-1-CR-hvKP highlights the urgent need for effective preventive strategies against aggravated accumulation of resistance genes.

<sup>†</sup>Qian Wang and Yue Liu contributed equally to this work.

\*Correspondence:

Mingju Hao  
haomingju@163.com  
Yingying Hao  
haoyyjie@163.com

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Keywords** *Bla*<sub>KPC</sub>, *Bla*<sub>NDM</sub>, ST11-KL64, CR-hvKP, IncC-type plasmid

## Introduction

*Klebsiella pneumoniae* is known to cause both community-acquired and nosocomial infections globally and is regarded as one of the most prevalent and clinically relevant opportunistic pathogens [1]. With limited treatment options, carbapenem-resistant *K. pneumoniae* (CRKP) poses an enormous public health threat owing to increasing antibiotic resistance as a result of the acquisition of multiple mobile genetic elements [2].

Since 2017, China has experienced a concerning outbreak of ST11 carbapenem-resistant hypervirulent *K. pneumoniae* (ST11-CR-hvKP), associated with high mortality rates [3]. A nationwide survey found ST11-KL64 *K. pneumoniae* as the most prevalent CRKP clone in China, with *bla*<sub>KPC-2</sub>, *bla*<sub>NDM-1</sub>, and *bla*<sub>OXA-48</sub> being the predominant carbapenemase genes among ST11-CRKP [4]. The emergence of the ST11-KL64 CR-hvKP clone in China is primarily attributable to the continuous coevolution of plasmids carrying hypervirulence and carbapenem resistance genes. These variants potentially develop in two directions: (i) CR-hvKP, an hvKP that acquires a plasmid encoding carbapenemase, and (ii) hypervirulent carbapenem-resistant *K. pneumoniae* (hv-CRKP), a CRKP strain that acquires a virulence plasmid. Initially, pLVPK-like virulence plasmids were described as nonconjugative; however, recent studies revealed that pLVPK-like virulence plasmids could be transferred from hvKP strains to ST11-CRKP with the help of a self-transferable IncF plasmid in various modes [5].

Of note, ST11-CR-hvKP co-harboring *bla*<sub>KPC</sub> or *bla*<sub>NDM</sub> has been increasingly reported, thereby necessitating urgent control measures. Nonetheless, evolutionary and dissemination strategies remain to be elucidated. A previous study based on the phylogenetic analysis of genomes from public databases showed that the formation of the superbug co-carrying NDM and KPC mainly emerged from a KPC-2-CRKP progenitor, which later acquired another highly transferable *bla*<sub>NDM-1</sub> plasmid [2]. However, further genomic studies and experiments are required to verify this hypothesis. The present study aimed to investigate the mechanisms underlying the resistance and virulence of CR-hvKP isolates collected from a Chinese hospital, with a focus on *bla*<sub>KPC</sub> and *bla*<sub>NDM</sub> dual-positive hvKP strains.

## Materials and methods

### Bacterial strains

Five non-duplicate ST11-KL64 CR-KP isolates co-producing *bla*<sub>KPC</sub> and *bla*<sub>NDM</sub> were collected from Shandong Provincial Hospital between January 2017 and June 2020. Permission to report cases was obtained from patients or their representatives.

### Antimicrobial susceptibility testing

Antimicrobial susceptibility testing was conducted using the VITEK-2 Compact system (bioMérieux, France) in accordance with the Clinical and Laboratory Standards Institute guidelines. The minimum inhibitory concentrations (MICs) of tigecycline and colistin were determined using the broth dilution method and interpreted in accordance with the US Food and Drug Administration standards and the European Committee on Antimicrobial Susceptibility Testing ([http://www.eucast.org/clinical\\_breakpoints](http://www.eucast.org/clinical_breakpoints)) guidelines, respectively.

### Conjugation assay

The transfer ability of KPC and NDM harboring plasmids was assessed using JNKPN26 as the representative donor and *Escherichia coli* J53<sup>Azi<sup>R</sup></sup> as the recipient strain. A second round of conjugation assay was conducted using transconjugants as donor strains and EC600 as the recipient strain to test the mobilization of the hybrid plasmids. Conjugation was performed in liquid Luria–Bertani (LB) medium incubated at 37 °C without shaking. The *bla*<sub>KPC-2</sub>- and/or *bla*<sub>NDM-1</sub>-bearing transconjugants were selected on China Blue lactose agar supplemented with sodium azide and ceftazidime. In order to improve the screening efficiency of transconjugants carrying KPC, the *bla*<sub>NDM</sub> expression was suppressed by adding 0.1 mM EDTA to the screening plate. For all transconjugants, polymerase chain reaction (PCR) detection combined with XbaI and S1 pulsed-field gel electrophoresis (S1-PFGE) was employed to validate the presence of *bla*<sub>NDM-1</sub> and/or *bla*<sub>KPC-2</sub> harboring the plasmids. The conjugation frequency was determined by calculating the ratio of transconjugants to recipients based on the colony forming unit (CFU) count on serial dilution plates with matching antibiotics.

### Plasmid stability assay

Hybrid plasmid stability was assessed as previously described, with moderate modifications [2]. Briefly, isolates were cultured in LB broth at 37 °C with shaking

(200 rpm) and serially passaged for 7 days (approximately 140 generations) with 1:1000 dilutions in antibiotic-free LB broth. Cultures were serially diluted and plated on Mueller Hinton agar plates without antibiotics every 24 h for 7 days. Two pairs of specific primers flanking the fusion site were designed to validate the fusion state of plasmids carrying *bla*<sub>NDM-1</sub> and *bla*<sub>KPC-2</sub>. Furthermore, 48 bacterial colonies were randomly selected, and PCR was performed every 24 h. The primers used in the assay are listed in Additional file 3: Table S1.

### Whole-genome sequencing (WGS) and bioinformatics analysis

Genomic DNA sequencing was conducted by Novogene Co., Ltd. (Beijing, China) using both Illumina HiSeq and Nanopore platforms. Sequence assembly and annotation were conducted following the methodology outlined in previous research [6]. Samtools was used to assess the coverage [7].

Orthologous gene groups were identified using OrthoFinder version 2.3.12 [8]. Multiple sequence alignment was performed using MAFFT [9], and phylogenetic reconstruction was conducted using Gubbins [10]. Based on the sample separation time, a sample time-evolution relationship was constructed using BactDating [11]. Transmission tree inference was conducted using the Bayesian program TransPhylo [12].

### *Galleria mellonella* infection model

The virulence of CRKPs was examined using a *G. mellonella* infection model. Larvae weighing approximately 300 mg (Tianjin Huiyude Biotech Company, Tianjin, China) were kept in the dark at approximately 10 °C for backup. Overnight cultures of *K. pneumoniae* strains were washed with phosphate-buffered saline and further adjusted to concentrations of  $1 \times 10^5$ ,  $1 \times 10^6$ , and  $1 \times 10^7$  CFU/mL. ATCC13883 and NTUH-K2044 were used as low- and high-virulence controls, respectively. Each group comprised eight *G. mellonella* larvae. The survival rate of *G. mellonella* was recorded every 12 h, and all experiments were performed in triplicate. The results for the *G. mellonella* model were analyzed using Kaplan–Meier survival curves and log-rank tests.

### Serum-killing assay

In vitro virulence was evaluated using a serum-killing assay, as described previously [13]. Briefly, 25 µL of bacterial suspension (at a concentration of  $1 \times 10^6$  CFU/mL) was added to 75 µL of mixed serum from 10 healthy humans for co-culture in a microtiter plate. After 0, 1, 2, and 3 h, the plates were inoculated with the mixture, and the number of viable bacteria was determined. The test was independently performed at least three times,

and the percentage of CFU counts was characterized as the result of serum resistance, which was graded from 1 to 6. With normal human serum, a strain was generally considered to be resistant if it could attain grades 5–6, as previously described [13]. Unpaired two-sided Student's t-test was performed for the strains, and data were presented as means  $\pm$  standard deviation (SD).

## Results

### Clinical and strain features

Patients with KPC-NDM-CR-KP infections were admitted to Shandong Provincial Hospital's ICU from January to April 2019. Most had severe pneumonia, and one had postoperative pulmonary infection. Patients developed symptoms like pulmonary edema, pleural effusion, sputum, and shortness of breath, requiring mechanical ventilation. The strains were isolated within 3–42 days of admission. Patients 1 and 2 died due to severe pneumonia, sepsis, and multi-organ failure, while patient 4 discontinued treatment after septic shock. The other three improved with antibiotic therapy (Table 1, Fig. 1). All strains were highly resistant to  $\beta$ -lactam antibiotics, including carbapenems, but susceptible to tigecycline. JNKPN26 isolated from patient 3 was resistant to polymyxin (MIC > 64 mg/L) (Table 2).

Phylogenetic analysis of these KPs and eleven other CRKPs from the hospital showed that five KPC-NDM-CR-KPs and NDM-CR-KP JNKPN30 formed a separate clade (Subclade Ib of Clade I) (Fig. 2). Six strains had identical PFGE profiles, indicating the same clone. This clade appeared to evolve between 2018 and 2019, possibly due to acquiring a *bla*<sub>NDM</sub>-positive plasmid. Genetic analysis of JNKPN23, JNKPN26 and JNKPN30 revealed a chromosome (~5470 kb) and three plasmids per isolate, revealing nearly 99.7% coverage more than 200 $\times$  in depth: a virulence plasmid (~220 kb), a KPC-positive plasmid (~120 kb), and an NDM-positive plasmid (~140 kb), as confirmed by S1-PFGE. High similarity was observed among the KPC-encoding IncFII/IncR plasmid, which co-harboring *bla*<sub>TEM-1</sub>, *bla*<sub>CTX-M-65</sub> and *bla*<sub>SHV-12</sub> resistance genes, with more than 99% identity. The IncC plasmid carried *bla*<sub>NDM</sub> and several AMR genes including *bla*<sub>CMY-6</sub>, *sul1*, *emrE*, *aadA16*, *dfxA27*, and *arr-3*, and exhibited a high similarity to others, with 100% coverage and 100% identity. JNKPN30 lacked *bla*<sub>KPC</sub> gene, but an IncFII/IncR-type plasmid was identified within it. WGS showed the IncFII/IncR plasmid lost a 25-kb fragment (including  $\Delta$ ISKpn6-*bla*<sub>KPC-2</sub>-ISKpn27 and IS26-*bla*<sub>SHV-12</sub>-TnAs1 resistance units, as well as  $\Delta$ Tn21) (Fig. 3). Notably, *mgrB* deletion was observed in the colistin-resistant strain JNKPN26 (Additional file 1: Fig. S1).

All six strains harbored a 200–220-kb pLVPK-like virulence plasmid, identified as IncFIB/IncHI1B, carrying

**Table 1** Clinical characteristics of patients with KPC-2-NDM-1-CR-hvKP

Variables/patients	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5
Strain	JNKPN23	JNKPN24	JNKPN26	JNKPN29	JNKPN31
Clinical characteristics					
Age	70	70	44	56	69
Gender	Female	Female	Female	Female	Female
City	JINAN	JINAN	JINAN	JINAN	JINAN
Ward	ICU	ICU	ICU	ICU	ICU
Underlying conditions	Right oophorectomy, myocardial ischemia, chronic ischemic heart disease, respiratory failure	Hypertension, diabetes, dyslipidemia	Hepatic cyst, splenic cyst	Esophageal squamous cell carcinoma	Hypertension, diabetes, UTI
Invasive procedures					
Mechanical ventilation	Yes	Yes	Yes	Yes	Yes
Drainage catheters	Yes	Yes	Yes	Yes	Yes
Surgery	Yes	Yes	Yes	Yes	Yes
Date of specimen collection	2019/2/5	2019/2/15	2019/2/28	2019/3/6	2019/3/15
Infection type	Severe pneumonia, sepsis	Severe pneumonia, sepsis	Severe pneumonia	Severe pneumonia	Severe pneumonia
Prior antibiotic usage within 30 days	Yes	Yes	Yes	Yes	Yes
Empirical antimicrobial usage	CSL + POL + TZP + TGC	TZP + TGC	CSL + TZP + CIP + LZD + POL + ETS	CXM + CSL + ETS + LZD + MIN	ETS + TZP + LEV
Clinical presentations					
Temperature (max °C)	39.2 °C	38.6 °C	39.5 °C	39.1 °C	38.2 °C
WBC (×10 <sup>9</sup> /L)	27.83	29.49	28.73	20.93	16.04
Therapeutic antimicrobial usage	TOB + IPM + TGC	TZP + TGC + POL	LZD + POL + MEM + LEV + TZP + ETS	LZD + MIN	LEV
Clinical outcomes					
Days of mechanical ventilation	6	12	51	10	7
Duration of ICU stay (days)	6	12	56	29	30
Outcome	Death	Death	Survived	Death	Survived


CSL cefoperazone/sulbactam, POL polymyxin, TZP piperacillin/tazobactam, IPM/CS imipenem/cilastatin, TOB tobramycin, TGC tigecycline, LZD linezolid, CIP ciprofloxacin, MEM meropenem, LEV levofloxacin, CXM cefuroxime sodium, MIN minocycline, ETS etimicin sulfate, UTI urinary tract infection

canonical virulence factors such as the regulator of the mucoid phenotype (*rmpA* and *rmpA2*), salmochelin (*iroBCDN*), aerobactin (*iucABCD* and *iutA*), and siderophore salmochelin (*iroBCDE* and *iroN*). The plasmid lacked antibiotic resistance genes and was devoid of T4SS, making it unable to undergo conjugative transfer. The *G. mellonella* virulence assay revealed that KPC-NDM-CR-KP JNKPN26 exhibited significantly higher virulence than ATCC13883 but showed similar virulence to the positive control NTUH-K2044. Furthermore, serum bactericidal assays indicated that KPC-NDM-CR-KP strains exhibited serum resistance, with a survival

rate of approximately 78% after 60 min of incubation with pooled human serum. These phenotypic findings confirmed that KPC-NDM-CR-KP strains were hyper-virulent (Fig. 4).

#### Mobilization of the pJNKPN26-KPC plasmid with the help of the conjugative pJNKPN26-NDM plasmid

Conjugation assays demonstrated that the NDM-1-positive plasmid, pJNKPN26-NDM, could efficiently self-transfer to *E. coli* J53 with a high frequency of  $3.56 \times 10^{-3}$ . In contrast, KPC-positive plasmid had a much lower transfer frequency of  $1.8 \times 10^{-9}$ . Interestingly, all the




**Fig. 1** Epidemiology of *Klebsiella pneumoniae* outbreak cases. Colored text and bars represent the source from which the bacterial species were isolated, whereas colored triangles represent the time at which the bacterial species were isolated. CRKP: carbapenem-resistant *Klebsiella pneumoniae*; CRPA: carbapenem-resistant *Pseudomonas aeruginosa*; CRAB: carbapenem-resistant *Acinetobacter baumannii*; *S. aureus*: *Staphylococcus aureus*; *E. coli*: *Escherichia coli*; *B. circulans*: *Bacillus circulans*; *C. striatum*: *Corynebacterium striatum*

**Table 2** Microbiological characteristics of clinical isolated

Isolate	Carbapenemase	MLST	MIC (µg/mL)													
			TZP	CAZ	CRO	FEP	ATM	ETP	IPM	AMK	GEN	CIP	LVX	SXT	POL	TGC
JNKPN01	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≤ 1	1	0.5
JNKPN03	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≤ 1	0.5	2
JNKPN05	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≤ 2	≥ 16	≥ 4	≥ 8	≤ 1	1	1
JNKPN15	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≤ 1	1	0.5
JNKPN23	KPC-2; NDM-1	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≥ 16	1	0.5
JNKPN24	KPC-2; NDM-1	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≥ 16	0.5	1
JNKPN26	KPC-2; NDM-1	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≥ 16	> 64	1
JNKPN28	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≤ 1	0.5	0.5
JNKPN29	KPC-2; NDM-1	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≥ 16	0.5	1
JNKPN30	NDM-1	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≥ 16	0.5	2
JNKPN31	KPC-2; NDM-1	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≥ 16	1	1
JNKPN34	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≤ 1	1	1
JNKPN37	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≤ 1	1	2
JNKPN38	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≤ 1	0.5	0.75
JNKPN49	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≥ 64	≥ 16	≥ 4	≥ 8	≥ 16	0.5	0.25
JNKPN58	KPC-2	ST11	≥ 128	≥ 64	≥ 64	≥ 64	≥ 64	≥ 32	≥ 16	≤ 2	≤ 1	≥ 4	≥ 8	≤ 1	0.5	1

MIC minimal inhibitory concentrations, CRO ceftriaxone, FEP cefepime, CAZ ceftazidime, ATM aztreonam, TZP piperacillin-tazobactam, ETP ertapenem, IMP imipenem, SXT trimethoprim-sulfamethoxazole, AMK amikacin, GEN gentamicin, CIP ciprofloxacin, LVX levofloxacin, TGC tigecycline, POL polymyxin



**Fig. 2** Hospital outbreak evolution analysis of CO-NDM-KPC-CRKP. The simulated outbreak evolution analysis of CRKP strains isolated in one health set. A phylogenetic tree was constructed using the sample separation time to infer evolutionary relationships among clinical isolates (BioProject PRJNA792451). The features of CO-NDM-KPC-CRKP associated with our hospital outbreak are highlighted in bold

KPC-positive transconjugants selected from plates containing EDTA, also carried the NDM bearing IncC plasmids (Table 3).

The IncC plasmid harbored a complete conjugative transfer-related module, including an *oriT* region, a relaxase of the MobH family, a type IV coupling protein (T4CP), and a *tra* gene cluster that coded for type IV T4SS. In contrast, relaxase and T4CP were absent in the pJNKPN26-KPC plasmid, suggesting that the KPC plasmid transfer relied on assistance from the IncC plasmid.

The plasmid state of transconjugants coharboring NDM and KPC was assessed with S1-PFGE. In addition to NDM and KPC located on separate plasmids, similar to the clinical strain (~140 kb and ~120 kb), two extra S1 patterns were detected in the conjugates. Briefly, the transconjugants co-harboring NDM and KPC displayed three different S1-PFGE profiles: (i) J-JNKPN26-1, which carried two independent plasmids (approximately 140 kb and 120 kb); (ii) J-JNKPN26-2 and J-JNKPN26-3, which contained a novel large plasmid (approximately 270 kb) and two smaller plasmids (approximately 140 kb and 120 kb, respectively); and (iii) J-JNKPN26-4, which possessed a novel large plasmid (approximately 270 kb)


that was smaller than the novel plasmids found in J-JNKPN26-2 and J-JNKPN26-3 (Fig. 5, Additional file 2: Fig. S2).

In pattern (i), pJ-JNKPN26-1\_NDM and pJ-JNKPN26-1\_KPC were almost identical with NDM and KPC harboring the plasmids in the clinical strain; nevertheless, inverted repeats (IRs) at the 5'-end repeat of *TnAsI* were only present within *dnaQ* in pJ-JNKPN26-1\_NDM. Thus, the recombinant junction mediated by *TnAsI* was supposed to occur (Fig. 6).

In pattern (ii), three plasmids existed in the transconjugant J-JNKPN26-2—namely, pJ-JNKPN26-2\_HNK (270,446 bp, GenBank accession number: OR041627), pJ-JNKPN26-2\_NDM (144,158 bp), and pJ-JNKPN26-2\_KPC (126,207 bp). The transconjugant J-JNKPN26-3 harbored three plasmids—namely, pJ-JNKPN26-3\_HNK (270,443 bp, GenBank accession number: OR041628), pJ-JNKPN26-3\_NDM (144,239 bp) and pJ-JNKPN26-3\_KPC (126,207 bp).

In pattern (iii), one large plasmid was identified in the transconjugant J-JNKPN26-4 with a size of 267,313 bp (Fig. 3). Comparative genetic analysis revealed that pJ-JNKPN26-4\_HNK (GenBank accession





**Fig. 4** Analysis of the virulence and serum resistance of *Klebsiella pneumoniae* strains. **A** Survival rates of *Galleria mellonella* larvae injected with *Klebsiella pneumoniae* strains. Survival data were plotted using the Kaplan–Meier method, and the groups were compared using the log-rank test. Hypervirulent *K. pneumoniae* NTUH-K2044 and classic *K. pneumoniae* ATCC13883 strains were used as lower-virulence comparators. **B** Evaluation of the serum resistance of ST11 *Klebsiella pneumoniae* strains. Data are presented as mean (SD). The reference *K. pneumoniae* strain ATCC 13883 and the hypervirulent strain NTUH-K2044 were used as controls for the susceptible (ATCC 13883) and resistant (NTUH-K2044) grades, respectively. Significant differences (two-tailed unpaired t-test) in viable counts (3 h) of NTUH-K2044 and ATCC 13883 cells are shown in the table. \*Unpaired t-test was not used because identical viable counts were recorded. **C** Virulence gene matrix of *Klebsiella pneumoniae* strains. The presence and absence of virulence genes are represented by dark blue and light blue boxes, respectively. Red triangles indicate virulence genes located in the plasmids. NTUH-K2044 and ATCC13883 are publicly available *K. pneumoniae* strains

facilitated by mobile genetic elements, even at a short time after the onset of the outbreak [15]. Previously reported strains with coexisting KPC and NDM were often sporadic [16]. As these strains have been increasingly reported worldwide, their stability, transmission capacity, and mechanism of action have drawn considerable attention [2, 17]. Recently, KPC-2-NDM-1-CRKP had been shown to be stable in the environment and could be nosocomially transmitted among patients [18]. Limited treatment choices, particularly for those carrying a typical hypervirulence plasmid, have led to a higher fatality rate in infected patients and have become a severe clinical challenge [2]. In this study, among six patients with nosocomial infections caused by KPC-2-NDM-1-CRKP, two died, whereas one discontinued treatment. All

six strains harbored a 200–220-kb pLVPK-like virulence plasmid exhibiting a hypervirulent phenotype similar to the canonical hvKP NUTH-K2044 strain.

Colistin-based combination therapy is a commonly used last-resort agent for treating severe CRKP infections [19]. In this study, a colistin-resistant strain, JNKPN26, with a PFGE profile identical to that of other KPC-2-NDM-1-CRKPs, was isolated from patient 3. Comparative genetic analysis was conducted to identify the deletion of the *mgrB* region and its microevolution in vivo. We retrospectively analyzed the medical records of patient 3. After ten day’s high-dose colistin therapy, the colistin-resistant JNKPN26 strain was isolated from the patient’s sputum. In this case, colistin exposure was suspected to be an independent risk factor for the



**Table 3** Microbiological characteristics of transconjugants

Isolate	Carbapenemase	MIC ( $\mu\text{g/mL}$ )											
		CRO	CAZ	ATM	TZP	ETP	IMP	SXT	AK	GN	CIP	LEV	FOS
J-JNKPN23	NDM-1	$\geq 64$	$\geq 64$	4	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 16$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN23-1	NDM-1, KPC-2	$\geq 64$	$\geq 64$	16	64	$\geq 8$	$\geq 16$	$\geq 320$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN24	NDM-1	$\geq 64$	$\geq 64$	4	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 16$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN26-1	NDM-1, KPC-2	$\geq 64$	$\geq 64$	4	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 320$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN26-2	NDM-1, KPC-2	$\geq 64$	$\geq 64$	4	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 320$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN26-3	NDM-1, KPC-2	$\geq 64$	$\geq 64$	4	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 320$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN26-4	NDM-1, KPC-2	$\geq 64$	$\geq 64$	4	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 320$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN26-5	NDM-1	$\geq 64$	$\geq 64$	4	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 16$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN29	NDM-1	$\geq 64$	$\geq 64$	2	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 16$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN30	NDM-1	$\geq 64$	$\geq 64$	4	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 16$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN31	NDM-1	$\geq 64$	$\geq 64$	$\geq 64$	$\geq 128$	$\geq 8$	$\geq 16$	$\geq 16$	$\geq 64$	$\geq 16$	$\leq 0.25$	$\leq 0.25$	$\leq 32$
J-JNKPN58	KPC-2	$\geq 64$	$\geq 64$	$\geq 64$	$\geq 128$	$\geq 8$	$\geq 16$	$\leq 1$	$\leq 2$	$\leq 1$	$\leq 0.25$	$\leq 0.25$	$\leq 32$

MIC minimal inhibitory concentrations, CRO ceftriaxone, CAZ ceftazidime, ATM aztreonam, TZP piperacillin–tazobactam, ETP ertapenem, IMP imipenem, SXT trimethoprim–sulfamethoxazole, AMK amikacin, GEN gentamicin, CIP ciprofloxacin, LVX levofloxacin


emergence of colistin resistance *in vivo*. These strains may have experienced rapid evolution and nosocomial outbreaks under antibiotic screening pressure.

Based on the retrospective genomic surveillance of CRKP strains at our hospital, our phylogenetic analysis suggested that the KPC-2-NDM-1-CRKP strains likely originated from the progenitor KPC-2-CRKP that acquired a highly transferable *bla*<sub>NDM-1</sub> plasmid. This finding is consistent with previous studies that conducted genomic analyses using publicly available complete genome sequences [2].

We attempted to experimentally assess the stability and transmission of carbapenemase-carrying plasmids, specifically the pJNKPN26-KPC plasmid, but failed to obtain transconjugants with this plasmid alone. However, pJNKPN26-KPC could be transferred accompanied by the co-transfer of *bla*<sub>NDM</sub> harboring the IncC plasmid. According to a previous study, only 8.0% of IncFII-IncR KPC-2-encoding plasmids from CRKP can be transferred to *E. coli* [20]. In this study, analysis of pJNKPN26-KPC by oriTfinder revealed that the relaxase gene and T4CP were absent, establishing that the IncFII-IncR-type plasmid employed an inactive type IV secretion system (T4SS). Further S1-PFGE and WGS revealed that the pJNKPN26-KPC plasmid was transferred through complete integration into the self-transferable IncC-type plasmid pJNKPN26-NDM to form a novel hybrid plasmid. Cointegration formed by the copy-in route was mediated by TnAs1 and IS26. TnAs1 or IS26 from pJNKPN26-KPC interrupted the IncC plasmid targeting at random sites, such as *dnaQ*, *uvrD*, and *ISEcp1*, followed by a replication step that duplicated themselves. Interestingly, the hybrid resolved back into two individual

plasmids during the recombination event mediated by TnAs1. These findings suggested that NDM harboring the plasmid acted as a helper plasmid for mobilization of the KPC-bearing plasmid.

We further examined the genetic architecture and hereditary patterns of the hybrid plasmids mediated by the two mobile elements. The first route was a combination mediated by IS26 on pJNKPN26-KPC inserted into *ISEcp1* on the NDM-bearing IncC-type plasmid pJNKPN26-NDM. Cointegration was experimentally determined to be stable without dissociation after 7 days of passage. The other route is mediated by TnAs1 in pJNKPN26-KPC inserted into *dnaQ* or *uvrD* in pJNKPN26-NDM. In contrast to IS26 mediated recombination, all three cointegrates mediated by TnAs1 can be dissociated after transfer or passage, resulting in more flexibility during the shuttle process of the mobile element. IS26 has been widely reported to play a critical role in the dissemination of antibiotic resistance genes in gram-negative bacteria, which have been shown to form cointegrates both by a copy-in mechanism involving one insertion sequence (IS) and by a targeted conservation mechanism involving two ISs [21]. In addition, the Tn3 family is ubiquitous in bacteria, molding host genomes via a paste-and-copy mechanism [22]. It has rarely been reported that TnAs1 carries the entire large plasmid and acts as a transposition module through the copy-in mode. TnAs1 contains a transposase (*tnpA*) with an integration function, a resolvase (*tnpR*) with a dissociation function, and an *res* site containing both promoters and IRs at both ends. We speculated that *tnpR* in TnAs1 mainly contributes to the dissociation of the integration intermediate. Among the co-integration events, TnAs1 was detected at




**Fig. 5** Mobilization of pJNKPN26-NDM and pJNKPN26-KPC. **A** XbaI and S1-PFGE of *K. pneumoniae* JNKPN26, as well as their corresponding transconjugants. Differently colored triangles denote different plasmids: virulence plasmid pJNKPN26-Vir (blue triangles), IncFII/IncR plasmid pJNKPN-KPC (yellow triangles), IncC plasmid pJNKPN-NDM (green triangles), and novel hybrid plasmid (orange triangles). **B** Schematic representation of a round of conjugation assays and the state of hybrid plasmid after passage. Rounded rectangles of the same color represent the same strains. Blue, red, and yellow circles denote pJNKPN26-NDM or its derivatives, pJNKPN26-KPC or its derivatives, and pJNKPN26-Vir, respectively

a higher frequency than IS26, suggesting that *TnAs1* may be a more active driver of translocatable units.

IncC plasmids have been widely studied because of their large size, conjugative transfer capability, pan-host prevalence worldwide, and ability to carry multiple drug resistance genes, especially CMY and *bla*<sub>NDM</sub> [23]. NDM harboring IncC plasmids are widespread among a variety of Enterobacteriaceae strains in many countries [24]. IncC-type plasmids accounted for 52.4%

of NDM-producing CRKP isolates in Pakistan [25]. In China, IncC-type plasmids are essential contributors to the dissemination of NDM, accounting for 12.28% of all *bla*<sub>NDM</sub> harboring the plasmids in *K. pneumoniae* [26]. Recently, a *bla*<sub>NDM-1</sub>-bearing cointegrated plasmid from a clinical *Salmonella* Lomita strain was generated from IS 26 mediated integration of an IncX3 and IncC plasmid. In addition to co-transference by forming hybrid plasmids, insertion sequence-mediated rearrangements



**Fig. 6** Genetic structures of the conjugative hybrid resistance plasmid fusion regions. **A** Alignment of hybrid resistance plasmids pJ-JNKPN26-1-NDM, pJ-JNKPN26-2\_HNK, pJ-JNKPN26-3\_HNK, and pJ-JNKPN26-4\_HNK with parental plasmids from JNKPN26. Black, yellow, red, cyan-blue, and other colored arrows indicate other function protein, conjugation transfer, resistance gene, IS insertion site, and mobile element protein, respectively. The evidence for the occurrence of the fusion event in transconjugant J-JNKPN261-1 was the emergence of TnAs1 on pJ-JNKPN26-1-NDM. **C** The resistance genes are indicated by rectangles. A target site and subsequent 8-bp duplications are indicated by a vertical flag. The relative frequencies of IS26- and TnAs1-mediated reactions are indicated by blue and green arrows, respectively. Proposed mechanisms of plasmid fusion. **i** pJ-JNKPN26-1\_NDM and pJNKPN26-2\_HNK: TnAs1 attacked the *dnaQ* gene of pJNKPN26-NDM, leading to the formation of fusion plasmids through a replicative transposition mechanism. **ii** pJ-JNKPN26-3\_HNK: TnAs1 interrupted the *uvrD* gene of pJNKPN26-NDM, leading to the reverse insertion of pJNKPN26-KPC into pJ-JNKPN26-1-NDM forming plasmid pJ-JNKPN26-3\_HNK. **iii** pJ-JNKPN26-4\_HNK: formation of cointegration was mediated by IS26 interrupting ISEcp1

may promote the diversity and rapid evolution of the IncC genome [27]. Further studies should investigate the fitness cost of excellent compatibility and transferability after the insertion of a large sequence.

### Conclusions

In conclusion, the present study investigated the evolutionary pathway based on a phylogenetic analysis of a cluster of highly homologous CRKPs isolated from one health set.

Our results verified that KPC-2-NDM-1-CRKP formed from a KPC-2-CRKP strain that acquired a highly transferable *bla*<sub>NDM-1</sub> plasmid in silico. Unfortunately, not all the strains isolated from related patients and environment were collected and sequenced. It's impossible to determine the origin of *bla*<sub>NDM</sub> bearing plasmid in the real world. More well planned sample and sequencing are needed for further confirmation. Interestingly, KPC harboring the IncFII/IncR-type plasmid could be transferred by the IncC-type plasmid via the formation of a large hybrid plasmid mediated by IS26 or TnAs1. A better understanding of the mechanisms underlying and triggering co-integration may facilitate the development of interventional measures to curb the formation and dissemination of such elements.

### Abbreviations

CR-hvKP	Carbapenem-resistant hypervirulent <i>Klebsiella pneumoniae</i>
hv-CRKP	Hypervirulent carbapenem-resistant <i>K. pneumoniae</i>
CRKP	Carbapenem-resistant <i>K. pneumoniae</i>
WGS	Whole-genome sequencing
MICs	Minimum inhibitory concentrations
CFU	Colony forming unit
PCR	Polymerase chain reaction
T4SS	Type IV secretion system
ISs	Insertion sequences
IRs	Inverted repeats
PFGE	Pulsed-field gel electrophoresis
CZA	Ceftazidime/avibactam

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12941-024-00686-3>.

**Additional file 1: Figure S1.** Comparative analysis of mgrB-related region. Alignment of mgrB-related region in colistin-resistant strain JNKPN26 with colistin-susceptible strain JNKPN30.

**Additional file 2: Figure S2.** Original PFGE Pattern.

**Additional file 3: Table S1.** The primers used in this study.

### Author contributions

Yingying Hao and Mingju Hao contributed to experiment conception and design. Qian Wang and Yue Liu conducted bioinformatics analysis and wrote the paper, prepared the tables and figures. Yuanyuan Bai, Xinglun Lu, and Zhen Song performed data analysis. Ran Chen, Yan Jin and Yueling Wang carried out the bacteria identification. The corresponding author is responsible for submitting a competing interest statement on behalf of all authors of the paper.

### Funding

The study was supported by grants from National Natural Science Foundation of China (81902119) and Shandong Province Natural Science Foundation (ZR2021MH214).

### Availability of data and materials

Data will be made available on request.

### Declarations

#### Ethics approval and consent to participate

Ethics committee approval of this study was granted by the institutional review board of the Shandong Provincial Hospital, and informed consent from the patient was obtained.

### Consent for publication

Informed consent was obtained from all the participants in this study as well as the co-authors.

### Competing interests

The authors have no relevant financial or non-financial interests to disclose.

### Author details

<sup>1</sup>Department of Clinical Laboratory, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan, China. <sup>2</sup>Department of Clinical Laboratory, Shandong Provincial Hospital, Cheeloo College of Medicine, Shandong University, Jinan, Shandong, China. <sup>3</sup>Shandong Medicine and Health Key Laboratory of Laboratory Medicine, Department of Clinical Laboratory Medicine, The First Affiliated Hospital of Shandong First Medical University & Shandong Provincial Qianfoshan Hospital, Jinan, China.

Received: 4 January 2024 Accepted: 21 March 2024

Published online: 29 March 2024

### References

- Munoz-Price LS, Poirel L, Bonomo RA, Schwaber MJ, Daikos GL, Cormican M, Cornaglia G, Garau J, Gniadkowski M, Hayden MK, Kumarasamy K, Livermore DM, Maya JJ, Nordmann P, Patel JB, Paterson DL, Pitout J, Villegas MV, Wang H, Woodford N, Quinn JP. Clinical epidemiology of the global expansion of *Klebsiella pneumoniae* carbapenemases. *Lancet Infect Dis*. 2013. [https://doi.org/10.1016/S1473-3099\(13\)70190-7](https://doi.org/10.1016/S1473-3099(13)70190-7).
- Gao H, Liu Y, Wang R, Wang Q, Jin L, Wang H. The transferability and evolution of NDM-1 and KPC-2 co-producing *Klebsiella pneumoniae* from clinical settings. *EBioMedicine*. 2020. <https://doi.org/10.1016/j.ebiom.2019.102599>.
- Gu D, Dong N, Zheng Z, Lin D, Huang M, Wang L, Chan EW, Shu L, Yu J, Zhang R, Chen S. A fatal outbreak of ST11 carbapenem-resistant hypervirulent *Klebsiella pneumoniae* in a Chinese hospital: a molecular epidemiological study. *Lancet Infect Dis*. 2018. [https://doi.org/10.1016/S1473-3099\(17\)30489-9](https://doi.org/10.1016/S1473-3099(17)30489-9).
- Zhao J, Liu C, Liu Y, Zhang Y, Xiong Z, Fan Y, Zou X, Lu B, Cao B. Genomic characteristics of clinically important ST11 *Klebsiella pneumoniae* strains worldwide. *J Glob Antimicrob Resist*. 2020. <https://doi.org/10.1016/j.jgar.2020.03.023>.
- Xu Y, Zhang J, Wang M, Liu M, Liu G, Qu H, Liu J, Deng Z, Sun J, Ou HY, Qu J. Mobilization of the nonconjugative virulence plasmid from hypervirulent *Klebsiella pneumoniae*. *Genome Med*. 2021. <https://doi.org/10.1186/s13073-021-00936-5>.
- Hao Y, Zhao X, Zhang C, Bai Y, Song Z, Lu X, Chen R, Zhu Y, Wang Y. Clonal dissemination of clinical carbapenem-resistant *Klebsiella pneumoniae* isolates carrying fosA3 and bla (KPC-2) coharboring plasmids in Shandong, China. *Front Microbiol*. 2021. <https://doi.org/10.3389/fmicb.2021.771170>.
- Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM. Twelve years of SAMtools and BCFtools. *Gigascience*. 2021. <https://doi.org/10.1093/gigascience/giab008>.
- Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol*. 2019. <https://doi.org/10.1186/s13059-019-1832-y>.
- Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*. 2013. <https://doi.org/10.1093/molbev/mst010>.
- Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris SR. Rapid phylogenetic analysis of large samples of recombinant bacterial whole genome sequences using Gubbins. *Nucleic Acids Res*. 2014. <https://doi.org/10.1093/nar/gku1196>.
- Didelot X, Croucher NJ, Bentley SD, Harris SR, Wilson DJ. Bayesian inference of ancestral dates on bacterial phylogenetic trees. *Nucleic Acids Res*. 2018. <https://doi.org/10.1093/nar/gky783>.
- Didelot X, Fraser C, Gardy J, Colijn C. Genomic infectious disease epidemiology in partially sampled and ongoing outbreaks. *Mol Biol Evol*. 2017. <https://doi.org/10.1093/molbev/msw275>.

13. Mei YF, Liu PP, Wan LG, Liu Y, Wang LH, Wei DD, Deng Q, Cao XW. Virulence and genomic feature of a virulent *Klebsiella pneumoniae* sequence type 14 strain of serotype K2 HARBORING bla(NDM-5) in China. *Front Microbiol*. 2017. <https://doi.org/10.3389/fmicb.2017.00335>.
14. Huang J, Yi M, Yuan Y, Xia P, Yang B, Liao J, Dang Z, Luo S, Xia Y. Emergence of a fatal ST11-KL64 tigecycline-resistant hypervirulent *Klebsiella pneumoniae* clone cocarrying bla(NDM) and bla(KPC) in plasmids. *Microbiol Spectr*. 2022. <https://doi.org/10.1128/spectrum.02539-22>.
15. Zhang X, Ouyang J, He W, Zeng T, Liu B, Jiang H, Zhang Y, Zhou L, Zhou H, Liu Z, Liu L. Co-occurrence of rapid gene gain and loss in an interhospital outbreak of carbapenem-resistant hypervirulent ST11-K64 *Klebsiella pneumoniae*. *Front Microbiol*. 2020. <https://doi.org/10.3389/fmicb.2020.579618>.
16. Huang Y, Li J, Wang Q, Tang K, Cai X, Li C. Detection of carbapenem-resistant hypervirulent *Klebsiella pneumoniae* ST11-K64 co-producing NDM-1 and KPC-2 in a tertiary hospital in Wuhan. *J Hosp Infect*. 2023. <https://doi.org/10.1016/j.jhin.2022.09.014>.
17. Wei DD, Wan LG, Liu Y. Draft genome sequence of an NDM-1- and KPC-2-coproducing hypervirulent carbapenem-resistant *Klebsiella pneumoniae* strain isolated from burn wound infections. *Genome Announc*. 2018. <https://doi.org/10.1128/genomeA.00192-18>.
18. Boralli C, Paganini JA, Meneses RS, Mata C, Leite EMM, Schurch AC, Paganelli FL, Willems RJL, Camargo I. Characterization of bla(KPC-2) and bla(NDM-1) plasmids of a *K. pneumoniae* ST11 outbreak clone. *Antibiotics*. 2023. <https://doi.org/10.3390/antibiotics12050926>.
19. Tang M, Li J, Liu Z, Xia F, Min C, Hu Y, Wang H, Zou M. Clonal transmission of polymyxin B-resistant hypervirulent *Klebsiella pneumoniae* isolates coharboring bla(NDM-1) and bla(KPC-2) in a tertiary hospital in China. *BMC Microbiol*. 2023. <https://doi.org/10.1186/s12866-023-02808-x>.
20. Zou H, Zhou Z, Berglund B, Zheng B, Meng M, Zhao L, Zhang H, Wang Z, Wu T, Li Q, Li X. Persistent transmission of carbapenem-resistant, hypervirulent *Klebsiella pneumoniae* between a hospital and urban aquatic environments. *Water Res*. 2023. <https://doi.org/10.1016/j.watres.2023.120263>.
21. Harmer CJ, Hall RM. IS 26 family members IS 257 and IS 1216 also form cointegrates by copy-in and targeted conservative routes. *mSphere*. 2020;5(1):10–128.
22. Lima-Mendez G, Oliveira Alvarenga D, Ross K, Hallet B, Van Melderden L, Varani AM, Chandler M. Toxin-antitoxin gene pairs found in Tn3 family transposons appear to be an integral part of the transposition module. *MBio*. 2020. <https://doi.org/10.1128/mBio.00452-20>.
23. Hancock SJ, Phan MD, Roberts LW, Vu TNM, Harris PNA, Beatson SA, Schembri MA. Characterization of DtrJ as an IncC plasmid conjugative DNA transfer component. *Mol Microbiol*. 2021. <https://doi.org/10.1111/mmi.14697>.
24. Xu J, Zhao Z, Ge Y, He F. Unravelling the genome sequence of NDM-1 and KPC-2 co-producing *Klebsiella pneumoniae* ST11 isolated from a bloodstream infection. *J Glob Antimicrob Resist*. 2020. <https://doi.org/10.1016/j.jgar.2020.01.021>.
25. Qamar MU, Ejaz H, Walsh TR, Shah AA, Al Farraj DA, Alkufeidy RM, Alkubaisi NA, Saleem S, Jahan S. Clonal relatedness and plasmid profiling of extensively drug-resistant New Delhi metallo- $\beta$ -lactamase-producing *Klebsiella pneumoniae* clinical isolates. *Future Microbiol*. 2021. <https://doi.org/10.2217/fmb-2020-0315>.
26. Zeng Z, Lei L, Li L, Hua S, Li W, Zhang L, Lin Q, Zheng Z, Yang J, Dou X. In silico characterization of blaNDM-harboring plasmids in *Klebsiella pneumoniae*. *Front Microbiol*. 2022. <https://doi.org/10.3389/fmicb.2022.1008905>.
27. Li R, Xie M, Liu L, Huang Y, Wu X, Wang Z, Chan EWC, Chen S. Characterisation of a cointegrate plasmid harbouring bla(NDM-1) in a clinical *Salmonella* Lomita strain. *Int J Antimicrob Agents*. 2020. <https://doi.org/10.1016/j.ijantimicag.2019.09.021>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.