

RESEARCH

Open Access

Spermine reverses lipopolysaccharide-induced memory deficit in mice

Pâmella Karina Santana Frühaufer¹, Rafael Porto Ineu¹, Lediane Tomazi¹, Thiago Duarte¹, Carlos Fernando Mello^{1,2} and Maribel Antonello Rubin^{1,3*}

Abstract

Background: Lipopolysaccharide (LPS) induces neuroinflammation and memory deficit. Since polyamines improve memory in various cognitive tasks, we hypothesized that spermine administration reverses LPS-induced memory deficits in an object recognition task in mice. The involvement of the polyamine binding site at the N-methyl-D-aspartate (NMDA) receptor and cytokine production in the promnesic effect of spermine were investigated.

Methods: Adult male mice were injected with LPS (250 µg/kg, intraperitoneally) and spermine (0.3 to 1 mg/kg, intraperitoneally) or ifenprodil (0.3 to 10 mg/kg, intraperitoneally), or both, and their memory function was evaluated using a novel object recognition task. In addition, cortical and hippocampal cytokines levels were measured by ELISA four hours after LPS injection.

Results: Spermine increased but ifenprodil decreased the recognition index in the novel object recognition task. Spermine, at doses that did not alter memory (0.3 mg/kg, intraperitoneally), reversed the cognitive impairment induced by LPS. Ifenprodil (0.3 mg/kg, intraperitoneally) reversed the protective effect of spermine against LPS-induced memory deficits. However, spermine failed to reverse the LPS-induced increase of cortical and hippocampal cytokine levels.

Conclusions: Spermine protects against LPS-induced memory deficits in mice by a mechanism that involves GluN2B receptors.

Keywords: Ifenprodil, Memory, Neuroinflammation, NMDA receptor object recognition, Polyamines, Spermine

Background

It has been shown that intraperitoneal injection of lipopolysaccharide (LPS), a cell-wall component of gram-negative bacteria, induces neuroinflammation, hippocampal cell loss, cognitive impairment, learning deficits and even β -amyloid plaque generation in the hippocampus [1,2], constituting a valid experimental model to study the physiological, behavioral, and emotional aspects of sickness behavior [3]. A number of studies have shown that upon exposure to LPS, microglia become activated and produce proinflammatory mediators such as cytokines, chemokines, prostanooids, and reactive oxygen species. Current evidence indicates that these products are key mediators of the neuroinflammatory

process and contribute to LPS-induced neuronal damage and subsequent cognitive loss [1,4-8]. In line with this view, it has been shown that LPS administration impairs contextual fear conditioning [8-10] and spatial memory [2,11-14] and avoidance learning [15-19] in rats and mice. In addition, LPS decreases the preference for a novel object in a novel object recognition test in mice [20]. The comprehension of the mechanisms by which such a deficit occurs may unveil additional (or ratify already-known) regulatory mechanisms of memory consolidation and retrieval in this pathologic condition, thereby guiding the search for novel therapeutic strategies or compounds to mitigate these deficits.

The N-methyl-D-aspartate (NMDA) receptors seem to be particularly susceptible to the neuroinflammatory challenge, since inflammation decreases the expression of GluN1 [21], GluN2A and GluN2B [22] subunits, and NMDA-dependent long-term potentiation [23] in the hippocampus. This finding is in agreement with previous

* Correspondence: maribel.rubin@gmail.com

¹Graduation Program in Pharmacology, Center of Health Sciences, Federal University of Santa Maria, Santa Maria, RS 97105-900, Brazil

³Department of Biochemistry and Molecular Biology, Natural and Exact Sciences Center, Federal University of Santa Maria, Camobi, CEP: 97105900 Santa Maria, RS, Brazil

Full list of author information is available at the end of the article

reports that neuroinflammation decreases total NMDA receptors (NR1 immunoreactivity) in the hippocampus and entorhinal cortex [24] and that LPS-treated animals present decreased MK801 binding [25]. Accordingly, the partial NMDA receptor agonist D-cycloserine prevents the deleterious effects of LPS [26] and closed head injury [27] on memory consolidation. Therefore, it sounds possible that other positive allosteric modulators of the NMDA receptor attenuate LPS-induced cognitive deficits.

Polyamines, such as putrescine, spermidine, and spermine, allosterically activate NMDA receptors by binding at the lower lobe of the N-terminal domain of GluN1 and GluN2B dimer interface [28]. Functionally, polyamines are involved in growth and differentiation, but they also regulate a broad array of cellular functions in both neurons and inflammatory cells [29-31]. Numerous reports have indicated that polyamines improve memory in several tasks and attenuate memory deficits induced by different amnesic agents [32-41]. In fact, agonists and antagonists of the polyamine binding site at the NMDA receptor respectively facilitate and impair memory in various tasks [33,34,36,39,40,42,43], and the sequential activation of protein kinase C (PKC) and PKA/CREB (protein kinase A/cAMP response element-binding protein) pathways in the hippocampus has been implicated in the promnesic effect of polyamines [44,45]. However, it has recently been described that interruption of the NMDA receptor modulation by polyamines reverses $A\beta_{25-35}$ -induced memory impairment in mice in a novel object recognition task [46], suggesting that the role of polyamines in memory may vary in physiological and pathological conditions.

The assumption that the effects of polyamines on memory depend on physiological or pathological conditions is in agreement with clinical and experimental data that support a beneficial role for NMDA antagonists in Alzheimer's disease [46,47]. However, one might ask whether this applies to every pathological or neuroinflammatory condition. Therefore, we investigated the effect of spermine on LPS-induced impairment of memory in a novel object recognition test. In addition, we investigated whether spermine alters LPS-induced increase of cortical and hippocampal cytokine levels in mice, since both anti- and proinflammatory activity have also been reported for these aliphatic amines [29,48-51].

Methods

Animals

Adult male Swiss mice approximately 12 weeks old (30 to 35 g) provided by the Animal Center of Universidade Federal de Santa Maria, were used for the behavioral experiments. The animals had free access to water and food (Guabi, Santa Maria, Rio Grande do Sul, Brazil), and were maintained in a humidity- and temperature-controlled room ($22 \pm 2^\circ\text{C}$) with a 12-hour light-dark

cycle. Behavioral experiments were conducted in a sound-attenuated and air-regulated room, where the animals were habituated for 1 hour prior to experiments. Behavioral tests were conducted during the light phase of the cycle (between 9:00 a.m. and 5:00 p.m.). All animal procedures were carried out in accordance with Brazilian law no. 11.794/2008, which is in agreement with the Policies on the Use of Animals and Humans in Neuroscience research, revised and approved by the Society for Neuroscience Research in January 1995 and with the Institutional and National regulations for animal research (process 068/2011). All experimental protocols were designed with the aim of keeping the number of animals used to a minimum, as well as their suffering.

Drug administration

Lipopolysaccharide (*Escherichia coli*, serotype 055:B5), spermine (N, N'-bis (3-aminopropyl) 1,4-butanediamine) and ifenprodil (α -(4-hydroxyphenyl)- β -methyl-4-benzyl-1-piperidineethanol tartrate salt) were obtained from Sigma (St. Louis, MO, USA). All drug solutions were prepared daily in saline (0.9% NaCl) and injections were performed intraperitoneally in a 10 ml/kg injection volume. Doses were selected based on previous studies [8], and pilot experiments.

Behavioral testing

Novel object recognition task

A novel object recognition task was carried out as described previously [46]. The task was performed in a $30 \times 30 \times 30$ cm wooden chamber, with walls painted black, a front wall made of Plexiglas and a floor covered with ethyl vinyl acetate sheet. A light bulb, hanging 60 cm above the behavioral apparatus, provided constant illumination of about 40 lux, and an air-conditioner provided constant background sound isolation. The objects used were plastic mounting bricks, each with different shapes and colors, but the same size. Throughout the experiments, objects were used in a counterbalanced manner. Animals had not previously displayed a preference for any of the objects. Chambers and objects were cleaned with 30% ethanol immediately before and at the end of each behavioral evaluation. The task consisted of habituation, training, and testing sessions, each lasting 8 minutes. In the first session, mice were individually habituated to the behavioral apparatus and then returned to their home cages. Twenty-four hours later, the animals were subjected to a training session in which the animals were exposed to two of the same objects (object A), and the exploration time was recorded with two stopwatches. Exploration was recorded when the animal touched or reached the object with the nose at a distance of less than 2 cm. Climbing or sitting on the object was not considered exploration. The test session was carried out 24 hours

after training. Mice were placed back in the behavioral chamber and one of the familiar objects (object A) was replaced by a novel object (object B). The times spent exploring the familiar and the novel object were recorded. The discrimination index was then calculated, taking into account the difference of time spent exploring the new and familiar objects, using the formula:

$$\left[\frac{(T_{\text{novel}} - T_{\text{familiar}})}{(T_{\text{novel}} + T_{\text{familiar}})} \right] \times 100(\%)$$

The discrimination index was used as a memory parameter.

Open field

Immediately after the object recognition test session, the animals were transferred to a 30 cm × 30 cm open field, with the floor divided into four squares. During the 5-min open field session, the number of crossing and rearing responses was recorded. The open field was used to identify motor disabilities, which might influence the object recognition performance.

Quantification of cytokines

Cytokine quantification was assessed by ELISA using commercial kits for mouse IFN- γ , TNF- α , IL-1 β , IL-6, and IL-10 (eBIO-SCIENCE, San Diego, CA, USA), according to the manufacturer's instructions. The presence and concentration of the cytokines were determined by the intensity of the color measured by spectrometry in a micro ELISA reader.

Statistical analysis

Statistical analyses were performed using Student's *t* test, one, two or three-way analysis of variance (ANOVA) followed by the Bonferroni post-hoc analysis. A value of $P < 0.05$ was considered significant. *F* and *P* values are shown only if $P < 0.05$.

Experimental design

Experiment 1

This experiment was designated to investigate the effect of LPS on the object recognition task performance. Animals were habituated and trained, as described. Immediately after training, the animals were injected with saline or LPS (250 $\mu\text{g}/\text{kg}$). Twenty-four hours after training, the animals were subjected to the novel object recognition test session and open field as described.

Experiment 2

This experiment was designated to investigate the effect of spermine or ifenprodil on the object recognition task performance. Animals were habituated and trained as described. Immediately after training, the animals were injected with saline, spermine (0.1 to 10 mg/kg) or

ifenprodil (0.3 to 10 mg/kg). Twenty-four hours after training, the animals were subjected to the novel object recognition test session and open field, as described.

Experiment 3

This experiment was designed to investigate the involvement of polyamine binding sites in the impairment of memory induced by LPS. Animals were habituated and trained as described. Immediately after training, the animals were injected with saline or LPS (250 $\mu\text{g}/\text{kg}$) and 5 minutes later with saline or the polyaminergic agonist, spermine (at doses that have no effect *per se* on memory, 0.3 mg/kg, as determined by the dose-effect curve in different flanks. Twenty-four hours after training, the animals were subjected to the novel object recognition test session and open field, as described.

Experiment 4

This experiment was designed to investigate the involvement of polyamine binding sites on the NMDA receptor in the reversal of the LPS-induced impairment of memory by spermine on the object recognition task performance. Animals were habituated and trained as described. Immediately after training, the animals were injected with saline or LPS (250 $\mu\text{g}/\text{kg}$), 5 min later they were injected with saline or the polyaminergic agonist, spermine (0.3 mg/kg, the dose that reverses the LPS-induced impairment of memory), and after another 5 minutes they were injected with saline or ifenprodil (at doses that have no effect *per se* on memory, 0.3 mg/kg, as determined by the dose-effect curve) in different flanks. Twenty-four hours after training, the animals were submitted to the novel object recognition test and open field, as described.

Experiment 5

This experiment was designed to investigate whether spermine prevents LPS-induced increases in levels of proinflammatory cytokines, IL-1 β , IL-6, TNF- α , interferon- γ (ITF- γ), and anti-inflammatory cytokine, IL-10. Immediately after the training session of the object recognition task, animals received saline, LPS (250 $\mu\text{g}/\text{kg}$), spermine (0.3 mg/kg) or a combination of LPS (250 $\mu\text{g}/\text{kg}$) and spermine (0.3 mg/kg). Four hours after drug administration, the animals were anesthetized with ketamine (100 mg/kg) and xylazine (10 mg/kg) and transcardially perfused with cold saline. The cerebral cortex and the hippocampi were dissected and homogenized in appropriated buffer (PBS containing 1 mM ethylenediamine tetraacetic acid (EDTA), 0.1 mM phenylmethylsulfonyl fluoride and 0.5% BSA, pH 7.4). Samples were centrifuged at 25,000 *g* for 10 min and the supernatant was used to measure IL-1 β , IL-6, TNF- α , ITF- γ , and IL-10 levels, which were corrected for total protein content. Protein concentration

was determined for each brain region by the Bradford method [52] enabling cytokine levels to be expressed as pg/mg protein. The cytokine quantification was assessed as described.

Results



LPS decreased the recognition index in novel object recognition task

Figure 1 shows the effect of the post-training intraperitoneal administration of LPS or saline on recognition index in the object recognition task. Statistical analysis (Student's *t* test) revealed that LPS (250 µg/kg) decreased the recognition index ($t_{14} = 3.086$; $P < 0.01$; $\eta^2 = 0.4$), compared with the control group, indicating a memory impairment in LPS-treated animals.

Spermine increased and ifenprodil decreased the recognition index on novel object recognition task

Figure 2A shows the effect of post-training administration of spermine (0.1 to 10 mg/kg) on the recognition index in the object recognition task. Statistical analysis (one-way ANOVA) revealed a significant effect of spermine ($F_{5,44} = 3.76$; $P < 0.05$; $\eta^2 = 0.29$). Post-hoc analysis revealed that spermine (1 mg/kg) increased the recognition index, indicating that spermine improved memory.

Figure 2B shows the effect of post-training administration of ifenprodil (0.3 to 10 mg/kg) on the recognition index in the object recognition task. Statistical analysis



(one-way ANOVA) revealed a significant effect of ifenprodil ($F_{3,17} = 7.28$; $P < 0.005$; $\eta^2 = 0.56$). Post-hoc analysis revealed that ifenprodil (10 mg/kg) decreased the recognition index, indicating that ifenprodil impaired memory.

Spermine attenuates LPS-induced discrimination impairments on novel object recognition task

Figure 3 shows the effect of post-training administration of spermine on LPS-induced impairment of memory in the novel object recognition task. Statistical analysis (two-way



ANOVA) revealed a significant treatment (saline or LPS) versus polyamine agonist (saline or spermine) interaction ($F_{1,56} = 8.10$; $P < 0.01$; $\eta^2 = 0.12$), revealing that spermine reversed the impairment of memory induced by LPS.

Ifenprodil reverses the effect of spermine in LPS-treated animals

Figure 4 shows the effect of ifenprodil (0.3 mg/kg) on the spermine-induced reversal of the impairment of memory induced by LPS in the object recognition task. Statistical analysis (three-way ANOVA) revealed a significant treatment (saline or LPS) versus polyamine agonist (saline or spermine) versus NMDA receptor antagonist (saline or ifenprodil) interaction ($F_{1,110} = 4.45$; $P < 0.05$; $\eta^2 = 0.038$). Post-hoc analysis (Bonferroni) revealed that ifenprodil reversed the effect of spermine in attenuating the impairment of memory induced by LPS on the novel object recognition task.

Treatments did not alter crossing responses.

Since the recognition index might be affected by locomotor alterations unrelated to the mnemonic component of the task, we monitored the number of crossing responses in the open field task. Treatments did not alter locomotor activity measured by crossing responses (data not shown). Therefore, the reported effects in this study are unlikely to be associated with changes in locomotion and coordination.



Spermine did not reverse LPS-induced increase of proinflammatory cytokines levels

Figures 5 and 6 show the effect of post-training administration of spermine and LPS in the levels of proinflammatory and anti-inflammatory cytokines in the hippocampus and cerebral cortex respectively. Statistical analysis (two-way ANOVA) revealed a significant effect of treatment (saline or LPS) on IL-1 β ($F_{1,15} = 40.01$; $P < 0.001$; $\eta^2 = 0.72$), IL-6 ($F_{1,15} = 52.43$; $P < 0.001$; $\eta^2 = 0.77$), TNF- α ($F_{1,15} = 27.64$; $P < 0.001$; $\eta^2 = 0.64$) and IL-10 ($F_{1,15} = 9.26$; $P < 0.01$; $\eta^2 = 0.38$) levels in the hippocampus (Figure 5), and on IL-1 β ($F_{1,15} = 37.74$; $P < 0.001$; $\eta^2 = 0.71$), IL-6 ($F_{1,15} = 47.43$; $P < 0.001$; $\eta^2 = 0.75$) and TNF- α ($F_{1,15} = 30.43$; $P < 0.001$; $\eta^2 = 0.67$), IL-10 ($F_{1,5} = 8.78$; $P < 0.05$; $\eta^2 = 0.36$) levels in the cerebral cortex (Figure 6). Both LPS and spermine alter the levels of IL-1 β ($F_{1,15} = 4.78$; $P < 0.05$; $\eta^2 = 0.24$), IL-6 ($F_{1,15} = 4.92$; $P < 0.05$; $\eta^2 = 0.24$), and IL-10 ($F_{1,15} = 4.76$; $P < 0.05$; $\eta^2 = 0.24$) levels in the hippocampus (Figure 5), and on IL-1 β ($F_{1,15} = 6.14$; $P < 0.05$; $\eta^2 = 0.29$), IL-6 ($F_{1,15} = 4.92$; $P < 0.05$; $\eta^2 = 0.24$) and IL-10 ($F_{1,5} = 5.95$; $P < 0.05$; $\eta^2 = 0.28$) levels in the cerebral cortex (Figure 6).

Discussion

This study showed that a polyaminergic agonist, spermine, reverses post-training bacterial endotoxin-induced memory impairment in a novel object recognition task. In addition, it showed that spermine, at doses higher than those capable of reversing the deleterious effect of LPS,




Figure 5 (See legend on next page.)

(See figure on previous page.)

Figure 5 Effects of post-training administration of spermine and LPS in the levels of proinflammatory and anti-inflammatory cytokines in hippocampus. Immediately after the training session of the object recognition task, animals received saline, LPS (250 µg/kg, intraperitoneally), spermine (0.3 mg/kg, intraperitoneally) or a combination of LPS (250 µg/kg, intraperitoneally) and spermine (0.3 mg/kg, intraperitoneally), and were euthanized 4 hours after injections to measure the levels of (A) IL-1β, (B) IL-6, (C) TNF-α, (D) ITF-γ, and (E) IL-10 in the hippocampus. **P* < 0.05 between pooled post-training saline (SAL-SAL and SAL-SPM groups) and pooled post-training LPS (LPS-SAL and LPS-SPM groups) by Bonferroni's *t* test. #*P* < 0.05 between pooled post-training saline (SAL-SAL and LPS-SAL groups) and pooled post-training spermine (SAL-SPM and LPS-SPM groups). Data are the mean ± standard error of the mean of 4 or 5 animals per group. LPS, lipopolysaccharide; SAL, saline; SPM, spermine.

increases the recognition index in the novel object recognition task. It also revealed that ifenprodil decreases the recognition index in the novel object recognition task, and a non-effective dose of ifenprodil reverses the memory-improving effect of spermine in LPS-treated animals, providing pharmacological evidence that the promnesic effect of spermine involves the polyamine binding site at the NMDA receptor. However, spermine failed to reverse the LPS-induced increase of cytokine levels in the cerebral cortex or hippocampus.

The currently described ability of spermine to reverse LPS-induced memory deficits is in agreement with a number of studies that have separately shown that while LPS disrupts [1,8,13,15,53], polyamines improve memory [32-39,54].

Several mechanisms have been implicated in LPS-induced alterations of neural functions. It has been suggested that memory impairment is triggered by direct stimulation of toll-like receptor 4 (TLR₄) by LPS. This TLR₄ activation recruits myeloid differentiation adaptor protein (MyD88) and ultimately activates the nuclear factor κB signaling pathway, increasing the production of proinflammatory cytokines by macrophages [55], microglia [56-58] and astrocytes [56,59]. The LPS-induced activation of glial cells results in the release of neurotoxic substances, including nitric oxide, glutamate, cytotoxic cytokines, and superoxide radicals [8,56,60,61], the suppression of neurotrophic factor secretion, and impaired neuroplasticity and, consequently, learning and memory [8,9,11,17,62,63]. In addition, TLR₄ activates Src family kinases, which phosphorylate GluN2B subunit or NMDA receptor and enhance GluN2B-dependent Ca²⁺ influx [64,65], promoting excitotoxicity. Indeed, LPS induces progressive and cumulative neuronal loss over time [66-68].

Conversely, there is a large body of evidence indicating that the promnesic effects of polyamines involve the activation of NMDA receptors, which have been critically implicated in learning and memory processes [36,40,44,45,69-71]. In the current study, ifenprodil, a noncompetitive GluN2B-containing NMDA receptor antagonist, not only impaired the consolidation of the memory of novel object recognition task, but also reversed (at a dose that did not alter memory *per se* in our study) the improving effect of spermine on the

memory of LPS-treated animals, supporting a role for NMDA receptors in this effect of spermine [72-74]. Interestingly, these results are in agreement with the study by Kranjac and colleagues [26], who have shown that partial NMDA receptor agonist D-cycloserine rescues memory consolidation following systemic bacterial endotoxin exposure. Furthermore, Velloso and colleagues [41] have found that post-training intrastriatal administration of spermine reverses the recognition memory deficits in the novel object recognition task induced by quinolinic acid, a model of Huntington's disease. It is worth noting that spermine improved memory *per se* in the novel object recognition task. To our knowledge, this is the first study showing that spermine improves memory. The finding that ifenprodil prevents the promnesic effect of spermine tempts us to propose that it may involve the same molecular targets proposed for spermidine [33,34,36,38]. However, further studies are necessary to clarify this point.

We have found that while LPS increased IL1-β, IL-6, TNF-α, and IFN-γ levels, it decreased IL-10 levels in the cerebral cortex and hippocampus. This is in agreement with previous studies that have shown that systemic administration of this endotoxin causes neuroinflammation [56-58]. Moreover, spermine increased IL1-β and IL-6 levels in the cerebral cortex and hippocampus *per se*, indicating that it facilitates inflammation, but to a much lesser degree than LPS. Interestingly, spermine also decreased the anti-inflammatory cytokine IL-10 in both cerebral structures, further supporting a proinflammatory role for this polyamine in our experimental conditions. These results are in line with other studies [51,75] which show that the spermine facilitates inflammation *in vivo*. Accordingly, it has been shown that polyamines promote macrophage influx into the murine central nervous system following pathogenic insult, in an *in vivo* model of secondary central nervous system inflammation [51]. Soulet and Rivest [75] have shown that systemic LPS increases ornithine decarboxylase expression throughout the central nervous system and that this precedes an increase in the expression of proinflammatory molecules TLR₂ and TNF-α, in mice. Moreover, treatment with a difluoromethylornithine, an irreversible inhibitor of ornithine decarboxylase, decreases LPS-induced TNF-α and TLR₂ expression, proving evidence that the LPS-induced increase of proinflammatory molecules is polyamine-dependent.




Figure 6 (See legend on next page.)

(See figure on previous page.)

Figure 6 Effects of post-training administration of spermine and LPS in the levels of proinflammatory and anti-inflammatory cytokines in cerebral cortex. Immediately after the training session of the object recognition task, animals received saline, LPS (250 µg/kg, intraperitoneally), spermine (0.3 mg/kg, intraperitoneally) or a combination of LPS (250 µg/kg, intraperitoneally) and spermine (0.3 mg/kg, intraperitoneally), and were euthanized 4 hours after injections to measure the levels of (A) IL-1β, (B) IL-6, (C) TNF-α, (D) ITF-γ, and (E) IL-10 in the cerebral cortex. * $P < 0.05$ between pooled post-training saline (SAL-SAL and SAL-SPM groups) and pooled post-training LPS (LPS-SAL and LPS-SPM groups) by Bonferroni's *t* test. # $P < 0.05$ between pooled post-training saline (SAL-SAL and LPS-SAL groups) and pooled post-training spermine (SAL-SPM and LPS-SPM groups). Data are the mean ± standard error of the mean of 4 or 5 animals per group. LPS, lipopolysaccharide; SAL, saline; SPM, spermine.

The intracerebral injection of spermine prior to LPS also increases the number of cells expressing both TNF-α and TLR₂ in the central nervous system [75]. Interestingly, LPS exposure stimulates IL-1β release from astrocytes (*in vitro*) through a mechanism that requires NMDA receptor stimulation [76].

We hypothesized that polyamines could decrease LPS-induced cognitive impairment by interfering in cytokine levels because: (1) LPS, at doses that cause cognitive impairment, sequentially increases brain IL-6, IL-1β, and TNF-α mRNA levels in the hippocampus [77,78] and frontal cortex [79]; (2) LPS-induced upregulation of IL-1β and TNF-α mRNA in hippocampal tissue of IL-6^(+/+) mice is absent in IL-6^(-/-) mice, which are also refractory to the LPS-induced impairment in working memory [77]. In addition, overexpression of TNF-α in neurons or glial cells impairs passive avoidance memory [80]; (3) Pharmacological treatments that decrease the levels of TNF-α, TNF-α receptor 1, and NF-κB p65 phosphorylation also decrease LPS-induced memory deficits [81,82], though conventional TNF^(-/-) knockout mice present cognitive dysfunction [83,84]. A recent review summarized the effect of TNF-α, IL-6, and IL-1 on learning and memory [85], where the authors suggest that TNF-α and its receptors might mediate the disrupting effect of LPS on learning and memory. Therefore, considering the existing evidence supporting a role for cytokines, particularly TNF-α, on LPS-induced cognitive impairment, we thought that the memory-improving effects of spermine could involve a decrease of cytokine levels in the hippocampus.

It has recently been described that LPS reduces the number of excitatory synapses in the hippocampus and cerebral cortex, leading to synaptic deficits [86] that may underlie LPS-induced cognitive deficits. In fact, LPS induces neurotoxic substance release and suppression of neurotrophic factors secretion, which are known to increase neuroplasticity and, consequently, learning and memory. Since polyamines increase excitatory activity [28], it is possible that spermine actions involve a compensatory increase of excitatory transmission in LPS-treated animals. However, specific studies have to be performed, to clarify this point.

Conclusions

Spermine improves memory *per se* and attenuates LPS-induced memory impairment. Our results also suggest

that spermine protects against LPS-induced memory impairment by mechanisms that involve the polyamine binding site at the NMDA receptor, since it is reversed by ifenprodil. Spermine, however, does not prevent LPS-induced increase of proinflammatory molecules, suggesting that the effect of spermine on memory does not involve anti-inflammatory mechanisms.

Abbreviations

ANOVA: analysis of variance; BSA: bovine serum albumin; CREB: cAMP response element-binding protein; EDTA: ethylenediamine tetraacetic acid; ELISA: enzyme-linked immunosorbent assay; IL-1β: interleukin-1β; IL-6: interleukin-6; IL-10: interleukin-10; ITF-γ: interferon-γ; LPS: lipopolysaccharide; MyD88: myeloid differentiation adaptor protein; NMDA: N-methyl-D-aspartate; PBS: phosphate buffered saline; PKA: protein kinase A; PKC: protein kinase C; TNF-α: tumor necrosis factor-α; TLR: toll-like receptor.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PKSF, RPI, LT, and TD conceived and designed the experiments. PKSF and LT performed the behavioral experiments. TD performed the cytokine quantification assay. PKSF, RPI, CFM, and MAR analyzed the data. CFM, MAR, and PKSF wrote the paper. All authors read and approved the final manuscript.

Acknowledgements

This study was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico-CNPq (306164/2010-8, 481664/2010-6, 476551/2009-9) and Fundação de Amparo à Pesquisa do Rio Grande do Sul-FAPERGS. CFM, RPI, and MAR are recipients of CNPq fellowships. PKSF, LT, and TD are recipients of Coordenação de Aperfeiçoamento de Pessoal de Nível Superior-CAPES fellowships.

Author details

¹Graduation Program in Pharmacology, Center of Health Sciences, Federal University of Santa Maria, Santa Maria, RS 97105-900, Brazil. ²Department of Physiology and Pharmacology, Center of Health Sciences, Federal University of Santa Maria, Santa Maria, RS 97105-900, Brazil. ³Department of Biochemistry and Molecular Biology, Natural and Exact Sciences Center, Federal University of Santa Maria, Camobi, CEP: 97105900 Santa Maria, RS, Brazil.

Received: 24 July 2014 Accepted: 11 December 2014

Published online: 09 January 2015

References

1. Lee JW, Lee YK, Yuk DY, Choi DY, Ban SB, Oh KW, et al. Neuro-inflammation induced by lipopolysaccharide causes cognitive impairment through enhancement of beta-amyloid generation. *J Neuroinflammation*. 2008;5:37.
2. Shaw KN, Commins S, O'Mara SM. Lipopolysaccharide causes deficits in spatial learning in the watermaze but not in BDNF expression in the rat dentate gyrus. *Behav Brain Res*. 2001;124:47-54.
3. Benson S, Kattoor J, Wegner A, Hammes F, Reidick D, Grigoleit JS, et al. Acute experimental endotoxemia induces visceral hypersensitivity and altered pain evaluation in healthy humans. *Pain*. 2012;153:794-9.
4. Block ML, Zecca L, Hong JS. Microglia-mediated neurotoxicity: uncovering the molecular mechanisms. *Nat Rev Neurosci*. 2007;8:57-69.

5. Beishuizen A, Thijs LG. Endotoxin and the hypothalamo-pituitary-adrenal (HPA) axis. *J Endotoxin Res.* 2003;9:3–24.
6. Turrin NP, Gayle D, Ilyin SE, Flynn MC, Langhans W, Schwartz GJ, et al. Pro-inflammatory and anti-inflammatory cytokine mRNA induction in the periphery and brain following intraperitoneal administration of bacterial lipopolysaccharide. *Brain Res Bull.* 2001;54:443–53.
7. Gabellec MM, Griffais R, Fillion G, Haour F. Expression of interleukin 1 α , interleukin 1 β and interleukin 1 receptor antagonist mRNA in mouse brain: regulation by bacterial lipopolysaccharide (LPS) treatment. *Brain Res Mol Brain Res.* 1995;31:122–30.
8. Kranjac D, McLinden KA, Deodati LE, Papini MR, Chumley MJ, Boehm GW. Peripheral bacterial endotoxin administration triggers both memory consolidation and reconsolidation deficits in mice. *Brain Behav Immun.* 2012;26:109–21.
9. Pugh CR, Kumagawa K, Fleshner M, Watkins LR, Maier SF, Rudy JW. Selective effects of peripheral lipopolysaccharide administration on contextual and auditory-cue fear conditioning. *Brain Behav Immun.* 1998;12:212–29.
10. Bilbo SD, Biedenkapp JC, Der-Avakian A, Watkins LR, Rudy JW, Maier SF. Neonatal infection-induced memory impairment after lipopolysaccharide in adulthood is prevented via caspase-1 inhibition. *J Neurosci.* 2005;25:8000–9.
11. Sparkman NL, Martin LA, Calvert WS, Boehm GW. Effects of intraperitoneal lipopolysaccharide on Morris maze performance in year-old and 2-month-old female C57BL/6 J mice. *Behav Brain Res.* 2005;159:145–51.
12. Terrando N, Rei Fidalgo A, Vizcaychipi M, Cibelli M, Ma D, Monaco C, et al. The impact of IL-1 modulation on the development of lipopolysaccharide-induced cognitive dysfunction. *Crit Care.* 2010;14:R88.
13. Zarifkar A, Choopani S, Ghasemi R, Naghdi N, Maghsoudi AH, Maghsoudi N, et al. Agmatine prevents LPS-induced spatial memory impairment and hippocampal apoptosis. *Eur J Pharmacol.* 2010;634:84–8.
14. Lee YJ, Choi DY, Choi IS, Kim KH, Kim YH, Kim HM, et al. Inhibitory effect of 4-O-methylhonokiol on lipopolysaccharide-induced neuroinflammation, amyloidogenesis and memory impairment via inhibition of nuclear factor-kappaB *in vitro* and *in vivo* models. *J Neuroinflammation.* 2012;9:35.
15. Sparkman NL, Kohman RA, Garcia AK, Boehm GW. Peripheral lipopolysaccharide administration impairs two-way active avoidance conditioning in C57BL/6 J mice. *Physiol Behav.* 2005;85:278–88.
16. Kohman RA, Tarr AJ, Sparkman NL, Day CE, Paquet A, Akkaraju GR, et al. Alleviation of the effects of endotoxin exposure on behavior and hippocampal IL-1 β by a selective non-peptide antagonist of corticotropin-releasing factor receptors. *Brain Behav Immun.* 2007;21:824–35.
17. Tarr AJ, McLinden KA, Kranjac D, Kohman RA, Amaral W, Boehm GW. The effects of age on lipopolysaccharide-induced cognitive deficits and interleukin-1 β expression. *Behav Brain Res.* 2011;217:481–5.
18. Lin GH, Lee YJ, Choi DY, Han SB, Jung JK, Hwang BY, et al. Anti-amyloidogenic effect of thiacecronone through anti-inflammation *in vitro* and *in vivo* models. *J Alzheimers Dis.* 2012;29:659–76.
19. Choi DY, Lee JW, Lin G, Lee YK, Lee YH, Choi IS, et al. Obovatol attenuates LPS-induced memory impairments in mice via inhibition of NF- κ B signaling pathway. *Neurochem Int.* 2012;60:68–77.
20. Miwa M, Tsuboi M, Noguchi Y, Enokishima A, Nabeshima T, Hiramatsu M. Effects of betaine on lipopolysaccharide-induced memory impairment in mice and the involvement of GABA transporter 2. *J Neuroinflammation.* 2011;8:153.
21. Harre EM, Galic MA, Mouihate A, Noorbakhsh F, Pittman QJ. Neonatal inflammation produces selective behavioural deficits and alters N-methyl-D-aspartate receptor subunit mRNA in the adult rat brain. *Eur J Neurosci.* 2008;27:644–53.
22. Ma J, Choi BR, Chung C, Min SS, Jeon WK, Han JS. Chronic brain inflammation causes a reduction in GluN2A and GluN2B subunits of NMDA receptors and an increase in the phosphorylation of mitogen-activated protein kinases in the hippocampus. *Mol Brain.* 2014;7:33.
23. Min SS, Quan HY, Ma J, Han JS, Jeon BH, Seol GH. Chronic brain inflammation impairs two forms of long-term potentiation in the rat hippocampal CA1 area. *Neurosci Lett.* 2009;456:20–4.
24. Rosi S, Ramirez-Amaya V, Hauss-Wegrzyniak B, Wenk GL. Chronic brain inflammation leads to a decline in hippocampal NMDA-R1 receptors. *J Neuroinflammation.* 2004;1:12.
25. Biegona A, Alvarado M, Budinger TF, Grossman R, Hensley K, West MS, et al. Region-selective effects of neuroinflammation and antioxidant treatment on peripheral benzodiazepine receptors and NMDA receptors in the rat brain. *J Neurochem.* 2002;82:924–34.
26. Kranjac D, Koster KM, Kahn MS, Eimerbrink MJ, Womble BM, Cooper BG, et al. Peripheral administration of D-cycloserine rescues memory consolidation following bacterial endotoxin exposure. *Behav Brain Res.* 2013;243:38–43.
27. Yaka R, Biegona A, Grigoriadis N, Simeonidou C, Grigoriadis S, Alexandrovich AG, et al. D-cycloserine improves functional recovery and reinstates long-term potentiation (LTP) in a mouse model of closed head injury. *FASEB J.* 2007;21:2033–41.
28. Mony L, Zhu S, Carvalho S, Paoletti P. Molecular basis of positive allosteric modulation of GluN2B NMDA receptors by polyamines. *EMBO J.* 2011;30:3134–46.
29. Zhang M, Wang H, Tracey KJ. Regulation of macrophage activation and inflammation by spermine: a new chapter in an old story. *Crit Care Med.* 2000;28:N60–6.
30. Gilad GM, Gilad VH. Astroglia growth retardation and increased microglia proliferation by lithium and ornithine decarboxylase inhibitor in rat cerebellar cultures: cytotoxicity by combined lithium and polyamine inhibition. *J Neurosci Res.* 2007;85:594–601.
31. Tabor CW, Tabor H. Polyamines. *Annu Rev Biochem.* 1984;53:749–90.
32. Kishi A, Ohno M, Watanabe S. Spermidine, a polyamine site agonist, attenuates working memory deficits caused by blockade of hippocampal muscarinic receptors and mGluRs in rats. *Brain Res.* 1998;793:311–4.
33. Rubin MA, Boemo RL, Jurach A, Rojas DB, Zanolta GR, Obregon AD, et al. Intra-hippocampal spermidine administration improves inhibitory avoidance performance in rats. *Behav Pharmacol.* 2000;11:57–61.
34. Rubin MA, Stiegemeier JA, Volkweis MA, Oliveira DM, Fenili AC, Boemo RL, et al. Intra-amygdala spermidine administration improves inhibitory avoidance performance in rats. *Eur J Pharmacol.* 2001;423:35–9.
35. Mikolajczak P, Okulicz-Kozaryn I, Kaminska E, Niedopad L, Polanska A, Gebka J. Effects of acamprosate and some polyamine site ligands of NMDA receptor on short-term memory in rats. *Eur J Pharmacol.* 2002;444:83–96.
36. Rubin MA, Berlese DB, Stiegemeier JA, Volkweis MA, Oliveira DM, dos Santos TL, et al. Intra-amygdala administration of polyamines modulates fear conditioning in rats. *J Neurosci.* 2004;24:2328–34.
37. Tadano T, Hozumi S, Yamadera F, Murata A, Nijima F, Tan-No K, et al. Effects of NMDA receptor-related agonists on learning and memory impairment in olfactory bulbectomized mice. *Methods Find Exp Clin Pharmacol.* 2004;26:93–7.
38. Berlese DB, Sauzem PD, Carati MC, Guerra GP, Stiegemeier JA, Mello CF, et al. Time-dependent modulation of inhibitory avoidance memory by spermidine in rats. *Neurobiol Learn Mem.* 2005;83:48–53.
39. Camera K, Mello CF, Ceretta AP, Rubin MA. Systemic administration of polyaminergic agents modulate fear conditioning in rats. *Psychopharmacology (Berl).* 2007;192:457–64.
40. Ribeiro DA, Mello CF, Signor C, Rubin MA. Polyaminergic agents modulate the reconsolidation of conditioned fear. *Neurobiol Learn Mem.* 2013;104:9–15.
41. Velloso NA, Dalmolin GD, Gomes GM, Rubin MA, Canas PM, Cunha RA, et al. Spermine improves recognition memory deficit in a rodent model of Huntington's disease. *Neurobiol Learn Mem.* 2009;92:574–80.
42. Ceretta AP, Camera K, Mello CF, Rubin MA. Arcaine and MK-801 make recall state-dependent in rats. *Psychopharmacology (Berl).* 2008;201:405–11.
43. da Rosa MM, Mello CF, Camera K, Ceretta AP, Ribeiro DA, Signor C, et al. Opioid mechanisms are involved in the disruption of arcaine-induced amnesia by context pre-exposure. *Neurobiol Learn Mem.* 2012;97:294–300.
44. Guerra GP, Mello CF, Bochi GV, Pazini AM, Fachineto R, Dutra RC, et al. Hippocampal PKA/CREB pathway is involved in the improvement of memory induced by spermidine in rats. *Neurobiol Learn Mem.* 2011;96:324–32.
45. Guerra GP, Mello CF, Bochi GV, Pazini AM, Rosa MM, Ferreira J, et al. Spermidine-induced improvement of memory involves a cross-talk between protein kinases C and A. *J Neurochem.* 2012;122:363–73.
46. Gomes GM, Dalmolin GD, Bar J, Karpova A, Mello CF, Kreutz MR, et al. Inhibition of the polyamine system counteracts β -amyloid peptide-induced memory impairment in mice: involvement of extrasynaptic NMDA receptors. *PLoS One.* 2014;9:e99184.
47. Rosi S, Vazdarjanova A, Ramirez-Amaya V, Worley PF, Barnes CA, Wenk GL. Memantine protects against LPS-induced neuroinflammation, restores behaviorally-induced gene expression and spatial learning in the rat. *Neuroscience.* 2006;142:1303–15.
48. Zhang M, Borovikova LV, Wang H, Metz C, Tracey KJ. Spermine inhibition of monocyte activation and inflammation. *Mol Med.* 1999;5:595–605.
49. Zhang M, Caragine T, Wang H, Cohen PS, Botchkina G, Soda K, et al. Spermine inhibits proinflammatory cytokine synthesis in human

- mononuclear cells: a counterregulatory mechanism that restrains the immune response. *J Exp Med*. 1997;185:1759–68.
50. Bussiere FI, Chaturvedi R, Cheng Y, Gobert AP, Asim M, Blumberg DR, et al. Spermine causes loss of innate immune response to *Helicobacter pylori* by inhibition of inducible nitric-oxide synthase translation. *J Biol Chem*. 2005;280:2409–12.
 51. Puntambekar SS, Davis DS, Hawel 3rd L, Crane J, Byus CV, Carson MJ. LPS-induced CCL2 expression and macrophage influx into the murine central nervous system is polyamine-dependent. *Brain Behav Immun*. 2011;25:629–39.
 52. Bradford MM. Rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*. 1976;72:248–54.
 53. Lee B, Sur B, Park J, Kim SH, Kwon S, Yeom M, et al. Ginsenoside Rg3 alleviates lipopolysaccharide-induced learning and memory impairments by anti-inflammatory activity in rats. *Biomol Ther (Seoul)*. 2013;21:381–90.
 54. Meyer RC, Knox J, Purwin DA, Spangler EL, Ingram DK. Combined stimulation of the glycine and polyamine sites of the NMDA receptor attenuates NMDA blockade-induced learning deficits of rats in a 14-unit T-maze. *Psychopharmacology (Berl)*. 1998;135:290–5.
 55. Borowski T, Kokkinidis L, Merali Z, Anisman H. Lipopolysaccharide, central *in vivo* biogenic amine variations, and anhedonia. *Neuroreport*. 1998;9:3797–802.
 56. Smith JA, Das A, Ray SK, Banik NL. Role of pro-inflammatory cytokines released from microglia in neurodegenerative diseases. *Brain Res Bull*. 2012;87:10–20.
 57. Nakajima K, Kohsaka S. Microglia: activation and their significance in the central nervous system. *J Biochem*. 2001;130:169–75.
 58. Van Dam AM, Bauer J, Tilders FJ, Berkenbosch F. Endotoxin-induced appearance of immunoreactive interleukin-1 β in ramified microglia in rat brain: a light and electron microscopic study. *Neuroscience*. 1995;65:815–26.
 59. Li C, Zhao R, Gao K, Wei Z, Yin MY, Lau LT, et al. Astrocytes: implications for neuroinflammatory pathogenesis of Alzheimer's disease. *Curr Alzheimer Res*. 2011;8:67–80.
 60. Fine SM, Angel RA, Perry SW, Epstein LG, Rothstein JD, Dewhurst S, et al. Tumor necrosis factor α inhibits glutamate uptake by primary human astrocytes. Implications for pathogenesis of HIV-1 dementia. *J Biol Chem*. 1996;271b:15303–6.
 61. Espey MG, Kustova Y, Sei Y, Basile AS. Extracellular glutamate levels are chronically elevated in the brains of LP-BMS5-infected mice: a mechanism of retrovirus-induced encephalopathy. *J Neurochem*. 1998;71:2079–87.
 62. Yirmiya R, Goshen I. Immune modulation of learning, memory, neural plasticity and neurogenesis. *Brain Behav Immun*. 2011;25b:181–213.
 63. Thomson LM, Sutherland RJ. Systemic administration of lipopolysaccharide and interleukin-1 β have different effects on memory consolidation. *Brain Res Bull*. 2005;67:24–9.
 64. Yu XM, Askalan R, Keil 2nd GJ, Salter MW. NMDA channel regulation by channel-associated protein tyrosine kinase Src. *Science*. 1997;275:674–8.
 65. Viviani B, Bartsaghi S, Gardoni F, Vezzani A, Behrens MM, Bartfai T, et al. Interleukin-1 β enhances NMDA receptor-mediated intracellular calcium increase through activation of the Src family of kinases. *J Neurosci*. 2003;23:8692–700.
 66. Cunningham C. Microglia and neurodegeneration: the role of systemic inflammation. *Glia*. 2013;61:71–90.
 67. Gyoneva S, Davalos D, Biswas D, Swanger SA, Garnier-Amblard E, Loth F, et al. Systemic inflammation regulates microglial responses to tissue damage *in vivo*. *Glia*. 2014;62:1345–60.
 68. Qin L, Wu X, Block ML, Liu Y, Breese GR, Hong JS, et al. Systemic LPS causes chronic neuroinflammation and progressive neurodegeneration. *Glia*. 2007;55:453–62.
 69. Castellano C, Cestari V, Ciamei A. NMDA receptors and learning and memory processes. *Curr Drug Targets*. 2001;2:273–83.
 70. Morris RG. NMDA receptors and memory encoding. *Neuropharmacology*. 2013;74:32–40.
 71. Newcomer JW, Krystal JH. NMDA receptor regulation of memory and behavior in humans. *Hippocampus*. 2001;11:529–42.
 72. Paoletti P, Bellone C, Zhou Q. NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity and disease. *Nat Rev Neurosci*. 2013;14:383–400.
 73. Rock DM, Macdonald RL. Polyamine regulation of N-methyl-D-aspartate receptor channels. *Annu Rev Pharmacol Toxicol*. 1995;35:463–82.
 74. Williams K, Romano C, Dichter MA, Molinoff PB. Modulation of the NMDA receptor by polyamines. *Life Sci*. 1991;48:469–98.
 75. Soulet D, Rivest S. Polyamines play a critical role in the control of the innate immune response in the mouse central nervous system. *J Cell Biol*. 2003;162:257–68.
 76. Gerard F, Hansson E. Inflammatory activation enhances NMDA-triggered Ca²⁺ signalling and IL-1 β secretion in primary cultures of rat astrocytes. *Brain Res*. 2012;1473:1–8.
 77. Sparkman NL, Buchanan JB, Heyen JR, Chen J, Beverly JL, Johnson RW. Interleukin-6 facilitates lipopolysaccharide-induced disruption in working memory and expression of other proinflammatory cytokines in hippocampal neuronal cell layers. *J Neurosci*. 2006;26:10709–16.
 78. Czerniawski J, Guzowski JF. Acute neuroinflammation impairs context discrimination memory and disrupts pattern separation processes in hippocampus. *J Neurosci*. 2014;34:12470–80.
 79. Bossu P, Cutuli D, Palladino I, Caporali P, Angelucci F, Laricchiuta D, et al. A single intraperitoneal injection of endotoxin in rats induces long-lasting modifications in behavior and brain protein levels of TNF- α and IL-18. *J Neuroinflammation*. 2012;9:101.
 80. Fiore M, Angelucci F, Alleva E, Branchi I, Probert L, Aloe L. Learning performances, brain NGF distribution and NPY levels in transgenic mice expressing TNF-alpha. *Behav Brain Res*. 2000;112:165–75.
 81. Gong QH, Wang Q, Pan LL, Liu XH, Xin H, Zhu YZ. S-propargyl-cysteine, a novel hydrogen sulfide-modulated agent, attenuates lipopolysaccharide-induced spatial learning and memory impairment: involvement of TNF signaling and NF- κ B pathway in rats. *Brain Behav Immun*. 2011;25:110–9.
 82. Belarbi K, Jopson T, Tweedie D, Arellano C, Luo W, Greig NH, et al. TNF- α protein synthesis inhibitor restores neuronal function and reverses cognitive deficits induced by chronic neuroinflammation. *J Neuroinflammation*. 2012;9:23.
 83. Baune BT, Wiede F, Braun A, Gollidge J, Arolt V, Koerner H. Cognitive dysfunction in mice deficient for TNF- and its receptors. *Am J Med Genet B Neuropsychiatr Genet*. 2008;147B:1056–64.
 84. Camara ML, Corrigan F, Jaehne EJ, Jawahar MC, Anscorn H, Koerner H, et al. TNF- α and its receptors modulate complex behaviours and neurotrophins in transgenic mice. *Psychoneuroendocrinology*. 2013;38:3102–14.
 85. Donzis EJ, Tronson NC. Modulation of learning and memory by cytokines: signaling mechanisms and long term consequences. *Neurobiol Learn Mem*. 2014;115C:68–77.
 86. Moraes CA, Santos G, Spohr TC, D'Avila JC, Lima FR, Benjamim CF, Bozza FA, Gomes FC. Activated microglia-induced deficits in excitatory synapses through IL-1 β : implications for cognitive impairment in sepsis. *Mol Neurobiol*. 2014. doi: 10.1007/s12035-014-8868-5.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

