

RESEARCH

Open Access

Generalized Gronwall-Bellman-type discrete inequalities and their applications

Qinghua Feng^{1,2*}, Fanwei Meng² and Yaoming Zhang¹

* Correspondence: fqhua@sina.com
¹School of Science, Shandong University of Technology, Zibo, 255049 Shandong, China
Full list of author information is available at the end of the article

Abstract

In this paper, some new nonlinear Gronwall-Bellman-type discrete inequalities are established, which can be used as a handy tool in the research of qualitative and quantitative properties of solutions of certain difference equations. The established results generalize some of the recent results obtained by Cheung and Ma, respectively.

Mathematics Subject Classification (2010) 26D15

Keywords: Discrete inequalities, Difference equations, Explicit bounds, Qualitative analysis, Quantitative analysis

1 Introduction

In the past few years, the research of Gronwall-Bellman-type finite difference inequalities has been paid much attention by many authors, which play an important role in the study of qualitative as well as quantitative properties of solutions of difference equations, such as boundedness, stability, existence, uniqueness, continuous dependence and so on. Many difference inequalities have been established (see [1]-[11] and the references therein). But in the analysis of some certain difference equations, the bounds provided by the earlier inequalities are inadequate and it is necessary to seek some new discrete inequalities in order to obtain a diversity of desired results. Our aim in this paper is to establish some new nonlinear Gronwall-Bellman-type discrete inequalities, which provide new bounds for unknown functions lying in these inequalities. We will illustrate the usefulness of the established results by applying them to study the boundedness, uniqueness, and continuous dependence on initial data of solutions of certain difference equations. Our results generalize some of the results in [1,2].

Throughout this paper, \mathbf{R} denotes the set of real numbers and $\mathbf{R}_+ = [0, \infty)$, while \mathbf{Z} denotes the set of integers. The definition domain and the image of a function f are denoted by $Dom(f)$ and $Im(f)$, respectively. $I := [m_0, \infty) \cap \mathbf{Z}$ and $J := [n_0, \infty) \cap \mathbf{Z}$ are two fixed lattices of integral points in \mathbf{R} . Let $\Omega := I \times J \subset \mathbf{Z}^2$, and $\wp(\Omega)$ denotes the set of all \mathbf{R} -valued functions on Ω , while $\wp_+(\Omega)$ denotes the set of all \mathbf{R}_+ -valued functions on Ω . For the sake of convenience, we extend the domain of definition of each function in $\wp(\Omega)$ and $\wp_+(\Omega)$ trivially to the ambient space \mathbf{Z}^2 . So, for example, a function in $\wp(\Omega)$ is regarded as a function defined on \mathbf{Z}^2 with support in $\wp(\Omega)$. As usual, the collection of all continuous functions of a topological space X into a topological space Y will be denoted by $C(X, Y)$. Finally, the partial difference operators Δ_1 and Δ_2 on u

$\in \wp(\mathbb{Z}^2)$ are defined as $\Delta_1 u(m, n) = u(m + 1, n) - u(m, n)$, $\Delta_2 u(m, n) = u(m, n + 1) - u(m, n)$.

2 Main results

Theorem 1 Let $\Omega_{(s, t)} = ([m_0, s] \times [n_0, t]) \cap \Omega$, $(s, t) \in \Omega$. Suppose $u(m, n)$, $a(m, n)$, $k(m, n)$, $b(m, n) \in \wp_+(\Omega)$, and a, k are nondecreasing in every variable. $\eta, \phi \in C(\mathbb{R}_+, \mathbb{R}_+)$, and η are strictly increasing, while ϕ is nondecreasing with $\phi(r) > 0$ for $r > 0$. If for $(m, n) \in \Omega$, $u(m, n)$, satisfies the following inequality

$$\eta(u(m, n)) \leq a(m, n) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \phi(u(s, t)), \quad (1)$$

then for $(m, n) \in \Omega_{(m_1, n_1)}$, we have

$$u(m, n) \leq \eta^{-1} \left\{ G^{-1} \left[G(a(m, n)) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \right] \right\}, \quad (2)$$

where

$$G(z) = \int_{z_0}^z \frac{1}{\phi(\eta^{-1}(z))} dz, z \geq z_0 > 0, \quad (3)$$

and m_1, n_1 are chosen so that for $(m, n) \in \Omega_{(m_1, n_1)}$,

$$G(a(m, n)) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \in \text{Dom}(G^{-1}). \quad (4)$$

Proof Given $(X, Y) \in \Omega_{(m_1, n_1)}$, and let $(m, n) \in \Omega_{(X, Y)}$. Then considering a, k are both nondecreasing, from (1), we have

$$\eta(u(m, n)) \leq a(X, Y) + k(X, Y) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \phi(u(s, t)), \quad (m, n) \in \Omega_{(X, Y)} \quad (5)$$

Let the right side of (5) be $v(m, n)$. Then

$$u(m, n) \leq \eta^{-1}(v(m, n)), \quad (m, n) \in \Omega_{(X, Y)}, \quad (6)$$

and

$$\begin{aligned} \Delta_1 v(m, n) &= v(m + 1, n) - v(m, n) = k(X, Y) \sum_{t=n_0}^{n-1} b(m, t) \phi(u(m, t)) \\ &\leq k(X, Y) \sum_{t=n_0}^{n-1} b(m, t) \phi(\eta^{-1}(v(m, t))) \leq k(X, Y) \phi(\eta^{-1}(v(m, n - 1))) \sum_{t=n_0}^{n-1} b(m, t). \end{aligned} \quad (7)$$

On the other hand, according to the Mean-Value Theorem for integrals, there exists ξ such that $v(m, n) \leq \xi \leq v(m + 1, n)$, and

$$\begin{aligned} \Delta_1 G(v(m, n)) &= G(v(m + 1, n)) - G(v(m, n)) \\ &= \int_{v(m, n)}^{v(m+1, n)} \frac{1}{\phi(\eta^{-1}(z))} dz = \frac{\Delta_1 v(m, n)}{\phi(\eta^{-1}(\xi))} \leq \frac{\Delta_1 v(m, n)}{\phi(\eta^{-1}(v(m, n)))}. \end{aligned} \quad (8)$$

Combining (7) and (8), we obtain

$$\Delta_1 G(v(m, n)) \leq k(X, Y) \frac{\varphi(\eta^{-1}(v(m, n-1)))}{\varphi(\eta^{-1}(v(m, n)))} \sum_{t=n_0}^{n-1} b(m, t) \leq k(X, Y) \sum_{t=n_0}^{n-1} b(m, t). \quad (9)$$

Setting $m = s$, and a summary on both sides of (9) with respect to s from m_0 to $m - 1$ yields

$$\sum_{s=m_0}^{m-1} \Delta_1 G(v(s, n)) \leq k(X, Y) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t), \quad (10)$$

that is,

$$G(v(m, n)) - G(v(m_0, n)) \leq k(X, Y) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t). \quad (11)$$

Considering G is strictly increasing, and $v(m_0, n) = a(X, Y)$, it follows

$$v(m, n) \leq G^{-1} \left[G(a(X, Y)) + k(X, Y) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \right], \quad (m, n) \in \Omega_{(X, Y)}. \quad (12)$$

Combining (6) and (12), we have

$$u(m, n) \leq \eta^{-1} \left\{ G^{-1} \left[G(a(X, Y)) + k(X, Y) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \right] \right\}, \quad (m, n) \in \Omega_{(X, Y)}. \quad (13)$$

Take $m = X, n = Y$ in (13), yields

$$u(X, Y) \leq \eta^{-1} \left\{ G^{-1} \left[G(a(X, Y)) + k(X, Y) \sum_{s=m_0}^{X-1} \sum_{t=n_0}^{Y-1} b(s, t) \right] \right\}. \quad (14)$$

Since $(X, Y) \in \Omega_{(m_1, n_1)}$ are selected arbitrarily, then in fact (14) holds for $\forall (m, n) \in \Omega_{(m_1, n_1)}$, that is

$$u(m, n) \leq \eta^{-1} \left\{ G^{-1} \left[G(a(m, n)) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \right] \right\}, \quad (m, n) \in \Omega_{(m_1, n_1)},$$

which is the desired result. \square

Remark 1 If we take $a(m, n) \equiv c, k(m, n) \equiv 1, \eta(u) = u^\alpha$ in Theorem 2.1, where $c \geq 0, \alpha > 0$ are constants, then Theorem 2.1 reduces to [1, Theorem 2.1].

Corollary 1 Under the conditions of Theorem 2.1, if for $(m, n) \in \Omega, u(m, n)$ satisfies the following inequality

$$\eta(u(m, n)) \leq a(m, n) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \eta(u(s, t)), \quad (15)$$

then for $(m, n) \in \Omega_{(m_1, n_1)}$, we have

$$u(m, n) \leq \eta^{-1}(a(m, n)e^{J(m, n)}), \quad (16)$$

where

$$J(m, n) = k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t). \tag{17}$$

Remark 2 In Corollary 2.2, if we take η for different functions, then we have various bounds for $u(m, n)$. For example, if we take $\eta(u) = u$, then we obtain $u(m, n) \leq a(m, n)e^{J(m, n)}$, while $u(m, n) \leq C^{\frac{1}{p}} e^{\frac{J(m, n)}{p}}$, if we take $\eta(u) = u^p$, $a(m, n) \equiv C$, $k(m, n) \equiv 1$.

Following in a same manner as the proof of Theorem 2.1, we can obtain the following three theorems.

Theorem 2 Let $\Omega_{(s, t)} = ([m_0, s] \times [t, \infty)) \cap \Omega$, $(s, t) \in \Omega$. Suppose u, a, k, b, η, ϕ are defined the same as in Theorem 2.1. If for $(m, n) \in \Omega$, $u(m, n)$ satisfies the following inequality

$$\eta(u(m, n)) \leq a(m, n) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n+1}^{\infty} b(s, t)\phi(u(s, t)), \tag{18}$$

then for $(m, n) \in \Omega_{(m_1, n_1)}$, we have

$$u(m, n) \leq \eta^{-1} \left\{ G^{-1} \left[G(a(m, n)) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n+1}^{\infty} b(s, t) \right] \right\}, \tag{19}$$

where G is defined as in Theorem 2.1, and m_1, n_1 are chosen so that for $G(a(m, n)) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n+1}^{\infty} b(s, t) \in \text{Dom}(G^{-1})$, $G(a(m, n)) + k(m, n) \sum_{s=m_0}^{m-1} \sum_{t=n+1}^{\infty} b(s, t) \in \text{Dom}(G^{-1})$.

Theorem 3 Let $\Omega_{(s, t)} = ([s, \infty) \times [n_0, t]) \cap \Omega$, $(s, t) \in \Omega$. Suppose u, a, k, b, η, ϕ are defined the same as in Theorem 2.1. If for $(m, n) \in \Omega$, $u(m, n)$ satisfies the following inequality

$$\eta(u(m, n)) \leq a(m, n) + k(m, n) \sum_{s=m+1}^{\infty} \sum_{t=n_0}^{n-1} b(s, t)\phi(u(s, t)), \tag{20}$$

then for $(m, n) \in \Omega_{(m_1, n_1)}$, we have

$$u(m, n) \leq \eta^{-1} \left\{ G^{-1} \left[G(a(m, n)) + k(m, n) \sum_{s=m+1}^{\infty} \sum_{t=n_0}^{n-1} b(s, t) \right] \right\}, \tag{21}$$

where G is defined as in Theorem 2.1, and m_1, n_1 are chosen so that for $G(a(m, n)) + k(m, n) \sum_{s=m+1}^{\infty} \sum_{t=n_0}^{n-1} b(s, t) \in \text{Dom}(G^{-1})$, $G(a(m, n)) + k(m, n) \sum_{s=m+1}^{\infty} \sum_{t=n_0}^{n-1} b(s, t) \in \text{Dom}(G^{-1})$.

Theorem 4 Let $\Omega_{(s, t)} = ([s, \infty) \times [t, \infty)) \cap \Omega$, $(s, t) \in \Omega$. Suppose u, a, k, b, η, ϕ are defined the same as in Theorem 2.1. If for $(m, n) \in \Omega$, $u(m, n)$ satisfies the following inequality

$$\eta(u(m, n)) \leq a(m, n) + k(m, n) \sum_{s=m+1}^{\infty} \sum_{t=n+1}^{\infty} b(s, t)\phi(u(s, t)), \tag{22}$$

then for $(m, n) \in \Omega_{(m_1, n_1)}$, we have

$$u(m, n) \leq \eta^{-1} \left\{ G^{-1} \left[G(a(m, n)) + k(m, n) \sum_{s=m+1}^{\infty} \sum_{t=n+1}^{\infty} b(s, t) \right] \right\}, \quad (23)$$

where G is defined as in Theorem 2.1, and m_1, n_1 are chosen so that for $G(a(m, n)) + k(m, n) \sum_{s=m+1}^{\infty} \sum_{t=n+1}^{\infty} b(s, t) \in \text{Dom}(G^{-1})$, $G(a(m, n)) + k(m, n) \sum_{s=m+1}^{\infty} \sum_{t=n+1}^{\infty} b(s, t) \in \text{Dom}(G^{-1})$.

In the following theorem, we will study a class of Volterra-Fredholm type difference inequality.

Theorem 5 Suppose u, η, ϕ are defined as in Theorem 2.1, $a \in \wp_+(\Omega)$, M, N, C are constants, and $M \in [m_0, \infty) \cap \mathbf{Z}$, $N \in [n_0, \infty) \cap \mathbf{Z}$, $C \geq 0$. If for $(m, n) \in \Omega$, $u(m, n)$ satisfies the following inequality

$$\eta(u(m, n)) \leq C + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t)\phi(u(s, t)) + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t)\phi(u(s, t)), \quad (24)$$

then we have

$$u(m, n) \leq \eta^{-1} \left\{ G^{-1} \left\{ G \left(\tilde{G}^{-1} \left[\sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t) \right] \right) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t) \right\} \right\}, \quad (m, n) \in \Omega, \quad (25)$$

provided that $\tilde{G}(z) = G(2z - C) - G(z)$ is strictly increasing for $z \geq C$, where G is defined as in (3).

Proof Suppose $C > 0$, and let the right side of (24) be $v(m, n)$. Then

$$u(m, n) \leq \eta^{-1}(v(m, n)), \quad (m, n) \in \Omega, \quad (26)$$

and

$$v(m_0, n) = C + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t)\phi(u(s, t)). \quad (27)$$

Furthermore,

$$\begin{aligned} \Delta_1 v(m, n) &= \sum_{t=n_0}^{n-1} a(m, t)\phi(u(m, t)) \\ &\leq \sum_{t=n_0}^{n-1} a(m, t)\phi(\eta^{-1}(v(m, t))) \leq \phi(\eta^{-1}(v(m, n-1))) \sum_{t=n_0}^{n-1} a(m, t). \end{aligned} \quad (28)$$

On the other hand, according to the Mean-Value Theorem for integrals, there exists ξ such that $v(m, n) \leq \xi \leq v(m+1, n)$, and

$$\begin{aligned} \Delta_1 G(v(m, n)) &= G(v(m+1, n)) - G(v(m, n)) \\ &= \int_{v(m, n)}^{v(m+1, n)} \frac{1}{\phi(\eta^{-1}(z))} dz = \frac{\Delta_1 v(m, n)}{\phi(\eta^{-1}(\xi))} \leq \frac{\Delta_1 v(m, n)}{\phi(\eta^{-1}(v(m, n)))}. \end{aligned} \quad (29)$$

Combining (28) and (29), we obtain

$$\Delta_1 G(v(m, n)) \leq \frac{\varphi(\eta^{-1}(v(m, n-1)))}{\varphi(\eta^{-1}(v(m, n)))} \sum_{t=n_0}^{n-1} a(m, t) \leq \sum_{t=n_0}^{n-1} a(m, t). \tag{30}$$

Setting $m = s$, and a summary on both sides of (30) with respect to s from m_0 to $m - 1$ yields

$$\sum_{s=m_0}^{m-1} \Delta_1 G(v(s, n)) \leq \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t), \tag{31}$$

that is,

$$G(v(m, n)) - G(v(m_0, n)) \leq \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t). \tag{32}$$

Considering G is strictly increasing, then it follows

$$v(m, n) \leq G^{-1} \left[G(v(m_0, n)) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t) \right], \quad (m, n) \in \Omega. \tag{33}$$

Take $m = M, n = N$ in (33), we obtain

$$v(M, N) \leq G^{-1} \left[G(v(m_0, N)) + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t) \right]. \tag{34}$$

So,

$$\begin{aligned} 2v(m_0, n) - C &= C + 2 \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t) \varphi(u(s, t)) = v(M, N) \\ &\leq G^{-1} \left[G(v(m_0, N)) + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t) \right] = G^{-1} \left[G(v(m_0, n)) + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t) \right], \end{aligned} \tag{35}$$

that is,

$$G(2v(m_0, n) - C) - G(v(m_0, n)) \leq \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t), \tag{36}$$

which is rewritten as

$$\tilde{G}(v(m_0, n)) \leq \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t). \tag{37}$$

Since \tilde{G} is strictly increasing, then furthermore we have

$$v(m_0, n) \leq \tilde{G}^{-1} \left[\sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t) \right]. \tag{38}$$

Combining (26), (33), and (38), we obtain the desired inequality.

If $C = 0$, we can substitute C with $\varepsilon > 0$ in the proof above and then after letting $\varepsilon \rightarrow 0$, we obtain the desired result. \square

Remark 3 If we take $\eta(u) = u$ in Theorem 2.6, then Theorem 2.6 reduces to [2, Theorem 2.1].

Corollary 2 Under the conditions of Theorem 2.6, if $C > 0$, and for $(m, n) \in \Omega$, $u(m, n)$ satisfies the following inequality

$$\eta(u(m, n)) \leq C + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t)\eta(u(s, t)) + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t)\eta(u(s, t)), \quad (39)$$

then we have

$$u(m, n) \leq \eta^{-1} \left\{ \frac{C}{2 - e^{J(M, N)}} e^{J(m, n)} \right\}, \quad (m, n) \in \Omega, \quad (40)$$

provided that $e^{J(M, N)} < 2$, where $J(m, n) = \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t)$.

Proof From Theorem 2.6, considering $\eta = \phi$, we have

$$G(z) = \int_{z_0}^z \frac{1}{z} dz = \ln z - \ln z_0 \quad (41)$$

and

$$\tilde{G}(z) = G(2z - C) - G(z) = \ln\left(2 - \frac{C}{z}\right). \quad (42)$$

Combining (25), (41) and (42), we can deduce the desired result. \square

Remark 4 In Corollary 2.7, if we take η for different functions, then we have various bounds for $u(m, n)$. For example, if we take $\eta(u) = u$, then we obtain $u(m, n) \leq \frac{C}{2 - e^{J(M, N)}} e^{J(m, n)}$. If we take $\eta(u) = u^p$, then we obtain $u(m, n) \leq \left\{ \frac{C}{2 - e^{J(M, N)}} e^{J(m, n)} \right\}^{\frac{1}{p}}$.

Theorem 6 Suppose u, η, ϕ, a, C, G are defined as in Theorem 2.6. Define $\tilde{J}(z) = G(2z - C) - G(z) - \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t)$. Furthermore, assume \tilde{J} is increasing, and $\exists \xi \geq C$ such that $\tilde{J}(\xi) = 0$. If for $(m, n) \in \Omega$, $u(m, n)$ satisfies (24), then

$$u(m, n) \leq \eta^{-1} \left\{ G^{-1} \left[G(\xi) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t) \right] \right\}, \quad (m, n) \in \Omega. \quad (43)$$

Proof Let the right side of (24) be $v(m, n)$. Then

$$u(m, n) \leq \eta^{-1}(v(m, n)), \quad (m, n) \in \Omega. \quad (44)$$

Similar to the process of (26)-(32), we deduce

$$v(m, n) \leq G^{-1} \left[G(v(m_0, n)) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t) \right], \quad (m, n) \in \Omega, \quad (45)$$

and

$$G(2v(m_0, n) - C) - G(v(m_0, n)) \leq \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t), \quad (46)$$

that is, $\tilde{J}(v(m_0, n)) \leq 0$. Considering $\tilde{J}(\xi) = 0$, then $\tilde{J}(v(m_0, n)) \leq \tilde{J}(\xi)$. Since \tilde{J} is increasing, then

$$v(m_0, n) \leq \xi. \tag{47}$$

Combining (45) and (47), we obtain

$$v(m, n) \leq G^{-1} \left[G(\xi) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t) \right], \quad (m, n) \in \Omega. \tag{48}$$

Then by (44) and (48), we obtain the desired inequality. \square

Remark 5 If we take $\eta(u) = u$ in Theorem 2.8, then Theorem 2.8 reduces to [2, Theorem 2.1].

3 Applications

In this section, we will present some applications for the established results above, and show that they are useful in the study of boundedness, uniqueness, and continuous dependence of solutions of certain difference equations.

Example 1 Consider the following difference equation

$$\Delta_{12} e^{u(m,n)} = F(m, n, u(m, n)) \tag{49}$$

with the initial condition

$$u(m, n_0) = \ln \left[e^{f(m)} + 1 \right], u(m_0, n) = \ln \left[e^{g(n)} + 1 \right], f(m_0) = g(n_0) = 0, \tag{50}$$

where $u \in \wp_+(\Omega)$, $F : \Omega \times \mathbf{R}_+ \rightarrow \mathbf{R}$, $f : I \rightarrow \mathbf{R}$, $g : J \rightarrow \mathbf{R}$.

Theorem 7 Suppose $u(m, n)$ is a solution of (49) and (50), and $|F(m, n, u)| \leq b(m, n) u^p$, $e^{f(m)} + e^{g(n)} \leq C$, where p, C are constants, and $p > 1, C > 0, b \in \wp_+(\Omega)$, then for $(m, n) \in \Omega_{(m_1, n_1)}$, we have

$$u(m, n) \leq \ln \left\{ G^{-1} \left[G(C) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \right] \right\}, \tag{51}$$

where

$$G(z) = \int_{z_0}^z \frac{1}{(\ln z)^p} dz, z \geq z_0 > 0, \tag{52}$$

$\Omega_{(m_1, n_1)} = ([m_0, m_1] \times [n_0, n_1]) \cap \Omega$, and m_1, n_1 are chosen so that for $G(C) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \in \text{Dom}(G^{-1})$, $G(C) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) \in \text{Dom}(G^{-1})$.

Proof The equivalent form of (49) and (50) can be denoted by

$$e^{u(m,n)} = e^{f(m)} + e^{g(n)} + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} F(s, t, u(s, t)). \tag{53}$$

So,

$$e^{u(m,n)} \leq e^{f(m)} + e^{g(n)} + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} |F(s, t, u(s, t))| \leq C + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) u^p(s, t). \tag{54}$$

Then, a suitable application of Theorem 2.1 (with $\eta(u) = e^\alpha$, $\phi(u) = u^p$) to (54) yields the desired result. \square

Lemma 1 For any $u_1, u_2 \in \mathbf{R}^+$, we have

$$|u_1 - u_2| \leq |e^{u_1} - e^{u_2}|. \tag{55}$$

Proof Without loss of generality, we assume $u_1 \geq u_2$, and let u_2 be fixed. Define $f(z) = (e^z - e^{u_2}) - (z - u_2)$. Then we have $f(z) = e_z - 1 \geq 0$ for $z \geq u_2 \geq 0$, which shows $f(z)$ is nondecreasing on $[u_2, \infty)$. So $f(u_1) \geq f(u_2) = 0$, that is, $e^{u_1} - e^{u_2} \geq u_1 - u_2$, which shows $|u_1 - u_2| \leq |e^{u_1} - e^{u_2}|$. \square

Theorem 8 Assume u_1, u_2 are two solutions of (49) and (50), and $|F(m, n, u_1) - F(m, n, u_2)| \leq b(m, n)|u_1 - u_2|, \forall (m, n) \in \Omega$, where $b \in \wp_+(\Omega)$, then $u_1 \equiv u_2$, that is, (49) and (50) has at most one solution.

Proof Since u_1, u_2 are two solutions of (49) and (50), then from (49), (50), and (53), we have

$$e^{u_1(m,n)} = e^{f(m)} + e^{g(n)} + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} F(s, t, u_1(s, t)) \tag{56}$$

and

$$e^{u_2(m,n)} = e^{f(m)} + e^{g(n)} + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} F(s, t, u_2(s, t)). \tag{57}$$

By (56) and (57) and Lemma 3.2, we deduce

$$\begin{aligned} & \left| e^{u_1(m,n)} - e^{u_2(m,n)} \right| \\ &= \left| \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} [F(s, t, u_1(s, t)) - F(s, t, u_2(s, t))] \right| \\ &\leq \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} |F(s, t, u_1(s, t)) - F(s, t, u_2(s, t))| \\ &\leq \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) |u_1(s, t) - u_2(s, t)| \\ &\leq \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) |e^{u_1(s,t)} - e^{u_2(s,t)}|. \end{aligned} \tag{58}$$

A suitable application of Corollary 2.2 to (58) yields $|e^{u_1(m,n)} - e^{u_2(m,n)}| \leq 0$, which implies $u_1 \equiv u_2$, and the proof is complete.

The following theorem deals with the continuous dependence of the solution of (49) and (50) on the function F and the initial value $f(m), g(n)$. \square

Theorem 9 Assume $u(m, n)$ is the solution of (49) and (50), $|F(m, n, u_1) - F(m, n, u_2)| \leq b(m, n)|u_1 - u_2|, \forall (m, n) \in \Omega$, where $b \in \wp_+(\Omega)$, $|e^{f(m)} - \tilde{e}^{f(m)} + e^{g(n)} - \tilde{e}^{g(n)}| \leq \varepsilon$, where $\varepsilon > 0$ is a constant. Furthermore, assume $\bar{u} \in \wp_+(\Omega), \bar{u} \in \wp_+(\Omega)$ is the solution of the following difference equation

$$\Delta_{12} e^{\bar{u}(m,n)} = \bar{F}(m, n, \bar{u}(m, n)) \tag{59}$$

with the initial condition

$$\bar{u}(m, n_0) = \ln[e^{\bar{f}(m)} + 1], \bar{u}(m_0, n) = \ln[e^{\bar{g}(n)} + 1], \bar{f}(m_0) = \bar{g}(n_0) = 0, \quad (60)$$

where $\bar{F} : \Omega \times \mathbf{R}_+ \rightarrow \mathbf{R}$, $\bar{f} : I \rightarrow \mathbf{R}$, $\bar{g} : J \rightarrow \mathbf{R}$, then

$$|u(m, n) - \bar{u}(m, n)| \leq 2\varepsilon e^{J(m,n)}, \quad (61)$$

where

$$J(m, n) = \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t). \quad (62)$$

Proof From (59) and (60), we deduce

$$e^{\bar{u}(m,n)} = e^{\bar{f}(m)} + e^{\bar{g}(n)} + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} \bar{F}(s, t, \bar{u}(s, t)). \quad (63)$$

Then by a combination of (53) and (63), we deduce

$$\begin{aligned} & \left| e^{u(m,n)} - e^{\bar{u}(m,n)} \right| \\ & \leq \left| e^{f(m)} - e^{\bar{f}(m)} + e^{g(n)} - e^{\bar{g}(n)} \right| + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} |F(s, t, u(s, t)) - \bar{F}(s, t, \bar{u}(s, t))| \\ & \leq \varepsilon + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} |F(s, t, u(s, t)) - F(s, t, \bar{u}(s, t)) + F(s, t, \bar{u}(s, t)) - \bar{F}(s, t, \bar{u}(s, t))| \\ & \leq \varepsilon + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} (|F(s, t, u(s, t)) - F(s, t, \bar{u}(s, t))| + |F(s, t, \bar{u}(s, t)) - \bar{F}(s, t, \bar{u}(s, t))|) \quad (64) \\ & \leq 2\varepsilon + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} |F(s, t, u(s, t)) - F(s, t, \bar{u}(s, t))| \\ & \leq 2\varepsilon + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) |u(s, t) - \bar{u}(s, t)| \\ & \leq 2\varepsilon + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} b(s, t) |e^{u(s,t)} - e^{\bar{u}(s,t)}|. \end{aligned}$$

After a suitable application of Corollary 2.2 to (64), we obtain $|e^{u(m,n)} - e^{\bar{u}(m,n)}| \leq 2\varepsilon e^{J(m,n)}$. So from Lemma 3.2, it follows $|u(m, n) - \bar{u}(m, n)| \leq |e^{u(m,n)} - e^{\bar{u}(m,n)}| \leq 2\varepsilon e^{J(m,n)}$, which is the desired result. \square

Example 2 Consider the following Volterra-Fredholm sum-difference equation

$$u^p(m, n) = f^p(m) + g^p(n) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} F(s, t, u(s, t)) + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} F(s, t, u(s, t)) \quad (65)$$

with the initial condition

$$u(m, n_0) = f(m), u(m_0, n) = g(n), f(m_0) = g(n_0) = 0, \quad (66)$$

where $u \in \wp(\Omega)$, $F : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$, $f : I \rightarrow \mathbf{R}$, $g : J \rightarrow \mathbf{R}$, $p \geq 1$ is an odd number.

Theorem 10 Suppose $u(m, n)$ is a solution of (65) and (66), and $|F(m, n, u)| \leq a(m, n) |u|^p$, $|f^p(m) + g^p(n)| \leq C$, where $C > 0$ is a constant, and $a \in \wp_+(\Omega)$, then we have

$$|u(m, n)| \leq \left\{ \frac{C}{2 - e^{J(M, N)}} e^{J(m, n)} \right\}^{\frac{1}{p}} \tag{67}$$

provided $e^{J(M, N)} < 2$, where

$$J(m, n) = \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t). \tag{68}$$

Proof From (65) and (66), we have

$$\begin{aligned} |u(m, n)|^p &\leq |f^p(m) + g^p(n)| + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} |F(s, t, u(s, t))| + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} |F(s, t, u(s, t))| \\ &\leq C + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t) |u(s, t)|^p + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t) |u(s, t)|^p \end{aligned} \tag{69}$$

Then a suitable application of Corollary 2.7 to (69) yields the desired result. \square

Theorem 11 Assume $|F(m, n, u_1) - F(m, n, u_2)| \leq a(m, n) |u_1^p - u_2^p|, \forall (m, n) \in \Omega$, where $a \in \wp_+(\Omega)$, and $e^{J(M, N)} < 2$, where $J(m, n) = \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t)$, then (65) and (66) has at most one solution.

Proof Suppose u_1, u_2 are two solutions of (65) and (66), then from (65) and (66), we have

$$\begin{aligned} &|u_1^p(m, n) - u_2^p(m, n)| \\ &= \left| \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} [F_1(s, t, u(s, t)) - F_2(s, t, u(s, t))] + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} [F_1(s, t, u(s, t)) - F_2(s, t, u(s, t))] \right| \\ &\leq \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} |F_1(s, t, u(s, t)) - F_2(s, t, u(s, t))| + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} |F_1(s, t, u(s, t)) - F_2(s, t, u(s, t))| \\ &\leq \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t) |u_1^p(s, t) - u_2^p(s, t)| + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} a(s, t) |u_1^p(s, t) - u_2^p(s, t)|. \end{aligned} \tag{70}$$

A suitable application of Corollary 2.7 to (70) yields $|u_1^p(m, n) - u_2^p(m, n)| \leq 0$, that is, $u_1^p(m, n) = u_2^p(m, n), \forall (m, n) \in \Omega$. Since p is an odd number, then $u_1(m, n) = u_2(m, n)$, which implies (65) and (66) has at most one solution.

Similar to Theorem 3.4, we have the following result showing the continuous dependence of the solution of (65) and (66) on the function F and the initial data $f(m), g(n)$. \square

Theorem 12 Assume $u(m, n)$ is the solution of (65) and (66), $|F(m, n, u_1) - F(m, n, u_2)| \leq a(m, n) |u_1^p - u_2^p|, \forall (m, n) \in \Omega$, where $a \in \wp_+(\Omega)$, $|f^p(m) - \bar{f}^p(m) + g^p(n) - \bar{g}^p(n)| \leq \varepsilon$, where $\varepsilon > 0$ is a constant. Furthermore, assume $\bar{u} \in \wp_+(\Omega), \bar{u} \in \wp_+(\Omega)$ is the solution of the following difference equation

$$\bar{u}^p(m, n) = \bar{f}^p(m) + \bar{g}^p(n) + \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} \bar{F}(s, t, \bar{u}(s, t)) + \sum_{s=m_0}^{M-1} \sum_{t=n_0}^{N-1} \bar{F}(s, t, \bar{u}(s, t)) \tag{71}$$

with the initial condition

$$\bar{u}(m, n_0) = \bar{f}(m), \bar{u}(m_0, n) = \bar{g}(n), \bar{f}(m_0) = \bar{g}(n_0) = 0, \tag{72}$$

where $\bar{F} : \Omega \times \mathbf{R} \rightarrow \mathbf{R}$, $\bar{f} : I \rightarrow \mathbf{R}$, $\bar{g} : J \rightarrow \mathbf{R}$, then

$$|u^p(m, n) - \bar{u}^p(m, n)| \leq \frac{2\varepsilon}{2 - e^{J(M, N)}} e^{J(m, n)}, \quad (m, n) \in \Omega, \quad (73)$$

provided that $e^{J(M, N)} < 2$, where $J(m, n) = \sum_{s=m_0}^{m-1} \sum_{t=n_0}^{n-1} a(s, t)$.

The proof for Theorem 3.7 is similar to Theorem 3.4, and we omit it here.

4 Competing interests

The authors declare that they have no competing interests.

5 Authors' contributions

QF carried out the main part of this article. All authors read and approved the final manuscript.

Acknowledgements

This work is supported by National Natural Science Foundation of China (Grant No. 10571110), Natural Science Foundation of Shandong Province (ZR2009AM011 and ZR2010AZ003) (China) and Specialized Research Fund for the Doctoral Program of Higher Education (20103705110003)(China). The authors thank the referees very much for their careful comments and valuable suggestions on this paper.

Author details

¹School of Science, Shandong University of Technology, Zibo, 255049 Shandong, China ²School of Mathematical Sciences, Qufu Normal University, Qufu, 273165 Shandong, China

Received: 4 March 2011 Accepted: 2 September 2011 Published: 2 September 2011

References

1. Cheung, WS, Ren, JL: Discrete non-linear inequalities and applications to boundary value problems. *J Math Anal Appl.* **319**, 708–724 (2006). doi:10.1016/j.jmaa.2005.06.064
2. Ma, QH: Some new nonlinear Volterra-Fredholm-type discrete inequalities and their applications. *J Comput Appl Math.* **216**, 451–466 (2008). doi:10.1016/j.cam.2007.05.021
3. Pachpatte, BG: On some new inequalities related to certain inequality arising in the theory of differential equations. *J Math Anal Appl.* **189**, 128–144 (1995). doi:10.1006/jmaa.1995.1008
4. Pang, PYH, Agarwal, RP: On an integral inequality and discrete analogue. *J Math Anal Appl.* **194**, 569–577 (1995). doi:10.1006/jmaa.1995.1318
5. Ma, QH: N-independent-variable discrete inequalities of Gronwall-Ou-lang type. *Ann Differ Equ.* **16**, 813–820 (2000)
6. Meng, FW, Li, WN: On some new nonlinear discrete inequalities and their applications. *J Comput Appl Math.* **158**, 407–417 (2003). doi:10.1016/S0377-0427(03)00475-8
7. Cheung, WS: Some discrete nonlinear inequalities and applications to boundary value problems for difference equations. *J Differ Equ Appl.* **10**, 213–223 (2004). doi:10.1080/10236190310001604238
8. Cheung, WS, Ren, JL: Discrete nonlinear inequalities and applications to boundary value problems. *J Math Anal Appl.* **319**, 708–724 (2006). doi:10.1016/j.jmaa.2005.06.064
9. Ma, QH, Cheung, WS: Some new nonlinear difference inequalities and their applications. *J Comput Appl Math.* **202**, 339–351 (2007). doi:10.1016/j.cam.2006.02.036
10. Meng, FW, Ji, DH: On some new nonlinear discrete inequalities and their applications. *J Comput Appl Math.* **208**, 425–433 (2007). doi:10.1016/j.cam.2006.10.024
11. Ma, QH: Estimates on some power nonlinear Volterra-Fredholm type discrete inequalities and their applications. *J Comput Appl Math.* **233**, 2170–2180 (2010). doi:10.1016/j.cam.2009.10.002

doi:10.1186/1029-242X-2011-47

Cite this article as: Feng et al.: Generalized Gronwall-Bellman-type discrete inequalities and their applications. *Journal of Inequalities and Applications* 2011 **2011**:47.