

Research

Open Access

Early biomarkers and potential mediators of ventilation-induced lung injury in very preterm lambs

Megan J Wallace*¹, Megan E Probyn¹, Valerie A Zahra¹, Kelly Crossley¹, Timothy J Cole², Peter G Davis³, Colin J Morley³ and Stuart B Hooper¹

Address: ¹Department of Physiology, Monash University, Melbourne, Victoria, Australia, ²Department of Biochemistry and Molecular Biology, Monash University, Melbourne, Victoria, Australia and ³Newborn Research, Royal Women's Hospital, Melbourne, Victoria, Australia

Email: Megan J Wallace* - megan.wallace@med.monash.edu.au; Megan E Probyn - m.probyn@uq.edu.au; Valerie A Zahra - valerie.zahra@med.monash.edu.au; Kelly Crossley - kelly.crossley@med.monash.edu.au; Timothy J Cole - tim.cole@med.monash.edu.au; Peter G Davis - pgd@unimelb.edu.au; Colin J Morley - colin@morleys.net; Stuart B Hooper - stuart.hooper@med.monash.edu.au

* Corresponding author

Published: 10 March 2009

Received: 28 November 2008

Respiratory Research 2009, **10**:19 doi:10.1186/1465-9921-10-19

Accepted: 10 March 2009

This article is available from: <http://respiratory-research.com/content/10/1/19>

© 2009 Wallace et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Bronchopulmonary dysplasia (BPD) is closely associated with ventilator-induced lung injury (VILI) in very preterm infants. The greatest risk of VILI may be in the immediate period after birth, when the lungs are surfactant deficient, still partially filled with liquid and not uniformly aerated. However, there have been very few studies that have examined this immediate post-birth period and identified the initial injury-related pathways that are activated. We aimed to determine if the early response genes; connective tissue growth factor (*CTGF*), cysteine rich-61 (*CYR61*) and early growth response 1 (*EGR1*), were rapidly induced by VILI in preterm lambs and whether ventilation with different tidal volumes caused different inflammatory cytokine and early response gene expression.

Methods: To identify early markers of VILI, preterm lambs (132 d gestational age; GA, term ~147 d) were resuscitated with an injurious ventilation strategy (V_T 20 mL/kg for 15 min) then gently ventilated (5 mL/kg) for 15, 30, 60 or 120 min ($n = 4$ in each). To determine if early response genes and inflammatory cytokines were differentially regulated by different ventilation strategies, separate groups of preterm lambs (125 d GA; $n = 5$ in each) were ventilated from birth with a V_T of 5 (VG5) or 10 mL/kg (VG10) for 135 minutes. Lung gene expression levels were compared to levels prior to ventilation in age-matched control fetuses.

Results: *CTGF*, *CYR61* and *EGR1* lung mRNA levels were increased ~25, 50 and 120-fold respectively ($p < 0.05$), within 30 minutes of injurious ventilation. VG5 and VG10 caused significant increases in *CTGF*, *CYR61*, *EGR1*, *IL-1*, *IL-6* and *IL-8* mRNA levels compared to control levels. *CTGF*, *CYR61*, *IL-6* and *IL-8* expression levels were higher in VG10 than VG5 lambs; although only the *IL-6* and *CYR61* mRNA levels reached significance.

Conclusion: *CTGF*, *CYR61* and *EGR1* may be novel early markers of lung injury and mechanical ventilation from birth using relatively low tidal volumes may be less injurious than using higher tidal volumes.

Introduction

The lungs of very preterm infants have an immature distal airway structure, with a thick air/blood barrier and a small surface area for gas-exchange. They are surfactant deficient because undifferentiated epithelial cells predominate with few type II alveolar cells. As a result, very preterm infants often require respiratory support in the minutes following birth. Although essential for survival, mechanical ventilation of very preterm infants is closely associated with a high risk of developing bronchopulmonary dysplasia (BPD). BPD is characterised by a simplification of airways, a cessation of alveolarisation, hypercellularity, variable fibrosis and capillary dysplasia [1].

Ventilator induced lung injury (VILI) in preterm infants is associated with many different forms of mechanical ventilation [2-7]. The inflammation that results from VILI is thought to play an important role in the pathogenesis of BPD. VILI promotes the recruitment of inflammatory cells such as neutrophils and macrophages and induces many pro-inflammatory cytokines, transcription factors and growth factors leading to abnormal lung development [8,9]. These factors include interleukin (IL)-1 β , IL-6, IL-8, IL-10, tumour necrosis factor (TNF)- α , transforming growth factor (TGF)- β ₁, nuclear factor (NF)- κ B and interferon- γ [8,10-13]. Although these factors are elevated in response to VILI, a detectable increase can take many hours or days [14], making it difficult to define the initial injury-related pathways involved [9,15]. Identifying the initial injury pathways is critical as the greatest risk of injury may be during the period immediately after birth when the lungs are partially liquid-filled, are surfactant deficient and are not uniformly aerated [16-18]. However, it is unclear whether the above factors are reliable markers of lung injury in studies that are of short duration e.g. investigations of the neonatal resuscitation period.

One of the histological hallmarks of BPD is hypercellularity of the lung [1] and we have recently demonstrated that VILI rapidly stimulates lung cell proliferation in the immature lung [19]. The early response genes connective tissue growth factor (CTGF), cysteine-rich 61 (CYR61) and early growth response factor 1 (EGR1) are known to promote cell proliferation [20,21] and we have recently shown that they are rapidly activated in response to a fetal lung growth stimulus [22]. Previous studies have also demonstrated that these genes are activated in response to lung injury in adults [23-27], but their role in VILI in the preterm neonate is unknown. Thus, our first aim was to investigate whether these early response genes are activated within 15 min-2 h of an injurious insult to the lungs of preterm lambs, before pathological changes to the lung have occurred. To determine their usefulness as early markers of lung injury, we compared their change in expression with changes in the expression of the inflam-

mation genes *IL-1*, *IL-6*, *IL-8* and *TGF- β ₁*, TNF- α protein levels and NF- κ B activity, which have previously been associated with VILI in neonates [8,11,13]. Our second aim was to determine if the mRNA levels of these genes could differentiate between ventilation strategies likely to induce only a mild degree of VILI. To address that aim we determined the mRNA levels of *CTGF*, *CYR61*, *EGR1*, *IL-1*, *IL-6* and *IL-8* in preterm lambs resuscitated from birth using tidal volumes of 5 or 10 mL/kg. Based on the known roles of CTGF, CYR61 and EGR1, it is possible that their aberrant expression contributes to abnormal lung development in very preterm infants destined to develop BPD.

Methods

Animal experiments

Delivery and ventilation of lambs

All experimental procedures on animals were approved by the Monash University Animal Ethics Committee. Pregnant Merino \times Border Leicester ewes at 125 or 132 days of gestational age (GA; term is \sim 147 d) were anaesthetised and the fetal head and neck were exposed for catheterisation and intubation. The fetus was then delivered and ventilated as described below for 135 min. Arterial blood samples were collected every 5 min for the first 15 min and then every 10 min until the end of the experiment. The peak inspiratory pressure (PIP), positive end expiratory pressure (PEEP), mean airway pressure (P_{aw}), tidal volume (V_T), inspiratory and expiratory times, ventilation rate, arterial blood pressure and heart rate were recorded using a data acquisition system (PowerLab, ADInstruments Pty. Ltd., Castle Hill, NSW, Aust.). The alveolar-arterial oxygen difference (AaDO₂) was calculated using the equation: $(P_{barometric} - P_{H_2O}) \times FiO_2 - (PaCO_2/0.8) - PaO_2$. Control fetuses at the same gestational ages were used to indicate the levels of gene expression prior to ventilation.

Time-course for the activation of early response genes caused by injurious ventilation (IV)

Preterm lambs delivered at 132 d gestation ($n = 16$) were resuscitated and mechanically ventilated from birth using a Dräger "Babylog 8000+" (Dräger Medical, Lubeck, Germany). For the first 15 min after birth, lambs were ventilated with an injurious ventilation (IV) protocol, consisting of a tidal volume (V_T) of 20 mL/kg in the absence of a PEEP. After 15 min, lambs were ventilated using a V_T of 5 mL/kg and 8 cmH₂O PEEP for a further 15 (LI 15), 30 (LI 30), 60 (LI 60) or 120 (LI 120) mins ($n = 4$ for each group).

Affect of tidal volume on the activation of early response genes

Preterm lambs delivered at 125 d GA were resuscitated and mechanically ventilated using the Dräger "Babylog 8000+" set to deliver a guaranteed V_T of either 5 (VG5) or 10 (VG10) mL/kg with 8 cmH₂O of PEEP for 135 min

from birth (15 minute resuscitation stabilisation period followed by 2 h of ventilation; n = 5 in each group). The ventilation settings and experimental protocol for these studies have been described previously [28].

Post-mortem examination and tissue collection

At the end of each experiment lambs were humanely killed with an overdose of sodium pentobarbitone (i.v.). The lungs were removed, weighed and the left bronchus was ligated. The left lung was cut into small sections and snap frozen in liquid nitrogen for analysis of *CTGF*, *CYR61*, *EGR1*, *IL-1*, *IL-6*, *IL-8* and *TGF- β* ₁ mRNA levels, active NF- κ B levels and TNF- α protein concentrations. The right lung was fixed via the airways, using 4% paraformaldehyde at 20 cmH₂O for light microscopy.

Tissue analysis

Active NF- κ B protein levels

NF- κ B protein activity was measured in lung tissue using an electromobility gel-shift assay. Lung nuclear proteins were extracted [29] from lung tissue and the protein concentration was determined using a BioRad DC Protein Assay kit (Sigma Aldrich, Australia). Nuclear protein (8 μ g) was incubated on ice for 20 min with 2 μ l binding buffer (100 mM HEPES, 50 mM MgCl₂, 50% glycerol, 10 mM EDTA, 500 mM potassium glutamate), 1 μ l DTT, 1 μ l poly dIdC and 1 μ l of a double stranded ³²P- κ B DNA probe containing the cognate κ B motif (5'-AGTTGAG-GGGACTTCC-3'; total volume 20 μ l). Samples were then electrophoresed for 2 h at 110 V at room temperature in a 5% non-denaturing polyacrylamide (19:1 Acrylamide:Bis-acrylamide) gel with 0.5 \times TBE buffer. The gel was then dried onto Whatmann 3 mm chromatography paper in a gel drier (Speed Gel SG210D, Savant Instruments, USA) and exposed to a storage phosphor screen for 24 – 48 h at room temperature. The relative levels of active NF- κ B bound to the κ B motif were quantified by measuring the total integrated density of each band using Image-Quant software (Molecular Dynamics, Sunnyvale, CA). To compare values from different electromobility gel-shift assays, values from each treatment group were expressed as a percentage of the mean value obtained from the same age-matched control fetuses that were run on all blots for the each experiment.

TNF- α protein concentration

The concentration of TNF- α in lung tissue was measured using a modified antibody-sandwich method of the enzyme-linked immunosorbent assay [30]. Tissue samples were homogenised in 1 \times PBS and centrifuged at 2,500 rpm for 20 min. Supernatant, plasma or standards (50 μ l) were incubated overnight in a 96-well microtitre plate precoated with 50 μ l of TNF- α mouse ascites monoclonal antibody (diluted 1:250 in 3 mM NaN₃, 20 mM Na₂CO₃, 30 mM NaHCO₃) and blocked with 1% skim milk powder in PBS. Plates were washed five times in PBS with 20% Tween 20 (Wash buffer), then incubated for 2 h with 50 μ l of rabbit anti-TNF- α polyclonal antisera (1:500 dilution in 0.001 M PBS/5%BSA). The plates were then washed with buffer and incubated for 1 h with 50 μ l of sheep anti-rabbit horseradish peroxidase (diluted 1:1000 in 0.01 M PBS/5% BSA). The plates were then washed, 100 μ l tetramethyl benzidine/dimethyl sulphoxide was added and the plates were incubated for 10 – 15 min in the dark before the colour reaction was stopped using 0.5 M sulphuric acid. An automatic plate reader (Original Labsystems Multiskan RC, USA) measured the absorbance (at 450 nm) and the levels of TNF α in each sample were determined by interpolation of the standard curve.

TGF- β ₁ gene expression

TGF- β ₁ mRNA levels in lung tissue were quantified by Northern Blot analysis as previously described [31]. The total integrated density of the *TGF- β* ₁ mRNA transcript was divided by the total integrated density of the 18S rRNA band for that sample to account for minor differences in total RNA loading between lanes. As a result, the band densities are presented as a ratio of the 18S rRNA band density and, therefore, have no units.

Quantitative real-time polymerase chain reaction

EGR1, *CTGF*, *CYR61*, *IL-1*, *IL-6* and *IL-8* mRNA levels in lung tissue were measured using quantitative real-time polymerase chain reaction (qRT-PCR). The primers used for amplification of these genes, the gene accession numbers and the regions amplified are shown in Table 1. Total RNA was extracted, DNase-treated and 1 μ g was reverse transcribed into cDNA (M-MLV Reverse Transcriptase, RNase H Minus, Point Mutant Kit; Promega, Madison,

Table 1: Primers used for quantitative real-time PCR

Gene	GenBank Accession #	Nucleotides amplified	Upstream primer 5'-3'	Downstream primer 5'-3'
<i>EGR1</i>	DQ239634	444–532	AGGGTCACTGTGGAAGGTC	GCAGCTGAAGTCAAAGGAA
<i>CTGF</i>	DQ239672	407–469	TATAGCTCCAGCGACAGCTC	ACGAACTTGACTCAGCCTCA
<i>CYR61</i>	DQ239628	286–354	ATCGTCCAAACAACCTCGTG	GGTAACGCGTGTGGAGATAC
<i>IL-1</i>	NM_001009465	353–473	CGATGAGCTTCTGTGTGATG	CTGTGAGAGGAGGTGGAGAG
<i>IL-6</i>	NM_001009392	598–705	CGCAAAGGTTATCATCATCC	CCCAGGAACTACCAATCA
<i>IL-8</i>	NM_001009401	438–520	CCTCAGTAAAGATGCCAATGA	TGACAAACCCTACACCAGACC
<i>18S</i>	X01117	1495–1673	GTCTGTGATGCCCTTAGATGTC	AAGCTTATGACCCGCACTTAC

WI). qRT-PCR was performed using a Mastercycler[®] ep gradient S realplex real-time PCR system (Eppendorf, Germany) using 20 µl reactions, containing 1 µl cDNA template (1.5 µg/µl for *IL-6*, 1 µg/µl for *IL-1*, *IL-8* and *CTGF*, 500 ng/µl for *EGR1* and 200 ng/µl for *CYR61* and 18S), 1 µl of each forward and reverse primer (10 µM for *IL-1*, *IL-6*, *IL-8*, *CYR61* and 18S and 4 µM for *CTGF* and *EGR1*), 10 µl SYBR green (Platinum[®] SYBR[®] Green qPCR SuperMix-UDG; Invitrogen Life Technologies, Carlsbad, CA) and 7 µl of nuclease-free water. The thermal profile used to amplify the PCR products included an initial 2 min incubation at 95°C, followed by 35–40 cycles of; denaturation at 95°C for 3 sec, annealing at 59°C (*IL-1*, *IL-8* and *EGR1*) or 60°C (*IL-6*, *CTGF* and *CYR61*) for 20 sec and elongation at 72°C for 20 sec. The fluorescence was recorded after each 72°C step. Dissociation curves were performed to ensure that a single PCR product had been amplified for each primer pair. Each sample was measured in triplicate and a control sample, containing no template, was included in each run. A threshold value (C_T value) for each sample was determined. Minor differences in the amount of cDNA template added to each reaction were adjusted by subtracting the C_T value for 18S from the C_T value for the gene of interest (ΔC_T). To enable comparisons between assays, a calibrator sample (in quadruplicate) was run in each assay. The average C_T value for the calibrator sample was subtracted from the ΔC_T of each sample ($\Delta\Delta C_T$). The mRNA levels of genes of interest were normalized using the equation $2^{-\Delta\Delta C_T}$ and the results were expressed relative to the mean mRNA levels of the gene of interest in non-ventilated control fetuses.

Light microscopy and immunohistochemistry for *EGR1* and *CYR61*
Each lobe of each right lung was cut into 5 mm slices. Every second slice was subdivided into 3 sections and 6 sections were chosen at random from each lobe, cut into ~1 cm × 1 cm sections and embedded in paraffin. Paraffin blocks were randomly selected and 5 µm sections were incubated at 60°C for 2 h, deparaffinised in xylene, rehydrated using graded alcohol washes and washed in PBS and either stained with Haematoxylin and Eosin (H&E) or treated further for immunohistochemistry. Sections used for immunohistochemistry were then boiled in sodium citrate (0.01 M, pH 6.0) for 20 mins (in a microwave, on high) to enhance antigen retrieval. Sections were then washed in PBS (*CYR61* 2 × 5 min; *EGR1* 3 × 5 min) and incubated (*CYR61* 5 min; *EGR1* 30 min) in hydrogen peroxide (3%) to block endogenous peroxidase activity. They were then rinsed in water (*CYR61* only), washed in PBS and incubated in blocking/permeabilisation buffer (10% normal goat serum and 0.1% TritonX-100 in 0.05 M TrisHCl for *CYR61* sections or 25% normal goat serum and 5% BSA in 0.05 M TrisHCl for *EGR1* sections) in a humidity chamber (*CYR61* 30 min; *EGR1* 45 min, at room temp). The sections were then incubated with the primary antibodies (*CYR61* Cat# sc-13100; *EGR1* Cat# sc-

189, Santa Cruz Biotechnology, California, USA) diluted in DAKO antibody diluent (*CYR61*, diluted 1:150; *EGR1* diluted 1:200) for either 90 min at room temperature (*CYR61*) or overnight at 4°C (*EGR1*). Sections were then washed in PBS (0.1% Tween-20) for 5 mins (×3) and incubated with a biotinylated secondary antibody (goat anti-rabbit diluted 1:700; Vector laboratories, Burlingame, CA) in PBS/0.1% Tween 20 (*CYR61*) or Dako antibody diluent (*EGR1*) for 1 hour at room temperature. The sections were again washed in PBS (0.1% Tween 20) for 5 mins (×3) before the secondary antibody was detected using the Vectastain ABC detection kit (Vector laboratories). The sections were washed, dehydrated and permanently mounted. Sections that lacked the primary antibodies or the secondary antibody were also included.

Sections were viewed under a light microscope and images were captured at a magnification of 1000× using a digital camera. Analysis was performed on images using ImagePro Plus (Media Cybernetics, MD) on 5 fields of view per section using 3 randomly chosen sections (from different regions of the lungs). For each field of view, the area of tissue positively stained for *EGR1* or *CYR61* was measured and expressed as a percentage of the total area of tissue. The percentage of stained tissue for each lamb was then averaged for each experimental group. Analysis was performed on the alveolar region of the lung, taking care to avoid areas containing major airways or blood vessels.

Data analysis

Data are expressed as the mean ± SEM with the level of statistical significance set at $p < 0.05$. PaCO₂, pHa, SaO₂, FiO₂ and PIP were analysed using a 2-way repeated measures ANOVA. The immunohistochemistry data was analysed by a nested ANOVA. The relative amounts of active NF-κB (all three bands summed) and the mRNA levels of *TGF- β 1*, *CTGF*, *CYR61*, *EGR1*, *IL-6*, *IL-8* and *IL-1* were compared between groups using one-way ANOVA. Significant differences indicated by ANOVA were subjected to a least significant difference post-hoc test to identify differences between individual time points and treatment groups.

Results

Activation of early response genes following IV

All blood gas and ventilation parameters were similar in the four groups of lambs exposed to 15 mins of IV immediately after birth (LI 15, LI 30, LI 60, LI 120). Thus, only data from the lambs ventilated for 2 hrs after the 15 min IV protocol (LI 120) are presented in Fig 1.

Blood gas parameters

Throughout the 135 min experimental period, the SaO₂ remained at or higher than 95% (Fig. 1). The FiO₂ was initially reduced from 0.60 ± 0.18 to 0.27 ± 0.03 at the end




Figure 1
Blood gas parameters following 15 minutes of injurious ventilation. The alveolar-arterial difference in oxygenation (AaDO₂) (A), oxygen saturation (SaO₂) (B), fraction of inspired oxygen (FiO₂) (C), arterial pH (pHa) (D) and partial pressure of CO₂ in arterial blood (PaCO₂) (E) in preterm lambs at 132 days of gestation resuscitated at birth using an injurious ventilation strategy then ventilated gently for 120 minutes. Values are mean ± SEM. The black bar indicates 15 min of ventilation with 20 mL/kg V_T and 0 cmH₂O of positive end-expiratory pressure. The asterisks (*) represent values significantly different (p < 0.05) to the initial (5 min) time point.

of the 15 min IV period (V_T 20 mL/Kg, 0 cmH₂O PEEP), but it was necessary to gradually increase the FiO_2 to a maximum of 0.47 ± 0.13 at 70 mins after completion of the IV period. The AaDO₂ was significantly reduced from 739.6 ± 213.1 mmHg to 285.9 ± 38.0 mmHg by the end of the 15 min IV period and then remained at this level for the duration of the experiment. During the 15 min IV period, the PaCO₂ and pH_a remained unchanged at 15 ± 1 mmHg and 7.66 ± 0.02 , respectively. However, during the remainder of the experimental period, the PaCO₂ gradually increased, reaching a maximum of 64 ± 6 mmHg, and the pH_a gradually decreased, reaching a minimum of 7.18 ± 0.04 (Fig. 1).

Ventilation parameters

During 15 min of IV, the PIP required to administer a V_T of 20 mL/kg (in the absence of PEEP) decreased ($p < 0.02$) from 54 ± 2 cmH₂O at 3 min after birth to 47 ± 3 cmH₂O by the end of the 15 min IV period. Within 10 min of change in ventilation strategy, the PIP required to deliver a V_T of 5 mL/kg with 8 cmH₂O PEEP was reduced ($p < 0.001$) to 32 ± 1 cmH₂O. The required PIP did not change further during the remainder of the 120 min ventilation period. However, because of the increasing PaCO₂ and decreasing pH, it was necessary to gradually increase the ventilation rate from 36.3 ± 6.6 breaths/min at the end of the 15 min IV period to 87.1 ± 18.5 breaths/min at the completion of the experiment. As a result, the mean airway pressure at the end of the 15 min IV period was similar to that at completion of the experiment (15.2 ± 0.5 vs 15.6 ± 0.6 cmH₂O).

Indicators of lung injury

The level of active NF-κB within lung tissue did not significantly change for up to 2 h following 15 min of IV; the levels were similar at 15 ($78.2 \pm 7.9\%$), 30 ($93.2 \pm 27.0\%$), 60 ($109.9 \pm 22\%$) and 120 ($70.4 \pm 23.3\%$) min after IV compared with values prior to ventilation measured in age-matched control fetuses ($100.0 \pm 5.8\%$). Similarly, TGF-β₁ mRNA levels in lung tissue were similar at 15 ($96.4 \pm 2.0\%$), 30 ($99.7 \pm 4.2\%$), 60 ($98.3 \pm 14.1\%$) and 120 ($99.1 \pm 13.6\%$) minutes after IV, compared with the levels before ventilation in age-matched control fetuses ($100.0 \pm 3.8\%$). TNF-α protein levels could not be detected in plasma or tissue homogenates in ventilated lambs or in unventilated age-matched control fetuses.

IV induced a large and sustained increase in *IL-1*, *IL-6* and *IL-8* mRNA levels; 28.3 ± 16.6 , 25.6 ± 13.9 and 74.1 ± 20.4 fold increase respectively ($p < 0.05$), compared with pre-ventilation control values, within 15 mins of completing IV (Fig 2). Although *IL-1* mRNA levels had returned to control levels at 120 mins after completion of the IV period, *IL-6* and *IL-8* mRNA levels remained significantly elevated ($p < 0.05$) at 11.0 ± 3.2 and 42.8 ± 11.3 fold,

respectively, above pre-ventilation control values at this time (Fig 2).

IV also induced a time-dependent increase in mRNA levels for *CTGF*, *EGR1* and *CYR61*. The expression levels of all three genes were significantly higher ($p < 0.05$) at every time point after IV, than the pre-ventilation mRNA levels in age-matched control fetuses. *CTGF* mRNA levels increased 15.5 ± 3.8 fold at 15 mins and increased further to 24.4 ± 2.1 fold the control values at 30 mins after the IV period. *CTGF* mRNA levels in lung tissue then declined to 10.9 ± 2.7 fold at 60 mins and to 7.8 ± 1.5 fold of the control values at 120 mins after the IV period (Fig. 3A). Compared with the values prior to ventilation in age-matched control fetuses, *EGR1* and *CYR61* mRNA levels increased by 123.7 ± 7.0 and 51.3 ± 11.4 fold, respectively, at 15 mins after the IV period. *EGR1* and *CYR61* mRNA levels in lung tissue then declined to 43.9 ± 8.8 and 29.1 ± 4.3 fold above control values at 30 mins, to 13.8 ± 4.1 and 13.7 ± 3.5 fold at 60 mins, and to 11.1 ± 2.7 and 5.6 ± 1.5 fold, respectively, at 120 mins after the IV period (Fig. 3A).

The increase in *CYR61* and *EGR1* gene expression was reflected by a gradual, but marked, increase in the percentage of lung tissue stained positive for these proteins (Fig 3B); representative histological sections immunostained for *CYR61* and *EGR1* are shown in Figure 4. The percentage of lung tissue labelled positive for the *CYR61* and *EGR1* proteins increased from 3.0 ± 1.4 and $11.2 \pm 1.2\%$ before ventilation in control fetuses to 16.8 ± 2.9 and $31.1 \pm 1.6\%$, respectively ($p < 0.05$), at 2 hours after IV (Fig. 3B). Sections of lung tissue that lacked the primary antibodies or the secondary antibody showed no evidence of staining. *CTGF* protein levels could not be determined as none of the commercial antibodies tested recognised ovine *CTGF*.

Affect of tidal volume on the activation of early response genes

Blood gas and ventilation parameters and indices of lung injury

The blood gas and ventilation parameters for these studies have been presented in detail previously [28]. The co-efficient of variation of the delivered V_T was $6.5 \pm 0.3\%$. The PIP and P_{aw} delivered to VG10 lambs was significantly higher ($p < 0.05$) than the PIP and P_{aw} delivered to VG5 lambs throughout the 15 minute resuscitation and 2 h ventilation period (Fig 5). PaCO₂ values were significantly lower ($p < 0.05$) in the VG10 group than the VG5 group throughout the 15 minute resuscitation period and 2 h ventilation period. pH_a values were significantly higher ($p < 0.05$) in lambs ventilated at 10 mL/kg compared with lambs ventilated at 5 mL/kg during the resuscitation period but were not different from the 5 mL/kg lambs during the 2 hour ventilation period. The SaO₂ and AaDO₂




Figure 2
***IL-1*, *-6* and *-8* mRNA levels following injurious ventilation.** *IL-1*, *IL-6* and *IL-8* mRNA levels (mean \pm SEM) in pre-term lamb lungs at 132 days of gestation resuscitated at birth using an injurious ventilation (IV) strategy for 15 minutes, then ventilated gently for 15–120 minutes. Values are expressed as a fold change relative to values in unventilated age-matched control fetuses (T = 0 values). *IL-6* and *IL-8* mRNA levels were significantly higher than the levels in unventilated control fetuses ($p < 0.05$) at all timepoints after the IV period. *IL-1* mRNA levels were significantly higher than the levels in unventilated control fetuses at 15, 30 and 60 minutes after the IV period.

were similar in both groups (Fig 5). Two of the VG10 lambs developed pneumothoraces and the experiments were terminated (just prior to the planned end of the ventilation period). Subpleural air leaks were also observed in three of the VG10 lambs. None of the VG5 lambs developed pneumothoraces and only one developed a subpleural air leak. At least three H&E stained tissue sections from three different regions of the lung from each lamb were closely examined under the light microscope for evidence of lung injury. All lung tissue sections from lambs ventilated with 10 mL/kg showed substantial and consistent evidence of hyaline membranes, cellular debris and epithelial cell detachment in the bronchioles and terminal airspaces of the lungs (Fig 6). In contrast, there was substantial variation within and between the lungs of the lambs ventilated with 5 mL/kg. Hyaline membranes in VG5 lambs were rare and minor in comparison to VG10 lambs and while epithelial cell detachment was a common finding (Fig 6) in all VG5 lambs, there was substantial regional variation. Hyaline membranes and epithelial cell detachment were not observed in lungs from control fetuses.

Indicators of lung inflammation

TNF α protein levels were not detectable and active NF- κ B levels and TGF- β_1 mRNA levels within lung tissue were not

altered by either of the ventilation procedures (data not shown).

The mRNA levels for *IL-1*, *IL-6* and *IL-8* in lung tissue were significantly increased in both groups of ventilated lambs, compared to the levels prior to ventilation measured in age-matched control fetuses ($p < 0.001$; Fig 7). The increase in *IL-1* mRNA levels was similar in VG5 (35.1 ± 12.0 fold) and VG10 (31.5 ± 9.9 fold) lambs and were greater than control levels (1.0 ± 0.3 ; $p < 0.001$). However, the increase in *IL-6* was significantly greater in VG10 (116.9 ± 44.6 fold) lambs compared to VG5 lambs (28.9 ± 4.8 fold, $p < 0.05$), both of which were significantly higher than the levels before ventilation in control fetuses (1.0 ± 0.3 ; $p < 0.001$). The increase in *IL-8* mRNA levels was also greater in the VG10 lambs (92.2 ± 52.4 fold) than in the VG5 lambs (32.8 ± 8.7 fold) and both groups were significantly higher than control levels (1.0 ± 0.4 ; $p < 0.001$), however, due to the large degree of variation between lambs the differences between the two ventilated groups were not statistically significant.

The lung mRNA levels of *EGR1*, *CYR61* and *CTGF* were also significantly increased ($p < 0.01$) in both ventilated groups of lambs, compared to the levels before ventilation in age-matched control fetuses (Fig 7). The fold increase in *EGR1* mRNA levels relative to control levels (1.0 ± 0.2 ; $p < 0.001$) was similar in VG5 (14.8 ± 2.6 fold) and VG10 (14.6 ± 2.5 fold) lambs. The fold increase in *CYR61* mRNA levels was greater in the VG10 (21.2 ± 4.9 fold; $p < 0.01$) lambs than in the VG5 treated lambs (8.8 ± 1.4 fold) and both were significantly greater than the levels prior to ventilation in control fetuses (1.0 ± 0.1 ; $p < 0.01$). The increase in mRNA levels for *CTGF* was also greater in the VG10 (11.8 ± 4.1 fold) lambs than in the VG5 treated lambs (6.5 ± 1.1 fold) but the difference between the ventilated groups failed to reach statistical significance. Both groups of ventilated fetuses had significantly higher *CTGF* mRNA levels than the control fetuses (1.0 ± 0.4 ; $p < 0.001$).

Discussion

Ventilator-induced lung injury (VILI) is closely associated with BPD in very preterm infants [1] and is thought to trigger an inflammatory response which results in abnormal lung development. However, the specific mechanisms by which mechanical ventilation causes lung injury in very preterm infants are largely unknown, as are the pathways resulting in the abnormal lung development that characterise BPD. We have recently demonstrated that VILI in the immature lung induces a rapid increase in distal lung cell proliferation [19] which is consistent with the fibroblast proliferation seen in infants with BPD [1] We have also identified a number of early response genes (*CTGF*, *EGR1* and *CYR61*) that regulate cell proliferation and are thought to play a role in normal lung develop-




Figure 3
CTGF, CYR61 and EGR1 mRNA levels following injurious ventilation. (A) *CTGF*, *CYR61* and *EGR1* lung mRNA levels and (B) the percentage of tissue staining positive for *CYR61* and *EGR1* protein in preterm lambs at 132 days of gestation resuscitated at birth using an injurious ventilation (IV) strategy for 15 minutes, then ventilated gently for 15–120 minutes. All values are mean ± SEM and expressed as a fold change relative to values in unventilated age-matched control fetuses (T = 0 values). The mRNA levels of *CTGF*, *CYR61* and *EGR1* were significantly higher ($p < 0.05$) than the levels prior to ventilation (T = 0), at all time points after IV. The asterisks (*) indicate protein levels of *CYR61* and *EGR1* that were significantly higher ($p < 0.05$) than the levels before ventilation measured in age-matched control fetuses.




Figure 4
EGR1 and CYR61 protein levels in lung tissue following injurious ventilation. Lung tissue sections stained for EGR1 and CYR61 proteins using immunohistochemical techniques. The lung tissue sections shown are representative of the sections from unventilated age-matched control fetuses and preterm lambs at 2 hours after a 15 minute period of injurious ventilation (IV). The brown stain represents lung tissue containing the EGR1 or CYR61 protein. Slides incubated without the primary or secondary antibodies did not show any evidence of brown staining (data not shown).

ment [22]. As these genes are also involved in adult lung injury and disease [24-27], we investigated their activation following VILI in preterm lambs. We found that *CTGF*, *EGR1* and *CYR61* expression is rapidly increased in a time-dependent manner in response to VILI in very preterm lambs and that *CTGF*, *CYR61*, *IL-6* and *IL-8* are differentially expressed during high and low tidal volume ventilation strategies. Thus, it is possible that the abnormal lung development that follows VILI, is explained at least in part by the abnormally high expression of these genes. Furthermore, the reduction in pneumothoraces and sub-pleural air-leaks, the histological evidence of lung injury and our gene expression findings indicate that volume-controlled mechanical ventilation (with PEEP) from birth, using a low tidal volume (5 mL/kg) was less injurious than using a tidal volume of 10 mL/kg.


A primary aim of this study was to determine the degree and rapidity of increase in expression of *CTGF*, *CYR61* and *EGR1* following injurious ventilation, in comparison to that of inflammatory factors that have previously been associated with VILI in neonates [8,11,32]. In the present study TNF α protein was not detectable, while NF- κ B activity and *TGF-1* mRNA levels did not change within 2 hr of VILI, suggesting that these proteins and genes do not form part of the very early response to lung injury in very preterm lambs. In contrast, the increases in *IL-1*, *IL-6* and *IL-*

8 after injurious ventilation support the findings of other studies that have also found these inflammatory cytokines are increased at 2-3 h after injurious ventilation from birth [32,33]. Our study extends those findings to demonstrate that *IL-1*, *IL-6*, *IL-8*, *CTGF*, *CYR61* and *EGR1* all responded very rapidly (within 15 minutes of an injurious resuscitation period) and to levels substantially higher (25-125 fold) than those in unventilated controls. These data suggest that the cascade of events leading to lung inflammation and lung remodelling can be rapidly initiated during the immediate resuscitation period after birth. The abnormally high expression levels of these genes was not only limited to resuscitation with high tidal volumes without PEEP, but also occurred in response to ventilation regimens similar to those commonly used for preterm infants.


CYR61 and *CTGF* are members of the CCN protein family which in mammals consists of 6 proteins (*CYR61*, *CTGF*, nephroblastoma-overexpressed1; *NOV1* and the Wnt-induced secreted proteins; *WISP-1*, *WISP-2* and *WISP-3*; [34]). The CCN family are secreted matricellular proteins that form interactions between the extracellular matrix and cell adhesion molecules, leading to diverse cellular responses including cell proliferation, extracellular matrix production, angiogenesis, adhesion, migration, apoptosis and growth arrest [34].

CTGF induces lung fibroblast proliferation, myofibroblast differentiation [35] and the expression of collagen and other extracellular molecules [34]. *CTGF* has increased expression (0.3 fold) in fetal sheep lungs undergoing accelerated lung growth [22] and *CTGF* knockout mice die at birth of respiratory failure due to defects in the rib cage and pulmonary hypoplasia [36]. Although these data indicate that *CTGF* is important for normal lung growth, abnormally elevated levels of *CTGF* expression are also implicated in the pathogenesis of adult human lung diseases such as idiopathic pulmonary fibrosis [24] and chronic obstructive pulmonary disease [26]. In the adult mouse, bleomycin-induced pulmonary fibrosis [23] and hyperoxia-induced lung injury [25], also exhibit elevated *CTGF* mRNA levels. As fibroblast proliferation, myofibroblast differentiation, hypercellularity and pulmonary fibrosis are commonly associated with VILI in very preterm infants [1] and fetal sheep [19], it is possible that abnormally high *CTGF* expression following VILI (~25 fold in the current study) may contribute to the pathogenesis of BPD.

CYR61 is structurally and functionally similar to *CTGF* and also acts as an early response gene. *CYR61* acts synergistically with other growth factors to potentiate their mitogenic effects on endothelial, epithelial and fibroblast cells [20,37] as well as to promote collagen and cartilage

**Figure 5**

Blood gas and ventilator parameters during VG5 and VG10 ventilation strategies. Arterial pH (pHa) (A), partial pressure of CO₂ in arterial blood (PaCO₂) (B), alveolar-arterial oxygen difference (AaDO₂) (C), peak inspiratory pressure (PIP) (D) and mean airway pressure (P_{aw}) (E) in preterm lambs mechanically ventilated from birth at 125 days of gestation. Lambs were mechanically ventilated with either 5 (VG5) or 10 (VG10) mL/kg. Values are mean ± SEM and the asterisks represent values significantly different ($p < 0.05$) between VG5 and VG10.

**Figure 6**

Histological evidence of lung injury in lambs ventilated with VG5 and VG10 ventilation strategies. Representative haematoxylin and eosin stained lung tissue sections in preterm lambs mechanically ventilated from birth at 125 d of gestation with a tidal volume of 10 mL/kg (**A**) or 5 mL/kg (**B**) and unventilated control fetuses (**C**). Hyaline membranes are shown with arrows and detached epithelial cells are shown with arrowheads.




Figure 7
Interleukin-1 , -6 and -8, EGR1, CYR61 and CTGF mRNA levels in control fetuses and following VG5 and VG10 ventilation strategies. IL-1 , IL-6 and IL-8 (A) and EGR1, CYR61 and CTGF (B) mRNA levels in unventilated age-matched control fetuses and in preterm lambs mechanically ventilated from birth at 125 days of gestation with either 5 (VG5) or 10 (VG10) mL/kg. The values are mean \pm SEM and expressed as a fold-change relative to the mean levels in unventilated control fetuses. The asterisks (*) represents values significantly greater ($p < 0.001$) than values before ventilation measured in age-matched control fetuses. The hash (#) represents values significantly greater than those in the VG5 lambs ($p < 0.05$).

production [34]. Depending on the cellular milieu, the primary role of *CYR61* is thought to be the regulation of angiogenesis by promoting the proliferation of endothelial cells and the production of angiogenic molecules such as vascular endothelial growth factor [38,39]. Interestingly, *CYR61* also up-regulates the expression of inflammatory genes, including *IL-1*, as well as modulators of the extracellular matrix such as proteases and their inhibitors [40]. Similar to *CTGF*, *CYR61* expression is also increased (~0.3 fold) in fetal sheep lungs undergoing accelerated growth [22] and abnormally high levels of *CYR61* have been implicated in the pathogenesis of chronic obstructive pulmonary disease [26] in humans as well as lung injury in adult rodents induced by hyperoxia [25] or volutrauma [27]. Based on its known roles, the large and rapid increase in *CYR61* expression (~50 fold in the current study) may contribute to the abnormal lung pathology caused by VILI via several mechanisms. It may contribute to the hypercellularity and fibrosis by directly stimulating the proliferation of fibroblasts and epithelial cells and may upset the normal balance of angiogenic factors, contributing to dysmorphic capillary growth. It may also contribute to the sustained inflammation and abnormal tissue repair that can occur in response to VILI and is an antecedent of BPD in very preterm infants. Our results indicate that increased *CYR61* expression may play a key role in initiating the cascade of events caused by VILI, as *CYR61* protein levels in lung tissue were increased 6-fold within two hours of VILI.

EGR1 is a transcription factor that is rapidly expressed by diverse stimuli that induce growth, differentiation and apoptosis [41]. *EGR1* up-regulates the expression of cell cycle regulatory proteins, growth factors, cytokines such as *IL-1 β* , *TNF α* and *TGF β* and other transcription factors including itself and matrix proteins [21,42-45]. *EGR1* is up-regulated in the fetal sheep during accelerated lung growth [22] and in hemi-pneumonectomy induced compensatory lung growth in adult mice suggesting that it may play a role in regulating normal lung growth [46]. However, *EGR1* expression is also increased by volutrauma in the adult rat lung [47] and it plays a pivotal role in the response to pulmonary ischaemia-reperfusion injury in the adult mouse [48]. In humans it has been implicated in the pathogenesis of chronic obstructive pulmonary disease [26,49] and vascular pathologies where it can cause vascular lesions, suppress the growth of damaged endothelial cells and modulate vascular tone [reviewed in [43]]. These roles for *EGR1*, suggest the high levels of its expression induced by VILI (~125 fold in the current study), may contribute to abnormal lung development by its ability to induce cell proliferation, impair vascular development, produce matrix proteins and induce cytokines that promote inflammation.

Regardless of whether *CTGF*, *CYR61* and *EGR1* are critical mediators of abnormal lung development caused by VILI, they are likely to be early markers of lung injury. All three genes were very rapidly elevated in response to the injurious ventilation strategy. More importantly, when taken together, the expression levels of *IL-6*, *IL-8*, *CTGF* and *CYR61* appeared to differentiate between ventilation strategies causing different degrees of lung injury. Expression levels of all four genes were lowest in lambs mechanically ventilated with a tidal volume of 5 mL/kg and were higher in lambs mechanically ventilated with 10 mL/kg that exhibited gross and histological evidence of lung injury. In contrast, *EGR1* and *IL-1* appeared not to be sufficiently sensitive to detect any differences between the ventilation strategies. Although the 135 minute ventilation period did not allow time for changes in lung structure to manifest histologically, other evidence indicated that VG10 lambs incurred more lung injury than VG5 lambs. This evidence included the presence of hyaline membranes, detached epithelial cells, red blood cells in the distal lung parenchyma, the presence of blood stained tracheal aspirates, the production of pneumothoraces and subpleural air leaks, and the high PIP required to achieve the tidal volume of 10 mL/kg [28].

Conclusion

The current international guidelines for neonatal resuscitation (ILCOR) provide little guidance on the most appropriate resuscitation techniques that minimise lung injury in the immediate newborn period when the lungs are partially liquid-filled and not uniformly aerated. Our data indicate that VILI during the immediate newborn period can rapidly (within 15 mins) initiate changes in gene expression which are abnormal and likely to potentiate inflammation and to promote abnormal lung development. Furthermore, our studies indicate that resuscitation and mechanical ventilation at birth with relatively high tidal volumes is potentially more injurious than with relatively low tidal volumes. We also conclude that *CTGF*, *CYR61*, *EGR1*, *IL-1 β* , *IL-6* and *IL-8* are likely to be useful biomarkers of VILI in the newborn, particularly in studies of short duration.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MW identified *EGR1*, *CTGF* and *CYR61* as likely candidate genes, oversaw the molecular and histological component of the analyses and prepared the manuscript. MP performed the animal experiments and the *TGF- β* ₁, *TNF- α* and *NF- κ B* analyses. VZ performed the real-time PCR and immunohistochemical analyses. KC supervised the animal experiments and TC provided intellectual input into

the studies and provided editorial assistance with the manuscript. CM, PD and SH designed and supervised the animal experiments, obtained funding for the project and provided intellectual input and editorial assistance with the manuscript.

Acknowledgements

We thank Alison Thiel, Foula Sozo, Peter Dargaville and Naomi McCallion for technical assistance. This study was supported by the National Health and Medical Research Council of Australia.

References

- Coalson JJ: **Pathology of new bronchopulmonary dysplasia.** *Semin Neonatal* 2003, **8**:73-81.
- Gerstmann DR, Minton SD, Stoddard RA, Meredith KS, Monaco F, Bertrand JM, et al.: **The Provo multicenter early high-frequency oscillatory ventilation trial: improved pulmonary and clinical outcome in respiratory distress syndrome.** *Pediatrics* 1996, **98**:1044-1057.
- Stewart TE, Meade MO, Cook DJ, Granton JT, Hodder RV, Lapinsky SE, et al.: **Evaluation of a ventilation strategy to prevent barotrauma in patients at high risk for acute respiratory distress syndrome.** *N Engl J Med* 1998, **338**:355-361.
- Keszler M, Modanlou HD, Brudno DS, Clark FI, Cohen RS, Ryan RM, et al.: **Multicenter controlled clinical trial of high-frequency jet ventilation in preterm infants with uncomplicated respiratory distress syndrome.** *Pediatrics* 1997, **100**:593-599.
- Tsuno K, Prato P, Kolobow T: **Acute lung injury from mechanical ventilation at moderately high airway pressures.** *J Appl Physiol* 1990, **69**:956-961.
- Parker JC, Hernandez LA, Peevy KJ: **Mechanisms of ventilator-induced lung injury.** *Crit Care Med* 1993, **21**:131-143.
- Dreyfuss D, Saumon G: **Ventilator-induced lung injury: lessons from experimental studies.** *Am J Respir Crit Care Med* 1998, **157**:294-323.
- Bose CL, Dammann CE, Laughon MM: **Bronchopulmonary dysplasia and inflammatory biomarkers in the premature neonate.** *Arch Dis Child Fetal Neonatal Ed* 2008, **93**:F455-F461.
- Kotecha S, Chan B, Azam N, Silverman M, Shaw RJ: **Increase in interleukin-8 and soluble intercellular adhesion molecule-1 in bronchoalveolar lavage fluid from premature infants who develop chronic lung disease.** *Arch Dis Child Fetal Neonatal Ed* 1995, **72**:F90-F96.
- Pugin J, Dunn I, Jolliet P, Tassaux D, Magnenat JL, Nicod LP, et al.: **Activation of human macrophages by mechanical ventilation in vitro.** *Am J Physiol* 1998, **275**:L1040-L1050.
- Kotecha S: **Cytokines in chronic lung disease of prematurity.** *Eur J Pediatr* 1996, **2**:S14-S17.
- Held HD, Boettcher S, Hamann L, Uhlig S: **Ventilation-induced chemokine and cytokine release is associated with activation of nuclear factor-kappaB and is blocked by steroids.** *Am J Respir Crit Care Med* 2001, **163**:711-716.
- Coalson JJ, Winter VT, Siler-Khodr T, Yoder BA: **Neonatal chronic lung disease in extremely immature baboons.** *Am J Respir Crit Care Med* 1999, **160**:1333-1346.
- Groneck P, Gotze-Speer B, Oppermann M, Eiffert H, Speer CP: **Association of pulmonary inflammation and increased microvascular permeability during the development of bronchopulmonary dysplasia: a sequential analysis of inflammatory mediators in respiratory fluids of high-risk preterm neonates.** *Pediatrics* 1994, **93**:712-718.
- Ambalavanan N, Novak ZE: **Peptide Growth Factors in Tracheal Aspirates of Mechanically Ventilated Preterm Neonates.** *Pediatr Res* 2003, **53**:240-244.
- Bjorklund LJ, Ingimarsson J, Curstedt T, Larsson A, Robertson B, Werner O: **Lung recruitment at birth does not improve lung function in immature lambs receiving surfactant.** *Acta Anaesthesiol Scand* 2001, **45**:986-993.
- Bjorklund LJ, Ingimarsson J, Curstedt T, John J, Robertson B, Werner O, et al.: **Manual ventilation with a few large breaths at birth compromises the therapeutic effect of subsequent surfactant replacement in immature lambs.** *Pediatr Res* 1997, **42**:348-355.
- Wada K, Jobe AH, Ikegami M: **Tidal volume effects on surfactant treatment responses with the initiation of ventilation in preterm lambs.** *J Appl Physiol* 1997, **83**:1054-1061.
- Allison BJ, Crossley KJ, Flecknoe SJ, Davis PG, Morley CJ, Harding R, et al.: **Ventilation of the very immature lung in utero induces injury and BPD-like changes in lung structure in fetal sheep.** *Pediatr Res* 2008:387-392.
- Brigstock DR: **The connective tissue growth factor/cysteine-rich 61/nephroblastoma overexpressed (CCN) family.** *Endocr Rev* 1999, **20**:189-206.
- Molnar G, Crozat A, Pardee AB: **The immediate-early gene Egr-1 regulates the activity of the thymidine kinase promoter at the G0-to-G1 transition of the cell cycle.** *Mol Cell Biol* 1994, **14**:5242-5248.
- Sozo F, Wallace MJ, Zahra VA, Filby CE, Hooper SB: **Gene expression profiling during increased fetal lung expansion identifies genes likely to regulate development of the distal airways.** *Physiol Genomics* 2006, **24**:105-113.
- Lasky JA, Ortiz LA, Tonthat B, Hoyle GW, Corti M, Athas G, et al.: **Connective tissue growth factor mRNA expression is upregulated in bleomycin-induced lung fibrosis.** *Am J Physiol* 1998, **275**:L365-L371.
- Pan LH, Yamauchi K, Uzuki M, Nakanishi T, Takigawa M, Inoue H, et al.: **Type II alveolar epithelial cells and interstitial fibroblasts express connective tissue growth factor in IPF.** *Eur Respir J* 2001, **17**:1220-1227.
- Perkowski S, Sun J, Singhal S, Santiago J, Leikauf GD, Albelda SM: **Gene expression profiling of the early pulmonary response to hyperoxia in mice.** *Am J Respir Cell Mol Biol* 2003, **28**:682-696.
- Ning W, Li CJ, Kaminski N, Feghali-Bostwick CA, Alber SM, Di YP, et al.: **Comprehensive gene expression profiles reveal pathways related to the pathogenesis of chronic obstructive pulmonary disease.** *Proc Natl Acad Sci USA* 2004, **101**:14895-14900.
- Ma SF, Grigoryev DN, Taylor AD, Nonas S, Sammani S, Ye SQ, et al.: **Bioinformatic identification of novel early stress response genes in rodent models of lung injury.** *Am J Physiol Lung Cell Mol Physiol* 2005, **289**:L468-L477.
- Probyn ME, Hooper SB, Dargaville PA, McCallion N, Harding R, Morley CJ: **Effects of tidal volume and positive end-expiratory pressure during resuscitation of very premature lambs.** *Acta Paediatr* 2005, **94**:1764-1770.
- Deryckere F, Gannon F: **A one-hour minipreparation technique for extraction of DNA-binding proteins from animal tissues.** *Biotechniques* 1994, **16**:405.
- Egan PJ, Rothel JS, Andrews AE, Seow HF, Wood PR, Nash AD: **Characterization of monoclonal antibodies to ovine tumor necrosis factor-alpha and development of a sensitive immunoassay.** *Vet Immunol Immunopathol* 1994, **41**:259-274.
- Lines A, Nardo L, Phillips ID, Possmayer F, Hooper SB: **Alterations in lung expansion affect surfactant protein A, B and C mRNA levels in fetal sheep.** *Am J Physiol* 1999, **276**:L239-L245.
- Hillman NH, Moss TJM, Kallapur SG, Bachurski C, Pillow JJ, Polglase GR, et al.: **Brief, large tidal volume ventilation initiates lung injury and a systemic response in fetal sheep.** *Am J Respir Crit Care Med* 2007, **176**:575-581.
- Naik AS, Kallapur SG, Bachurski CJ, Jobe AH, Michna J, Kramer BW, et al.: **Effects of ventilation with different positive end-expiratory pressures on cytokine expression in the preterm lamb lung.** *Am J Respir Crit Care Med* 2001, **164**:494-498.
- Brigstock DR: **The CCN family: a new stimulus package.** *J Endocrinol* 2003, **178**:169-175.
- Grotendorst GR, Duncan MR: **Individual domains of connective tissue growth factor regulate fibroblast proliferation and myofibroblast differentiation.** *FASEB J* 2005, **19**:729-738.
- Baguma-Nibasheka M, Kablar B: **Pulmonary hypoplasia in the connective tissue growth factor (Ctgf) null mouse.** *Dev Dyn* 2008, **237**:485-493.
- Sakamoto S, Yokoyama M, Aoki M, Suzuki K, Kakehi Y, Saito Y: **Induction and function of CYR61 (CCN1) in prostatic stromal and epithelial cells: CYR61 is required for prostatic cell proliferation.** *Prostate* 2004, **61**:305-317.
- Zhou D, Herrick DJ, Rosenbloom J, Chaqour B: **Cyr61 Mediates the Expression of VEGF, (alpha)v Integrin and (alpha) Actin Genes through Cytoskeletal-Based Mechanotransduction**

- Mechanisms in Bladder Smooth Muscle Cells.** *J Appl Physiol* 2005, **98**:2344-2354.
39. Kubota S, Takigawa M: **CCN family proteins and angiogenesis: from embryo to adulthood.** *Angiogenesis* 2007, **10**:1-11.
 40. Chen CC, Mo FE, Lau LF: **The angiogenic factor Cyr61 activates a genetic program for wound healing in human skin fibroblasts.** *J Biol Chem* 2001, **276**:47329-47337.
 41. Lee SL, Sadovsky Y, Swirnonoff AH, Polish JA, Goda P, Gavrilina G, et al.: **Luteinizing hormone deficiency and female infertility in mice lacking the transcription factor NGFI-A (Egr-1).** *Science* 1996, **273**:1219-1221.
 42. Fu M, Zhu X, Zhang J, Liang J, Lin Y, Zhao L, et al.: **Egr-1 target genes in human endothelial cells identified by microarray analysis.** *Gene* 2003, **315**:33-41.
 43. Silverman ES, Collins T: **Pathways of Egr-1-mediated gene transcription in vascular biology.** *Am J Pathol* 1999, **154**:665-670.
 44. Reynolds PR, Cosio MG, Hoidal JR: **Cigarette smoke-induced Egr-1 upregulates proinflammatory cytokines in pulmonary epithelial cells.** *Am J Respir Cell Mol Biol* 2006, **35**:314-319.
 45. Khachigian LM, Collins T: **Inducible expression of Egr-1-dependent genes. A paradigm of transcriptional activation in vascular endothelium.** *Circ Res* 1997, **81**:457-461.
 46. Landesberg LJ, Ramalingam R, Lee K, Rosengart TK, Crystal RG: **Upregulation of transcription factors in lung in the early phase of postpneumonectomy lung growth.** *Am J Physiol Lung Cell Mol Physiol* 2001, **281**:L1138-L1149.
 47. Copland IB, Kavanagh BP, Engelberts D, McKerlie C, Belik J, Post M: **Early changes in lung gene expression due to high tidal volume.** *Am J Respir Crit Care Med* 2003, **168**:1051-1059.
 48. Yan SF, Fujita T, Lu J, Okada K, Shan ZY, Mackman N, et al.: **Egr-1, a master switch coordinating upregulation of divergent gene families underlying ischemic stress.** *Nat Med* 2000, **6**:1355-1361.
 49. Zhang W, Yan SD, Zhu A, Zou YS, Williams M, Godman GC, et al.: **Expression of Egr-1 in late stage emphysema.** *Am J Pathol* 2000, **157**:1311-1320.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

