

RESEARCH ARTICLE

Open Access

# Hypochlorous acid and hydrogen peroxide-induced negative regulation of *Salmonella enterica* serovar Typhimurium *ompW* by the response regulator ArcA

Eduardo H Morales<sup>1†</sup>, Iván L Calderón<sup>1†</sup>, Bernardo Collao<sup>1</sup>, Fernando Gil<sup>1</sup>, Steffen Porwollik<sup>2</sup>, Michael McClelland<sup>2,3</sup> and Claudia P Saavedra<sup>1\*</sup>

## Abstract

**Background:** Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and hypochlorous acid (HOCl) are reactive oxygen species that are part of the oxidative burst encountered by *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) upon internalization by phagocytic cells. In order to survive, bacteria must sense these signals and modulate gene expression. Growing evidence indicates that the ArcAB two component system plays a role in the resistance to reactive oxygen species. We investigated the influx of H<sub>2</sub>O<sub>2</sub> and HOCl through OmpW and the role of ArcAB in modulating its expression after exposure to both toxic compounds in *S. Typhimurium*.

**Results:** H<sub>2</sub>O<sub>2</sub> and HOCl influx was determined both *in vitro* and *in vivo*. A *S. Typhimurium ompW* mutant strain ( $\Delta ompW$ ) exposed to sub-lethal levels of H<sub>2</sub>O<sub>2</sub> and HOCl showed a decreased influx of both compounds as compared to a wild type strain. Further evidence of H<sub>2</sub>O<sub>2</sub> and HOCl diffusion through OmpW was obtained by using reconstituted proteoliposomes. We hypothesized that *ompW* expression should be negatively regulated upon exposure to H<sub>2</sub>O<sub>2</sub> and HOCl to better exclude these compounds from the cell. As expected, qRT-PCR showed a negative regulation in a wild type strain treated with sub-lethal concentrations of these compounds. A bioinformatic analysis in search for potential negative regulators predicted the presence of three ArcA binding sites at the *ompW* promoter region. By electrophoretic mobility shift assay (EMSA) and using transcriptional fusions we demonstrated an interaction between ArcA and one site at the *ompW* promoter region. Moreover, qRT-PCR showed that the negative regulation observed in the wild type strain was lost in an *arcA* and in *arcB* mutant strains.

**Conclusions:** OmpW allows the influx of H<sub>2</sub>O<sub>2</sub> and HOCl and is negatively regulated by ArcA by direct interaction with the *ompW* promoter region upon exposure to both toxic compounds.

## Background

Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and hypochlorous acid (HOCl) are reactive oxygen species that are part of the oxidative burst encountered by *S. Typhimurium* upon internalization by phagocytic cells. Under acidic conditions, such as those found inside the phagosome, H<sub>2</sub>O<sub>2</sub> is generated spontaneously by the reaction of two superoxide anion (O<sub>2</sub><sup>-</sup>) molecules [1]. Moreover, *S. Typhimurium* encodes both

periplasmic and cytoplasmic superoxide dismutases that catalyze O<sub>2</sub><sup>-</sup> dismutation to generate H<sub>2</sub>O<sub>2</sub> and molecular oxygen [2-4]. HOCl is produced by the action of myeloperoxidase (MPO) in a reaction that depends on H<sub>2</sub>O<sub>2</sub>, Cl<sup>-</sup> and acidic conditions [5,6]. Taken together, H<sub>2</sub>O<sub>2</sub> and HOCl react with thiol and heme groups, copper and iron salts generating the reactive hydroxyl radical (OH<sup>•</sup>). As a consequence, they produce lipid peroxidation, chlorination of tyrosine residues, oxidation of iron centers, protein cross linking and DNA damage [5-8].

In order to enter Gram negative bacteria, H<sub>2</sub>O<sub>2</sub> and HOCl must be able to cross the outer membrane (OM) and even though several biological membranes are

\* Correspondence: csaavedra@unab.cl

†Equal contributors

<sup>1</sup>Laboratorio de Microbiología Molecular, Facultad Ciencias Biológicas, Universidad Andres Bello, Santiago, Chile

Full list of author information is available at the end of the article

permeable to H<sub>2</sub>O<sub>2</sub>, studies in *E. coli* and *Saccharomyces cerevisiae* showed that this compound cannot diffuse freely [9,10]. For HOCl, diffusion through the OM is reported to be limited [11]. One possibility for H<sub>2</sub>O<sub>2</sub> and HOCl influx through the OM is diffusion through porins. In this context, we recently reported that OmpD, *S. Typhimurium* most abundant OM porin, allows H<sub>2</sub>O<sub>2</sub> diffusion [12]. OM porins are organized as homo-trimers (classic porins) or monomers (small porins) forming aqueous channels that allow the influx of hydrophilic solutes with a molecular weight ≤ 600 Da [13]. Classic porins, including OmpC and OmpF, form β-barrels with 12–22 transmembrane segments while small porins (OmpW) are composed of 8–10 [14,15]. The crystal structure of OmpW from *E. coli* revealed that it forms an 8-stranded β-barrel and functions as an ion channel in lipid bilayers [16,17]. In *Vibrio cholerae*, OmpW was described as an immunogenic 22 KDa protein [18] and its expression is altered by factors such as temperature, salinity, nutrient availability and oxygen levels [19]. Additionally, several studies show that porins are regulated by ROS. Due its oxidant nature and diffusion through the OM, regulation of porin expression must be tightly regulated as a mechanism of controlling OM permeability. Accordingly, *S. Typhimurium* *ompD* and *ompW* expression is regulated in response to H<sub>2</sub>O<sub>2</sub> and paraquat [12,20], respectively, and *S. Enteritidis* and *Typhimurium* exposure to HOCl results in lower levels of *ompD*, *ompC* and *ompF* transcripts [21].

The cellular response to oxidative stress is regulated at the transcriptional level by activating the SoxRS and OxyR regulons in response to O<sub>2</sub><sup>-</sup> and H<sub>2</sub>O<sub>2</sub>, respectively [22,23], however, several studies have provided evidence for a role of the ArcAB two component system in the resistance to ROS induced damage [12,24–26]. ArcA is essential for *S. Enteritidis*, *Typhimurium* and *E. coli* resistance to ROS [24,26,27]. ArcB is a sensor member of the histidine kinase family that is anchored to the inner membrane [28]. In response to oxygen availability, ArcB autophosphorylates in an ATP dependant intramolecular reaction at position His-292 [29,30] and transfers the phosphate group to the cytoplasmic response regulator ArcA [31–33], which binds to promoter regions regulating gene expression [34,35]. ArcB activity is regulated in response to oxygen conditions by the redox state of both the ubiquinone and menaquinone pools [29,36–38]. However, recent studies in *E. coli* show that the system is regulated by the degree of aerobiosis but not by the redox state of the ubiquinone pool, challenging the idea that the system is inhibited by oxidized quinones [39].

In this work we provide further evidence of the role of the ArcAB two component system in the response to ROS under aerobic conditions and show that this system mediates regulation of *ompW* expression in response to a novel signal, HOCl. First we demonstrate, both *in vivo*

and *in vitro*, that OmpW mediates diffusion of H<sub>2</sub>O<sub>2</sub> and HOCl and that exposure of *S. Typhimurium* to these compounds results in a negative regulation of *ompW*. By EMSA and using transcriptional fusions, we demonstrate that the global regulator ArcA binds to the *ompW* promoter region. Furthermore, we show that *ompW* negative regulation observed in wild type cells treated with H<sub>2</sub>O<sub>2</sub> and HOCl was not retained in an *arcA* or *arcB* mutant strain, indicating that the ArcAB two component system mediates *ompW* negative regulation in response to H<sub>2</sub>O<sub>2</sub> and HOCl. These results further expand our knowledge in both the mechanisms of ROS resistance and the role of ArcAB in this process.

## Results and discussion

### The OmpW porin facilitates H<sub>2</sub>O<sub>2</sub> and HOCl diffusion through the OM and reconstituted proteoliposomes

Hydrogen peroxide and hypochlorous acid are ROS generated by phagocytic cells and in order to enter Gram-negative bacteria they must be able to cross the OM. Even though several biological membranes are permeable to H<sub>2</sub>O<sub>2</sub>, studies in *E. coli* and *S. cerevisiae* demonstrate that this compound cannot diffuse freely [9,10]. Additionally, the dielectric properties of H<sub>2</sub>O<sub>2</sub> are comparable to those of water and this compound has a slighter larger dipolar moment, further limiting its diffusion through the OM lipid bilayer. For HOCl, diffusion through the OM is also reported to be limited [11]. Therefore, H<sub>2</sub>O<sub>2</sub> and HOCl must be channeled through the lipid bilayer and one possibility is the influx through porins. We recently demonstrated that the most abundant OM protein in *S. Typhimurium*, OmpD, allows H<sub>2</sub>O<sub>2</sub> diffusion and is regulated by ArcAB [12]. Little is known about the diffusion of HOCl, but genetic evidence has suggested that in *E. coli* porins might be used as entry channels for hypothiocyanate ions (OSCN<sup>-</sup>), a molecule with a similar chemical structure generated by lactoperoxidase using thiocyanate and H<sub>2</sub>O<sub>2</sub> as an oxidant [40]. In one study, *ompC* and *ompF* knockout mutants showed an increased resistance to OSCN<sup>-</sup>, however, a direct role of porins in mediating HOCl diffusion was not evaluated.


To assess whether OmpW allows the diffusion of H<sub>2</sub>O<sub>2</sub> and HOCl, scopoletin and dihydrorhodamine (DHR)-123 probes, respectively, were used to measure uptake of both toxic compounds separately in a wild type, Δ*ompW* and a genetically complemented Δ*ompW* (pBAD-*ompW*) strain as described in methods. The Δ*ompW* strain showed an increase in extracellular fluorescence levels after exposure to H<sub>2</sub>O<sub>2</sub> and HOCl resulting in higher extra/intracellular ratios (24 and 4-fold, respectively) as compared to the wild type strain, indicating that in the absence of OmpW the influx of both toxic compounds is decreased. Genetic complementation of Δ*ompW* resulted in nearly identical

levels of both extra and intracellular fluorescence as those observed in the wild type strain, suggesting that OmpW is necessary for H<sub>2</sub>O<sub>2</sub> and HOCl uptake (Figure 1A and C). Even though OmpW appears as a direct responsible for the influx of the compounds, a pleiotropic effect cannot be ruled out at this point because the absence of OmpW in the mutant strain could be producing a remodeling of the membrane organization.


To establish a direct contribution of OmpW in H<sub>2</sub>O<sub>2</sub> and HOCl transport, we used reconstituted proteoliposomes. OmpW-proteoliposomes showed a decrease in H<sub>2</sub>O<sub>2</sub> and HOCl extra/intraliposomal ratios (3.5 and 5-fold respectively) when compared to free liposomes (Figure 1B and D). Proteoliposomes with *S. Typhimurium* OmpA porin were used as a negative control as previously described [12]. As expected, OmpA-proteoliposomes showed similar levels to those of free liposomes, indicating that OmpW facilitates H<sub>2</sub>O<sub>2</sub> and HOCl uptake.

Since OmpW channels both toxic compounds across the lipid bilayer, we hypothesized that a  $\Delta ompW$  strain should be more resistant to both toxic compounds when compared to the wild type strain. As shown in Figure 2, exposure of  $\Delta ompW$  to H<sub>2</sub>O<sub>2</sub> 4 mM or HOCl 5 mM

resulted in an increase in the number of colony forming units (CFU) after 60 min of treatment. However, at longer periods the CFU count between strains 14028s and  $\Delta ompW$  was similar. At 30 min post-treatment with either of the toxic compounds, strain  $\Delta ompW$  showed an increase from  $1 \times 10^6$  CFU/ml to approximately  $6 \times 10^7$  CFU/ml. In contrast, the CFU/ml count for strain 14028s remained almost unaltered at  $1 \times 10^6$ , resulting in a 1.5-log<sub>10</sub>-fold increase in growth for  $\Delta ompW$ . A similar result was observed after 60 min of treatment where the *ompW* mutant strain showed an increase from  $6 \times 10^7$  to  $1.5 \times 10^9$  CFU/ml while the wild type strain changed from  $1 \times 10^6$  to  $8 \times 10^7$  CFU/ml. Our results suggest that the absence of OmpW in the mutant strain represents an advantage at short time points due to a decreased permeability towards both H<sub>2</sub>O<sub>2</sub> and HOCl. At longer periods, OM permeability should be reduced because exposure to both toxic compounds results in a negative regulation of *S. Typhimurium* porins including OmpD, OmpC and OmpF [12,21]. One important possibility that cannot be ruled out at this time is that in the  $\Delta ompW$  strain, the expression of other porins or the OM lipid composition might be altered, therefore changing OM



**Figure 1** OmpW facilitates H<sub>2</sub>O<sub>2</sub> and HOCl diffusion through the outer membrane and reconstituted proteoliposomes. **A** and **C**. H<sub>2</sub>O<sub>2</sub> and HOCl levels were measured indirectly by specific fluorescence assays in the wild type (14028s), mutant ( $\Delta ompW$ ) and genetically complemented strains ( $\Delta ompW/pBAD-ompW$  + arabinose). Exponentially growing cells were exposed to H<sub>2</sub>O<sub>2</sub> (**A**) or NaOCl (**C**) for 5 min and fluorescence was determined in the extracellular (extra) and intracellular fractions. **B** and **D**. Free liposomes (L), proteoliposomes reconstituted with *S. Typhimurium* OmpW (PL OmpW) or OmpA (PL OmpA) proteins were incubated with H<sub>2</sub>O<sub>2</sub> (**B**) or NaOCl (**D**) for 5 min and fluorescence was determined in the extraliposomal (extra) and intraliposomal fractions. AU indicates arbitrary units. Values represent the average of four independent experiments  $\pm$  SD.



permeability. For example, a study conducted in *E. coli* showed that an *ompC* knockout mutant had increased levels of OmpA [40], however, changes in permeability were not evaluated. Furthermore, this has not been evaluated in a *S. Typhimurium* or *E. coli*  $\Delta ompW$  strain.

Our data supports the proposed model where OmpW allows the influx of small polar molecules, like  $H_2O_2$  and HOCl. The crystal structure of OmpW from *E. coli* revealed that the cross-section of the barrel has approximate dimensions of  $17 \times 12 \text{ \AA}$  along the length of the barrel and although the interior of the channel has a hydrophobic character, the observed single channel activities shows that polar molecules traverse the barrel [17]. Taken together, these results provide biochemical and genetic evidence indicating that both toxic compounds are channeled through OmpW. From our knowledge, this is the first direct evidence of HOCl diffusion through porins. Furthermore, preliminary analyses indicate that  $H_2O_2$  and HOCl channeling is common for *S. Typhimurium* OmpD, OmpC and OmpF porins (unpublished data).


#### Hydrogen peroxide and hypochlorous acid exposure results in *ompW* negative regulation

Since the OmpW porin channels  $H_2O_2$  and HOCl through the OM and exposure to these molecules is detrimental to bacteria, we hypothesized that *ompW* should be negatively regulated when *S. Typhimurium* is exposed to  $H_2O_2$  and HOCl. To study this effect, wild type *S. Typhimurium* cells were grown to mid-log phase, exposed to  $H_2O_2$  or HOCl and *ompW* mRNA levels were measured by qRT-PCR. As seen in Figure 3, exposure to  $H_2O_2$  and HOCl resulted in

lower levels of *ompW* transcripts ( $0.27 \pm 0.04$  and  $0.156 \pm 0.079$ , respectively) relative to control untreated cells. In agreement with our results of *ompW* negative regulation, similar results were observed by Wang et al. (2010) who showed that *S. Enteritidis* and Typhimurium cells exposed to HOCl results in modulation of *ompD*, *ompC*, *ompF* (negatively) and *ompA* (positively) expression. Furthermore, Calderón et al. (2011) demonstrated that the *S. Typhimurium* *ompD* gene is negatively regulated in response to  $H_2O_2$ . Therefore, our and all the published data suggest that in the presence of  $OCl^-$  or  $H_2O_2$  there might be a general lowering in the concentration of porins in the outer membrane, in order to diminish the permeability. To assess the specificity of our assay, we evaluated *ompD*, *ompC* and *arcB* transcript levels as positive (*ompD* and *ompC*) and negative controls (*arcB*). The *arcB* gene was used as a negative control based on our microarray analysis which shows that it remains unaltered under these conditions and between strains 1408s and  $\Delta arcA$  (unpublished data). Our results indicate that after exposure to both toxic compounds, *arcB* transcript levels remain unchanged while those of *ompD* and *ompC* are lowered as compared to untreated cells (Figure 3). Therefore, all the evidence indicates that OM permeability is tightly regulated in response to ROS and could represent a novel mechanism of resistance when bacteria are exposed to these toxic compounds.

#### ArcA binds the *ompW* promoter region

In addition to the *soxRS* and *oxyR* systems, several studies have provided evidence that the ArcAB two component system plays an important role in the resistance to




ROS induced damage. For example, ArcA is essential for *S. Enteritidis* and Typhimurium resistance to ROS [24,27] and *E. coli* mutant strains of the sensor ArcB and the regulator ArcA, show an increased susceptibility to H<sub>2</sub>O<sub>2</sub> [26]. However, neither of these studies identified genes directly regulated by the system under oxidative stress. We recently demonstrated that ArcA negatively regulates the expression of *S. Typhimurium ompD* after H<sub>2</sub>O<sub>2</sub> exposure by direct interaction with its promoter region [12]. To determine if ArcA mediates *ompW* down-regulation in response to H<sub>2</sub>O<sub>2</sub> and HOCl, a search for putative ArcA binding sites at the *ompW* promoter region was performed using Virtual Footprint 3.0 [41]. The analysis predicted the presence of three ArcA binding sites (ABS) located at positions -61 to -70 (ABS-1, forward orientation), -230 to -239 (ABS-2) and -286 to -295 (ABS-3, both in reverse orientation) relative to the experimentally determined transcription start site [42]. Comparison with the extended core region 5'-GTTAATTAATGTTA-3' described by Evans *et al.* (2011) further revealed that only ABS-1 presented a high degree of identity (14 out of 15 nucleotides) with the consensus sequence. To confirm or rule out a direct interaction between ArcA and the predicted binding sites, deletions of the promoter region were generated by PCR (schematized in Figure 4B) and used to perform non-radioactive EMSAs with ArcA and phosphorylated

ArcA (ArcA-P). The purity of the protein was assessed by PAGE and ArcA was the dominant product. Electrophoretic mobility shift with ArcA-P was only observed when incubated with fragments that included ABS-1 (Figure 4C and D, W1 and W4). No shifts were observed in fragments that include both ABS-2 and ABS-3 (W3, even at three-fold excess) or control fragments that did not include any ABS (W2 and W5). Non-phosphorylated ArcA only generated electrophoretic mobility shifts at higher concentrations (over 1200 nM) where the negative controls were also retarded as a result of non-specific binding (Figure 3E). Taken together our bioinformatic and EMSA analyses indicate that ArcA-P binds to the *ompW* promoter region at a site located between positions -80 and -41 and suggests that this site is ABS-1 which is located between positions -70 to -55.

#### Evaluating ArcA binding site 1 (ABS-1) functionality

To further confirm that ABS-1 (Figure 4A) was the functional ArcA binding site mediating *ompW* negative regulation in response to ROS, we constructed transcriptional fusions of the *ompW* promoter region. We generated two different fusions which included the whole promoter from positions +1 to -600, with respect to the translation start site. One construction contained the native promoter (pompW-lacZ) while substitutions that mutated ABS-1 (shown in red and underlined,



**Figure 4 ArcA binding to the *ompW* promoter region.** **A.** *S. Typhimurium ompW* promoter region. Black and red boxes indicate predicted ArcA binding sites. -10 and -35 boxes are underlined. The transcription start site is shown in bold and indicated as +1. The translation start site is underlined and in red. The consensus ArcA binding site is shown under the promoter sequence. **B.** Schematic representation of the *ompW* promoter region. Positions relative to the transcription start site are indicated. ArcA binding sites are indicated as in the text. PCR products used in EMSAs are shown and names of each fragment are indicated. **C, D and E.** EMSA of the *ompW* promoter region. A 3-fold excess (60 ng) of fragments W2 and W3 were incubated with W1 (**C**) and the fragment W4 was incubated with W5 (**D**) and increasing amounts of phosphorylated ArcA as indicated on the top of each gel. (**E**) W1, W2 and W3 were incubated with increasing amounts of non-phosphorylated ArcA.

Figure 5A) were included in the second construction (pompW/ABS1-lacZ). The constructions were transformed into the wild type strain and  $\beta$ -galactosidase activity was measured in response to treatment with H<sub>2</sub>O<sub>2</sub> and HOCl.


The activity of the constructions was compared to the untreated 14028s strain with the wild type fusion. Treatment of this strain with H<sub>2</sub>O<sub>2</sub> and HOCl resulted in lower activity levels ( $0.58 \pm 0.008$  and  $0.53 \pm 0.095$ , respectively), in agreement with qRT-PCR experiments. However, a 5 nucleotide substitution of the most conserved residues at ABS-1 site (pompW/ABS1-lacZ) resulted in no regulation after exposure to either of the toxic compounds ( $1.09 \pm 0.104$  and  $0.93 \pm 0.061$ ), indicating that they are relevant for the transcriptional activity of *ompW* in response to H<sub>2</sub>O<sub>2</sub> and HOCl (Figure 5B). Furthermore, these results are in agreement with EMSAs which indicate that ArcA only binds to fragments containing ABS-1.

#### The ArcAB two component system mediates *ompW* negative regulation


To establish a direct relationship between *ompW* negative regulation and ArcA-P binding to its promoter region, *ompW* expression was evaluated by qRT-PCR in a  $\Delta$ *arcA* strain exposed to H<sub>2</sub>O<sub>2</sub> and HOCl. The negative regulation observed in the wild type strain was not retained in an *arcA* mutant treated with either of the

toxic compounds and *ompW* transcript levels were similar as those observed in untreated cells. Genetic complementation of  $\Delta$ *arcA* restored the negative regulation observed in wild type cells exhibiting lower *ompW* mRNA levels ( $0.161 \pm 0.068$  and  $0.488 \pm 0.027$ , respectively) as compared to untreated cells (Figure 6A and C). Growth of the genetically complemented strain in the presence of glucose (non-induction) resulted in similar *ompW* mRNA levels between treated and untreated cells (data not shown). As controls, we measured *ompD*, *ompC* and *arcB* transcript levels after exposure to H<sub>2</sub>O<sub>2</sub> and HOCl in a  $\Delta$ *arcA* strain. Transcript levels of *ompD* were measured since its expression is regulated by ArcA under ROS conditions [12]. Our results indicate that neither *ompD* or *arcB* transcript levels were decreased after exposure to H<sub>2</sub>O<sub>2</sub> or HOCl while those of *ompC* remained regulated in a  $\Delta$ *arcA* strain treated with either of the toxic compounds (Figure 6A), confirming that ArcA mediates *ompD* regulation under ROS conditions and showing that the expression of *ompC* is ArcA independent and regulated by different mechanisms which remain unsolved to the date, and are under study in our laboratory. Furthermore, our bioinformatic analyses in search for ArcA motifs predicted binding sites in the promoter regions of *ompW* and *ompD*, but not for *ompC* ([12], data not shown).

To determine whether the negative regulation by ArcA was dependant on its cognate sensor ArcB, *ompW*



**Figure 5 Evaluating ArcA binding site 1 (ABS-1) functionality at the *ompW* promoter. (A)** Schematic representation of substitutions generated at the *ompW* promoter. Substituted bases are in red, underlined and shown below the core ArcA binding sequence. Black box indicates ABS-1. -35 is indicated. **(B)** Expression of the wild type and mutagenized regulatory region of *ompW* in *S. Typhimurium*. Strain 14028s was transformed with the reporter plasmids pompW-lacZ (wild type) or pompW/ABS1-lacZ (ABS-1 mutated). Cells were grown to OD ~ 0.4 and treated with H<sub>2</sub>O<sub>2</sub> 1.5 mM or NaOCl 530  $\mu$ M for 20 min and  $\beta$ -galactosidase activity was measured. Values represent the average of three independent experiments  $\pm$  SD.



**Figure 6** ArcAB-dependent expression of *ompW*. *ompW*, *ompD*, *ompC*, *arcB* and *arcA* mRNA levels were measured by qRT-PCR in a (A)  $\Delta arcA$ , (B)  $\Delta arcB$  and (C)  $\Delta arcA/pBAD-arcA$  and  $\Delta arcB/pBAD-arcB$ . *arcB* and *arcA* were used as negative controls in (A) and (B), respectively. Exponentially growing cells were treated with H<sub>2</sub>O<sub>2</sub> 1.5 mM or NaOCl 530 μM for 20 min and transcript levels were measured. Genetically complemented cells were grown in the presence of arabinose 1 mM. Control cells received no treatment. 16S rRNA levels were used for normalization. Values represent the average of three independent experiments ± SD.

mRNA levels were evaluated in a  $\Delta arcB$  strain. In contrast to the negative regulation observed in wild type cells, *ompW* mRNA levels were further increased in a  $\Delta arcB$  strain after exposure to HOCl ( $3.37 \pm 1.09$ ). Transcript levels after treatment with H<sub>2</sub>O<sub>2</sub> were similar as those observed in untreated cells (Figure 6B). One possibility for this result is that in the absence of ArcA, ArcB

might phosphorylate (*i.e.* ArcB-OmpR, [43]) one or more response regulators, either unspecifically or due to cross-talk, which could bind to the promoter region and therefore prevent binding of positive regulators like SoxS, which has been demonstrated to regulate *ompW* and is up-regulated in response to HOCl [20,44]. This could result in constant *ompW* transcript levels as shown in Figure 6A. On the other hand, in the absence of ArcB no phosphorylation occurs and SoxS or other positive regulator(s) might have free accessibility to the *ompW* promoter and therefore increase its expression (Figure 6B), although this possibility has not been evaluated in this study. Genetic complementation of  $\Delta arcB$  restored the negative regulation observed in wild type cells exposed to H<sub>2</sub>O<sub>2</sub> and HOCl ( $0.19 \pm 0.04$  and  $0.24 \pm 0.11$ , respectively, Figure 6C). The *ompD* and *ompC* transcripts levels remained down-regulated after exposure to H<sub>2</sub>O<sub>2</sub> and HOCl in the  $\Delta arcB$  strain, while the negative control *arcA* remained unaltered (Figure 6B).

The ArcA regulon in anaerobically grown *S. Typhimurium* was recently determined [27]. Interestingly, neither *ompD* nor *ompW* expression was down-regulated in an ArcA dependant manner, suggesting that the ArcA regulon under anaerobic and aerobic ROS conditions could be different. Even in *E. coli*, *ompW* expression is suggested to be regulated by FNR in response to oxygen availability [39]. The difference between the ArcA regulons under aerobic and ROS conditions might be explained by studies suggesting that the mechanism of ArcA activation under aerobic conditions is different from those classically described. *E. coli* mutant strains in residue H-717 of ArcB are able to phosphorylate and activate ArcA through the transfer of the phosphate group from residue His-292 under aerobic conditions [45] and Loui *et al.* (2009) suggested that H<sub>2</sub>O<sub>2</sub> resistance is independent of ArcA phosphorylation at residue Asp-54. To the date, the detailed molecular mechanism of ArcAB activation in response to ROS remains unsolved. Therefore, further experiments to unveil the molecular mechanism by which the *S. Typhimurium* ArcAB two component system is activated are needed and under way in our laboratory.

### Conclusion

We provide both genetic and biochemical evidence indicating that the OM porin OmpW mediates the influx of H<sub>2</sub>O<sub>2</sub> and HOCl. The results revealed that the *S. Typhimurium* *ompW* gene is negatively regulated upon exposure to both toxic compounds. Furthermore, we demonstrate that the response regulator ArcA mediates *ompW* negative regulation in response to H<sub>2</sub>O<sub>2</sub> and HOCl via a direct interaction with the upstream region of *ompW*. Taken together, with our previous observation that OmpD mediates influx of H<sub>2</sub>O<sub>2</sub> and is negatively regulated by ArcA in response

to H<sub>2</sub>O<sub>2</sub>, these results further expand our knowledge regarding the coordinated regulatory mechanisms of ROS resistance and the role of ArcAB in this process.

## Methods

### Bacterial strains and growth conditions

Bacterial strains used in this work are listed in Table 1. Cells were grown aerobically with agitation in LB medium at 37°C. Solid media consisted of agar (20 g l<sup>-1</sup>) and plates were incubated at 37°C. Dilutions (1:100) of overnight cultures were used to initiate growth. When necessary, growth media was supplemented with the appropriate antibiotics (see below).

### Strain construction and genetic complementation

*S. Typhimurium arcB* gene was interrupted by gene disruption as previously described [46]. Strain 14028s (wild type) harboring plasmid pKD46 was grown in the presence of arabinose (10 mM) and ampicillin (100 µg ml<sup>-1</sup>) to OD<sub>600</sub> ~ 0.4, made electrocompetent and transformed with a PCR product generated with plasmid pKD3 as template and primers 5' ATTGGGTATTATGTGC-GAAGTTGTGGTGAAGGAATCCTCTTGTAGGCTGGA

GCTGCTTCG 3' (WarcBF) and 5' GGTGTTGGCGCAG-TATTCGCGCACCCCGGTCAAACCGGGGCATATGAA-TATCCTCCTTAG 3' (WarcBR). Transformants were selected on LB plates supplemented with chloramphenicol (20 µg ml<sup>-1</sup>) and confirmed by PCR using primers 5' GCTACGCATATTTCGCACAA 3' (arcBF) and 5' GCGCCTTTGACATCATCATA 3' (arcBR).

Genetic complementation of the  $\Delta arcB$  strain was performed using plasmid pBAD-*arcB*. To generate this plasmid, *S. Typhimurium arcB* gene was amplified by PCR using primers 5' ATGAAGCAAATTCGTATGCTG 3' (pBADarcBF) and 5' TCATTTTTTTTCCGCGTTTGC-CACCC 3' (pBADarcBR) and cloned into plasmid pBAD-TOPO TA<sup>®</sup> (Invitrogen) according to manufacturer's instructions. Insertion was verified by DNA sequencing.

### Bacterial survival after exposure to oxidative stress

Bacteria were cultured in 5 ml of LB medium at 37°C overnight with shaking. Antibiotics were added as appropriate. 1:1000 dilutions of the overnight cultures were grown in 25 ml to OD ~ 0.4 and H<sub>2</sub>O<sub>2</sub> 4 mM or NaOCl 5 mM (final concentration) were added. In all the assays the cultures were grown aerobically at 250 rpm. Aliquots

**Table 1 Bacterial strains used in this study**

Strain	Relevant characteristic(s)	Source
<i>S. Typhimurium</i>		
14028s	wild type strain	G. Mora
14028s/pompW-lacZ	14028s transformed with a derivative of plasmid pLacZ-Basic carrying the <i>ompW</i> promoter (nt -600 to +1)	This work
14028s/pompW/ABS1-lacZ	14028s transformed with a derivative of plasmid pLacZ-Basic carrying the <i>ompW</i> promoter (nt -600 to +1) with substitution GTTAA to TCCGG into position -70 to -66	This work
$\Delta ompW$	<i>ompW::kan</i>	C. Saavedra
$\Delta ompW$ /pBAD- <i>ompW</i>	$\Delta ompW$ strain complemented with pBAD vector carrying the <i>S. Typhimurium ompW</i> gene	C. Saavedra
$\Delta arcA$	<i>arcA::cam</i>	[12]
$\Delta arcA$ / pBAD- <i>arcA</i>	$\Delta arcA$ strain complemented with pBAD vector carrying the <i>S. Typhimurium arcA</i> gene	[12]
$\Delta arcB$	<i>arcB::cam</i>	This work
$\Delta arcB$ / pBAD- <i>arcB</i>	$\Delta arcB$ strain complemented with pBAD vector carrying the <i>S. Typhimurium arcB</i> gene	This work
<i>E. coli</i> Top10	F- <i>mcrA</i> $\Delta(mrr-hsdRMS-mcrBC)$ $\Phi 80lacZ\Delta M15$ $\Delta lacX74$ <i>recA1</i> <i>araD139</i> $\Delta(ara-leu)7697$ <i>galJ</i> <i>galK</i> <i>rpsL</i> (Str <sup>R</sup> ) <i>endA1</i> <i>nupG</i>	Invitrogen
Top10 pBAD- <i>ompW</i>	Top10 transformed with the pBAD vector carrying the <i>S. Typhimurium ompW</i> gene	C. Saavedra
Top10 pBAD- <i>ompA</i>	Top10 transformed with the pBAD vector carrying the <i>S. Typhimurium ompA</i> gene	C. Saavedra
Top10 pBAD- <i>arcB</i>	Top10 transformed with the pBAD vector carrying the <i>S. Typhimurium arcB</i> gene	This work
BL21 pET-TOPOArcA	BL21(DE3) transformed with the pET-TOPO101ArcA vector carrying the <i>S. Typhimurium arcA</i> gene	[12]



of cultures were withdrawn at the different time points, diluted and plated in triplicate. Bacterial cultures were enumerated by counting the number of CFU after overnight incubation to determine the bacterial concentrations.

### Construction of transcriptional fusions with reporter gene *lacZ*

The native *ompW* promoter region from positions +1 to -600 (with respect to the translation start) site was amplified by PCR with primers *ompW\_pLacZ\_-600F\_ATG* 5' CGGGGTACCCCGATATCGAAAATTCGCG 3' and *ompW\_pLacZ\_-1R\_ATG* 5' CCCAAGCTTACCCGCTC-CATCGTTATGGT 3' using genomic DNA from *S. Typhimurium* (strain 14028s). The restriction sites (*KpnI* and *HindIII*, respectively) at the ends of the DNA fragment were introduced by the PCR primers (underlined sequences) and digested with the corresponding enzymes. The digested PCR product was cloned into the multiple cloning site (MCS) of the  $\beta$ -galactosidase reporter vector pLacZ-Basic (GenBank accession no. U13184), Clontech, generating plasmid *pompW-lacZ*. To generate plasmid *pompW/ABS1-lacZ*, primers *ompW\_pLacZ\_-600F\_ATG* with *Mut\_sit\_arcAR* 5' TGTTCTTATAATGCGGAATT-TATTGATCCAG 3' and *ompW\_pLacZ\_-1R\_ATG* with *Mut\_sit\_arcAF* 5' CTGGATCAATAAATTCGGAAT-TATAAGAACA 3' were used to generate overlapping PCR products spanning the whole length of the *ompW* promoter. Mutation of ABS-1 was generated by incorporating substitutions in primers *Mut\_sit\_arcAF* and *Mut\_sit\_arcAR* (underlined sequences). The resulting PCR products were used as templates in a second reaction with primers *ompW\_pLacZ\_-600F\_ATG* and *ompW\_pLacZ\_-1R\_ATG* to generate the mutated *ompW* promoter, which was digested and cloned into the MCS of plasmid pLacZ-Basic. Constructions were confirmed by DNA sequencing. The generated constructs were transformed into wild type strain 14028s. To evaluate activity, cells at  $OD_{600} \sim 0.4$  were grown for 20 min in the presence of  $H_2O_2$  (1.5 mM) or NaOCl (530  $\mu$ M). Control cells received no treatment.  $\beta$ -galactosidase activity was determined as previously described [20].

### Protein purification

His-tagged ArcA used in EMSAs was purified as previously described [12]. Briefly *E. coli* BL21 cells harboring plasmid pET-TOPO-*arcA* were grown in 500 ml of LB medium supplemented with ampicillin (100  $\mu$ g ml<sup>-1</sup>) to  $OD_{600} \sim 0.4$  and protein overexpression was carried out by adding 1 mM IPTG and further growth for 6 h. Protein was purified by affinity chromatography as described by Georgellis *et al.*, (1997).

Outer membrane proteins used in proteoliposomes were purified as described by Calderón *et al.* (2011). *E. coli* Top10 cells carrying pBAD-*ompA* or pBAD-*ompW* were

grown in 500 ml to  $OD_{600} \sim 0.6$  at 37°C and overexpression was performed for 5 h in the presence of 1 mM arabinose. His-tagged porins were purified by affinity chromatography using HisTrap HP columns (Amersham) according to the manufacturer's instructions.

Plasmid pBAD-*ompW* was generated amplifying the coding region of *S. Typhimurium ompW* by PCR using primers 5' ATGAAAAAATTTACAGTGGC 3' (pBAD-*ompWF*) and 5' GAAACGATAGCCTGCCGAG 3' (pBAD-*ompWR*) and cloned into plasmid pBAD-TOPO TA<sup>®</sup> (Invitrogen) according to the manufacturer's instructions. Insertion was verified by DNA sequencing.

### RNA isolation and *ompW* mRNA detection

Overnight cultures were diluted (1:100) and cells were grown to  $OD_{600} \sim 0.4$ . Genetically complemented cells ( $\Delta$ *arcA*/pBAD-*arcA* and  $\Delta$ *arcB*/pBAD-*arcB*) were grown in the presence of arabinose (1 mM) and ampicillin (100  $\mu$ g ml<sup>-1</sup>). At this point,  $H_2O_2$  (1.5 mM) or NaOCl (530  $\mu$ M) was added and cells were grown for 20 min. Control cells received no treatment. After exposure to the toxic compounds, 4 ml were withdrawn from the culture and used to extract total RNA using GenElute Total RNA purification Kit<sup>®</sup> (Sigma). Total RNA treatment with DNase I and cDNA synthesis was performed as previously described [19].

Relative quantification of *ompW* mRNA was performed using Brilliant II SYBR Green QPCR Master Reagent Kit and the Mx3000P detection system (Stratagene). 16S rRNA was used for normalization. Specific primers were 5' ATGAAAAAATTTACAGTGG 3' (RT*ompWF*) and 5' GAAACGATAGCCTGCCGA 3' (RT*ompWR*) for the *ompW* gene; 5' GTAGAATTCCAGGTGTAGCG 3' (16SF) and 5' TTATCACTGGCAGTCTCCTT 3' (16SR) for 16S rRNA gene (16S). The reaction mixture was carried out in a final volume of 20  $\mu$ l containing 1  $\mu$ l of diluted cDNA (1:1000), 0.24  $\mu$ l of each primer (120 nM), 10  $\mu$ l of 10 x Master Mix, 0.14  $\mu$ l of diluted ROX (1:200) and 8.38  $\mu$ l of  $H_2O$ . The reaction was performed under the following conditions: 10 min at 95°C followed by 40 cycles of 30 s at 95°C, 30 s at 53°C and 45 s at 72°C. Finally a melting cycle from 53 to 95°C was performed to check for amplification specificity. Amplification efficiency was calculated from a standard curve constructed by amplifying serial dilutions of RT-PCR products for each gene. These values were used to obtain the fold change in expression for the gene of interest normalized with 16S levels according to [47]. Experiments were performed in three biological and technical replicates.

### DNA binding assays

Non-radioactive EMSAs were performed as described [48]. Briefly, increasing amounts of purified ArcA (phosphorylated and unphosphorylated) were incubated

with 20 or 60 ng of PCR product(s) in binding buffer (100 mM Tris-Cl [pH 7.4], 100 mM KCl, 10 mM MgCl<sub>2</sub>, 10% glycerol, and 2 mM dithiothreitol) for 20 min at 30°C. Reaction mixtures were immediately loaded on prerun 4% native polyacrylamide gels. The DNA-protein complexes were visualized by ethidium bromide staining. PCR fragments used in EMSAs were generated by PCR using reverse primer 5' ACCCGCTCCATCGTTATGGT 3' (ompWR) in combination with 5' GAGCAGACAAATATTTGCAT 3' (300WF) or 5' TATTAGATCACTTATTACTT 3' (170WF) to generate fragments W1 and W2, respectively. Fragment W3 was generated using primers 300WF and 5' GATCCA-GATTAATTTAGAAC 3'. Fragments W4 and W5 were generated by using reverse primer 5' AATTTTTTCA-TACCCGCTCC 3' in combination with primers 5' CCTATAACCAGGATTTTCAA 3' and 170WF, respectively. ArcA phosphorylation was carried out as described by Linch and Lin (1996). Briefly purified ArcA was incubated with 50 mM disodium carbamoyl phosphate (Sigma) in a buffer containing 100 mM Tris-Cl (pH 7.4), 10 mM MgCl<sub>2</sub>, 125 mM KCl, for 1 h at 30°C and used immediately in EMSA assays.

#### **In vivo and in vitro determination of hydrogen peroxide and hypochlorous acid diffusion**

*In vivo* diffusion of H<sub>2</sub>O<sub>2</sub> was assessed as previously described [12]. For HOCl detection, overnight cultures were diluted and cells were grown to OD<sub>600</sub> ~ 0.5. Two ml of bacterial culture were centrifuged for 5 min at 4500 x g and resuspended in 1 ml of 100 mM phosphate buffer (pH 7.2). A 200 µl aliquot was incubated for 5 min with 530 µM NaOCl and constant agitation. Following, cells were vacuum filtered using polycarbonate filters of 0.025 µm (Millipore) and pass through was collected (extracellular fraction). Bacteria retained in the filter were recovered with 1 ml of 50 mM phosphate buffer (pH 7.2) and disrupted by sonication (intracellular fraction). Both fractions (190 µl) were incubated separately with dihydrorhodamine-123 to a final concentration of 5 µM as previously described [49]. The fluorescent product, rhodamine-123, was measured by fluorescence detection with excitation and emission wavelengths of 500 and 536 nm, respectively. HOCl and H<sub>2</sub>O<sub>2</sub> uptake was determined as the extracellular/intracellular fluorescence ratio. The background fluorescence from a bacterial suspension not exposed to either of the toxic compounds was subtracted and results were normalized by protein concentration.

Proteoliposomes were prepared as described [50] with modifications [51]. For *in vitro* diffusion, proteoliposomes were incubated with 1.5 mM H<sub>2</sub>O<sub>2</sub> or 530 µM NaOCl for 5 min, vacuum filtered and pass through was recovered (extraliposomal fraction). Proteoliposomes were recovered from the filters with 2 ml of 50 mM phosphate buffer (pH 7.2) and disrupted by sonication (intraliposomal fraction).

Fluorescence was measured in both fractions as described above and H<sub>2</sub>O<sub>2</sub> or HOCl uptake was determined as the extraliposomal/intraliposomal fluorescence ratio.

#### **Acknowledgments**

This work was supported by grants from FONDECYT #1085131 and #1120384 (to CPS), Universidad Andres Bello DI-34-11/R (to CPS), POSTDOC FONDECYT 3095013 (to ILC) and Universidad Andres Bello DI-50-09/R (to ILC). EHM and BC received doctoral fellowships by CONICYT and MECESUP UAB0802 additionally to EHM. We would like to thank Nicolás Pacheco for his assistance in the UFC experiments. The authors have declared that no competing interests exist. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. Publication fees were covered by FONDECYT grant # 1120384 and from Universidad Andres Bello DI-34-11/R (to CPS).

#### **Author details**

<sup>1</sup>Laboratorio de Microbiología Molecular, Facultad Ciencias Biológicas, Universidad Andres Bello, Santiago, Chile. <sup>2</sup>The Vaccine Research Institute of San Diego, 3030 Bunker Hill Suite# 203, San Diego, CA 92109, USA. <sup>3</sup>Department of Pathology and Laboratory Medicine, D440 Medical Sciences I, University of California, Irvine, CA 92697, USA.

#### **Author's contributions**

EHM and CPS conceived the project. EHM, BC and ILC performed the experiments. FG and SPo conducted partial data analysis. EHM, ILC, MM and CPS wrote the paper. All authors read and approved the final manuscript.

Received: 2 February 2012 Accepted: 30 April 2012

Published: 30 April 2012

#### **References**

1. Fridovich I: *The biology of oxygen radicals*. *Science* 1978, **201**:875–880.
2. Hassett D, Cohen M: **Bacterial adaptation to oxidative stress: implications for pathogenesis and interaction with phagocytic cells**. *FASEB J* 1989, **3**:2574–2582.
3. Canvin J, Langford PR, Wilks KE, Kroll JS: **Identification of sodC encoding periplasmic [CuZn]-superoxide dismutase in Salmonella**. *FEMS Microbiol Lett* 1996, **136**:215–220.
4. Storz G, Imlay JA: **Oxidative stress**. *Curr Opin Microbiol* 1999, **2**:188–194.
5. Thomas E: **Myeloperoxidase: Hydrogen Peroxide, Chloride Antimicrobial System: Nitrogen-Chlorine Derivatives of Bacterial Components in Bactericidal Action Against Escherichia coli**. *Infect Immun* 1979, **23**:522–531.
6. Rosen H, Crowley J, Heinecke J: **Human Neutrophils Use the Myeloperoxidase-Hydrogen Peroxide-Chloride System to Chlorinate but Not Nitrate Bacterial Proteins during Phagocytosis**. *J Biol Chem* 2002, **277**:30463–30468.
7. Hampton M, Kettle A, Winterbourn C: **Inside the Neutrophil Phagosome: Oxidants, Myeloperoxidase and Bacterial Killing**. *Blood* 1998, **92**:3007–3017.
8. Imlay J: **Pathways of Oxidative Damage**. *Annu Rev Microbiol* 2003, **57**:395–418.
9. Seaver LC, Imlay JA: **Hydrogen peroxide fluxes and compartmentalization inside growing Escherichia coli**. *J Bacteriol* 2001, **183**:7182–7189.
10. Sousa-Lopes A, Antunes F, Cyrne L, Marinho HS: **Decreased cellular permeability to H<sub>2</sub>O<sub>2</sub> protects Saccharomyces cerevisiae cells in stationary phase against oxidative stress**. *FEBS Lett* 2004, **578**:152–156.
11. Leyer G, Johnson E: **Acid Adaptation Sensitizes Salmonella Typhimurium to Hypochlorous Acid**. *Appl Environ Microbiol* 1997, **63**:461–467.
12. Calderón IL, Morales E, Caro NJ, Chahuán CA, Collao B, Gil F, Villareal JM, Ipinza F, Mora GC, Saavedra CP: **Response regulator ArcA of Salmonella enterica serovar Typhimurium downregulates the expression of OmpD, a porin facilitating uptake of hydrogen peroxide**. *Res Microbiol* 2011, **162**:214–222.
13. Nikaido H: **Multidrug efflux pumps of gram-negative bacteria**. *J Bacteriol* 1996, **178**:5853–5859.
14. Shulz GE: **β-barrel membrane proteins**. *Curr Opin Struct Biol* 2000, **10**:443–447.
15. Klebba P: *The Porinologist*. *J. Bacteriol.* 2005, **187**:8232–8236.

16. Albrecht R, Zeth K, Soding J, Lupas A, Linke D: Expression, crystallization and preliminary X-ray crystallographic studies of the outer membrane protein OmpW from *Escherichia coli*. *Acta Cryst.* 2006, **62**:415–418.
17. Hong H, Patel DR, Tamm LK, Van den Berg B: The outer membrane protein OmpW forms an eight-stranded beta-barrel with a hydrophobic channel. *J Biol Chem* 2006, **281**:7568–7577.
18. Jalajakumari MB, Manning PA: Nucleotide sequence of the gene *ompW*, encoding a 22kDa immunogenic outer membrane protein of *Vibrio cholerae*. *Nucleic Acids Res* 1990, **18**:2180.
19. Bisweswar N, Nandy RK, Sarkar A, Ghose AC: Structural features, properties and regulation of the outer-membrane protein W (OmpW) of *Vibrio cholerae*. *Microbiology* 2005, **151**:2975–2986.
20. Gil F, Ipinza P, Fuentes J, Fumeron R, Villareal JM, Aspée A, Mora GC, Vásquez CC, Saavedra C: The *ompW* (porin) gene mediates methyl viologen (paraquat) efflux in *Salmonella enterica* serovar Typhimurium. *Res Microbiol* 2007, **158**:529–536.
21. Wang S, Phillippy A, Deng K, Rui X, Li Z, Tortorello ML, Zhang W: Transcriptomic Responses of *Salmonella enteric* Serovars Enteritidis and Typhimurium to Chlorine-Based Oxidative Stress. *Appl Environ Microbiol* 2010, **76**:5013–5024.
22. Christman MF, Morgan RW, Jacobson FS, Ames BN: Positive control of a regulon for defenses against oxidative stress and some heat-shock proteins in *Salmonella typhimurium*. *Cell* 1985, **41**:753–762.
23. Greenberg JT, Monach P, Chou JH, Josephy PD, Demple B: Positive control of a global antioxidant defense regulon activated by superoxide-generating agents in *Escherichia coli*. *Proc Natl Acad Sci* 1990, **87**:6181–6185.
24. Lu S, Killoran PB, Fang FC, Riley LW: The global regulator ArcA controls resistance to reactive nitrogen and oxygen intermediates in *Salmonella enterica* serovar Enteritidis. *Infect Immun* 2002, **70**:451–461.
25. Wong SM, Alugupalli KR, Ram S, Akerley BJ: The ArcA regulon and oxidative stress resistance in *Haemophilus influenzae*. *Mol Microbiol* 2007, **64**:1375–1390.
26. Loui C, Chang AC, Lu S: Role of the ArcAB two-component system in the resistance of *Escherichia coli* to reactive oxygen stress. *BMC Microbiol* 2009, **9**:183.
27. Evans MR, Fink RC, Vazquez-Torres A, Porwollik S, Jones-Carson J, McClelland M, Hassan HM: Analysis of the ArcA regulon in anaerobically grown *Salmonella enterica* sv. Typhimurium. *BMC Microbiol* 2011, **21**:11–58.
28. Iuchi S, Matsuda Z, Fujiwara T, Lin EC: The *arcB* gene of *Escherichia coli* encodes a sensor-regulator protein for anaerobic repression of the arc modulon. *Mol Microbiol* 1990, **4**:715–727.
29. Malpica R, Franco B, Rodriguez C, Kwon O, Georgellis D: Identification of a quinone-sensitive redox switch in the ArcB sensor kinase. *Proc Natl Acad Sci* 2004, **101**:13318–13323.
30. Peña-Sandoval G, Georgellis D: The ArcB Sensor Kinase of *Escherichia coli* Autophosphorylates by an Intramolecular Reaction. *J Bacteriol* 2010, **192**:1735–1739.
31. Iuchi S, Lin EC: Purification and phosphorylation of the Arc regulatory components of *Escherichia coli*. *J Bacteriol* 1992, **174**:5617–5623.
32. Georgellis D, Lynch AS, Lin EC: In vitro phosphorylation study of the arc two-component signal transduction system of *Escherichia coli*. *J Bacteriol* 1997, **179**:5429–5435.
33. Kwon O, Georgellis D, Lin EC: Phosphorelay as the Sole Physiological Route of Signal Transmission by the Arc Two-Component system of *Escherichia coli*. *J Bacteriol* 2000, **182**:3858–3862.
34. Lynch AS, Lin EC: Transcriptional control mediated by the ArcA two-component response regulator protein of *Escherichia coli*: characterization of DNA binding at target promoters. *J Bacteriol* 1996, **178**:6238–6249.
35. Jeon Y, Lee Y, Han J, Kim J, Hwang D: Multimerization of Phosphorylated and Non-phosphorylated ArcA is Necessary for the Response Regulator Function of the Arc Two-Component Signal Transduction System. *J Biol Chem* 2001, **276**:40873–40879.
36. Georgellis D, Kwon O, Lin EC: Quinones as the Redox Signal for the Arc Two-Component System of Bacteria. *Science* 2001, **292**:2314–2316.
37. Alexeeva S, Hellingwerf K, Mattos JT: Requirement of ArcA for Redox Regulation in *Escherichia coli* under Microaerobic but Not Anaerobic or Aerobic Conditions. *J Bacteriol* 2003, **185**:204–209.
38. Bekker M, Alexeeva S, Laan W, Sawers G, Mattos JT, Hellingwerf K: The ArcBA Two-Component System of *Escherichia coli* Is Regulated by the Redox State of both the Ubiquinone and the Menaquinone Pool. *J Bacteriol* 2010, **191**:746–754.
39. Rolfe MD, Ter Beek A, Graham AI, Trotter EW: Shahzad Asif HM, SysMO-SUMO, Sanguinetti G, Teixeira de Mattos J, Poole RK, Green J: Transcript Profiling and Inference of *Escherichia coli* K-12 ArcA Activity across the Range of Physiologically Relevant Oxygen Concentrations. *J Biol Chem* 2011, **286**:10147–10154.
40. De Spiegeleer P, Sermon J, Vanoirbeek K, Aertens A, Michiels CW: Role of porins in sensitivity of *Escherichia coli* to antibacterial activity of the lactoperoxidase enzyme system. *Appl Environ Microbiol* 2005, **71**:3512–3518.
41. Münch R, Hiller K, Grote A, Scheer M, Klein J, Schobert M, Jahn D: Virtual Footprint and PRODORIC: an integrative framework for regulon prediction in prokaryotes. *Bioinformatics* 2005, **21**:4187–4189.
42. Gil F, Hernández-Lucas I, Polanco R, Pacheco N, Collao B, Villareal JM, Nardocci G, Calva E, Saavedra CP: SoxS regulates the expression of the *Salmonella enterica* serovar Typhimurium *ompW* gene. *Microbiology* 2009, **155**:2490–2497.
43. Matsubara M, Kitaoka SI, Takeda SI, Mizuno T: Tuning of the porin expression under anaerobic growth conditions by his-to-Asp cross-phosphorelay through both the EnvZ-osmosensor and ArcB-anaerosensor in *Escherichia coli*. *Genes Cells* 2000, **5**:555–569.
44. Dukan S, Dadon S, Smulski DR, Belkin S: Hypochlorous Acid Activates the Heat Shock and soxRS Systems of *Escherichia coli*. *Appl Environ Microbiol* 1996, **62**:4003–4008.
45. Matsushika A, Mizuno T: A dual-signaling mechanism mediated by the ArcB hybrid sensor kinase containing the histidine-containing phosphotransfer domain in *Escherichia coli*. *J Bacteriol* 1998, **180**:3973–3977.
46. Datsenko KA, Wanner BL: One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci* 2000, **97**:6640–6645.
47. Pfaffl MW: A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 2001, **29**:e45.
48. Mika F, Hengge R: A two-component phosphotransfer network involving ArcB, ArcA, and RssB coordinates synthesis and proteolysis of  $\sigma^S$  (RpoS) in *E. coli*. *Genes Dev* 2005, **19**:2770–2781.
49. Rezk BM, Haenen G, van der Vijgh W, Bast A: Lipic Acid Protects Efficiently Only against a Specific Form of Peroxynitrite-induced Damage. *J Biol Chem* 2004, **279**:9693–9697.
50. Nikaido H, Rosenberg EY: Porin channels in *Escherichia coli*: studies with liposomes reconstituted from purified proteins. *J Bacteriol* 1983, **153**:241–252.
51. Cubillos MA, Lissi EA, Abuin EB: Kinetics of peroxidation of linoleic acid incorporated into DPPC vesicles initiated by the thermal decomposition of 2,2'-azobis(2-amidinopropane) dihydrochloride. *Chem Phys Lipids* 2001, **112**:41–46.

doi:10.1186/1471-2180-12-63

Cite this article as: Morales et al.: Hypochlorous acid and hydrogen peroxide-induced negative regulation of *Salmonella enterica* serovar Typhimurium *ompW* by the response regulator ArcA. *BMC Microbiology* 2012 **12**:63.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

