

Research article

Open Access

The gene for a lectin-like protein is transcriptionally activated during sexual development, but is not essential for fruiting body formation in the filamentous fungus *Sordaria macrospora*

Minou Nowrousian* and Patricia Cebula

Address: Lehrstuhl für Allgemeine und Molekulare Botanik, Ruhr-Universität Bochum, 44801 Bochum, Germany

Email: Minou Nowrousian* - minou.nowrousian@ruhr-uni-bochum.de; Patricia Cebula - patricia.cebula@gmx.de

* Corresponding author

Published: 03 November 2005

Received: 10 September 2005

BMC Microbiology 2005, 5:64 doi:10.1186/1471-2180-5-64

Accepted: 03 November 2005

This article is available from: <http://www.biomedcentral.com/1471-2180/5/64>

© 2005 Nowrousian and Cebula; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The filamentous fungus *Sordaria macrospora* forms complex three-dimensional fruiting bodies called perithecia that protect the developing ascospores and ensure their proper discharge. In previous microarray analyses, several genes have been identified that are downregulated in sterile mutants compared to the wild type. Among these genes was *tap1* (transcript associated with perithecial development), a gene encoding a putative lectin homolog.

Results: Analysis of *tap1* transcript levels in the wild type under conditions allowing only vegetative growth compared to conditions that lead to fruiting body development showed that *tap1* is not only downregulated in developmental mutants but is also upregulated in the wild type during fruiting body development. We have cloned and sequenced a 3.2 kb fragment of genomic DNA containing the *tap1* open reading frame and adjoining sequences. The genomic region comprising *tap1* is syntenic to its homologous region in the closely related filamentous fungus *Neurospora crassa*. To determine whether *tap1* is involved in fruiting body development in *S. macrospora*, a knockout construct was generated in which the *tap1* open reading frame was replaced by the hygromycin B resistance gene *hph* under the control of fungal regulatory regions. Transformation of the *S. macrospora* wild type with this construct resulted in a *tap1* deletion strain where *tap1* had been replaced by the *hph* cassette. The knockout strain displayed no phenotypic differences under conditions of vegetative growth and sexual development when compared to the wild type. Double mutants carrying the $\Delta tap1$ allele in several developmental mutant backgrounds were phenotypically similar to the corresponding developmental mutant strains.

Conclusion: The *tap1* transcript is strongly upregulated during sexual development in *S. macrospora*; however, analysis of a *tap1* knockout strain shows that *tap1* is not essential for fruiting body formation in *S. macrospora*.

Background

Fruiting body formation in ascomycetes is a complex process leading to the formation of a number of specialized cell types from a comparatively undifferentiated veg-

etative mycelium [1]. Recently, the molecular basis of this process has been investigated by forward and reverse genetics approaches, and a number of genes that are essential for fruiting body development have been identi-




Figure 1
Transcript levels of *tap1* are upregulated during sexual development. *tap1* transcript levels in the wild type under conditions of vegetative and sexual development were analyzed by real time PCR. For normalization, transcript levels of the SSUrRNAs were used as described previously [3]. Expression is given as fold induction (mean of two independent experiments, error bars indicate standard deviation) with transcript levels during vegetative growth set to 1. Real time PCR results were tested for significance of differential expression at a level of $p = 0.001$ using REST [24].

fied. However, a coherent picture of fungal multicellular development has yet to emerge [2].

One avenue towards a deeper understanding of developmental processes is by functional genomics analyses, e.g. microarray studies. Such approaches can help to identify genes that are regulated differentially during fruiting body development and are therefore candidates for further functional analysis. In a previous study, we have performed microarray analyses of fruiting body development in *S. macrospora* [3]. This ascomycete is homothallic and produces fruiting bodies called perithecia within seven days under laboratory conditions. *S. macrospora* is a close relative of *N. crassa*, but in contrast to *N. crassa*, it does not produce any asexual spores. Therefore, changes of gene expression patterns during sexual development are not superimposed by changes related to asexual sporulation. We have previously analyzed gene expression in three developmental mutants of *S. macrospora*, and have identified a number of genes that are downregulated in the sterile mutants when compared with the wild type [3]. One of these genes is *tap1* (transcript associated with perithecial

development, formerly known as SMU5651 [3]). *tap1* transcript levels are downregulated in the three developmental mutants *pro1*, *pro11* and *pro22* as well as in all three double mutants which led us to speculate that the gene might be involved in sexual development in *S. macrospora*. In addition to this intriguing expression pattern, the derived TAP1 amino acid sequence shows homology to lectins from other filamentous fungi with the highest similarity to lectins isolated from fruiting bodies of several basidiomycetes [3]. It has long been speculated that lectins play a role in fungal development; however, as no mutants in lectin-encoding genes from fruiting body-producing fungi have been analyzed to date, definite proof for this hypothesis is lacking [4,5]. The only known fungal lectin mutant is a strain of *Arthrobotryx oligospora* in which the lectin gene *aol* was deleted [6]. This mutant does not exhibit any phenotypical differences from the wild type under all conditions investigated, but as no sexual cycle is known for *A. oligospora*, the question whether *aol* might be involved in fruiting body development could not be addressed. Here, we present data on the expression of the *S. macrospora tap1* gene as well as the characterization of a *tap1* knockout strain to address the role of a putative lectin in fungal sexual development.

Results

Expression of *tap1* during sexual development in *S. macrospora*

Previously, we reported that *tap1* is downregulated in several developmental mutants when compared with the wild type [3]. In that analysis, all strains were grown under conditions that allowed sexual development, i.e. in floating culture. To investigate whether *tap1* expression in the wild type is associated with sexual development, we compared transcript levels of *tap1* in sexually developing mycelia with vegetative mycelia. For this purpose, the wild type was grown in floating or in submerged culture where it develops either fruiting bodies or vegetative mycelium, respectively. Analysis of *tap1* expression by quantitative real time PCR revealed that *tap1* transcript levels are upregulated nearly 60-fold in mycelia undergoing sexual development (Figure 1). These data further confirm that *tap1* expression is linked with fruiting body development in *S. macrospora*.

Sequence analysis of the *S. macrospora tap1* gene

For previous expression analyses, part of the *tap1* open reading frame was amplified from *S. macrospora* genomic DNA by PCR [3]. For further characterization of *tap1*, we have isolated a *tap1*-containing cosmid clone from an indexed cosmid library of *S. macrospora* [7]. A 3.2 kb restriction fragment carrying the *tap1* open reading frame and adjacent regions was subcloned and sequenced. The fragment carries the uninterrupted *tap1* open reading frame of 429 bp encoding a predicted polypeptide of 143




Figure 2

Synteny between *S. macrospora* (*S.m.*) and *N. crassa* (*N.c.*) in the genomic region containing *tap1*. Coding regions are given as dark gray and intergenic regions as light gray boxes. Nucleic acid identities between the *S. macrospora* open reading frames and their *N. crassa* orthologues as well as between the intergenic sequences are indicated.

amino acids. In addition, the fragment contains part of another open reading frame downstream of *tap1* that was named *ibl1* (importin beta like 1) due to its similarity to an importin beta subunit. The sequenced region is syntenic to a region within the *N. crassa* genome that contains the open reading frames NCU05651.2 and NCU05650.2 that are orthologs of *tap1* and *ibl1* respectively (Figure 2). In a previous comparison of 85 protein-coding genes of *S. macrospora* and *N. crassa*, the average nucleic acid identity within coding regions was found to be close to 90 %, and even non-coding regions of these two closely related Pyrenomycetes can be readily aligned [8]. These findings are further supported by our analysis of the genomic region containing *tap1* (Figure 2).

The predicted TAP1 polypeptide was used for BLASTP [9] searches of the public databases, and a multiple alignment was constructed of TAP1 and several of its closest homologs from other fungi (Figure 3). The closest homolog is the predicted protein NCU05651.2 from *N. crassa*; however, interestingly, the second-best sequence identity is found in two lectins from the basidiomycetes *Xerocomus chrysenteron* and *Paxillus involutus*. Surprisingly, lectins and predicted lectin-like proteins from the ascomycetes *Podospora anserina*, *Fusarium graminearum*, and *A. oligospora* are less similar to the *S. macrospora* TAP1 even though these fungi are much more closely related to *S. macrospora* than the basidiomycetes (Figure 3). However, the (predicted) lectins from *P. anserina*, *F. graminearum*, and *A. oligospora* have a higher similarity compared to the

basidiomycete lectins than TAP1 does (Figure 3). These findings might indicate that TAP1 and its corresponding *N. crassa* ortholog, even though being clearly members of this fungal lectin family, have evolved rather fast compared to other ascomycete homologs.

Construction of a $\Delta tap1$ strain

A construct for generating a *tap1* deletion strain was generated. It consists of a hygromycin resistance cassette flanked by about 1 kb of sequences upstream and downstream of the *tap1* open reading frame (Figure 4). The construct was used to transform the *S. macrospora* wild type, and transformants were screened for homologous recombination by PCR with oligonucleotide pairs d1 and d2 as well as d3 and d4 (Figure 4). These primer combinations yield amplicons only in the case of homologous recombination. Among 50 primary transformants, one transformant was found that displayed the expected PCR fragments (data not shown). Ascospores from this transformant were isolated to purify the putative knockout strain, and Southern blot analysis of nine ascospore isolates was performed with probes for *tap1* and the Hygromycin resistance gene *hph*. For two of the ascospore isolates, T2.35 and T2.41, results are shown in Figure 5. As expected, the *tap1* probe hybridized to a 3.2 kb BglII fragment in genomic DNA of the wild type, whereas no hybridization signal was obtained with genomic DNA from the knockout strains T2.35 and T2.41. Conversely, the *hph* probe marked a 4.2 kb BglII fragment in the knockout strains and produced no signal with wild type




Figure 3
Multiple alignment of TAPI and lectins from filamentous fungi. The multiple alignment was created using CLUSTALX [25] with the following sequences: *S.m.*, *Sordaria macrospora* TAPI [this work, emb:CAH03681.2]; *N.c.*, *Neurospora crassa* NCU05651.2 [ref:XP_325506.1]; *X.c.*, *Xerococcus chrysenteneri* lectin [gb:AAL73236.1]; *P.i.*, *Paxillus involutus* LECA [gb:AAT91249.1]; *P.c.*, *Pleurotus cornucopiae* PCL-F2 [dbj:BAB63923.1]; *A.b.*, *Agaricus bisporus* ABL [sp:Q00022]; *P.a.*, *Podospora anserina* Pa5D0092 [emb:CAD60779.1]; *F.g.*, *Fusarium graminearum* FG07558.1 [gb:EAA76455.1]; *A.o.*, *Arthrotrichum oligospora* lectin [emb:CAA65781.1]. Jalview was used to visualize the alignment [26]. Amino acid residues conserved in at least eight of the nine sequences are given in dark blue, residues conserved in at least six sequences in medium blue, and residues present in at least four sequences in light blue. At the end of the alignment, amino acid identity in % is given for all sequences in pair-wise comparisons.




Figure 4
Strategy for the generation of a *tap1* knockout strain. The *tap1* open reading frame is shown in dark gray, flanking non-coding regions that are present in the knockout construct are given as white boxes, and adjoining regions not present in the knockout construct are shown in light gray. The Hygromycin B resistance cassette comprising the *hph* gene and the *trpC* promoter and 5' UTR is given in black. Positions of oligonucleotide primers d1 to d4 are indicated. For further information see text.

DNA. The 4.2 kb fragment is of the size expected in a strain where homologous recombination has taken place (Figure 4). The $\Delta tap1$ strains T2.35 and T2.41 were used for further analysis.

Phenotypic characterization of the $\Delta tap1$ strain in different genetic backgrounds

The *tap1* deletion strains T2.35 and T2.41 were analyzed with respect to their phenotype during the sexual cycle. Surprisingly, both were completely wild type-like in all aspects of fruiting body development, i.e. both *tap1* deletion strains produced mature perithecia in the same numbers and in the same amount of time as the wild type. Moreover, neither perithecial nor spore morphology were altered in any recognizable way (Figure 6). Fruiting body formation took place both in constant light as well as in constant darkness both on complete as well as minimal media, similar to that of the wild type. Ascospores from the knockout strains germinated readily and were black like wild type spores (Figure 6).

We then investigated whether there were any phenotypes unrelated to sexual development present in the mutant strains. Mycelial growth rates were determined as 24.7 (\pm

1.7) and 24.8 (\pm 1.4) mm per day for $\Delta tap1$ and wild type respectively; thus, indicating that mycelial growth is not altered in the knockout strains. Also, renewal of growth and fruiting body development after incubation at 4 °C or 37 °C, conditions that prevent growth and fruiting body formation, respectively, was not different in the mutants when compared to the wild type (data not shown). Also, wettability of mycelium and fruiting bodies was similar in wild type and mutants indicating that the hydrophobic coating of the mycelium was not altered in the knockout strains. As the lack of *tap1* did not cause any discernable phenotype, we wondered whether *tap1* might have a redundant function or whether its absence might be masked by the presence of other genes. We therefore crossed the $\Delta tap1$ allele into strains bearing mutations in other developmental genes, namely *pro1*, *pro11*, *pro22*, and *pro41*. The *pro1* mutant is defective in a gene encoding a transcription factor [10], the *pro11* mutant lacks a functional WD40 repeat protein [11], and *pro22* and *pro41* are mutants non-allelic to *tap1* or the other *pro* genes (Kück et al., unpublished data). *tap1* transcript levels are downregulated in mutants *pro1*, *pro11*, *pro22* [3], and *pro41* (Nowrousian, unpublished data). Thus, we speculated whether a complete lack of *tap1* would lead to





Figure 5

Southern blot analysis of $\Delta tap1$ strains. Genomic DNA from the wild type, from two different $\Delta tap1$ single spore isolates from the original knockout strain, and from double mutants of $\Delta tap1$ with pro1, pro41, pro11, and pro11 was hydrolyzed with *Bgl*II, and after gel electrophoresis, the Southern blot was probed with radio-labeled DNA fragments containing the open reading frames of *tap1* (A) and *hph* (B), respectively. Marker sizes in kb are provided on the right. Numbers in brackets are strain numbers for single spore isolates.

a more pronounced developmental phenotype, especially as several other developmental genes are also downregulated in the pro mutants [3]. We obtained double mutants with strains pro1, pro11, pro22, and pro41 and verified the presence of the $\Delta tap1$ allele in the double mutants by Southern blot analysis (Figure 5). All double mutant

strains were phenotypically similar to the respective single pro mutants in that they produced only protoperithecia and were therefore sterile. There were no differences in the number of protoperithecia produced by the double mutants when compared with the single mutants. Also, growth rates of the vegetative mycelium as well as overall

**Figure 6**

Δ tapI strains have a wild type-like phenotype. The wild type and Δ tapI single spore isolates T2.35 and T2.41 (as indicated above each column) were grown in Petri dishes on BMM solid medium [10, 21] for 6d (A) or 7d (B, C) at 25°C in constant light. A) Segments of petri dishes, black dots are individual fruiting bodies. B) Segments of ascus rosettes with mature (black-spored) and immature asci. Scale bar 100 μ m. C) Mature asci. Scale bar 20 μ m.

morphological appearance were the same (data not shown). For the crosses of the *tap1* knockout with the pro mutant strains and subsequent back-crosses, we analyzed a total of 17 full tetrads and 37 partial tetrads from crosses with different mutant strains and in all cases found the expected 1:1 segregation pattern for each of the single markers (hygromycin resistance and pro mutant phenotype, respectively). This is a further indication that deletion of *tap1* does not interfere with sexual development, and also shows that the process of generating the $\Delta tap1$ allele did not introduce further mutations into the strains that would cause any different segregation patterns. Overall, no phenotype for $\Delta tap1$ was found in any of the genetic backgrounds investigated.

Discussion

The *tap1*-encoded polypeptide from *S. macrospora* has significant homology to lectins and lectin-like protein from other fungi (Figure 3). The highest degree of amino acid identity is found in comparison with lectins that were isolated from fruiting bodies of several basidiomycetes [12-14]. Lectins are carbohydrate-binding proteins that are found in a variety of organisms [4,5]. On the basis of sequence homology, the *S. macrospora* TAP1 polypeptide can be included into a class of fungal lectins; however, whether it has lectin activity, i.e. whether it specifically binds carbohydrates, remains to be elucidated. Interestingly, TAP1 displays a greater sequence identity towards basidiomycete lectins than to lectins or putative lectins from ascomycetes with the (notable) exception of its closest homolog, the *N. crassa* protein NCU05651.2 (Figure 3). However, BLAST searches in the *N. crassa* genome [15] with the sequences of TAP1 as well as the lectin sequences from the other fungi used in our sequence comparisons yielded only NCU05651.2 as a significant result (data not shown). This finding indicates that the gene is present as a single copy in *N. crassa*, and that there is no other member of this lectin gene family present in the *N. crassa* genome. As *N. crassa* and *S. macrospora* are close relatives with highly syntenic genomes [8], it is likely that this is the case in *S. macrospora* as well. This observation is supported by the fact that only a single band in *S. macrospora* genomic DNA hybridizes with a *tap1* probe (Figure 5). Thus, it seems that this particular class of fungal lectins has evolved faster in *S. macrospora* and *N. crassa* compared to other ascomycetes, for which it still retains more similarity with its basidiomycete relatives (Figure 3). Another class of fungal lectins has been found in the basidiomycete *Coprinus cinereus*. These lectins bind galactose and are therefore called galectins, and two galectins from *C. cinereus* are specifically expressed during different stages of fruiting body formation [16,17]. However, BLAST searches for galectin homologs in the *N. crassa* genome yielded no significant results (data not shown) indicating that no galectin-like proteins exist in *N. crassa* or that their

sequences are too dissimilar to the *C. cinereus* sequences to be detected by sequence comparisons alone. Thus, the *N. crassa* NCU05651.2 gene and its *S. macrospora* ortholog *tap1* are so far the only genes encoding putative lectins that have been identified by sequence analysis in these two ascomycetes.

In fungi, most lectins have been isolated from basidiomycetes, especially from mushroom fruiting bodies, and it has been speculated that they play a role in fruiting body development [5,18,19]. However, as no lectin mutant in a fruiting body-producing fungus has been characterized to date, this hypothesis has not been verified experimentally. Previous investigations and the results presented here show that *tap1* transcript levels are closely correlated with fruiting body development in *S. macrospora*; therefore, we decided to construct a *tap1* knockout strain to analyze whether this putative lectin plays a role in fruiting body formation. A $\Delta tap1$ strain was generated by gene replacement, but the knockout strain has no discernable phenotype under all conditions investigated. This might indicate that *tap1* has indeed no function in vegetative growth or sexual development of *S. macrospora*; however, the possibility that *tap1* is needed under environmental conditions not tested in our experiments or that its function is redundant or that in the absence of *tap1*, another gene product can take its place, cannot be excluded. The latter effect is well known in other organisms, and it was, for example, tested in a large-scale analysis of yeast synthetic lethal interactions where it was found that genes involved in similar biological processes, but not necessarily in the same regulatory pathway, can buffer one another in single mutant backgrounds but show a phenotype in the double mutant strain [20]. To test whether any of the known developmental genes of *S. macrospora* show this kind of genetic interaction with *tap1*, we obtained double mutants of $\Delta tap1$ with strains bearing mutations in the developmental genes *pro1*, *pro11*, *pro22*, or *pro41*. However, all double mutant strains were phenotypically similar to the respective pro mutant strains. *tap1* transcript levels are downregulated in the pro mutants, and our analysis demonstrates that even the complete loss of *tap1* does not worsen the condition of the mutants. This leaves open the possibility that *tap1* is necessary in a different genetic background or under different environmental conditions.

With respect to fruiting body formation, our results show that the putative lectin-encoding gene *tap1* is not an essential gene for this developmental process. As mentioned previously, the only other known fungal lectin mutant is the *aol* mutant of the nematode-trapping fungus *A. oligospora* [6]. Similar to our findings, the *aol* mutant also has no phenotype under all conditions investigated. Since no sexual stages from *A. oligospora* are known, these obser-

Table 1: *S. macrospora* strains used in this study. All strains are single spore isolates and are kept in our laboratory collection. The *fus* allele that is present in some of the strains is a spore color marker (brown instead of black spores) but has no influence on growth or fertility.

| strain number | genotype | description |
|---------------|---------------------------|----------------------|
| S48977 | wild type | wild type |
| T2.35 | Δ tap1 | tap1 deletion strain |
| T2.41 | Δ tap1 | tap1 deletion strain |
| M8871 | pro1 | sterile mutant [10] |
| S24117 | pro11 | sterile mutant [11] |
| S22528 | pro22 | sterile mutant |
| S46357 | pro41 | sterile mutant |
| S62222 | Δ tap1, pro1 | sterile mutant |
| S62355 | Δ tap1, pro1 | sterile mutant |
| S62497 | Δ tap1, pro41 | sterile mutant |
| S63491 | Δ tap1, pro11, fus | sterile mutant |
| S63433 | Δ tap1, pro11, fus | sterile mutant |
| S62292 | Δ tap1, pro22, fus | sterile mutant |
| S62456 | Δ tap1, pro22, fus | sterile mutant |

vations do not include fruiting body formation; however, vegetative growth, conidiation, and nematode-trapping were unchanged in the aol mutant strain [6]. Thus, possible functions of this class of lectins and lectin-like proteins in filamentous fungi remain enigmatic.

Conclusion

tap1 expression is strongly associated with sexual development in *S. macrospora*. An analysis of the *tap1* gene and its surrounding genomic region revealed a high degree of sequence identity and overall synteny with the corresponding region in the genome of *N. crassa*. Sequence comparisons of TAP1 with lectins and lectin-like proteins from other fungi indicate that it is most closely related to lectins isolated from basidiomycete fruiting bodies. However, analysis of a *tap1* knockout strain shows that *tap1* is not essential for fruiting body formation nor vegetative growth in *S. macrospora*. This is the case for a Δ tap1 allele in an otherwise wild type genetic background as well as in combination with mutations in several developmental genes. Whether *tap1* has any function under growth conditions not investigated here, e.g. in a more natural setting, remains to be elucidated.

Methods

Strains and growth conditions

Details of the *S. macrospora* strains used in this study are provided in Table 1. Double mutants were obtained from crosses of single mutant strains. Double mutant genotypes were verified by crosses against single mutants. Unless stated otherwise, standard growth conditions and DNA-mediated transformation were as previously described [10,21]. For analysis of growth velocity, 30 cm long race tubes were filled with 15 ml of medium, inoculated at one

end and the growth front was marked every 24 h for 7 consecutive days. For RNA extraction from cultures developing fruiting bodies, *S. macrospora* was grown at 25 °C in constant light in floating culture as described [3]. For RNA extraction from vegetative mycelium, a mycelial plug of 0.7 cm in diameter from a Petri dish with liquid medium [3] was inoculated into an Erlenmeyer flask with 100 ml of liquid medium and shaken at 130 rpm.

RNA extraction and quantitative real time PCR

Extraction of total RNA and quantitative real time PCR were performed as described previously [3] with the following modifications: reverse transcription was performed with 400 U Superscript II (Invitrogen) and 0.33 mM dNTPs, and real time PCR was carried out in a DNA Engine Opticon 2 (MJ Research).

Identification of a cosmid clone carrying tap1 and analysis of the tap1 gene

An indexed *S. macrospora* cosmid library [7] was screened for *tap1* by PCR with oligonucleotides SMU5651-1 (5' CATCAACGACACCTCCGACACCC) and SMU5651-2 (5' CATCGGCCTGATAGAAGCTTGATCC). For a first round of screening, pooled DNA from 48 cosmid clones was used as a template, DNA from clones from positive pools was then subpooled and used for the next round of screening. This led to the isolation of cosmid D3 from pool VI518-614 that contains the *tap1* gene. A 3.2 kb BglII restriction fragment carrying *tap1* was subcloned from cosmid D3 into pBluescript II/KS+ (Stratagene). The insert of the resulting vector pPC24 was sequenced at MWG Biotech or GATC Biotech AG [emb:AJ781427.2].

Construction of a Δ tap1 strain

To create a *tap1* knockout construct for homologous recombination in *S. macrospora*, flanking regions upstream and downstream of the *tap1* open reading frame were amplified by PCR from *S. macrospora* genomic DNA using oligonucleotides SMU5651-BamHI (5' AGGATC-CGTGATTCTCATGCTGTGGAAGGAAGC) and SMU5651-NheI (5' AGCTAGCTTTGCGGTTTGGTT-GGGGGTTGGT) for the upstream region and oligonucleotides SMU5651-ApaI (5' AGGGCCCGTACTCGTCAGTGGGAAAGTGGGTGG) and SMU5651-SacI (5' AGAGCTCTATGCACTTGCTCCT-CAAGCGTCTC) for the downstream region, introducing restriction sites as indicated in the oligonucleotide names. Additionally, the hygromycin-resistance cassette consisting of the *hph* gene from *Escherichia coli* and the *trpC* promoter from *Aspergillus nidulans* was amplified from plasmid pCB1004 [22] using oligonucleotides Hph-ApaI (5' AGGGCCCTCAACGGAACCCTATTCCTTTGCC) and Hph-NheI (5' AGCTAGCAACTGATATTGAAGGAG-CATTTTGG). All PCR fragments were subcloned in pDrive (Qiagen) resulting in pDriveA (upstream region),

pDriveB (downstream region) and pDrivehph (hygromycin-resistance cassette). Sequences of inserts and orientation within the vector was verified by restriction analysis and sequencing (MWG Biotech AG). The PCR fragment containing the downstream region was obtained by ApaI/SacI digestion of pDriveB and cloned into ApaI/SacI digested vector pDriveA resulting in plasmid pDriveAB. The hygromycin-resistance cassette was obtained by digesting plasmid pDrivehph with NheI and ApaI and was cloned into plasmid pDriveAB hydrolyzed with NheI and ApaI resulting in the knockout plasmid pABXY. For transformation of *S. macrospora*, plasmid pABXY was digested with BamHI and SacI and the knockout cassette was obtained by gel elution. The knockout cassette was transformed into the *S. macrospora* wild type and primary transformants were screened for homologous integration by PCR. For this purpose, total DNA was prepared from the transformants according to [23], and PCR was performed with oligonucleotides d1 (5' CGATGGCTGTGA-GAAGTACTCGC) and d2 (5' TGCCTCCTCCGAGGCTGATAACCT) for the downstream region or d3 (5' CCGTGGGTAAGGTATCTCTGATG) and d4 (5' CACCGCCTGGACGACTAAACCAA) for the upstream region (Figure 4).

Authors' contributions

MN designed the study, performed expression analyses and isolation of the cosmid clone as well as part of the characterization of the knockout strain and wrote the manuscript. PC cloned and sequenced the *tap1* gene, made the knockout construct and the *tap1* knockout strain and carried out part of the characterization of the knockout strain.

Acknowledgements

The authors would like to thank Swenja Ellßel and Susanne Schlewinski for excellent technical assistance and Prof. Dr. Ulrich Kück for his generous support that enabled us to carry out this study as part of project A1 of research initiative SFB 480 funded by the Deutsche Forschungsgemeinschaft (DFG).

References

- Bistis GN, Perkins DD, Read ND: Different cell types in *Neurospora crassa*. *Fungal Genet Biol* 2003, **50**:17-19.
- Pöggeler S, Nowrousian M, Kück U: **Fruiting body development in ascomycetes**. In *The Mycota* / Edited by: Kües U, Fischer R. Berlin, Heidelberg: Springer; 2005 in press.
- Nowrousian M, Ringelberg C, Dunlap JC, Loros JJ, Kück U: **Cross-species microarray hybridization to identify developmentally regulated genes in the filamentous fungus *Sordaria macrospora***. *Mol Genet Genomics* 2005, **273**:137-149.
- Wang H, Ng TB, Ooi VEC: **Lectins from mushrooms**. *Mycol Res* 1998, **102**:897-906.
- Guillot J, Kanska G: **Lectins in higher fungi**. *Biochem Syst Ecol* 1997, **25**:203-230.
- Balogh J, Tunlid A, Rosen S: **Deletion of a lectin gene does not affect the phenotype of the nematode-trapping fungus *Arthrobotrys oligospora***. *Fungal Genet Biol* 2003, **39**:128-135.
- Pöggeler S, Nowrousian M, Jacobsen S, Kück U: **An efficient procedure to isolate fungal genes from an indexed cosmid library**. *J Microbiol Meth* 1997, **29**:49-61.
- Nowrousian M, Würtz C, Pöggeler S, Kück U: **Comparative sequence analysis of *Sordaria macrospora* and *Neurospora crassa* as a means to improve genome annotation**. *Fungal Genet Biol* 2004, **41**:285-292.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: **Gapped BLAST and PSI-BLAST: a new generation of protein database search programs**. *Nucleic Acids Res* 1997, **25**:3389-3402.
- Masloff S, Pöggeler S, Kück U: **The *proI+* gene from *Sordaria macrospora* encodes a C₆ zinc finger transcription factor required for fruiting body development**. *Genetics* 1999, **152**:191-199.
- Pöggeler S, Kück U: **A WD40 repeat protein regulates fungal cell differentiation and can be replaced functionally by the mammalian homologue striatin**. *Eukaryot Cell* 2004, **3**:232-240.
- Iijima N, Yoshino H, Ten LC, Ando A, Watanabe K, Nagata Y: **Two genes encoding fruit body lectins of *Pleurotus cornucopiae*: sequence similarity with the lectin of a nematode-trapping fungus**. *Biosci Biotechnol Biochem* 2002, **66**:2083-2089.
- Crenshaw R, Harper SN, Moyer M, Privalle LS: **Isolation and characterization of a cDNA clone encoding a lectin gene from *Agaricus bisporus***. *Plant Physiol* 1995, **107**:1465-1466.
- Trigueros V, Lougarre A, Ali-Ahmed D, Rahbé Y, Guillot J, Chavant L, Fournier D, Paquereau L: **Xerocomus chryseron lectin: identification of a new pesticidal protein**. *Biochim Biophys Acta* 2003, **1621**:2992-2298.
- Galagan J, Calvo S, Borkovich K, Selker E, Read N, Jaffe D, FitzHugh W, Ma L, Smirnov S, Purcell S, et al.: **The genome sequence of the filamentous fungus *Neurospora crassa***. *Nature* 2003, **422**:859-868.
- Boulianne RP, Liu Y, Aebi M, Lu BC, Kües U: **Fruiting body development in *Coprinus cinereus*: regulated expression of two galectins secreted by a non-classical pathway**. *Microbiology* 2000, **146**:1841-1853.
- Cooper DN, Boulianne RP, Charlton S, Farrell EM, Sucher A, Lu BC: **Fungal galectins, sequence and specificity of two isolectins from *Coprinus cinereus***. *J Biol Chem* 1997, **272**:1514-1521.
- Oguri S, Ando A, Nagata Y: **A novel developmental stage-specific lectin of the basidiomycete *Pleurotus cornucopiae***. *J Bacteriol* 1996, **178**:5692-5698.
- Kaneko T, Oguri S, Kato S, Nagata Y: **Developmental appearance of lectin during fruit body formation in *Pleurotus cornucopiae***. *J Gen Appl Microbiol* 1993, **39**:93-100.
- Tong AH, Lesage G, Bader GD, Ding H, Xu H, Xin X, Young J, Berriz GF, Brost RL, Chang M, et al.: **Global mapping of the yeast genetic interaction network**. *Science* 2004, **303**:808-813.
- Nowrousian M, Masloff S, Pöggeler S, Kück U: **Cell differentiation during sexual development of the fungus *Sordaria macrospora* requires ATP citrate lyase activity**. *Mol Cell Biol* 1999, **19**:450-460.
- Carroll AM, Sweigard JA, Valent B: **Improved vectors for selecting resistance to Hygromycin**. *Fungal Genet Newsl* 1994, **41**:22.
- Lecellier G, Silar P: **Rapid method for nucleic acid extraction from petri dish-grown mycelia**. *Curr Genet* 1994, **25**:122-123.
- Pfaffl MW, Horgan GW, Dempfle L: **Relative expression software tool (REST) for group-wise comparison and statistical analysis of relative expression results in real-time PCR**. *Nucleic Acids Res* 2002, **30**:e36.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG: **The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools**. *Nucleic Acids Res* 1997, **24**:4876-4882.
- Clamp M, Cuff J, Searle SM, Barton GJ: **The Jalview Java alignment editor**. *Bioinformatics* 2004, **12**:426-427.