

Research article

Open Access

The *Francisella* pathogenicity island protein IgIA localizes to the bacterial cytoplasm and is needed for intracellular growth

Olle M de Bruin, Jagjit S Ludu and Francis E Nano*

Address: The Department of Biochemistry and Microbiology, University of Victoria, Victoria, B.C., Canada

Email: Olle M de Bruin - debruin@uvic.ca; Jagjit S Ludu - jludu@uvic.ca; Francis E Nano* - fnano@uvic.ca

* Corresponding author

Published: 17 January 2007

Received: 11 August 2006

BMC Microbiology 2007, 7:1 doi:10.1186/1471-2180-7-1

Accepted: 17 January 2007

This article is available from: <http://www.biomedcentral.com/1471-2180/7/1>

© 2007 de Bruin et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: *Francisella tularensis* is a gram negative, facultative intracellular bacterium that is the etiological agent of tularemia. *F. novicida* is closely related to *F. tularensis* but has low virulence for humans while being highly virulent in mice. IgIA is a 21 kDa protein encoded by a gene that is part of an *iglABCD* operon located on the *Francisella* pathogenicity island (FPI).

Results: Bioinformatics analysis of the FPI suggests that IgIA and IgIB are components of a newly described type VI secretion system. In this study, we showed that IgIA regulation is controlled by the global regulators MglA and MglB. During intracellular growth IgIA production reaches a maximum at about 10 hours post infection. Biochemical fractionation showed that IgIA is a soluble cytoplasmic protein and immunoprecipitation experiments demonstrate that it interacts with the downstream-encoded IgIB. When the *igIB* gene was disrupted IgIA could not be detected in cell extracts of *F. novicida*, although IgIC could be detected. We further demonstrated that IgIA is needed for intracellular growth of *F. novicida*. A non-polar *igIA* deletion mutant was defective for growth in mouse macrophage-like cells, and *in cis* complementation largely restored the wild type macrophage growth phenotype.

Conclusion: The results of this study demonstrate that IgIA and IgIB are interacting cytoplasmic proteins that are required for intramacrophage growth. The significance of the interaction may be to secrete effector molecules that affect host cell processes.

Background

Francisella tularensis is the etiological agent of the severe, febrile disease tularemia. Although there have been rare isolates of *F. tularensis* in Australia, tularemia is mainly a disease of the Northern hemisphere that is spread by blood-sucking mosquitoes, flies, and ticks or acquired from contact with infected animals such as rabbits, rodents, and beavers [1]. Occasionally, local outbreaks of tularemia are associated with contact or consumption of contaminated natural water. In addition, *F. tularensis* is potentially a threat as a bioterrorist agent due to its high

infectivity and lethality when inhaled. *F. novicida* is highly related at the DNA level to *F. tularensis*, and serves as a model organism since it is very virulent in mice while being avirulent in humans.

F. tularensis is a gram-negative, facultative intracellular bacterium capable of survival and replication in macrophages [2]. A common virulence strategy of intracellular pathogens is to favorably modulate the intracellular milieu of hosts for their own benefit. In *Legionella pneumophila* a type IV secretion system (T4SS) delivers effectors

that allow the pathogen to replicate in ribosome-studded phagosomes that fail to fuse with lysosomes [3,4]. *Salmonella enterica* relies on a pathogenicity island-encoded type III secretion system (TTSS) to modify phagosome biogenesis [5,6], including inhibition of phago-lysosomal fusion [7] and the NADPH oxidase-mediated killing by host cells [5]. Other intracellular pathogens, such as *Listeria monocytogenes*, degrade the phagosomal membrane and escape into the cytoplasm to replicate freely [8]. *F. tularensis* initially resides in a phagosome which accumulates some late endosome markers. After about four hours most *F. tularensis* cells escape the phagosome and grow in the cytoplasm. [2,9-11]. Although an intact *iglC* gene is needed for *F. tularensis* to escape phagosomes, the role of *IglC* is unknown.

We recently described a *Francisella* pathogenicity island (FPI) harboring several genes necessary for intracellular growth. Four FPI genes, *iglABCD*, are organized in an apparent operon [12]. The production of *IglC* mRNA is in part dependent on *MglA* [13] which is thought to be a global regulator of virulence factors in *F. tularensis*. By analogy with its *Escherichia coli* homologue, *SspA*, *MglA* likely interacts with RNA polymerase to directly or indirectly alter transcription of several genes [14]. Disruption of *mglA* or *mglB* results in mutants that are severely attenuated for virulence [15]. *IglC* has been shown to be induced about four-fold during intracellular growth relative to broth growth and necessary for virulence [16-18], and it was recently demonstrated that inactivation of *iglC* and *mglA* result in mutants that remain in phagosomes that fuse with lysosomes [19,20]. Although an *iglA* transposon insertion mutant has been shown to be defective for intracellular growth, it could not be ruled out that the observed phenotype was due to interruption of transcription of downstream genes, including *iglC* [17].

In this study, we use *F. novicida* to investigate the properties of *IglA* and its role in *F. novicida* intracellular growth. *F. novicida* is particularly suited for these studies since, unlike *F. tularensis*, it contains only one copy of the FPI, and this simplifies the construction of mutants. Further, the biology of *F. novicida* growth in human macrophages is indistinguishable from that of *F. tularensis* strains [9,11], and thus *F. novicida* serves as a valid surrogate for virulent strains when studying basic aspects of *Francisella* intracellular growth. In this work we supply evidence that *IglA* is a cytoplasmic protein that interacts with *IglB*, and is required for intramacrophage growth.

Results

***IglAB* homologues in diverse bacteria are organized in a conserved gene cluster**

Homologues of *iglA* and *iglB* exist in several bacterial species that are either animal or plant pathogens or plant

symbionts [12] but there are no known homologues of *iglC* or *iglD*. *IglAB* homologues in *Vibrio cholerae*, *Salmonella enterica*, *Rhizobium leguminosarum*, and other bacteria are found in a cluster of genes encoding proteins known as IcmF-associated homologous proteins (IAHPs) [21-23]. Recently, it was demonstrated that this gene cluster encodes components of a proposed type VI secretion system (T6SS) in *Vibrio cholerae* [24].

In light of the emerging role of IAHP/T6SS in the secretion of proteins we re-examined the ORFs in the FPI to determine if components of a type VI secretion system may be present. Three essential components of a T6SS are a protein with an IcmF-motif and two linked genes that correspond to *iglA* and *iglB*. A BLASTP search revealed that an IcmF region was found as part of the C-terminal third of *PdpB* which aligned with the corresponding regions of proteins belonging to the IcmF conserved orthologous group (COG3523.2 with an E-value of 7×10^{-9}). The identification of *IglA* and *IglB* as members of COGs is much clearer. *IglA* has strong identity to members of COG3516 (E-value of 2×10^{-20}) and *IglB* has strong identity with COG3517 (E-value of 2×10^{-102}). Remarkably all of the relatives of *iglAB* are organized in the same order, and are always adjacent to each other on the chromosome. The *iglAB* genes together with an *icmF*-containing gene form the core set of genes that suggest the presence of a type VI secretion system. We also found through BLASTP analysis that the deduced product of an ORF 380 bp downstream of *pdpB* (shown as "vgr" in Fig. 1) shows a weak similarity (E-value 0.15) to the family of *vgr*-encoded proteins, such as *VgrG* [24] which is secreted by a T6SS in *V. cholerae*. *Vgr* proteins are hydrophilic proteins that contain valine-glycine repeats, and are found in a number of gram negative pathogens. Another ORF, 4587 bp downstream of *pdpB* show similarity (E-value, 0.0005) to proteins in COG3455 that includes the IAHP-associated protein *DotU*. The clustering of *iglAB* and the *icmF*-containing *pdpB* gene, together with two other IAHP-associated genes strongly suggests that the FPI carries a type VI secretion system.

***IglA* expression in an *mglAB* background**

Previously RT-PCR analysis of the level of *iglA*, *iglC* and *iglD* transcripts revealed a role of *MglA* in regulating expression of the *iglABCD* operon mRNA production [13]. We wished to test if *IglA* protein expression levels are depressed in mutant *mglA* and *mglB* backgrounds. Western immunoblot analysis of *IglA* in an *mglA* mutant and an *mglB* background revealed that *IglA* is not expressed at detectable levels in these strains (Fig. 2).

***IglA* expression during intramacrophage growth**

Previous studies provide evidence that *MglA* expression peaks at about 5 hours after infection of macrophages




Figure 1
Similarity of the FPI to other virulence gene clusters. Homologues of *Francisella* pathogenicity island proteins IglA and IglB are found on a conserved gene clusters known as IcmF associated homologous proteins (IAHP), which, in some cases encode a proposed type VI secretion system. In the FPI the IcmF motif appears at the C-terminus of PdpB. Downstream of the *pdpB* gene is an ORF designated "vgr" that encodes a protein with similarity the Vgr family, one of which is secreted by the proteins encoded by an IAHP cluster in *Vibrio cholerae*. Homologues of the *Legionella dotU* gene are often associated with IAHP clusters. A very weak similarity to *dotU* is seen in an ORF that is sixth downstream of *pdpB*.

[25], and that IglC expression is maximal at between 6 and 24 hours after infection [16]. To access the pattern of IglA expression during *F. novicida* infection of macrophages, we lysed J774 macrophages at various time points after infection with the wild type strain U112 and examined the lysates for IglA using immunoblotting. In our assays IglA was first detectable at 8 hours post-infection, peaked at 10 hours, and showed a decline by 12 hours (Fig. 3). In broth grown cultures IglA appeared to be maximally expressed at the late logarithmic phase of growth.

IglA is cytoplasmically located

Knowing the cellular localization of a protein can help lead to a hypothesis as to its biological role. To investigate the subcellular localization of IglA, we fractionated *F. novicida* U112 into soluble and membrane-associated fractions and determined the amount of IglA in each fraction by immunoblot analysis. The data from this experiment

revealed that IglA is exclusively a soluble protein (Fig 4). Although IglA lacks a signal peptide sequence, it could not be ruled out that IglA localizes to the periplasm by a novel mechanism. Therefore, we isolated the periplasmic contents from *F. novicida* and determined by immunoblotting that IglA does not localize to this compartment. We also failed to detect IglA in culture supernatant (data not shown). The data from these experiments strongly suggest that IglA is a cytoplasmic protein. In agreement with this, the IglA homologue in *Salmonella enterica* has been predicted to be localized to the cytoplasm [23].

IglA interacts with IglB in vivo

To investigate interactions of IglA with other *F. novicida* proteins we performed immunoprecipitations with anti-IglA antibody on soluble proteins. A co-precipitating protein with a relative molecular mass of approximately 60 was detected (Fig 5A). This protein band was excised and subjected to MALDI-TOF analysis, and the resulting peptide fragment masses were submitted to searches against predicted peptide fragments of prokaryotes in the MASCOT data bank. This analysis revealed that the only significant match was IglB from *F. novicida* (Fig. 5B). The relative molecular mass of the co-precipitated protein is consistent with this result as IglB is predicted to be 58 kDa. Immunoprecipitations performed with an *iglA* null strain did not result in the appearance of the 60 kDa band, nor did immunoprecipitations of U112 done with pre-immune serum. These results strongly suggest that IglA and IglB interact in the cytoplasm of *F. novicida*.



Figure 2
IglA regulation by MglA and MglB. Western blot showing lack of IglA in *mglA* and *mglB* mutants but present in the wild type strain U112. All samples were normalized to 6 µg protein per lane.

Supporting the hypothesis that IglA interacts with IglB is the finding that IglB mutants but not IglC mutants lack detectable IglA (see below, Fig. 8). Presumably a lack of




Figure 3
IgA expression in J774 macrophages. Western blot showing expression of IgA during infection of macrophages. J774 macrophages were infected with parent strain U112 (m.o.i 300:1) and lysed at the indicated time post infection. Loading was normalized according to the number of viable bacteria (CFU) in each sample as determined by plating on TSA-C plates. Lane J774, uninfected macrophages. TSB, broth grown U112 grown to indicated optical density (600 nm). All samples were normalized to 10⁷ CFU by viable counts. The macrophage cell lysates altered the appearance of the IgA bands, but control experiments showed that the cell lysates did not mask IgA reactivity with antibody.

association of IgA with IgB makes the former susceptible to degradation.

Deletion mutagenesis of *iglA* and complementation of the mutant strain

An *iglA* deletion mutant, ODB2, was constructed using a two-step integration-excision method (Fig. 6A). First, the PCR-amplified 1.5 kbp regions flanking *iglA* were joined so as to leave *iglB* intact, including its ribosome binding region. This recombinant construct was ligated to an




Figure 4
Subcellular localization of IgA. Anti-IgA was used to probe Western immunoblot of subcellular fractions of *F. novicida*. The sarkosyl insoluble fraction represents an enrichment of outer membrane protein and the sarkosyl soluble fraction contains largely inner membrane protein. Samples were prepared as outlined in Methods and normalized to 10 µg protein per lane before separation on a 12% SDS-PAGE gel. Results are representative of three independent experiments.




Figure 5
Co-immunoprecipitation of a 60 kDa protein with IgA. Panel A. Anti-IgA serum co-immunoprecipitates a circa 60 kDa soluble protein (arrow, lanes 1 and 4). The band is absent in control reactions with non-specific antibody (lane 3) and in immunoprecipitations with an *iglA* mutant (lane 2). Numbers shown indicate molecular mass standards. Results are representative of those of three experiments. Panel B. MALDI-TOF identified the 60 kDa protein as IgB. Underlined sequences indicate peptides identified by MALDI-TOF. The second and third regions each represent two peptides (break after the "R"). Of 25 queries submitted, 9 showed significant identity with rabbit heavy chain and 9 showed significant identity with IgB of *F. novicida*. No other significant hits were found in the MSDB 20060224 databank.

erythromycin resistance-*sacB* cassette and the ligation mixture was used to chemically transform *F. novicida* JLO to erythromycin resistance. The JLO strain is a derivative of U112 that has a deletion in one of its putative sucrose hydrolase genes, and is thus sensitive to sucrose when *sacB* is expressed. This strain behaves like wild type in our virulence assays (data not shown). An erythromycin resistant colony was grown and plated on agar media containing 10% sucrose which acts as a counter selective marker for the *sacB* gene. Sucrose sensitive strains were examined for loss of *iglA* by PCR (Fig. 6B). Attempts to genetically complement the Δ *iglA* strain by incorporating *iglA* into a *F. tularensis* plasmid pFNLT1 [26] failed, presumably because the over-expression of IgA was lethal to *F. novic-*

ida. Hence, an *in cis* complementation approach was devised, allowing *iglA* to be incorporated into the chromosome linked to a kanamycin resistance marker (Fig. 7A and 7B). The *iglA* deletion strain failed to produce IglA as determined by Western immunoblotting (Fig. 8). However, the Δ *iglA* strain retained expression of IglC at parental strain levels. *In cis* complementation of the Δ *iglA* strain resulted in a strain that regained partial expression of IglA. An insertion mutant of *iglB* gave a reduction in the amount of IglC that was made, and this is not surprising since many insertion mutation decrease the expression of downstream genes. Surprisingly, this same mutant lacked expression of IglA, suggesting that the co-expression of IglB is needed for expression of IglA or to prevent degradation of IglA. Disruption of *iglC* however, does not affect the amount of IglA detected (Fig. 8).

IglA is required for growth in the J774 macrophage cell line

Previous work has suggested that IglA is required for *F. novicida* intramacrophage growth and virulence; however, its role has never been unequivocally demonstrated. In order to assess the requirement for IglA expression in intramacrophage growth we used our defined deletion and complemented strains to infect a culture of the J774 macrophage cell line. The data shown in figure 9 illustrates that the Δ *iglA* strain is incapable of intramacrophage growth, as is the *iglC* negative strain, CG62. The Δ *iglA* strain that was complemented for IglA production partially regained its ability to grow in macrophages. The residual defect in intracellular growth is not unexpected since we showed that the expression of IglA was not at wild type levels. Δ *iglA* replicated as the parental strain in broth (data not shown).

The Δ *iglA* strain has lowered virulence in chicken embryos

When the Δ *iglA* strain was used to infect chicken embryos it caused low mortality when compared to wild type *F. novicida* (Fig. 10). The wild type strain of *F. novicida* caused 100% mortality at day 5 post infection at an infecting dose of 600 CFU, whereas the Δ *iglA* strain caused only 14% mortality at day 6 with an infecting dose of 4,500 CFU (Fig. 10) or 50% mortality at day 6 with an infecting dose of 45,000 CFU (data not shown).

Discussion

There is growing evidence that the *iglABCD* operon is needed for *F. tularensis* intracellular growth and virulence and that the MglAB proteins are involved in regulating the expression of *iglABCD*. However, there is very little genetic and corresponding biochemical data demonstrating the roles of MglAB and IglAB and their corresponding homologues in other bacteria. For example, while it is clear that MglA plays a role in regulating the amount of *iglABCD* transcript it is unclear if the role precisely corresponds to that of the *E. coli* SspA protein. The data that exists for the




Figure 6
Deletion mutagenesis of *iglA*. Panel A. Diagram of steps used to construct an *iglA* deletion mutant. A fragment of *pdpD* was joined to *iglB* and these two fragments were ligated to an Em^R -*sacB* cassette. After transformation the recombinant construct integrated into the *F. novicida* chromosome. Plating the strain with the integrated fragment on sucrose selected for strains that had undergone an excision of the *sacB* and neighboring regions. Panel B. PCR confirmation of the deletion of *iglA*. The small arrows indicate the location of the primers used in the reactions.

functioning of SspA suggest that much of the regulation of stationary phase proteins occurs indirectly via the repression of H-NS, and that some of the effect of SspA is post-transcriptional [14].

There is also growing evidence that proteins encoded by IAHP clusters, of which IglAB homologues are important




Figure 7
In cis complementation of *iglA*. Panel A. Diagram of complementation scheme. A PCR amplicon containing the *iglA* and neighboring regions was ligated to a Km^R cassette and used to transform a \DeltaiglA strain. Integration of the recombinant construct resulted in a strain with a chromosomally-integrated *iglA*. Panel B. PCR reactions demonstrating the presence of *iglA* in the complemented strain. Arrows in lower part of diagram indicate the location of the PCR primers used in the reactions.

components, are involved in secretion of proteins from gram-negative bacteria [24,27]. There are approximately 30 homologues of *iglAB* and in every case the two genes are adjacent to each other and arranged in the same gene order. In this work we provided biochemical evidence that




Figure 8
An *iglA* mutant lacks the expression of a 21 kDa protein. Western blot showing the lack of an anti-IgIA serum reactive 21 kDa protein in the \DeltaiglA strain (top panel). Wild type levels of IgIC are retained in the \DeltaiglA strain (bottom panel). In contrast, the *iglA::Em* mutant lacks expression of IgIC. The expression of IgIC is threefold lower in an *iglB::Em* strain than in JLO and \DeltaiglA . Fluorescence intensity was used to quantify relative amounts of protein.

the IglAB proteins physically associate with each other and are localized to the cytoplasm. The surprising finding that inactivation of the *iglB* gene results in the disappearance of the IglA protein suggest that the presence of IglB is required for IglA to be stable.

IglA was first identified as a locus that when inactivated by a transposon insertion rendered *F. novicida* defective for growth in macrophages [17]. However, it could not be ruled out that the effect was due to interruption of transcription of downstream genes. In this report, we provide strong evidence that IglA is necessary for intracellular growth as a non-polar *iglA* deletion mutant was defective for growth in a mouse macrophage-like cell line. *In cis* complementation of the \DeltaiglA strain restored intramacrophage growth although the growth was slower than in the wild type strain. The *in cis* complementation strategy created two *iglA* promoter regions on the chromosome, one on either side of a kanamycin resistance cassette. It is conceivable that this results in aberrant regulation of *iglA* expression, which could explain why the growth of the complementation strain lags early during infection. We were unable to complement the *iglA* deletion mutant *in trans* with pFNLTP1::*iglA*, a high copy-derivative of an endogenous *Francisella* plasmid. Presumably, over-expression of IglA was lethal to *F. novicida*.

We hypothesize that IglA and IglB are cytoplasmic, chaperone-like proteins that are involved in secretion of viru-




Figure 9
IglA is required for intracellular growth. Growth of \DeltaiglA strain in J774 mouse macrophage-like cells. Filled squares, parental strain JLO; open squares, \DeltaiglA ; triangles, *in cis* complementation strain; diamonds, *iglC* transposon insertion mutant CG62. The experiments were done in triplicate and standard errors are shown by bars. This graph shows data from one of three independent experiments.

lence factors. Therefore, the biological significance of IglAB interaction may be to secrete *Francisella* effector molecules. In other pathogens, secretion of virulence proteins often requires interaction between two cytoplasmic proteins. For example, in *Yersinia pestis*, a complex composed of SycN and YscB function as chaperones for YopN [28], which is secreted to the cell surface [29]. Also, interaction of IcmS and IcmW is required for translocation of effector proteins via the Dot/Icm complex during *Legionella pneumophila* intracellular growth [30,31]. Hager *et al.* recently demonstrated protein secretion by *F. novicida* [32]. We did not observe any difference in secreted peptides between broth-grown wild type *F. novicida* and the \DeltaiglA strain by SDS-PAGE electrophoresis (data not shown). This observation is not surprising given the fact it has been demonstrated that secretion involving IAHPs is a highly regulated or an *in vivo*-induced process [27].

In summary, our results suggest that IglA and IglB are interacting cytoplasmic proteins that are required for intramacrophage growth. The significance of the interaction may be to secrete effector molecules that affect host cell processes.

Conclusion

The *Francisella* Pathogenicity Island harbors uncharacterized genes implicated in virulence. By constructing an in-frame deletion mutant we have shown that the FPI gene *iglA* is needed for intramacrophage growth. Biochemical characterization of IglA strongly suggests that it is a cytoplasmic protein that interacts physically with IglB. In addition, we provide data that show IglA is induced dur-




Figure 10
 \DeltaiglA mutant is less virulent in chicken embryos. Infection of chicken embryos with 600 CFU of wild type (JL0) *F. novicida* lead to death of 7/7 embryos in 5 days (Panel A), whereas infection with 4,500 CFU of the \DeltaiglA strain (ODB2) lead to the death of 1/7 embryos in 6 days (Panel B).

ing infection of macrophages. Bioinformatics analysis of the FPI suggests that it is similar to virulence loci that encode a protein secretion apparatus. We propose that IglA and IglB are chaperone-like proteins that are part of a secretion system in *F. novicida*.

Methods

Bacterial strains and culture conditions

All strains used in this work are listed in Table 1. *F. novicida* strains were grown in trypticase soy broth supplemented with 0.1% cysteine (TSBC) or on trypticase soy agar supplemented with 0.1% cysteine (TSAC) unless stated otherwise. Kanamycin (45 μ g/ml) or erythromycin (30 μ g/ml) or 10% sucrose were added as needed.

Subcellular fractionation

1000 ml of overnight *F. novicida* U112 culture was harvested and resuspended in 50 ml of cold phosphate buff-

Table 1: Strains and plasmids used in study.

Name	Phenotype/Relevant Characteristics	Reference or Source
U112	<i>Francisella novicida</i> prototype strain.	ATCC
JL0	U112, Δ sucrose hydrolase strain used to make deletion mutants.	Laboratory strain
ODB2	JL0, Δ iglA	This study
ODB7	U112, <i>iglA::Em^R</i>	This study
ODB1	U112, <i>iglB::Em^R</i>	This study
ODB5	Δ iglA/ <i>iglA::Km^R</i> , <i>in cis</i> complementation of <i>iglA</i> in strain ODB2	This study
CG62	U112, <i>iglC::TnMax2</i>	Gray et al. (2002)
GB2	U112, <i>mgIA</i>	Baron et al. (1998)
GB6	U112, <i>mgIB::mTn 10Km</i>	Baron et al. (1998)
DH5 α	F- Φ 80d <i>lacZ</i> Δ M15 Δ (<i>lacZYA-argF</i>)U169 <i>recA1 endA1 hsdR17</i> (<i>r_k⁻, m_k⁺</i>) <i>phoA supE44 thi-1 gyr A96 relA1 λ-</i>	Invitrogen
pCR2.1	Cloning vector, Amp ^R Km ^R	Invitrogen

ered saline (PBS). Cells were broken by repeated passage through a French Pressure cell (American Instruments Co, Silver Spring, MD) at 1200 PSI. Unbroken cells were removed by 20 min of centrifugation at 10,000 \times g at 4 °C, and a sample was taken as the total protein fraction. The lysate was subjected to ultracentrifugation (Beckman L8-70, rotor Type 45 Ti) for 1 hr at 100,000 \times g at 4 °C to pellet the membranes. The supernatant (soluble protein fraction) was removed, whereas the membrane pellet was resuspended in 2.5 ml of 1% Sarkosyl (Sigma). The sarkosyl soluble (inner membrane) and the sarkosyl insoluble (outer membrane) were separated by ultracentrifugation for 1 hr at 100,000 \times g at 4 °C in a Beckman TLA-100.3 ultramicrocentrifuge. The activity of the inner membrane-associated enzyme NADH oxidase was determined per mg of protein [33] for each of the fractions as a measure of the relative mixing of the different cell compartments. The soluble fraction contained 3%, the sarkosyl soluble membrane fraction 79% and the sarkosyl insoluble membrane fraction 18% of the NADH oxidase activity. In addition, we found that 90% of IglC was found in the soluble fraction (data not shown) and 10% was in the total membrane fraction. IglC could not be detected in the sarkosyl-soluble or sarkosyl-insoluble membrane fractions. As IglC has previously been shown to be a soluble protein [16], this served as another control of our fractionation experiment. Isolation of periplasmic proteins was performed as described by Nossal and Heppel [34].

Co-immunoprecipitation

500 μ l of soluble fraction was pre-cleared by incubation with 20 μ l protein-G/Agarose beads (40% slurry; EMB Bioscience, La Jolla, CA) and 10 μ g nonspecific antibody for 1 h at room temperature (RT). Beads and bound proteins were removed by centrifugation and the soluble fraction was incubated with 10 μ l rabbit anti-IglA serum or nonspecific antibody for 1 h at RT followed by addition of 75 μ l protein-G/Agarose beads and incubation 1 h at RT. Complexes were recovered by centrifugation, 6500 rpm, 3 min, and beads were washed three times with 150 mM

NaCl, 10 mM Na₂H₃PO₄, pH 7.2. After the final wash, complexes were resuspended in 30 μ l SDS-PAGE loading buffer and the sample was boiled for 5 min. Beads were removed by centrifugation and released proteins were separated on a 12% Sodium dodecyl sulphate-polyacrylamide electrophoresis (SDS-PAGE) gel. The immunoprecipitated material was examined by immunoblotting with anti-IglA to confirm that IglA was present (data not shown).

SDS-PAGE and Western blotting

To normalize the amount of protein added to each lane, the concentration of protein samples were determined by use of the BCA assay (Pierce). SDS-PAGE was performed according to standard techniques. Separated proteins were transferred onto a Trans Blot[®] nitrocellulose (BioRad) or Immobilon-FL (Millipore) membrane and blocked with 5% skim milk (Difco) in PBS. Anti-IglA, and anti-IglC antibody were used at dilutions of 1:4,000 and 1:500 respectively. To detect bound antibody blots were incubated with IRDye800DX-conjugated goat anti-rabbit or IRDye700DX-conjugated goat anti-rat immunoglobulin G (Rockland, Gilbertsville, Pa.) and visualized in a LiCor Odyssey imaging system.

MALDI-TOF

Following SDS-PAGE separation of proteins in-gel digestion with trypsin was carried out, and peptides extracted. 10 μ l of the peptide sample was loaded on to a C18 zip tip and washed three times in 10 μ l of 0.1% TFA and eluted with 2 μ l of 50% ACN and 0.1% TFA containing 10 mg/ml 4-hydroxy alpha cyanocinnamic acid. MALDI-TOF MS analysis of the peptides was carried out using a Voyager-DE STR (Applied Biosystems, Foster City, CA). Mass fingerprint analysis was done using Mascot (Matrix Science, UK).

Construction of *iglA* deletion mutant

IglA deletion mutant, ODB2, was constructed using a two-step integration-excision method. 1.5 kilobasepair (kbp) regions flanking *iglA* were amplified with primers *iglA* L-F

5' cgcggccgcagcaaaaatgctggaggtgt, *iglA* L-R 5' cctcgagcat-caaccttgaattgggatt, for the left-hand flanking region, and with primers *iglA* R-F 5' cctcgagctctgtgatgctgctgagtct, *iglA* R-R 5' cgcggccgcaataaccagccaggcttacc, for the right-hand flanking regions. These were cloned into plasmid pCR2.1 (Invitrogen) and verified by sequencing. The flanking regions were then joined by ligation. The flanking region construct was ligated to an erythromycin resistance-*sacB* cassette and the ligation mixture was used to chemically transform *F. novicida* JLO to erythromycin resistance as previously described [35]. The JLO strain (Ludu et al., unpublished data) is a derivative of the *F. novicida* U112 prototype strain that has a deletion in a sucrose hydrolase gene, and thus is sensitive to *sacB* expression in the presence of sucrose. An erythromycin resistant colony was grown and plated on TSAC containing 10% sucrose which acts as a counter selective marker for the *sacB* gene. Sucrose sensitive strains were examined for loss of *iglA* by PCR.

The *iglA* and *iglB* allelic replacement mutants, ODB7 and ODB1, were constructed as previously described [12]. Briefly, 1.5 kbp regions flanking *iglB* were PCR amplified with primers *iglB* L-F 5' cgcggccgcgaagaagataattcttctt-gaaaccg, *iglB* L-R 5' cctcgag attgcatacaaaaatctcttactt, *iglB* R-F 5' cctcgagtgactatagatactaggcttgaacca, *iglB* R-R 5' cgcggccgctcaaaaggctttggaaatcaa incorporating Xho I sites and ligated to an erythromycin resistance cassette with added Xho I sites. *F. novicida* U112 was transformed with the construct as previously described [35]. The same primers used for construction of ODB2 were used for ODB7.

In cis complementation

IglA and its promoter region were amplified with primers *IglA* int-L 5' CCCCTCGAGAGCCGTTTTCAATATTGGTTT and *IglA* int-R 5' CCCCTCGAGCAACTTCTGTAGATC-CCCCAAA incorporating added XhoI sites and ligated to a kanamycin resistance cassette carrying a *F. novicida* promoter (Ludu et al., unpublished data). The construct was used to transform ODB2 as previously described [35].

Macrophage infection assay

Macrophage infection assays were performed essentially as described previously [2]. Briefly, J774.1 mouse macrophage-like cells were infected with *F. novicida* strains at a multiplicity of infection of 50:1 (bacterium-to-macrophage), and monolayers were incubated for 2 h in Dulbecco's Modified Eagle Medium containing 10% fetal bovine serum (DMEM), washed five times in Dulbecco's Phosphate Buffered Saline (DPBS), and incubated at 37°C in 5% CO₂. Macrophages were lysed in 0.1% deoxycholate at 0, 24, 48 and 72 h post infection. To determine bacterial growth, lysed macrophages and culture supernatants were serially diluted in DPBS and plated on TSAC. As

F. novicida does not grow in DMEM, this allows for an adequate determination of intracellular growth [2].

Chicken embryo infections

Fertilized White Leghorn eggs were obtained from the University of Alberta Poultry Research Station. Seven-day old embryos were injected under the chorioallantoic membrane with various doses of 100 µl of *F. novicida* diluted in PBS as previously described [36]. The embryos were monitored for death for 6 days.

Authors' contributions

OMB performed all experiments, constructed *iglA* and *iglB* mutants and drafted the manuscript. JSL constructed the deletion of the sucrose hydrolase gene in *F. novicida*. FEN was the principal investigator and supervised the project. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by Grant Number 5R01 AI056215-02 from the National Institutes of Allergy and Infectious Diseases. OMB thanks Mike Roberts and Karen Cheung for many helpful and insightful discussions.

References

- Oyston PC, Sjostedt A, Titball RW: **Tularaemia: bioterrorism defence renews interest in *Francisella tularensis***. *Nat Rev Microbiol* 2004, **2**:967-978.
- Anthony LD, Burke RD, Nano FE: **Growth of *Francisella* spp. in rodent macrophages**. *Infect Immun* 1991, **59**:3291-3296.
- Vogel JP, Andrews HL, Wong SK, Isberg RR: **Conjugative transfer by the virulence system of *Legionella pneumophila***. *Science* 1998, **279**:873-876.
- Segal G, Purcell M, Shuman HA: **Host cell killing and bacterial conjugation require overlapping sets of genes within a 22-kb region of the *Legionella pneumophila* genome**. *Proc Natl Acad Sci U S A* 1998, **95**:1669-1674.
- Vazquez-Torres A, Xu Y, Jones-Carson J, Holden DW, Lucia SM, Din-auger MC, Mastroeni P, Fang FC: ***Salmonella* pathogenicity island 2-dependent evasion of the phagocyte NADPH oxidase**. *Science* 2000, **287**:1655-1658.
- Ochman H, Soncini FC, Solomon F, Groisman EA: **Identification of a pathogenicity island required for *Salmonella* survival in host cells**. *Proc Natl Acad Sci U S A* 1996, **93**:7800-7804.
- Uchiya K, Barbieri MA, Funato K, Shah AH, Stahl PD, Groisman EA: **A *Salmonella* virulence protein that inhibits cellular trafficking**. *Embo J* 1999, **18**:3924-3933.
- de Chastellier C, Berche P: **Fate of *Listeria monocytogenes* in murine macrophages: evidence for simultaneous killing and survival of intracellular bacteria**. *Infect Immun* 1994, **62**:543-553.
- Clemens DL, Lee BY, Horwitz MA: **Virulent and avirulent strains of *Francisella tularensis* prevent acidification and maturation of their phagosomes and escape into the cytoplasm in human macrophages**. *Infect Immun* 2004, **72**:3204-3217.
- Golovliov I, Baranov V, Krocova Z, Kovarova H, Sjostedt A: **An attenuated strain of the facultative intracellular bacterium *Francisella tularensis* can escape the phagosome of monocytic cells**. *Infect Immun* 2003, **71**:5940-5950.
- Santic M, Molmeret M, Abu Kwaik Y: **Modulation of biogenesis of the *Francisella tularensis* subsp. *novicida*-containing phagosome in quiescent human macrophages and its maturation into a phagolysosome upon activation by IFN-gamma**. *Cell Microbiol* 2005, **7**:957-967.
- Nano FE, Zhang N, Cowley SC, Klose KE, Cheung KK, Roberts MJ, Ludu JS, Letendre GW, Meierovics AI, Stephens G, Elkins KL: **A *Francisella tularensis* pathogenicity island required for intramacrophage growth**. *J Bacteriol* 2004, **186**:6430-6436.
- Lauriano CM, Barker JR, Yoon SS, Nano FE, Arulanandam BP, Hassett DJ, Klose KE: **MglA regulates transcription of virulence factors**

- necessary for *Francisella tularensis* intramacrophage and intramacrophage survival. *Proc Natl Acad Sci U S A* 2004, **101**:4246-4249.
14. Hansen AM, Qiu Y, Yeh N, Blattner FR, Durfee T, Jin DJ: **SspA is required for acid resistance in stationary phase by downregulation of H-NS in *Escherichia coli*.** *Mol Microbiol* 2005, **56**:719-734.
 15. Baron GS, Nano FE: **MglA and MglB are required for the intramacrophage growth of *Francisella novicida*.** *Mol Microbiol* 1998, **29**:247-259.
 16. Golovliov I, Ericsson M, Sandstrom G, Tarnvik A, Sjostedt A: **Identification of proteins of *Francisella tularensis* induced during growth in macrophages and cloning of the gene encoding a prominently induced 23-kilodalton protein.** *Infect Immun* 1997, **65**:2183-2189.
 17. Gray CG, Cowley SC, Cheung KK, Nano FE: **The identification of five genetic loci of *Francisella novicida* associated with intracellular growth.** *FEMS Microbiol Lett* 2002, **215**:53-56.
 18. Lai XH, Golovliov I, Sjostedt A: **Expression of IgIC is necessary for intracellular growth and induction of apoptosis in murine macrophages by *Francisella tularensis*.** *Microb Pathog* 2004, **37**:225-230.
 19. Lindgren H, Golovliov I, Baranov V, Ernst RK, Telepnev M, Sjostedt A: **Factors affecting the escape of *Francisella tularensis* from the phagolysosome.** *J Med Microbiol* 2004, **53**:953-958.
 20. Santic M, Molmeret M, Klose KE, Jones S, Kwai YA: **The *Francisella tularensis* pathogenicity island protein IgIC and its regulator MglA are essential for modulating phagosome biogenesis and subsequent bacterial escape into the cytoplasm.** *Cell Microbiol* 2005, **7**:969-979.
 21. Sexton JA, Miller JL, Yoneda A, Kehl-Fie TE, Vogel JP: ***Legionella pneumophila* DotU and lcmF are required for stability of the Dot/lcm complex.** *Infect Immun* 2004, **72**:5983-5992.
 22. Das S, Chaudhuri K: **Identification of a unique IAHP (lcmF associated homologous proteins) cluster in *Vibrio cholerae* and other proteobacteria through in silico analysis.** *In Silico Biol* 2003, **3**:287-300.
 23. Folkesson A, Lofdahl S, Normark S: **The *Salmonella enterica* subspecies I specific centisome 7 genomic island encodes novel protein families present in bacteria living in close contact with eukaryotic cells.** *Res Microbiol* 2002, **153**:537-545.
 24. Pukatzki S, Ma AT, Sturtevant D, Krastins B, Sarracino D, Nelson WC, Heidelberg JF, Mekalanos JJ: **Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system.** *Proc Natl Acad Sci U S A* 2006, **103**:1528-1533.
 25. Baron GS, Nano FE: **An erythromycin resistance cassette and mini-transposon for constructing transcriptional fusions to cat.** *Gene* 1999, **229**:59-65.
 26. Maier TM, Havig A, Casey M, Nano FE, Frank DW, Zahrt TC: **Construction and characterization of a highly efficient *Francisella* shuttle plasmid.** *Appl Environ Microbiol* 2004, **70**:7511-7519.
 27. Mougous JD, Cuff ME, Raunser S, Shen A, Zhou M, Gifford CA, Goodman AL, Joachimiak G, Ordonez CL, Lory S, Walz T, Joachimiak A, Mekalanos JJ: **A virulence locus of *Pseudomonas aeruginosa* encodes a protein secretion apparatus.** *Science* 2006, **312**:1526-1530.
 28. Day JB, Plano GV: **A complex composed of SycN and YscB functions as a specific chaperone for YopN in *Yersinia pestis*.** *Mol Microbiol* 1998, **30**:777-788.
 29. Forsberg A, Viitanen AM, Skurnik M, Wolf-Watz H: **The surface-located YopN protein is involved in calcium signal transduction in *Yersinia pseudotuberculosis*.** *Mol Microbiol* 1991, **5**:977-986.
 30. Ninio S, Zuckman-Cholon DM, Cambronnie ED, Roy CR: **The *Legionella* lcmS-lcmW protein complex is important for Dot/lcm-mediated protein translocation.** *Mol Microbiol* 2005, **55**:912-926.
 31. Coers J, Kagan JC, Matthews M, Nagai H, Zuckman DM, Roy CR: **Identification of lcm protein complexes that play distinct roles in the biogenesis of an organelle permissive for *Legionella pneumophila* intracellular growth.** *Mol Microbiol* 2000, **38**:719-736.
 32. Hager AJ, Bolton DL, Pelletier MR, Brittnacher MJ, Gallagher LA, Kaul R, Skerrett SJ, Miller SI, Guina T: **Type IV pili-mediated secretion modulates *Francisella* virulence.** *Mol Microbiol* 2006, **62**:227-237.
 33. McDonald MK, Cowley SC, Nano FE: **Temperature-sensitive lesions in the *Francisella novicida* valA gene cloned into an *Escherichia coli* msbA lpxK mutant affecting deoxycholate resistance and lipopolysaccharide assembly at the restrictive temperature.** *J Bacteriol* 1997, **179**:7638-7643.
 34. Nossal NG, Heppel LA: **The release of enzymes by osmotic shock from *Escherichia coli* in exponential phase.** *J Biol Chem* 1966, **241**:3055-3062.
 35. Mdluli KE, Anthony LS, Baron GS, McDonald MK, Myltseva SV, Nano FE: **Serum-sensitive mutation of *Francisella novicida*: association with an ABC transporter gene.** *Microbiology* 1994, **140** (Pt 12):3309-3318.
 36. Nix EB, Cheung KKM, Wang D, Zhang N, Burke RD, Nano FE: **Virulence of *Francisella* spp. in chicken embryos.** *Infect Immun* 2006, **74**:4809-4816.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

