

Research article

Open Access

Molecular and biochemical characterization of urease and survival of *Yersinia enterocolitica* biovar IA in acidic pH *in vitro*

Neeru Bhagat and Jugsharan S Viridi*

Address: Microbial Pathogenicity Laboratory, Department of Microbiology, University of Delhi South Campus, Benito Juarez Road, New Delhi - 110 021, India

Email: Neeru Bhagat - neeru_bhagat@yahoo.com; Jugsharan S Viridi* - viridi_dusc@rediffmail.com

* Corresponding author

Published: 17 December 2009

Received: 20 May 2009

BMC Microbiology 2009, 9:262 doi:10.1186/1471-2180-9-262

Accepted: 17 December 2009

This article is available from: <http://www.biomedcentral.com/1471-2180/9/262>

© 2009 Bhagat and Viridi; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: *Yersinia enterocolitica*, an important food- and water-borne enteric pathogen is represented by six biovars viz. IA, IB, 2, 3, 4 and 5. Despite the lack of recognized virulence determinants, some biovar IA strains have been reported to produce disease symptoms resembling that produced by known pathogenic biovars (IB, 2-5). It is therefore imperative to identify determinants that might contribute to the pathogenicity of *Y. enterocolitica* biovar IA strains. *Y. enterocolitica* invariably produces urease and the role of this enzyme in the virulence of biovar IB and biovar 4 strains has been reported recently. The objective of this work was to study genetic organization of the urease (*ure*) gene complex of *Y. enterocolitica* biovar IA, biochemical characterization of the urease, and the survival of these strains under acidic conditions *in vitro*.

Results: The *ure* gene complex (*ureABCEFGD*) of *Y. enterocolitica* biovar IA included three structural and four accessory genes, which were contiguous and was flanked by a urea transport (*yut*) gene on the 3' side. Differences were identified in *ure* gene complex of biovar IA strain compared to biovar IB and 4 strains. This included a smaller *ureB* gene and larger intergenic regions between the structural genes. The crude urease preparation exhibited optimal pH and temperature of 5.5 and 65°C respectively, and Michaelis-Menten kinetics with a K_m of 1.7 ± 0.4 mM urea and V_{max} of 7.29 ± 0.42 μ mol of ammonia released/min/mg protein. The urease activity was dependent on growth temperature and growth phase of *Y. enterocolitica* biovar IA, and the presence of nickel in the medium. The molecular mass of the enzyme was > 545 kDa and an isoelectric point of 5.2. The number of viable *Y. enterocolitica* biovar IA decreased significantly when incubated at pH 2.5 for 2 h. However, no such decrease was observed at this pH in the presence of urea.

Conclusions: The *ure* gene cluster of biovar IA strains though similar to biovar IB and 4 strains, exhibited important differences. The study also showed the ability of biovar IA strains of *Y. enterocolitica* to survive at highly acidic pH *in vitro* in the presence of urea.

Background

Yersinia enterocolitica, an important food- and water-borne human enteropathogen is known to cause a variety of gastrointestinal problems. Most commonly, it causes acute

diarrhea, terminal ileitis and mesenteric lymphadenitis [1]. Long-term sequelae following infection include reactive arthritis and erythema nodosum [1]. Blood transfusion associated septicemia due to *Y. enterocolitica* has been

reported to have high mortality [2]. Currently, *Y. enterocolitica* is represented by six biovars (1A, 1B, 2, 3, 4 and 5) and more than 30 distinct serovars. The virulence of known pathogenic biovars namely 1B and 2-5 is attributed to pYV (plasmid for *Yersinia* virulence) plasmid [3] and chromosomally borne virulence factors [4].

The biovar 1A strains however lack pYV plasmid and have generally been regarded as avirulent. But several clinical, epidemiological and experimental evidences indicate their potential pathogenicity [5]. Some biovar 1A strains have been reported to produce disease symptoms resembling that produced by pathogenic biovars [6,7]. These have been implicated in nosocomial [8] and food-borne [9] outbreaks and isolated from extra-intestinal sites [10]. The biovar 1A strains also invade epithelial cells [11,12], resist killing by macrophages [13] and carry virulence-associated genes such as *ystB* (enterotoxin), *inv* (invasin), *myfA* (fimbriae), *hreP* (subtilisin/kexin-like protease) and *tccC* (insecticidal-toxin like complex) [5,14]. In the past, enterotoxin has been thought to be the only major virulence factor produced by biovar 1A strains. Recently insecticidal-toxin complex [15] and flagella [16] have been identified as virulence factors of *Y. enterocolitica* biovar 1A strains. However the exact mechanisms underlying the pathogenesis by biovar 1A strains remains unclear and there is need to investigate the role of other putative virulence factors.

Urease (urea amidohydrolase; EC 3.5.1.5) has been implicated to play a role in the pathogenesis of many bacteria such as *Helicobacter pylori*, *Proteus mirabilis* and *Brucella abortus* [17-19]. *Y. enterocolitica* invariably produces urease which has been reported to enable biovar 1B and biovar 4 strains to survive in the acidic environment of the stomach [20,21]. However, the role of urease in the survival of biovar 1A strains has not been investigated. The objective of this study was to determine the genetic organization of urease (*ure*) gene cluster, factors affecting urease activity, and the survival of biovar 1A strain of *Y. enterocolitica* in acidic pH *in vitro*.

Methods

Bacterial strains and growth conditions

Y. enterocolitica biovar 1A (serovar O:6,30) isolated from the stools of a diarrheic patient and deposited with *Yersinia* National Reference Laboratory and WHO Collaborating Center, Pasteur Institute (Paris) under reference number IP27403 was used to characterize *ure* gene complex and the enzyme urease. The details of other *Y. enterocolitica* strains used in this study namely serovars, source of isolation, country of origin, reference laboratory accession numbers and clonal groups have been reported previously [22]. *Y. enterocolitica* 8081 (bio serovar 1B/O:8) was obtained from M. Skurnik (Haartman Institute, Hel-

sinki, Finland). *Y. enterocolitica* IP26329 (bio serovar 2/O:9), IP26249 (bio serovar 2/O:5,27), and IP134 (bio serovar 4/O:3) were obtained from E. Carniel (*Yersinia* National Reference Laboratory and WHO Collaborating Center, Pasteur Institute, France). All strains were grown overnight at 28°C in Luria broth (HiMedia, Mumbai, India).

DNA extraction, primers and Polymerase Chain Reaction

Genomic DNA was isolated from overnight grown cultures using DNeasy tissue kit (Qiagen GmbH) as reported earlier [14].

Urease gene sequences of *Y. enterocolitica* biovar 1B and biovar 4 with GenBank accession numbers [L24101](#)[23] and [Z18865](#)[24] respectively were used to design primers U1 and U2 using PrimerSelect 5.03 software (DNASTAR Inc., Madison, USA) such that the structural genes (*ureA*, *ureB*, *ureC*) may be amplified as one amplicon. As these primers failed to consistently amplify the *ureABC* region of biovar 1A strains, primers for amplification of each of the structural genes separately were designed from the following sequences in the database (accession numbers are given in parentheses): *Y. enterocolitica* biovar 1B ([L24101](#), [AM286415](#)), *Y. enterocolitica* biovar 4 ([Z18865](#)), *Y. aldovae* ([AY363680](#)), *Y. bercovieri* ([AY363681](#)), *Y. frederiksenii* ([AY363682](#)), *Y. intermedia* ([AY363683](#)), *Y. kristensenii* ([AY363684](#)), *Y. mollaretii* ([AY363685](#)), *Y. rohdei* ([AY363686](#)), *Y. pestis* ([AE017042](#), [AL590842](#), [AE009952](#), [AF095636](#)) and *Y. pseudotuberculosis* ([U40842](#), [BX936398](#)). These sequences were also used to design primers for *ure* accessory (*ureE*, *ureF*, *ureG*, *ureD*) and urea transport (*yut*) genes. The most conserved regions for each of the genes were identified using MegAlign (DNASTAR) or ClustalW version 1.83 (accessible at <http://www.ebi.ac.uk/tools/clustalW>).

Primer pairs - *ureA1-ureA2* (for *ureA*), *ureB1-ureB2* (for *ureB*), *ureC1-ureC2* (for *ureC*), *ureCE1-ureCE2* (for *ureC-ureE* and *ureE*), *ureF1-ureF2* (for *ureF*), *ureG1-ureG2* (for *ureG*), *ureD1-ureD2* (for *ureD*) and *yut1-yut2* (for *yut*) were designed from the conserved regions (Fig. 1) with PrimerSelect software. The sequences of the amplicons thus obtained (with strain IP27403) were used subsequently to design primers for the intergenic regions and a remaining part of the *ureC* gene. The intergenic regions between *ureA-ureB*, *ureB-ureC*, *ureE-ureF*, *ureF-ureG* and *ureD-yut* were amplified using primer pairs - *ureAB1-ureAB2*, *ureBC1-ureBC2*, *ureE1-ureE2*, *ureFG1-ureFG2* and *ureD3-ureD4* respectively and part of *ureC* gene by *ureC3-ureC4*. As *ureD* could not be amplified in biovar 1A strain with *ureD1-ureD2*, another primer pair *ureG1-ureD2* was used for amplification of the *ureG-ureD* intergenic region and *ureD* gene. The primers were synthesized

from Microsynth or Sigma Genosys. The details of the PCR primers and the target genes are given in Table 1.

PCRs for *ure* structural and accessory genes, intergenic regions and the *yut* gene were performed using a thermal cycler (MyCycler, Bio-Rad). The 25 μ l PCR reaction mixture contained 100 ng of genomic DNA, 2.5 μ l of 10 \times Taq buffer containing 1.5 mM MgCl₂, 2.5 μ l of 2 mM dNTP, 25 pmol of each primer, and 2 U of Taq DNA polymerase (New England BioLabs). The details of the conditions used for amplification are given in Table 1. After amplification, 10 μ l of the PCR product was resolved in 2% agarose gel in 1 \times Tris-acetate-ethylenediaminetetraacetic acid (TAE) buffer (40 mM Tris-HCl, 20 mM acetic acid, 1 mM EDTA, pH 8.0) at 70 V for 2 h. The gels were stained with ethidium bromide (0.5 μ g/ml) and photographed under UV-transillumination in a gel documentation system (Bio-Rad, CA). The 1 kb and 100 bp DNA ladders (New England BioLabs) served as molecular size markers.

Sequencing of PCR amplicons, ORF analysis and phylogenetic relationships

The PCR amplicons obtained above using the genomic DNA of *Y. enterocolitica* biovar 1A (strain IP27403) were extracted, purified using QIAquick Gel extraction kit (Qiagen) and sequenced directly in one or both directions (Microsynth, Balgach, Switzerland or LabIndia, Gurgaon, India). The sequences were analyzed, edited and compiled using Editseq and MegAlign of DNASTAR. Homology searches for nucleotide and deduced amino acid sequences were carried out by BLASTN and BLASTP respectively. The multiple nucleotide and protein sequence alignments were performed by MegAlign or ClustalW. The percent identity and similarity were calculated using MatGAT 2.02 [25]. The theoretical molecular weight and isoelectric point (pI) of urease structural and accessory proteins were determined by EditSeq (DNASTAR).

The open reading frames (ORFs) in the compiled *ure* gene cluster were identified using GeneMark [26], GeneMark.hmm [27], FGENESB [28] and the NCBI ORF finder [29] programs. All ORFs were checked further for homology to known protein sequences using BLASTX.

The relationship of urease structural and accessory protein sequences of biovar 1A strain of *Y. enterocolitica* to sequences available in GenBank were determined by constructing phylogenetic trees with the program MEGA 4.0 using the neighbor-joining algorithm. Bootstrap value for each node of the tree was calculated over 1,000 replicate trees.

PCR-Restriction fragment length polymorphism (PCR-RFLP) of urease genes

Primer pairs ureAB3-ureAB4 and ureC1-ureC4 were designed to amplify the 1,004 bp and 1,727 bp of *ureAB* and *ureC* genes respectively (Fig. 1). The biovar 1A strains were chosen such that each belonged to a different serovar, country, source of isolation, REP/ERIC-type [22] and VNTR01-type [30]. The PCR amplicon of *ureAB* was digested with *Hae*III and *Sau*96I while that of *ureC* was digested with *Rsa*I and *Sau*96I. The choice of the restriction enzymes was based on *in silico* restriction of the expected amplicons such that DNA fragments were amenable to separation by gel electrophoresis. Restriction enzymes were from New England BioLabs (*Rsa*I and *Hae*III) or Bangalore Genei (*Sau*96I). Ten microlitre of amplified DNA was digested with 2.5 U (*Hae*III and *Sau*96I) or 5 U (*Rsa*I) of restriction enzyme using appropriate buffer recommended by the manufacturer, in a total volume of 25 μ l at 37 °C overnight. The digested products were separated by electrophoresis in 2.5% agarose gel at 50 V for 5 h in TAE buffer. 100 bp ladder (New England BioLabs) was used as the molecular size standard. The gel was stained with ethidium bromide and examined under UV transillumination.

Growth and preparation of cell free extract

Y. enterocolitica strain IP27403 was grown overnight at 28 °C in 20 ml LB medium with shaking at 200 rpm. Cells were collected by centrifugation (9,000 \times g, 10 min, 4 °C), washed twice, and resuspended to 1.5 \times 10⁸ CFU/ml equivalent to 0.5 McFarland standard (A₆₀₀ = 0.1). These were diluted to 1.0 \times 10⁶ CFU/ml and 50 μ l of this suspension was inoculated into 50 ml of fresh LB medium, and incubated further (28 °C, shaking at 200 rpm). Samples were withdrawn at different time intervals up to 78 h and diluted in 20 mM sodium phosphate buffer (pH 7.0). 0.1 ml of the appropriate dilution was plated, in triplicate, on Luria agar and incubated overnight at 28 °C. The number of viable bacteria was recorded at different intervals and CFU/ml was calculated. The log₁₀CFU/ml was plotted against incubation time (in h).

For preparing lysate, cells grown in 50 ml LB medium were harvested by centrifugation, washed twice and resuspended in 2.5 ml of 20 mM sodium phosphate buffer (pH 7.0). Cells were disrupted by sonication with three cycles (2 s "pulse on" and 2 s "pulse off" for 2 min) at 25% intensity with Vibra-Cell (Sonics). The cell lysate was centrifuged at 18,000 \times g for 30 min at 4 °C to obtain cell-free extract. The supernatant was transferred to pre-chilled microcentrifuge tubes and used immediately for determination of urease activity. Protein concentration was estimated by Bradford [31] method using bovine serum albumin (Sigma) as standard.

Table 1: PCR amplification of urease structural (*ureA*, *ureB*, *ureC*) and the accessory (*ureE*, *ureF*, *ureG*, *ureD*) genes and the intergenic regions thereof, in *Y. enterocolitica* biovar IA strain.

Primer	Sequence (5' - 3')	Target	Accession no.	Region amplified	Amplicon length (bp)	PCR conditions (°C, s)*		
						Den	Ann	Ext
U1	GCAGCCGTTTGGTC	<i>ureA-ureC</i>	DQ350880	275...2896	2622	94, 60	62.0, 110	72, 110
U2	ACGG		AM286415	1075847...1078426	2580			
	CTATGCCACGCATC CCGACC		Z18865	325...2907	2582			
<i>ureA1</i>	GGAGGGCTTATGCA	<i>ureA</i>	DQ350880	1...161	161	94, 60	61.4, 60	72, 60
<i>ureA2</i>	GCTCACCCCAAG		AM286415	1075573...1075733	161			
	TTGCCATCTCTGGC CCCTTCCA		Z18865	51...211	161			
<i>ureAB1</i>	CAATGGAAGGGGCC	<i>ureA-ureB</i>	DQ350880	137...579	443	94, 60	60.3, 60	72, 60
<i>ureAB2</i>	AGAGATGG		AM286415	1075709...1076210	502			
	GTAAGCCGCAGCAC GGTCAAACCTC		Z18865	187...688	502			
<i>ureAB3</i>	GCAGCTCACCCAA	<i>ureA-ureB</i>	DQ350880	12...1015	1004	95, 60	56.9, 110	72, 60
<i>ureAB4</i>	GAGAAGTTGA		AM286415	1075584...1076608	1025			
	AATTTGAGGCATCT GTCGCTCCTT		Z18865	62...1086	1025			
<i>ureB1</i>	ATTGCAGAGGATTA	<i>ureB</i>	DQ350880	349...650	302	94, 60	60.0, 60	72, 60
<i>ureB2</i>	AAGCATGAGC		AM286415	1075920...1076281	362			
	AGCGAACTTCGGT TTCATCAC		Z18865	398...759	362			
<i>ureBC1</i>	TGCGGCTTACGGAA	<i>ureB-ureC</i>	DQ350880	570...1022	453	94, 60	60.3, 60	72, 60
<i>ureBC2</i>	AAAGGCTGAATA		AM286415	1076201...1076615	415			
	GCCGAGAAATTTGA GGCATCTGTCG		Z18865	679...1093	415			
<i>ureC1</i>	AAAGGAGCGACAGA	<i>ureC</i>	DQ350880	991...1749	759	94, 60	63.2, 60	72, 60
<i>ureC2</i>	TGCCTCAAAA		AM286415	1076584...1077342	759			
	GAAACCTGAATATC CATTTTCATCCGCCA T		Z18865	1062...1823	762			
<i>ureC3</i>	GGCTATAAAGTTCA	<i>ureC</i>	DQ350880	1661...2717	1057	94, 60	52.9, 60	72, 60
<i>ureC4</i>	CGAAGACTG		AM286415	1077254...1078310	1057			
	CAAAGAAATAGCGC TGGTTCA		Z18865	1735...2791	1057			
<i>ureC1</i>	AAAGGAGCGACAGA	<i>ureC</i>	DQ350880	991...2717	1727	94, 60	50.0, 60	72, 120
<i>ureC4</i>	TGCCTCAAAA		AM286415	1076584...1078310	1727			
	CAAAGAAATAGCGC TGGTTCA		Z18865	1062...2791	1730			
<i>ureCE1</i>	GCGCTGGATGACGG	<i>ureC-ureE, ureE</i>	DQ350880	2504...3552	1019	94, 60	61.0, 60	72, 60
<i>ureCE2</i>	TGTGAAAGAG		AM286415	1078097...1079082	986			
	ATGTAAGCCGGAGC CATGAGGTTTC							
<i>ureE1</i>	ACCATGATGGATTC	<i>ureE-ureF</i>	DQ350880	3364...3734	371	95, 60	58.7, 60	72, 60
<i>ureE2</i>	CGTGATGAGA		AM286415	1078894...1079270	377			
	GTGAAGGCCCCCGAC CGGCAGTACG							
<i>ureF1</i>	TGAATGCATCAGAT	<i>ureF</i>	DQ350880	3668...4304	637	95, 60	50.0, 60	72, 60
<i>ureF2</i>	CTGATTCGTA		AM286415	1079204...1079840	637			
	ACATCCACAATAGG GACATAAGA							

Table 1: PCR amplification of urease structural (*ureA*, *ureB*, *ureC*) and the accessory (*ureE*, *ureF*, *ureG*, *ureD*) genes and the intergenic regions thereof, in *Y. enterocolitica* biovar IA strain. (Continued)

<i>ureFG1</i>	CAATATGGCGTGGC	<i>ureF-ureG</i>	DQ350880	4132...4535	403	95, 60	55.7, 60	72, 60
<i>ureFG2</i>	GATGACAAT CCACCGGGCCACC AATACCAA		AM286415	1079668...1080070	401			
<i>ureG1</i>	GAATAGCCATTCAA	<i>ureG</i>	DQ350880	4474...5091	618	95, 60	51.3, 60	72, 60
<i>ureG2</i>	CCGATAAAC CGCATAATCATATC CACCAAC		AM286415	1080009...1080626	618			
<i>ureG1</i>	GAATAGCCATTCAA	<i>ureG-ureD, ureD</i>	DQ350880	4474...6099	1626	95, 60	50.4, 60	72, 120
<i>ureD2</i>	CCGATAAAC TTCCGGCAATGTCA CACCGAGAAT		AM286415	1080009...1081634	1626			
<i>ureD1</i>	AGCCAGAATATCGT	<i>ureD</i>	DQ350880	5146...6099	954	95, 60	50.0, 60	72, 60
<i>ureD2</i>	GGAAACTCCT TTCCGGCAATGTCA CACCGAGAAT		AM286415	1080681...1081634	954			
<i>ureD3</i>	TTGTTAACCCCAA	<i>ureD-yut</i>	DQ350880	5884...6416	533	95, 60	58.0, 60	72, 60
<i>ureD4</i>	AGAGCATCAT CTGCCGGATTCCCT TCGCCATAG		AM286415	1081419...1081950	532			
<i>Yut1</i>	CGCGGCTGTGCTCA	<i>yut</i>	AM286415	1081851...1082745	895	95, 60	50.0, 60	72, 60
<i>Yut2</i>	AGTC GTGCTGGCATCACA TCTTTATTAGG							

The primer details and the PCR conditions used are given.

[DQ350880](#): *Y. enterocolitica* IP27403 (bio serovar IA/O:6,30); [AM286415](#): *Y. enterocolitica* 8081 (bio serovar IB/O:8); [Z18865](#): *Y. enterocolitica* 6471/76 (bio serovar 4/O:3)

Nucleotides sequences in bold are different in biovar IA strain ([DQ350880](#))

*PCRs were performed with initial denaturation step of 94°C for 10 min, 30 cycles each of denaturation (Den), annealing (Ann) and extension (Ext) as indicated and a final extension of 10 min at 72°C

Urease assay

Urease activity in the cell extract was assayed by measuring release of ammonia from urea in the phenol-hypochlorite assay [32]. Briefly, extract containing 2 µg of protein was added to 100 mM citrate buffer (pH 5.5) containing 50 mM urea in 200 µl of final volume. The mixture was incubated at 37°C for 15 min. A similar volume of the extract

boiled for 10 min served as negative control. The reaction was terminated by the addition of 1.5 ml of solution containing 1% phenol and 0.005% sodium nitroprusside; this was followed by the addition of 1.5 ml solution containing 0.5% (w/v) NaOH and 0.044% (v/v) NaClO, and the contents were mixed well. Following incubation at 37°C for 30 min, the absorbance was measured at 625 nm




Figure 1

Organization of ure gene cluster of *Y. enterocolitica* biovar IA. Primers used for amplification of structural and accessory genes, and the intergenic regions thereof are indicated.

using a spectrophotometer (UV-1700 Pharmaspec; Shimadzu Scientific Instruments Inc., Columbia, Md.). Assays were carried out in triplicate and the amount of the ammonia released per minute was determined. The quantity of ammonia (in nmol) released was calculated from the calibration curve obtained from appropriate dilutions of freshly prepared NH_4Cl solution, which was determined to be linear between 20-500 nmol. Data are presented as specific activity of urease, defined as μmol of $\text{NH}_3/\text{min}/\text{mg}$ of protein. Stated values are the mean \pm standard deviation of triplicate determinations.

Biochemical characterization

The optimum pH for urease was determined by measuring activity at pH 1.5 to 7.5. The assays were carried out in 20 mM sodium phosphate (for pH 1.5, 2.5, 5.5, 6.0, 6.5, 7.0 and 7.5) and 100 mM citrate (for pH 3.0, 3.5, 4.0 and 5.5) buffers. The optimum temperature for urease was determined by incubating the extract containing enzyme with substrate at different temperatures (18-75°C) in the phenol-hypochlorite assay described above. The kinetic data (K_m and V_{max}) of urease were calculated from Lineweaver-Burk plot of the initial rate of hydrolysis of urea in citrate buffer (100 mM, pH 5.5). To determine the effect of growth temperature and growth phase on urease activity, extracts were prepared from cells grown in shaking at 28°C and 37°C in LB medium for different time intervals. To determine the effect of urea and nickel on production of urease, medium was supplemented with urea (16.7 mM) or NiCl_2 (up to 200 μM).

Native and SDS PAGE

Cell-free extracts from different biovars of *Y. enterocolitica* were electrophoresed on non-denaturing polyacrylamide gel [33]. Briefly, extract containing *ca.* 100 μg of protein was mixed with 1 \times tracking dye and loaded on 5% resolving gel in 380 mM Tris-HCl (pH 8.8) with 4% stacking gel in 63 mM Tris-HCl (pH 6.8) in a mini-Protein III apparatus (Bio-Rad). Samples were electrophoresed with Tris-Glycine (pH 8.4) as the running buffer at 70 V for 2 h at 4°C. The gel was removed and equilibrated with 5-10 changes of solution containing 0.02% cresol red and 0.1% EDTA until the entire gel turned yellow. After draining the solution, gel was flooded with 1.5% (w/v) solution of urea. The pink bands of urease were recorded by scanning (UMAX Astra 3600). Urease from jack bean (Sigma) was used as the marker.

SDS-PAGE was performed as per standard protocol [34]. Briefly, extract containing 25 μg of protein was boiled in reducing Laemmli sample buffer and separated on 12% polyacrylamide gel.

Isoelectric focusing (IEF)

IEF of the cell extract was carried out in 6% polyacrylamide gel containing 2% ampholyte of pH 3-10 (BioLyte Ampholyte, Bio-Rad). 3-5 μl of extract containing *ca.* 20-25 μg of protein was loaded on the gel and focused at 4°C using a Mini IEF cell (Bio-Rad) according to the manufacturer's instructions. After focusing, the gel was equilibrated with a solution containing 0.02% cresol red and 0.1% EDTA. Urease bands were visualized by superimposing the gel with Whatman No. 1 filter paper presaturated with cresol red-EDTA solution containing 1.5% urea. Urease appeared as pink band against a yellow background. Broad range IEF standard with pI 4.45-9.6 (Bio-Rad) was used as the pI marker to determine the isoelectric point of the urease.

Survival of *Y. enterocolitica* in acidic pH in vitro

The *in vitro* survival of *Y. enterocolitica* was performed by slight modification of the method reported earlier [35]. Briefly, ten microlitre of the bacterial suspension was added to 1 ml of 20 mM sodium phosphate (for pH 2.5 and 7.0) or 100 mM citrate (for pH 4.0) buffer with or without 3.4 mM urea in 0.6% NaCl, and prewarmed to 37°C to give an initial count of *ca.* 7.0 $\log_{10}\text{CFU}/\text{ml}$. The contents were mixed and incubated with shaking at 37°C for 2 h. At the end of the incubation, samples were removed and diluted serially in 20 mM sodium phosphate buffer (pH 7.0). 0.1 ml of an appropriate dilution was plated on LB agar to determine CFU/ml. At conclusion of each assay, the pH of the buffer was recorded. All assays were repeated at least thrice on separate occasions.

Statistical analysis

The mean and the standard deviation for each data set were calculated using Microsoft Excel 2003 software (Microsoft Corporation, Redmond, Wash.). Statistical significance was calculated using unpaired *t* test (Sigma Stat version 3.5). *p* value < 0.05 was considered significant.

Nucleotide sequence accession number

The nucleotide sequence data of *ure* gene complex and the *yut* gene reported in this paper have been deposited in GenBank database under accession numbers [DQ350880](#) and [EU527335](#) respectively.

Results

Characterization of urease genes

Primers U1 and U2 were designed to amplify the *ure* structural (*ureA*, *ureB*, *ureC*) genes of *Y. enterocolitica*. Although amplification was obtained with biovar 1B, 2 and 4 strains, these primers did not consistently amplify the *ure* structural genes of biovar 1A strains. Thus, new primers were designed to amplify each of the *ure* structural and accessory (*ureE*, *ureF*, *ureG*, *ureD*) genes separately, and the intergenic regions so as to encompass the entire urease

gene cluster of biovar 1A strain. Amplicons of expected sizes were obtained for all genes except *ureB* and the intergenic regions namely *ureA-ureB*, *ureB-ureC* and *ureC-ureE* (Table 1). The sequences thus obtained were analyzed for homology with sequences available in databases, edited and combined to obtain 7,180 bp sequence of *ure* gene cluster of biovar 1A strain (See Additional file 1 for *ure* gene cluster sequence).

Seven ORFs were identified in the *ure* gene cluster of *Y. enterocolitica* biovar 1A strain and designated as *ureA*, *ureB*, *ureC*, *ureE*, *ureF*, *ureG* and *ureD* (Fig. 1) as in the *ure* gene complex of *Y. enterocolitica* 8081 (biovar 1B, accession number [AM286415](#)). As with *Y. enterocolitica* 8081, *yut* gene which encodes a urea transport protein was present downstream of the *ure* gene cluster. All ORFs had ATG as the start codon except *ureG* where the start codon was GTG. These ORFs were preceded by ribosome-binding consensus sequence. Although *ure* gene cluster of biovar 1A strain was broadly similar to that of biovar 1B and bio-

var 4 strains, differences were identified. These were - smaller *ureB* gene and *ureA-ureB* intergenic region and larger *ureB-ureC* and *ureC-ureE* intergenic regions in biovar 1A strain (Table 2). The size of *ureB* gene of *Y. enterocolitica* biovar 1A was identical to *ureB* of *Y. aldovae*, *Y. bercovieri*, *Y. intermedia*, *Y. mollaretii* and exhibited higher nucleotide sequence identity to these species than to *Y. enterocolitica* biovar 1B or 4. The stop codon of *ureG* overlapped with the start codon of *ureD* gene. The G + C content of the urease gene cluster was 49.76% which was typical of *Y. enterocolitica* with G + C content of 47.27%.

The comparison of *Y. enterocolitica* biovar 1A *ure* genes and the deduced amino acid sequences with that of *Yersinia* spp. and other bacteria are given in Tables 2 and 3 respectively. Besides *Yersinia* species, the homologies of *ure* genes (upto 76% identity) and their deduced amino acid sequences (upto 86% identity and 95% similarity) were significant with ureases from *Photothabdus luminescens* and *Edwardsiella ictaluri*. The UreA, UreC and UreG

Table 2: Urease structural and accessory genes and the intergenic regions thereof, in *Y. enterocolitica* biovar 1A.

Gene/Intergene	Ye1A	Size (in bp) and % identity							
		YeO8	YeO3	Yers	Yps	Ype	Pl	Ei	Ka
Structural									
<i>ureA</i>	303	303 98	303 98	303 89-96	303 91	303 91	303 77	303 76	303 63
<i>ureB</i>	435	495 83	495 83	435-495* 78-89	435/477 74-82	435/477/528 67-81	390 63	423 62	321 48
<i>ureC</i>	1719	1719 96	1722 94	1719 87-91	1719 86	1719 82-86	1716 75	1719 76	1704 61
Accessory									
<i>ureE</i>	687	792/693 85/97	NA	681-720 83-91	696 82	696/705 80-82	597 59	678 58	477 44
<i>ureF</i>	687	687 98	NA	687 85-90	687 85-86	687 86	687 66	705 65	675 48
<i>ureG</i>	666	666/606 99/90	NA	666 88-92	663 84	663 85	636 78	630 73	618 59
<i>ureD</i>	984	978/984 96/97	NA	966-984 84-90	966 81	834/964/967 71-82	966 64	963 62	-
Intergenic region									
<i>ureAB</i>	54	53	53	53-65	10/52	0/10/52	91	46	9
<i>ureBC</i>	202	164	164	87-97/201-202*	89	89	67	42	0
<i>ureCE</i>	236	74/173	NA	133-204	294/295	286/295	74	57	9
<i>ureEF</i>	21	21	NA	20-24	21	21	0	0	1
<i>ureFG</i>	117	117/177	NA	58-102	125/126	126	52	14	8
<i>ureGD</i>	0	1/0	NA	0	0	0	0	6	-

Comparison with different *Yersinia* spp. and other bacteria.

The abbreviations correspond to following species with accession number(s) in parentheses. Ye1A: *Y. enterocolitica* bioserovar 1A/O:6,30 ([DQ350880](#)); YeO8: *Y. enterocolitica* bioserovar 1B/O:8 ([L24101](#), [AM286415](#)); YeO3: *Y. enterocolitica* bioserovar 4/O:3 ([Z18865](#)); Yers included *Y. aldovae* ([AY363680](#)), *Y. bercovieri* ([AY363681](#)), *Y. frederiksenii* ([AY363682](#)), *Y. intermedia* ([AY363683](#)), *Y. kristensenii* ([AY363684](#)), *Y. mollaretii* ([AY363685](#)), *Y. rohdei* ([AY363686](#)); Yps: *Y. pseudotuberculosis* ([U40842](#); [CP000720](#); [CP000950](#); [BX936398](#)); Ype: *Y. pestis* ([CP000901](#), [CP000308](#), [AL590842](#), [AE017042](#), [CP000305](#), [CP000668](#), [AF095636](#)); Pl: *Photothabdus luminescens* ([BX571866](#)); Ei: *Edwardsiella ictaluri* ([AY607844](#)); Ka: *Klebsiella aerogenes* ([M36068](#))

% identity is indicated in bold

0 indicates that the intergenic region had overlapping stop and start codons

**ureB* gene size was 435 bp (*Y. aldovae*, *Y. bercovieri*, *Y. intermedia*, and *Y. mollaretii*), 441 bp (*Y. rohdei*), 468 bp (*Y. frederiksenii*) and 495 bp (*Y. kristensenii*); *ureBC* intergenic region of 201-202 bp was present in *Y. aldovae* and *Y. intermedia*

Table 3: Urease structural and accessory proteins of *Y. enterocolitica* biovar 1A (Ye 1A).

	Gene	Gene product (aa)	Mol. mass (Da)*	pI*	% identity/% similarity							
					YeO8	YeO3	Yers	Yps	Ype	PI	Ei	Ka
Structural subunits												
UreA	<i>ureA</i>	100	11,048	5.29	99-100	100	97-100/100	100	100	79/95	86/95	60/82
UreB	<i>ureB</i>	144	15,854	9.06	84-85/85-86	85/86	84-99/85-99	86-94/88-97	78-94/79-97	60/72	61/73	36/47
UreC	<i>ureC</i>	572	61,026	5.64	99/100	95/97	97-99/99-100	97/99	93-97/95-99	83/91	86/94	58/73
Accessory proteins												
UreE	<i>ureE</i>	228	25,507	6.35	86-99	NA	91-95/95-97	94/97	92-93/96-97	55/69	50/70	27/39
UreF	<i>ureF</i>	228	25,040	6.41	100	NA	96-99/97-100	98/99	97-98/99	67/76	65/83	22/43
UreG	<i>ureG</i>	221	24,181	4.94	91-100	NA	98-100/99-100	96/97	96/97	86/91	86/91	54/71
UreD	<i>ureD</i>	327	36,592	6.61	93-98/95-99	NA	91-98/95-99	93/96	FS	64/77	59/71	-

Comparison with different *Yersinia* spp. and other bacteria.

The abbreviations correspond to following species with protein accession numbers for UreA, UreB, UreC, UreE, UreF, UreG and UreD in parentheses: Ye1A: *Y. enterocolitica* biovar 1A (ABC74582-ABC74585; ACA51855-ACA51857); YeO8: *Y. enterocolitica* O8 biovar 1B (AAA50994-AAA51000, CAL11049-CAL11055); YeO3: *Y. enterocolitica* O3 biovar 4 (CAA79314-AA79320); Yers included *Y. aldovae* (AAR15084-AAR15090); *Y. bercovieri* (AAR15092-AAR15098); *Y. frederiksenii* (AAR15100-AAR15106); *Y. intermedia* (AAR15108-AAR15114); *Y. kristensenii* (AAR15117-AAR15123); *Y. mollaretii* (AAR15126-AAR15132); *Y. rohdei* (AAR15135-AAR15141); Yps: *Y. pseudotuberculosis* (CAH22182-CAH22176, AAA87852-AAA87858, ACA67429-ACA67435); Ype: *Y. pestis* (ABG14357-ABG14363; CAL21284-CAL21289; AAS62666-AAS62671; AAM84812-AAM84817; ABG17479-ABG17485; ABP39996-ABP39990; AAC78632-AAC78638); PI: *Photobacterium luminescens* (CAE14464-CAE14470); Ei: *Edwardsiella ictaluri* (ABD93708-ABD93706, AAT42448-AAT42445); Ka: *Klebsiella aerogenes* (AAA25149-AAA25154); NA: Not available; FS: frameshift mutation

* Theoretical molecular mass and pI were determined with DNASTAR

proteins were most conserved among *Yersinia* spp. The estimated molecular weights, in Da, of the protein subunits were 11,048 (UreA), 15,854 (UreB), 61,026 (UreC), 25,507 (UreE), 25,040 (UreF), 24,181 (UreG) and 36,592 (UreD) (Table 3).

Phylogenetic analysis of urease structural and accessory proteins of *Y. enterocolitica* biovar 1A showed clustering with members of gamma-proteobacteria such as *P. luminescens* and *E. ictaluri* along with *Yersinia* spp. (See Additional files 2 and 3). These protein sequences were also related closely to members of alpha-proteobacteria like *Methylobacterium chloromethanicum*, *M. extorquens*, *M. populi* and *Brucella* spp. but were related distantly to other members of gamma-proteobacteria like *Klebsiella aerogenes*, *P. mirabilis* and *Escherichia coli*.

PCR-RFLP of ure genes

The regions constituting the structural genes namely *ureAB* and *ureC* were amplified in several *Y. enterocolitica* biovar 1A strains using primer pairs AB3-AB4 and C1-C4 respectively. Restriction digestion of *ureAB* region with *HaeIII* and *Sau96I* resulted in almost identical patterns among all biovar 1A strains (See Additional file 4). But,

differences were clearly evident in restriction profiles of *ureC* digested with *RsaI* and *Sau96I* (Fig. 2). With *RsaI*, strains belonging to clonal group A exhibited profile different from that of clonal group B strains. Thus, it may be inferred that sequence of urease gene in clonal group A strains is different from that of clonal group B strains.

The *HaeIII* and *Sau96I* restriction profiles of *ureAB* of biovar 1B, 2 and 4 strains were distinct from that of biovar 1A strains (See Additional file 4). As with *ureAB*, restriction patterns of *ureC* for these biovars were also quite distinct from biovar 1A strains (data not shown).

Biochemical characterization

The crude extract of urease of *Y. enterocolitica* biovar 1A strain was active over a pH range of 4.0-7.0. The maximum activity was observed at pH 5.5 (Fig. 3a). The enzyme was quite heat-stable as urease activity was recorded up to 65 °C but decreased progressively at higher temperature (Fig. 3b). The optimum temperature for urease activity was 65 °C (Fig. 3b). The urease exhibited Michaelis-Menten kinetics with K_m and V_{max} of 1.74 ± 0.4 mM urea and 7.29 ± 0.42 μmol of ammonia released/min/mg of protein respectively (data not shown).




Figure 2
PCR-RFLP of ureC. PCR-RFLP of *ureC* of *Y. enterocolitica* biovar 1A strains amplified with primers ureC1-ureC4, and restriction digested using (A) *RsaI* and (B) *Sau96I* enzymes. Lanes 1: IP27360, 2: IP27362, 3: IP27364, 4: IP27365, 5: IP26310, 6: IP26311, 7: IP26312, 8: IP26315, 9: IP27403, 10: IP27407, 11: IP27429, 12: IP27433, 13: IP27434, 14: IP26261, 15: IP26305, 16: E1281580, 17: IP26316, 18: E1281550, 19: IP26152, 20: P346, 21: P354, 22: P386, 23: P472, 24: IP27404, 25: IP27406, 26: IP27430, 27: IP27432, 28: IP27484, 29: IP26147, 30: IP26148, 31: E1281600, 32: IP27385, 33: IP27386, 34: IP27388, 35: IP27485, 36: STM 126, 37: 8660/90 STM 484, 38: 0310/90, 39: ST5 NF-O, 40: IP27879, 41: IP27873, 42: IP27950, 43: IP27985, 44: IP24121, 45: IP27648, 46: IP27210, 47: IP27149, M: Molecular mass marker (100 bp ladder, New England BioLabs). * Strains belonging to clonal group B are shown in lanes 10, 14, 15, 19, 21, 22, 28, 29, 31, 45, 46 and 47. Clonal group A strains are in other lanes. For clonal groups refer to [22].

Y. enterocolitica biovar 1A grown at 28°C (optimum temperature for growth) exhibited higher urease activity than that grown at 37°C (Fig. 3c). Irrespective of the growth temperature, stationary phase cells showed higher activity (Fig. 3c). The supplementation of growth medium (Luria broth) with 16.7 mM urea did not show significant difference in urease activity. However, supplementation with nickel chloride resulted in *ca.* 10-fold increase in the activity. 1 µM NiCl₂ was sufficient to induce urease activity as no significant increase in the activity was observed with further increase in concentration up to 200 µM (See Additional file 5).

On native PAGE, urease was observed as two bands with the major band having molecular weight > 545 kDa and a slowly-developing band above it (Fig. 4). The electrophoretic mobility of urease of *Y. enterocolitica* biovar 1A strain was shown to be different from that of biovar 1B, 2 and 4 strains though similar to the *Y. intermedia* urease. The isoelectric point of the crude extract urease was 5.2.

Survival of *Y. enterocolitica* in vitro

The ability of *Y. enterocolitica* biovar 1A strain to survive at pH 2.5, 4.0 and 7.0 *in vitro* was investigated. Strains belonging to other biovars were also studied concurrently.




Figure 3
Biochemical characterization of *Y. enterocolitica* biovar 1A urease. (a) optimal pH for urease activity (b) effect of temperature on urease activity and (c) effect of growth phase and growth temperature on urease production; growth curve of biovar 1A strain grown at 28°C is also shown. Data points represent mean of triplicate determinations. The error bars indicate standard deviation.




Figure 4
Non-denaturing PAGE showing urease activity of *Yersinia* spp. Lane 1: *Y. enterocolitica* IP27403 (1A/O:6,30); lane 2: *Y. enterocolitica* IP134 (4/O:3); lane 3: *Y. enterocolitica* IP26329 (2/O:9); lane 4: *Y. enterocolitica* IP26249 (2/O:5,27); lane 5: *Y. enterocolitica* 8081 (1B/O:8); lane 6: *Y. intermedia* IP27478 (serotype O:7,8-8); M: Jack bean urease [272 kDa (trimer) and 545 kDa (hexamer)]; BV: Biovar.

The biovar 1A strain survived at pH 4.0 and 7.0 for 2 h without significant differences in their viable counts (Fig. 5). However, no viable cells were recovered after 2 h at pH 2.5. In fact, the decrease in the viable counts at this pH was evident even within 5 min of incubation. The addition of 3.4 mM urea at pH 2.5 was sufficient to increase the survival of *Y. enterocolitica* biovar 1A equivalent to that observed at pH 4.0 and 7.0. Similar results were observed for other biovars also. The pH of the assay medium at the end of experiment was same as that at the start, suggesting that increased survival of *Y. enterocolitica* was not due to any significant change in the pH.

Discussion

The *ure* gene cluster of *Y. enterocolitica* biovar 1A strain included three structural (*ureA*, *ureB*, *ureC*) and four (*ureE*, *ureF*, *ureG*, *ureD*) accessory genes. The *yut* gene, which is required for transport of urea was present downstream of this cluster. Thus, the organization (*ureABCEFGD*) of *ure* gene cluster in *Y. enterocolitica* biovar 1A strain was similar to that reported for *Y. enterocolitica* biovar 1B, *P. luminescens* and *E. ictaluri* [23,36,37]. Similar organization has




Figure 5
Survival of *Y. enterocolitica* in vitro at different pH.
 Number of bacterial cells (\log_{10} CFU/ml) of *Y. enterocolitica* after incubation for 2 h at pH 2.5, 4.0 and 7.0 in the absence and presence (U) of 3.4 mM urea. The values are mean of three independent observations. The error bars indicate standard deviation.

been reported for other species such as *Streptococcus salivarius*, *Synechococcus* sp. WH7805, and *B. abortus ure-2* operon [19,38,39]. However, important differences were observed compared to urease genes of *Y. enterocolitica* biovar 1B and biovar 4 strains. These included differences in the size of *ureB* gene and the intergenic regions. Also, the restriction profiles of *ure* structural genes of biovar 1A strains were different from that of biovars 1B, 2 and 4. These observations indicated that RFLP of urease genes may be used to study the epidemiology of *Y. enterocolitica*.

The amino acid residues in the urease structural proteins namely UreA (γ subunit), UreB (β subunit) and UreC (α subunit) that are reported to have functional significance in *K. aerogenes* urease [40] were also conserved in *Y. enterocolitica* biovar 1A. The crystallographic [41] and genetic [40] analysis of *K. aerogenes* urease has shown that four histidine residues (His-134, -136, -246 and -272), an aspartate (Asp-360) and a carbamylated lysine (Lys-217) of UreC are involved in nickel metallocenter binding. All these amino acids were conserved at positions His-139, -141, -251, -277; Asp-365 and Lys-222 in UreC of *Y. enterocolitica* biovar 1A. Histidine residues in the α -subunit of *K. aerogenes* shown to be important for substrate binding (His-219) and catalysis (His-320) are present at positions 224 and 325 in α -subunit of biovar 1A [40]. The urease active-site consensus sequence (MVCHHLD) [42] deviated by two residues (MVCHNLN) in biovar 1A strain. Amino acid residues with functional significance including His-97 (UreA) and His-39, -41 (UreB) [40] were also

conserved in relative positions in *Y. enterocolitica* biovar 1A. The conservation of amino acids in *Y. enterocolitica* biovar 1A urease involved in coordination of nickel at active site, substrate binding and catalysis as seen in *K. aerogenes* urease, suggested similar quaternary structure of the two enzymes. UreE consisted of histidine-rich motif at carboxy terminus as in UreE of *K. aerogenes*, *B. abortus*, *Actinobacillus pleuropneumoniae*, *E. ictaluri* and *Synechococcus* [19,36,39,43,44]. A P-loop motif (GPVGSCKT), which contains ATP and GTP binding sites [45] and probably provides energy for Ni activation [46] was present at the amino terminus (positions 19-26) of UreG.

A pH optimum in the acidic range for urease produced by a neutrophile like *Y. enterocolitica* biovar 1A was similar to that reported for *Y. enterocolitica* biovars 1B and 4, and *Morganella morganii* [35,47]. Ureases with optima in the acidic range reportedly carried a phenylalanine seven residues towards N-terminus, and an asparagine one residue toward the C-terminus, from the catalytic site [35]. Both these residues are also present at respective positions in UreC of *Y. enterocolitica* biovar 1A. The maximal activity of urease at 65°C by *Y. enterocolitica* biovar 1A has also been reported for other bacteria [44]. A low K_m of *Y. enterocolitica* biovar 1A urease as in biovar 4 strains [47], indicated its high affinity for urea. This suggested that the enzyme might function quite normally in the gut despite low concentrations (1.7-3.4 mM) of the urea available there. Also, consistent with our observation, organisms which produce urease with low K_m have been reported to possess urea transport (*yut*) gene as seen in *S. salivarius*, *Lactobacillus fermentum*, *Bacillus* sp. strain TB-90 and *B. suis* [48].

The cultural conditions which affected production of urease by *Y. enterocolitica* biovar 1A included growth phase, growth temperature and availability of nickel ions. The expression of bacterial ureases is known to be either constitutive or induced by factors like low nitrogen, urea or pH [49]. The maximal urease activity during stationary phase of the growth and at 28°C as observed for *Y. enterocolitica* biovar 1A strain was consistent with that of biovar 4 strains [47]. In *Y. enterocolitica*, several other virulence factors such as invasins, Myf fibrillae and enterotoxin have also been reported to be regulated by growth phase and the growth temperature [50]. A 10-fold increase in urease activity following supplementation of growth medium with nickel was not accompanied by increase in the expression of urease structural proteins suggesting that increased activity was probably due to the activation of pre-existing apoenzyme. Nickel has been reported to regulate both expression and activity of urease in *H. pylori* [51]. *In silico* analysis of whole genome of *Y. enterocolitica* 8081 (biovar 1B) revealed two systems (*ureH* and *ynt*) for transport of nickel. It would be interesting to determine

the role of multiple nickel transport genes in urease activity and its regulation in *Y. enterocolitica*.

The M_w of *Y. enterocolitica* biovar 1A urease as assessed from native PAGE was > 545 kDa. The molecular mass of urease is known to vary from as low as 130 kDa in *B. suis* [52] to as high as 620 kDa in *Providencia rettgeri* or > 700 kDa in *M. morganii* [53]. The difference in the molecular mass of urease of *Y. enterocolitica* biovar 1A vis-à-vis *Y. enterocolitica* biovar 1B and biovar 4 seems to be due to difference in the size of UreB (β -subunit), which is smaller in the former and thus may account for its lower molecular mass. The isoelectric point (pI) of 5.2 of biovar 1A urease was close to that reported for *Proteus penneri* (pI = 5.1) and *H. pylori* (pI = 5.9) urease [33,54]. No data on molecular mass and isoelectric point of ureases produced by *Y. enterocolitica* strains belonging to other biovars has been reported.

The ability of *Y. enterocolitica* biovar 1A strains to survive at pH 2.5 *in vitro* in the presence of 3.4 mM urea implicated urease in their survival. This suggested the possible role urease might play in the survival of *Y. enterocolitica* biovar 1A under acidic conditions in the gut. However, this needs to be confirmed by comparison of wild type strain with an isogenic urease mutant. The role of urease in survival during transit through gut has been reported for *B. suis*, *B. abortus*, *H. pylori* and *E. ictaluri* [18,19,36,55,56]. Interestingly, the biovar 1A strains have also been reported to resist killing, and survive within macrophages [13]. It would therefore be worthwhile to determine the role urease may play in the survival of *Y. enterocolitica* biovar 1A strains in the acidic environment of phagolysosomes.

Conclusions

The *ure* gene cluster of *Y. enterocolitica* biovar 1A though broadly similar to that of biovar 1B and biovar 4 strains showed differences in structural (*ureB*) genes and the intergenic regions thereof. The kinetic data indicated that urease produced by *Y. enterocolitica* biovar 1A strain would be active at low concentration of urea typically present in the gut. The ability of biovar 1A strain to survive at acidic pH in the presence of urea suggested that urease might play role in their survival in the gut. This however needs to be corroborated using *ure* isogenic mutants.

Authors' contributions

NB carried out the experimental part of the study. JSV conceived and supervised the work. Both authors participated in interpretation of data and preparation of the final manuscript.

Additional material

Additional file 1

Nucleotide and deduced amino acid sequences of ure gene cluster of *Y. enterocolitica* biovar 1A. The nucleotide sequence of the ure gene cluster of *Y. enterocolitica* biovar 1A and the deduced amino acid sequences of the structural (A, B, and C) and accessory (E, F, G and D) proteins are shown. Putative ribosome binding site consensus sequences upstream of ureA, ureB, ureC, ureF and ureG are in bold face. Stop codons are indicated by an asterisk.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2180-9-262-S1.PDF>]

Additional file 2

Phylogenetic relationships of urease structural (UreA, UreB and UreC) proteins. Dendrograms showing phylogenetic relationships of *Yersinia* spp. including *Y. enterocolitica* biovar 1A and other bacterial species based on amino acid sequence of urease structural proteins (UreA, UreB and UreC). The trees were constructed by the neighbor joining method in MEGA 4.0 package. The bootstrap values presented at corresponding branches were evaluated from 1,000 replications. GenBank accession numbers are indicated for strains used in creating the dendrogram. The bar scale shows substitutions per site.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2180-9-262-S2.PDF>]

Additional file 3

Phylogenetic relationships of urease accessory (UreE, UreF UreG and UreD) proteins. Dendrograms showing phylogenetic relationships of *Yersinia* spp. including *Y. enterocolitica* biovar 1A and other bacterial species based on amino acid sequence of urease accessory proteins (UreE, UreF, UreG and UreD). The trees were constructed by the neighbor joining method in MEGA 4.0 package. The bootstrap values presented at corresponding branches were evaluated from 1,000 replications. GenBank accession numbers are indicated for strains used in creating the dendrogram. The bar scale shows substitutions per site.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2180-9-262-S3.PDF>]

Additional file 4

PCR-RFLP of ureAB of *Y. enterocolitica*. DNA was amplified with primers ureAB3-ureAB4 and restriction digested using (A) HaeIII and (B) Sau96I enzymes. Lanes 1: IP27403, 2: IP26305, 3: E1281550, 4: P346, 5: P472, 6: IP27387, 7: STM 126, 8: 0310/90, 9: IP27938, 10: IP27879, 11: IP27873, 12: IP24121, 13: IP134, 14: IP26329, 15: IP26249, 16: 8081. M: Molecular mass marker (100 bp ladder, New England Biolabs); BV: biovar.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2180-9-262-S4.PDF>]

Additional file 5

Effect of urea/nickel chloride on activity (A) and expression (B) of urease of *Y. enterocolitica*. Strain IP27403 was grown in Luria Broth (LB) or in LB supplemented with 16.7 mM urea (LB-urea) or NiCl₂ at 1 μM (Ni-1), 10 μM (Ni-10), 100 μM (Ni-100) and 200 μM (Ni-200) concentration. M: Medium range protein ladder (Bangalore Genei). Data points represent mean of triplicate determinations; error bars denote standard deviation.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2180-9-262-S5.PDF>]

Acknowledgements

This work was supported by research grants to JSV from Indian Council of Medical Research (ICMR) and Defence Research and Development Organization (DRDO) and Senior Research Fellowship to NB from Indian Council of Medical Research (ICMR), New Delhi. The financial assistance from University of Delhi to strengthen R & D doctoral research programme is also acknowledged gratefully.

References

- Bottone EJ: **Yersinia enterocolitica: overview and epidemiologic correlates.** *Microbes Infect* 1999, **1**:323-333.
- Leclercq A, Martin L, Vergnes ML, Ounnoughene N, Laran JF, Giraud P, Carniel E: **Fatal Yersinia enterocolitica biotype 4 serovar O:3 sepsis after red blood cell transfusion.** *Transfusion* 2005, **45**:814-818.
- Cornelis GR, Boland A, Boyd AP, Geuijen C, Iriarte M, Neyt C, Sory MP, Stainier I: **The virulence plasmid of Yersinia, an antihost genome.** *Microbiol Mol Biol Rev* 1998, **62**:1315-1352.
- Revell PA, Miller VL: **Yersinia virulence: more than a plasmid.** *FEMS Microbiol Lett* 2001, **205**:159-164.
- Tennant SM, Grant TH, Robins-Browne RM: **Pathogenicity of Yersinia enterocolitica biotype IA.** *FEMS Immunol Med Microbiol* 2003, **38**:127-137.
- Morris JG Jr, Prado V, Ferreccio C, Robins-Browne RM, Bordun AM, Cayazzo M, Kay BA, Levine MM: **Yersinia enterocolitica isolated from two cohorts of young children in Santiago, Chile: incidence of and lack of correlation between illness and proposed virulence factors.** *J Clin Microbiol* 1991, **29**:2784-2788.
- Burnens AP, Frey A, Nicolet J: **Association between clinical presentation, biogroups and virulence attributes of Yersinia enterocolitica strains in human diarrhoeal disease.** *Epidemiol Infect* 1996, **116**:27-34.
- Ratnam S, Mercer E, Picco B, Parsons S, Butler R: **A nosocomial outbreak of diarrheal disease due to Yersinia enterocolitica serotype O:5, biotype I.** *J Infect Dis* 1982, **145**:242-247.
- Greenwood MH, Hooper VL: **Excretion of Yersinia spp. associated with consumption of pasteurized milk.** *Epidemiol Infect* 1990, **104**:345-350.
- Bissett ML, Powers C, Abbott SL, Janda JM: **Epidemiologic investigations of Yersinia enterocolitica and related species: sources, frequency, and serogroup distribution.** *J Clin Microbiol* 1990, **28**:910-912.
- Grant T, Bennett-Wood V, Robins-Browne RM: **Identification of virulence-associated characteristics in clinical isolates of Yersinia enterocolitica lacking classical virulence markers.** *Infect Immun* 1998, **66**:1113-1120.
- Singh I, Virdi JS: **Interaction of Yersinia enterocolitica biotype IA strains of diverse origin with cultured cells in vitro.** *Jpn J Infect Dis* 2005, **58**:31-33.
- Grant T, Bennett-Wood V, Robins-Browne RM: **Characterization of the interaction between Yersinia enterocolitica biotype IA and phagocytes and epithelial cells in vitro.** *Infect Immun* 1999, **67**:4367-4375.
- Bhagat N, Virdi JS: **Distribution of virulence-associated genes in Yersinia enterocolitica biovar IA correlates with clonal groups and not the source of isolation.** *FEMS Microbiol Lett* 2007, **266**:177-183.
- Tennant SM, Skinner NA, Joe A, Robins-Browne RM: **Homologues of insecticidal toxin complex genes in Yersinia enterocolitica biotype IA and their contribution to virulence.** *Infect Immun* 2005, **73**:6860-6867.
- McNally A, La Ragione RM, Best A, Manning G, Newell DG: **An aflagellate mutant Yersinia enterocolitica biotype IA strain displays altered invasion of epithelial cells, persistence in macrophages, and cytokine secretion profiles in vitro.** *Microbiology* 2007, **153**:1339-1349.
- Jones BD, Lockett CV, Johnson DE, Warren JW, Mobley HL: **Construction of a urease-negative mutant of Proteus mirabilis: analysis of virulence in a mouse model of ascending urinary tract infection.** *Infect Immun* 1990, **58**:1120-1123.
- Marshall BJ, Barrett LJ, Prakash C, McCallum RW, Guerrant RL: **Urea protects Helicobacter (Campylobacter) pylori from the bactericidal effect of acid.** *Gastroenterology* 1990, **99**:697-702.
- Sangari FJ, Seoane A, Rodriguez MC, Agüero J, García Lobo JM: **Characterization of the urease operon of Brucella abortus and assessment of its role in virulence of the bacterium.** *Infect Immun* 2007, **75**:774-780.
- de Koning-Ward TF, Robins-Browne RM: **Contribution of urease to acid tolerance in Yersinia enterocolitica.** *Infect Immun* 1995, **63**:3790-3795.
- Gripenberg-Lerche C, Zhang L, Ahtonen P, Toivanen P, Skurnik M: **Construction of urease-negative mutants of Yersinia enterocolitica serotypes O:3 and O:8: role of urease in virulence and arthritogenicity.** *Infect Immun* 2000, **68**:942-947.
- Sachdeva P, Virdi JS: **Repetitive elements sequence (REP/ERIC)-PCR based genotyping of clinical and environmental strains of Yersinia enterocolitica biotype IA reveal existence of limited number of clonal groups.** *FEMS Microbiol Lett* 2004, **240**:193-201.
- de Koning-Ward TF, Ward AC, Robins-Browne RM: **Characterisation of the urease-encoding gene complex of Yersinia enterocolitica.** *Gene* 1994, **145**:25-32.
- Skurnik M, Batsford S, Mertz A, Schiltz E, Toivanen P: **The putative arthritogenic cationic 19-kilodalton antigen of Yersinia enterocolitica is a urease β-subunit.** *Infect Immun* 1993, **61**:2498-2504.
- Campanella JJ, Bitincka L, Smalley J: **MatGAT: an application that generates similarity/identity matrices using protein or DNA sequences.** *BMC Bioinformatics* 2003, **4**:29.
- GeneMark [http://exon.biology.gatech.edu/genemark_prok_gms_plus.cgi]
- GeneMark.hmm [http://exon.gatech.edu/gmhmm2_prok.cgi]
- FGENESB [<http://www.softberry.com/berry.phtml>]
- NCBI ORF finder [<http://www.ncbi.nlm.nih.gov/gorf/gorf.html>]
- Gulati P, Varshney RK, Virdi JS: **Multilocus variable number tandem repeat analysis as a tool to discern genetic relationships among strains of Yersinia enterocolitica biovar IA.** *J Appl Microbiol* 2009, **107**:875-884.
- Bradford M: **A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding.** *Anal Biochem* 1976, **72**:248-254.
- McGee DJ, May CA, Garner RM, Himpls JM, Mobley HL: **Isolation of Helicobacter pylori genes that modulate urease activity.** *J Bacteriol* 1999, **181**:2477-2484.
- Mobley HL, Jones BD, Penner JL: **Urease activity of Proteus penneri.** *J Clin Microbiol* 1987, **25**:2302-2305.
- Sambrook J, Fritsch EF, Maniatis T: **Molecular cloning: a laboratory manual.** Cold Spring Harbor Laboratory Press C.S.H., New York; 1989.
- Young GM, Amid D, Miller VL: **A bifunctional urease enhances survival of pathogenic Yersinia enterocolitica and Morganella morganii at low pH.** *J Bacteriol* 1996, **178**:6487-6495.
- Booth NJ: **The role of urease in the pathogenesis of Edwardsiella ictaluri.** In *Ph.D thesis Louisiana State University, Department of Pathobiological Sciences, Baton Rouge*; 2005.
- Heermann R, Fuchs TM: **Comparative analysis of the Photorhabdus luminescens and the Yersinia enterocolitica genomes: uncovering candidate genes involved in insect pathogenicity.** *BMC Genomics* 2008, **9**:40.
- Chen YY, Clancy KA, Burne RA: **Streptococcus salivarius urease: genetic and biochemical characterization and expression in a dental plaque streptococcus.** *Infect Immun* 1996, **64**:585-592.

39. Collier JL, Brahamsha B, Palenik B: **The marine cyanobacterium *Synechococcus* sp. WH7805 requires urease (urea amidohydrolase, EC 3.5.1.5) to utilize urea as a nitrogen source: molecular-genetic and biochemical analysis of the enzyme.** *Microbiology* 1999, **145**:447-459.
40. Park IS, Hausinger RP: **Site-directed mutagenesis of *Klebsiella aerogenes* urease: identification of histidine residues that appear to function in nickel ligation, substrate binding, and catalysis.** *Protein Sci* 1993, **2**:1034-1041.
41. Jabri E, Carr MB, Hausinger RP, Karplus PA: **The crystal structure of urease from *Klebsiella aerogenes*.** *Science* 1995, **268**:998-1004.
42. Todd MJ, Hausinger RP: **Identification of the essential cysteine residue in *Klebsiella aerogenes* urease.** *J Biol Chem* 1991, **266**:24327-24331.
43. Mulrooney SB, Hausinger RP: **Sequence of the *Klebsiella aerogenes* urease genes and evidence for accessory proteins facilitating nickel incorporation.** *J Bacteriol* 1990, **172**:5837-5843.
44. Bossé JT, MacInnes JI: **Genetic and biochemical analyses of *Actinobacillus pleuropneumoniae* urease.** *Infect Immun* 1997, **65**:4389-4394.
45. Saraste M, Sibbald PR, Wittinghofer A: **The P-loop: a common motif in ATP- and GTP-binding proteins.** *Trends Biochem Sci* 1990, **15**:430-434.
46. Moncrief MB, Hausinger RP: **Characterization of UreG, identification of a UreD-UreF-UreG complex, and evidence suggesting that a nucleotide-binding site in UreG is required for *in vivo* metallocenter assembly of *Klebsiella aerogenes* urease.** *J Bacteriol* 1997, **179**:4081-4086.
47. de Koning-Ward TF, Robins-Browne RM: **A novel mechanism of urease regulation in *Yersinia enterocolitica*.** *FEMS Microbiol Lett* 1997, **147**:221-226.
48. Sebbane F, Bury-Moné S, Cailliau K, Browaey-Poly E, De Reuse H, Simonet M: **The *Yersinia pseudotuberculosis* Yut protein, a new type of urea transporter homologous to eukaryotic channels and functionally interchangeable *in vitro* with the *Helicobacter pylori* Urel protein.** *Mol Microbiol* 2002, **45**:1165-1174.
49. Mobley HL, Island MD, Hausinger RP: **Molecular biology of microbial ureases.** *Microbiol Rev* 1995, **59**:451-480.
50. Straley SC, Perry RD: **Environmental modulation of gene expression and pathogenesis in *Yersinia*.** *Trends Microbiol* 1995, **3**:310-317.
51. van Vliet AH, Kuipers EJ, Waidner B, Davies BJ, de Vries N, Penn CW, Vandenbroucke-Grauls CM, Kist M, Bereswill S, Kusters JG: **Nickel-responsive induction of urease expression in *Helicobacter pylori* is mediated at the transcriptional level.** *Infect Immun* 2001, **69**:4891-4897.
52. Contreras-Rodriguez A, Quiroz-Limon J, Martins AM, Peralta H, Avila-Calderon E, Sriranganathan N, Boyle SM, Lopez-Merino A: **Enzymatic, immunological and phylogenetic characterization of *Brucella suis* urease.** *BMC Microbiol* 2008, **8**:121.
53. Jones BD, Mobley HL: **Genetic and biochemical diversity of ureases of *Proteus*, *Providencia*, and *Morganella* species isolated from urinary tract infection.** *Infect Immun* 1987, **55**:2198-2203.
54. Mobley HL, Cortesia MJ, Rosenthal LE, Jones BD: **Characterization of urease from *Campylobacter pylori*.** *J Clin Microbiol* 1988, **26**:831-836.
55. Bandara AB, Contreras A, Contreras-Rodriguez A, Martins AM, Dobrean V, Poff-Reichow S, Rajasekaran P, Sriranganathan N, Schurig GG, Boyle SM: ***Brucella suis* urease encoded by *ureI* but not *ure2* is necessary for intestinal infection of BALB/c mice.** *BMC Microbiol* 2007, **7**:57.
56. Thune RL, Fernandez DH, Benoit JL, Kelly-Smith M, Rogge ML, Booth NJ, Landry CA, Bologna RA: **Signature-tagged mutagenesis of *Edwardsiella ictaluri* identifies virulence-related genes, including a salmonella pathogenicity island 2 class of type III secretion systems.** *Appl Environ Microbiol* 2007, **73**:7934-7946.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

