

RESEARCH ARTICLE

Open Access

Overexpression of members of the microRNA-183 family is a risk factor for lung cancer: A case control study

Wangyu Zhu^{2†}, XiaoGuang Liu^{2†}, JianYing He², DongDong Chen², YanYan Hunag² and Yong Kui Zhang^{1*}

Abstract

Background: Lung cancer is the leading cause of cancer-related deaths worldwide. Early detection is considered critical for lung cancer treatment. MicroRNAs (miRNAs) have shown promise as diagnostic and prognostic indicators. This study was to identify specific miRNAs with diagnostic and prognostic value for patients with lung cancer, and to explore the correlation between expression profiles of miRNAs and patient survival.

Methods: Gene expression of members of the miR-183 family (miR-96, miR-182, and miR-183) were examined in 70 paired samples from lung cancer patients (primary cancer and non-cancerous tissues and sera), as well as 44 serum samples from normal volunteers and lung cancer cell lines by quantitative real-time reverse transcription polymerase chain reaction (RT-qPCR). The correlation between the expression of miRNAs in tissues, sera, and patient overall survival were also examined by log-rank and Cox regression analysis.

Results: Expression levels of members of the miR-183 family in lung cancer tumor and sera were higher than that of their normal counterparts. The miR-96 expression in tumors was positively associated with its expression in sera. Log-rank and Cox regression analyses demonstrated that high expression of tumor and serum miRNAs of the miR-183 family were associated with overall poor survival in patients with lung cancer.

Conclusions: Our results suggest that the expressions of miR-96, miR-182, and miR-183 in tumor and sera may be considered potential novel biomarkers for the diagnosis and prognosis of lung cancer.

Keywords: miRNA, diagnosis, prognosis, sera, RT-quantitative PCR, human lung cancer

Background

Lung cancer is the leading cause of cancer-related deaths worldwide [1], and the current pathologic staging is inadequate to predict outcome for patient treatment. The development of molecular target therapy has improved the management of patients with lung cancer who are at high risk of relapse following surgery. One of the most promising classes of molecular markers in tumor prognosis is the small noncoding RNAs, or microRNAs (miRNAs) [2-4]. MiRNAs, acting as oncogenes or tumor suppressors, have been shown to regulate the expression of hundreds of targeted genes at the

posttranscriptional level and are implicated in the pathogenesis and therefore prognosis of human cancers [5-9]. For non-small cell lung cancer (NSCLC) in particular, previous reports have indicated that miRNA expression patterns could be potential biomarkers used for diagnosis, prognosis, and personalized therapy [4,8-11]. In addition, some studies found that human serum or plasma contained large numbers of stable miRNAs and that the expression profiles of some specific circulating miRNAs could be useful in the diagnosis and prognosis of cancer [12-14].

In our study, we performed a miRNA expression array by quantitative real-time reverse transcription polymerase chain reaction (RT-qPCR) and identified the expression of members of the miR-183 family (miR-96, miR-182, and miR-183) when comparing primary NSCLC tumor with adjacent normal lung tissues. The miR-183

* Correspondence: zyk801801@126.com

† Contributed equally

¹Department of Cardio-Thoracic Surgery, Zhoushan Hospital of Zhejiang Province, Zhoushan, Zhejiang, 316004, 316004, China

Full list of author information is available at the end of the article

family is located on human chromosome 7 and members of this family have been identified as potential oncogenes in several tumor types, including medulloblastomas [15], breast cancer [16], prostate cancer [17], hepatocellular tumors [18], and colon cancer [19,20], as well as lung cancer [21,22]. However, the role that these miRNAs play in the diagnosis and prognosis of lung cancer patients remains unknown.

We compared the expression levels of the miR-183 family in lung cancer cell lines with normal lung cells, and in primary tumor tissues and sera from NSCLC patients with normal volunteers. Our results indicate for the first time that members of the miR-183 family expressed in tumors and sera may be potential biomarkers in the diagnosis and prognosis of human lung cancer.

Methods

Sample population

NSCLC and matched adjacent noncancerous tissues were collected from patients undergoing lung resection surgery from January 2008 to May 2008 at Zhoushan Hospital, Zhejiang Province, China. The details of patients were shown in Table 1. Eligible samples were obtained from primary lung cancer that had not received any preoperative radiotherapy or chemotherapy, and in which there was no co-existing disease. Moreover, we excluded tissue blocks of mixed histology, or specimens with insufficient tumor material. The subtypes included 36 squamous-cell carcinoma and 34 adenocarcinoma. All human materials were obtained with patients' informed consent and the Ethical Review Committee of Zhoushan Municipal Government of China approved this study.

Upon removal, the surgical specimens were immediately transported to the clinical pathology laboratory, where each sample was placed in a cryovial and flash-frozen in liquid nitrogen within 30 minutes and then stored at -80°C until analyzed. Sera from all NSCLC patients and healthy volunteers were also collected. There was no significant difference in gender or age between NSCLC patients and healthy volunteers. All cases were reviewed by two pathologists and diagnoses were confirmed according to the criteria recently established by the National Comprehensive Cancer Network (NCCN).

Cell lines and cell culture conditions

The following cell lines were cultured individually in RPMI-1640 medium (Gibco): A549, H1299, and SPC-A1 human lung adenocarcinoma; 95C and 95D human giant-cell lung carcinoma; NCI-H466 human small cell lung carcinoma; NCI-H460 human large-cell lung carcinoma; and human bronchial epithelia. In addition, SK-

Table 1 Clinico-pathological Characteristics of 70 Patients with NSCLC

| Characteristics | n |
|----------------------------------|-----------|
| Mean age | |
| < 60 | 34 |
| ≥60 | 36 |
| Sex | |
| Male | 56 |
| Female | 14 |
| Smoking | |
| Nonsmokers | 24 |
| Current smokers | 46 |
| Tumor size | |
| 0-3 cm | 18 |
| > 3 cm | 52 |
| Histological classification | |
| Adenocarcinoma | 34 |
| SCC | 36 |
| Invasion to lung membrane | |
| Yes | 17 |
| No | 53 |
| differentiation | |
| Mod-well | 46 |
| Poor | 24 |
| Lymph node | |
| Negative | 38 |
| Positive | 32 |
| Stage classification | |
| Stagel | 36 |
| Stagell,III and IV | 34 |

MES1 human squamous-cell lung carcinoma cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco). Both media were supplemented with 10% fetal bovine serum (Gibco), 2 mM L-glutamine, 100 IU/mL penicillin, and 100 mg/mL streptomycin. Cells were incubated in 5% CO₂ at 37°C.

RNA isolation

Total RNA isolated from 6 NSCLC and 6 matched adjacent noncancerous tissues by the Trizol (Invitrogen) method were prepared for miRNA microarray according to the manufacturer's instructions. miRNA isolated from cells, tissues, and sera were obtained using the miRNA Isolation Kit and mirVana PARIS Kit (Applied Biosystems, Foster City, CA, USA), according to the manufacturer's protocol. The specific sources of miRNAs were: 10⁶ to 10⁷ cells (described above), 100 mg of tissue from each of 70 NSCLC patients and matched adjacent noncancerous tissues, and circulating miRNAs from 600 μL of serum from each of 70 NSCLC patients and 44 normal volunteers. The RNA concentration was measured by a NanoDrop ND-1000 spectrophotometer (nm

readings: A260/280 > 2.0, A260/230 > 1.8; NanoDrop Technologies).

MicroRNA microarray and data analysis

Three to five micrograms of total RNA were labeled by ligating the fluorescent RNA-linker 5'-cytidine bisphosphate-cyanine 3 (pCp-Cy3) to the 3' end of miRNAs. Slides were then incubated with the labeled RNA and washed. Five hundred and nine miRNAs passed the initial screening criteria of the normalized median fluorescence signal. The images were analyzed using Spot-Reader software (Niles Scientific, Portola Valley, CA). The significance of differences in expression levels was assessed by a two-sided paired *t*-test within the significance analysis of microarrays (SAM). *P* < 0.05 indicated a significant difference.

RT-qPCR of miRNA derived from lung cancer cell lines and lung primary tissues

The amounts of the miRNAs in lung tumors, analyzed by microarray, were quantified by RT-qPCR using TaqMan MicroRNA Assay Kits (Applied Biosystems, USA). Briefly, the reverse transcription (RT) reaction was carried out with a TaqMan MicroRNA Reverse Transcription Kit (Applied Biosystems) according to the instruction of the protocol. One to ten nanograms of total RNA per 15 μ L RT reaction were processed at 16°C for 30 min, 42°C for 30 min, and 85°C for 5 min. Following the RT, quantitative real-time PCR was performed in an ABI 7500 Real-Time PCR system (Applied Biosystems) at 95°C for 10 min, followed by 40 cycles of 95°C for 15 s and 60°C for one minute.

The cycle threshold (Ct) values were calculated with SDS 2.0.1 software (Applied Biosystems). The average expression levels of miRNAs in tissues and sera were normalized with U6 small nuclear RNA (snRNA) and U48 snRNA using the $2^{-\Delta\Delta C_t}$ method [23]. The mean Ct value in three candidate miRNAs was calculated, excluding outliers (i.e., replicates with a Ct differing by > one cycle from the median). If $Ct_{U6, ave}$ and $Ct_{U48, ave}$ were not within 20 and 32 cycles, the assay was repeated. Samples with low U6 or U48 snRNA levels were not included in this study.

Statistical analyses

The statistical analyses were performed with Graphpad Prism 5.0 statistical software. The data were examined according to the degree of homogeneity. The paired *t*-test, unpaired *t*-test, or Mann-Whitney *U*-test was used to analyze the correlation between the miRNA expression levels and clinical-pathological features of the patients. The paired sample *t*-test was used to compare the differences in miRNA expression between lung tissues and sera. All data were expressed as the mean \pm

standard error of the mean (SEM). Binary logistic regression was used to assess associations between the miR-183 family and the clinical-pathological features of the patients. Survival analysis was performed with the Kaplan-Meier method, and the log-rank test was used to compare survival times between groups and was adjusted according to age and tumor stages. The Cox hazard regression model was used to analyze the miRNAs as risk factors for lung cancer. A probability (*P*)-value < 0.05 was considered statistically significant.

Results


Expression of the miR-183 family of miRNAs in lung tissues, sera, and cell lines

MicroRNA arrays were performed from 6 lung cancer and 6 matched adjacent normal lung tissues that contained 723 human miRNA probes (from version 10.0 of miRBase sequence database, <http://www.mirbase.org/>). A total of 94 miRNAs were differentially expressed between lung carcinoma and the normal counterpart. Among those, we selected for further study 3 cluster miRNAs, namely miR-96, miR-182, and miR-183, based on the magnitude of fold changes (10.77-, 10.75-, and 6.62-fold, respectively) and probability values (*P* = 0.00127, 0.00209 and 0.00190), as shown in our previous study [24].

We validated the three miRNAs in 70 pairs of lung carcinoma and the corresponding noncancerous lung tissues by RT-qPCR. We showed that, compared with the adjacent normal lung tissues, the levels of the mature forms of miR-96, miR-182, and miR-183 were significantly increased in lung cancer tissues (*P* < 0.0001 for each; Figure 1A). The highly expressed (> 2-fold changes) miR-96, miR-182 and miR-183 were observed in 42 (60.0%), 54 (77.1%) and 48 (68.6%) of the 70 samples, respectively.

Compared to the sera of normal volunteers, the expression levels of miR-96, miR-182, and miR-183 in the sera of NSCLC patients were significantly higher in 62 (88.6%), 38 (54.3%), and 32 (45.7%) of the 70 samples (*P* < 0.0001, = 0.0130, and = 0.0086, respectively; Figure 1B). A correlation was also found between the serum and tumor levels of miR-96 (*Pearson r* = 0.419, *P* = 0.0003), but not for miR-182 (*Pearson r* = 0.0464, *P* = 0.703) or miR-183 (*Pearson r* = 0.118, *P* = 0.332). This suggested that serum levels of miR-96, but not miR-182 and miR-183, might correspond to the levels of tumor miR-96 (Figure 2).

Finally, we tested levels of miRNAs in different types of lung cancer vs. normal cell lines. The results from RT-qPCR were expressed as fold changes in each miRNA in lung cancer cells relative to normal cells. As shown in Figure 3, the expression levels of miR-96 were higher in 8 lung cancer cell lines. High levels of miR-




182 were found in H466 and A549 cells, and high levels of miR-183 in H466 and SK-MES1. The highest expression levels for the three miRNAs were found in NCI-H466 cells. In addition, the expression levels of all three were higher in A549 and 95D cells than in H1299 and 95C cells.



Correlation between miR-183 family and clinical-pathological features of NSCLC

We also analyzed with the Mann-Whitney *U*-test the correlation between miRNAs and clinical-pathological features of NSCLC (including age, gender, smoking history, histological classification, lymph node metastasis, differentiation, and clinical-pathological stage) to understand better the potential role of these miRNAs in NSCLC development and progression. Our results

showed that overexpression of miR-183 in tumors was strongly associated with lymph node metastasis, invasion of the lung membrane, and advanced clinical stage of NSCLC ($P = 0.0086$, 0.0222 and 0.0478 , respectively). Overexpression of miR-182 in tumors was positively related to invasion of the lung membrane and tumor size > 3 cm ($P = 0.0140$ and 0.0464 , respectively). In sera, miR-182 overexpression was also positively related to invasion of the lung membrane and tumor size > 3 cm ($P = 0.0222$ and 0.0351 , respectively). Compared to lung adenocarcinoma, higher expression of miR-96, miR-182, and miR-183 in tumors, and miR-96 in sera, were found in squamous cell lung carcinoma ($P = 0.0216$, 0.0190 , 0.0042 , and 0.0310 , respectively; Figure 4).

We also analyzed the association between levels of the miR-183 family and clinical-pathological features with a





binary logistic regression model. Results showed that patients over 60 years old, or with squamous-cell lung carcinoma, had higher levels of tumor miR-96. Patients with squamous-cell lung carcinoma or tumor invasive to lung membrane had higher levels of tumor miR-183. Patients with poor differentiation had higher levels of miR-96, miR-182 and miR-183 in sera (Table 2).

Correlation between expression of miRNAs of the miR-183 family and overall survival of NSCLC patients

Next, we investigated whether the expressions of miR-96, miR-182, and miR-183 were correlated with overall survival of NSCLC patients. Specifically, we used the Kaplan-Meier method, log-rank test and univariate Cox hazard regression model to analyze the correlation between overall survival of NSCLC patients and the expressions of miR-96, miR-182, and miR-183 in tumors (medians of $2^{-\Delta\Delta C_t}$ values: 6.45, 6.25, and 4.44, respectively) and in sera (medians 5.98, 2.08, and 1.84). The results showed that high expressions of miR-96, miR-182 and miR-183 in tumors and sera were independently correlated with shorter overall survival of NSCLC patients (log-rank test: $P = 0.0039, 0.0027, 0.0053$, respectively in tumors, and $P = 0.0017, 0.0025, 0.0046$ in sera; Figure 5).

We also examined if the expressions of miR-96, miR-182, and miR-183 in tumors and sera were correlated with clinical-pathological features of the patients, with hazard ratios (adjusted for sex, age and tumor stage) of 9.637, 7.163, 8.616, $P = 0.005, 0.010, 0.005$ respectively in tumors and with hazard ratios (adjusted for sex, age and tumor stage) of 5.512, 5.327, 5.972 yielded $P = 0.027, 0.030, 0.022$ respectively (Table 3) in sera. This result suggested that miR-96, miR-182 and miR-

183 in tumor and sera independently contributed to the overall survival of lung cancer patients

Discussion


Lung carcinoma is one of the most common malignancies in the world, and is the leading cause of cancer death in men and women in the United States with a dismal five-year survival rate (< 15%) [1]. Although recent advances have been made in improving diagnosis and treatment strategies, the prognosis of lung cancer patients remains unchanged and early detection is critical in improving survival duration. Studies have shown that miRNAs play an important role in tumorigenesis and they have been considered potential biomarkers for early diagnosis and prognosis in a wide variety of human cancers [6,14,25,26]. More importantly, circulating miRNAs might act as noninvasive blood-based biomarkers for cancer diagnosis [14,27]. One recent report demonstrated that miRNAs in tissues and plasma could have a critical role as molecular predictors of lung cancer development and management outcome [28].

The upregulation of members of the miR-183 family (miR-96, miR-182, miR-183) has been reported in several types of cancers [15-20]. Consistent with our results, members of the miR-183 family were elevated in lung cancer, revealed by novel rank-based statistical methods using microarrays [21,22,29]. miR-96, a known onco-miRNA, targets the tumor suppressor gene forkhead box O3 (*FOXO3*; a transcription factor that plays important roles in tumorigenesis) by binding to the 3'-UTR of *FOXO3* mRNA and thereby mediating protection against apoptosis and promoting cell survival [16,30]. Studies have also reported that miR-27a, miR-96, miR-182, and miR-183 coordinately regulated the

Table 2 Binary logistic regression analysis for an association between the expression levels of the miR-183 family and clinical-pathological features of patients

| | subset | Exp (B) (95% confidence interval) | P-value |
|-----------------------------|---------------------------------------|--------------------------------------|---------------|
| Tumor miR-96 | High, > 6.45/low, < 6.45 | | |
| Age | < 60/≥60 | 2.857 (1.080-7.559) | 0.034* |
| Histological classification | adenocarcinoma/SCC | 3.667 (1.366-9.842) | 0.010* |
| Tumor miR-183 | High, > 4.44/low, < 4.44 | | |
| Histological classification | adenocarcinoma/SCC | 3.667 (1.366-9.842) | 0.010* |
| Invasion to lung membrane | Negative/positive | 3.750 (1.072-13.121) | 0.039* |
| Serum miR-96 | High, > 5.98/low, < 5.98 | | |
| Differentiation | poor/Mod-well | 0.321 (0.114-0.905) | 0.032* |
| Serum miR-182 | High, > 2.08/low, < 2.08 | | |
| Differentiation | poor/Mod-well | 0.178 (0.059-0.537) | 0.002* |
| Serum miR-183 | High, > 1.84/low, < 1.84 | | |
| Differentiation | poor/Mod-well | 0.321 (0.114-0.905) | 0.032* |

Binary logistic regression analysis. *Indicates a significant difference ($P < 0.05$).



expression of forkhead box O1 (*FOXO1*) by directly targeting the mRNA *FOXO1* 3'-UTR [31,32]. In addition, miR-183 functioned as an onco-miRNA, shown to target the tumor suppressor transcription factor early growth

response 1 (*EGRI*), and phosphatase and tensin homolog (*PTEN*) to promote tumor cell migration [19].

Consistent with other reports [6,19,32], we showed that compared to adjacent non-cancerous lung tissues

Table 3 Univariate Cox hazard regression analysis for prognostic factors

| | subset | Adjusted Hazard ratio (95% confidence interval) | P-value |
|----------------------------------|---------------------------------------|--|----------------|
| Gender | male/female | 0.031 (0.000-5.569) | 0.190 |
| Age | < 60/≥60 | 1.630 (0.564-4.716) | 0.367 |
| Smoking | nonsmoker/current smoker | 1.388 (0.306-6.297) | 0.671 |
| Tumor size | 0-3 cm/> 3 cm | 0.766 (0.239-2.455) | 0.654 |
| Histological classification | adenocarcinoma/SCC | 2.739 (0.781-9.604) | 0.115 |
| Invasion to lung membrane | Negative/positive | 1.000 (0.221-4.534) | 1.000 |
| Differentiation | Mod-well/poor | 0.757 (0.182-3.141) | 0.701 |
| Lymph node | Negative/positive | 1.279 (0.425-3.850) | 0.661 |
| Stage classification | stage I/stage II, III, IV | 1.001 (0.334-2.996) | 0.999 |
| Tumor miR-96 | High, > 6.45/low, < 6.45 | 9.637 (1.963-47.315) | 0.005 * |
| Tumor miR-182 | High, > 6.25/low, < 6.25 | 7.163 (1.597-32.138) | 0.010 * |
| Tumor miR-183 | High, > 4.44/low, < 4.44 | 8.616 (1.918-38.705) | 0.005* |
| Serum miR-96 | High, > 5.98/low, < 5.98 | 5.512 (1.219-24.915) | 0.027 * |
| Serum miR-182 | High, > 2.08/low, < 2.08 | 5.327 (1.181-24.030) | 0.030 * |
| Serum miR-183 | High, > 1.84/low, < 1.84 | 5.972 (1.289-27.658) | 0.022 * |

Univariate cox hazard regression analysis. *indicates a significant difference ($P < 0.05$).

and normal volunteers, members of the miR-183 family were highly expressed in lung cancer primary tissues and sera. Moreover, we found that the expression levels of miR-96, miR-182, and miR-183 were higher in A549 and 95D cells than in H1299 and 95C cells, respectively. This implies that the miR-183 family might participate in tumor metastasis [19,30,33,34]. Consistent with this, other studies demonstrated that overexpressed miR-96 and miR-182 in melanoma cell lines and breast cancer cell lines resulted in enhanced oncogenic properties as well as invasion and metastasis *in vitro* [30,33]. Sarver *et al.* [19] showed that a reduction in miR-183 led to a decrease in migration in colon cancer and synovial sarcoma cell lines. Abraham *et al.* [34] reported that overexpression of miR-183 in medullary thyroid cancer predicted lymph node metastasis.

We note that opposite results have also been reported. According to one study, expression of miR-96 decreased cell invasion and migration in pancreatic cancer [35]. Another report demonstrated that miR-182 suppressed cell proliferation of lung cancer cell line A549 *in vitro* and tumor growth *in vivo* through its interference with the target gene cortactin (*CTTN*) by epigenetic modification [36]. Others found that overexpression of miR-183 inhibited migration and invasion of lung cancer and breast cancer cells, by targeting the protein ezrin [22,37]. These discrepancies might be due to different cancer cell lines used, and suggest that miRNA might have distinct functions depending upon cell type [38].

Our results demonstrate an association between the expressions of miR-183 and miR-182 and metastasis, invasion, advanced clinical stage, and tumor sizes of NSCLC. Together with other reports, they imply that

miR-182 and miR-183 might play a role in lung cancer invasion and metastasis, and progression of NSCLC [28,30]. Moreover, in our study patients with high tumor or serum levels of miR-96, miR-182, and miR-183 had shorter overall post-operative survival times. The hazard ratios adjusted for gender, age and tumor stage indicated that these miRNAs could be used independently for evaluating the prognoses of NSCLC patients.

In addition, our results demonstrated that serum miRNAs might be used as potential biomarkers for diagnosis and prognosis of lung cancers. Under lung cancer condition, miRNAs entered the circulation mainly from tumor cells [27]. One study showed that miRNAs in tissue specimens were not detected by Solexa sequencing in sera [39]. By comparing the expressions of miR-96, miR-182 and miR-183 in tumors and sera, our results suggest that the levels of serum and tumor miR-96 were correlated, while those of miR-182 and miR-183 were not. These results are consistent with reports from Mitchell [27] and Hu [39]. The actual role and functions of these miRNAs need to be further investigated.

Conclusions

Our results show that members of the miR-183 family, found in tumors and sera, may play a role in the development of NSCLC and have potential as biomarkers in the diagnosis and prognosis of lung cancer. We further demonstrate a positive correlation between expression levels of miR-96 in tumors and sera from NSCLC patients. Further study needs to be done to define the true value of combined expression profiles of the miR-183 family in tumors and sera in the diagnosis and prognosis of lung cancer.

List of abbreviations

Ct: cycle threshold; **CTTN**: cactactin; **DMEM**: Dulbecco's modified Eagle's medium; **EGR1**: early growth response 1; **FOXO3**: forkhead box O3; **FOXO1**: forkhead box O1; **miRNA**: microRNA; **NCCN**: National Comprehensive Cancer Network; **NSCLC**: non-small cell lung cancer; **pCp-Cy3**: 5'-cytidine bisphosphate-cyanine 3; **PTEN**: phosphatase and tensin homolog; **RT**: reverse transcription; **qRT-PCR**: quantitative real-time reverse transcription polymerase chain reaction; **SAM**: significance analysis of microarrays; **SEM**: standard error of the mean; **snRNA**: small nuclear RNA.

Acknowledgements

This research was supported by a grant from the Zhejiang Provincial Natural Science Foundation of China (No. Y2101391), a grant from the Medical Bureau of Zhejiang Province (No. 2009A210), and grants from the Science and Technology Bureau of Zhoushan (No. 20081059, No.091042 and No.101137) and from the Medical Bureau of Zhoushan (No. 2009B03). The authors would like to thank Medjaden Bioscience Limited for assisting in the preparation of this manuscript.

Author details

¹Department of Cardio-Thoracic Surgery, Zhoushan Hospital of Zhejiang Province, Zhoushan, Zhejiang, 316004, 316004, China. ²Joint Laboratory of Immunogenomics, Zhoushan Hospital-BIG/CAS, Zhoushan, Zhejiang Province, 316004, China.

Authors' contributions

WYZ, XGL, and YKZ were involved in the conception and design of the study. WYZ, JYH, DDC, and YYH were involved in the provision of study material and patients. WYZ and XGL performed the data analysis and interpretation. WYZ wrote the manuscript. YKZ approved the final version. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 17 June 2011 Accepted: 15 September 2011

Published: 15 September 2011

References

- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ: **Cancer statistics, 2009.** *CA-Cancer J Clin* 2009, **59**:225-249.
- Jay C, Nemunaitis J, Chen P, Fulgham P, Tong AW: **miRNA profiling for diagnosis and prognosis of human cancer.** *DNA Cell Biol* 2007, **26**:293-300.
- Cummins J, Velculescu V: **Implications of micro-RNA profiling for cancer diagnosis.** *Oncogene* 2006, **25**:6220-6227.
- Yu SL, Chen HY, Chang GC, Chen CY, Chen HW, Singh S, Cheng CL, Yu CJ, Lee YC, Chen HS: **MicroRNA signature predicts survival and relapse in lung cancer.** *Cancer Cell* 2008, **13**:48-57.
- Calin GA, Croce CM: **MicroRNA-cancer connection: the beginning of a new tale.** *Cancer Res* 2006, **66**:7390-7394.
- Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, Visone R, Iorio M, Roldo C, Ferracin M: **A microRNA expression signature of human solid tumors defines cancer gene targets.** *Proc Natl Acad Sci* 2006, **103**:2257-2261.
- Rosell R, Wei J, Taron M: **Circulating microRNA signatures of tumor-derived exosomes for early diagnosis of non-small cell lung cancer.** *Clin Lung Cancer* 2009, **10**:8-9.
- Ventura A, Jacks T: **MicroRNAs and cancer: short RNAs go a long way.** *Cell* 2009, **136**:586-591.
- Yanaihara N, Caplen N, Bowman E, Seike M, Kumamoto K, Yi M, Stephens RM, Okamoto A, Yokota J, Tanaka T: **Unique microRNA molecular profiles in lung cancer diagnosis and prognosis.** *Cancer cell* 2006, **9**:189-198.
- Raponi M, Dossey L, Jatko T, Wu X, Chen G, Fan H, Beer DG: **MicroRNA classifiers for predicting prognosis of squamous cell lung cancer.** *Cancer Res* 2009, **69**:5776-5783.
- Galluzzi L, Morselli E, Vitale I, Kepp O, Senovilla L, Criollo A, Servant N, Paccard C, Hupe P, Robert T: **MiR-181a and miR-630 regulate cisplatin-induced cancer cell death.** *Cancer Res* 2010, **70**:1793-1803.
- Zhao H, Shen J, Medico L, Wang D, Ambrosone CB, Liu S, Creighton C: **A pilot study of circulating miRNAs as potential biomarkers of early stage breast cancer.** *PLoS One* 2010, **5**:876-887.
- Resnick KE, Alder H, Hagan JP, Richardson DL, Croce CM, Cohn DE: **The detection of differentially expressed microRNAs from the serum of ovarian cancer patients using a novel real-time PCR platform.** *Gynecol oncol* 2009, **112**:55-59.
- Chen X, Ba Y, Cai X, Yin Y, Wang K, Guo J, Zhang Y, Chen J, Guo X: **Characterization of microRNAs in serum: a novel class of biomarkers for diagnosis of cancer and other diseases.** *Cell Res* 2008, **18**:997-1006.
- Gokhale A, Kunder R, Goel A, Sarin R, Moiyadi A, Shenoy A, Mamidipally C, Noronha S, Kannan S, Shirsat N: **Distinctive microRNA signature of medulloblastomas associated with the WNT signaling pathway.** *J Cancer Res Ther* 2010, **6**:521-529.
- Lin P, Yu S, Yang P: **MicroRNA in lung cancer.** *Br J cancer* 2010, **103**:1144-1148.
- Schaefer A, Jung M, Mollenkopf HJ, Wagner I, Stephan C, Jentzmik F, Miller K, Lein M, Kristiansen G, Jung K: **Diagnostic and prognostic implications of microRNA profiling in prostate carcinoma.** *Int J Cancer* 2010, **126**:1166-1176.
- Pineau P, Volinia S, McLunkin K, Marchio A, Battiston C, Terris B, Mazzaferro V, Lowe SW, Croce CM, Dejean A: **miR-221 overexpression contributes to liver tumorigenesis.** *Proc Natl Acad Sci* 2010, **107**:264-269.
- Sarver AL, Li L, Subramanian S: **MicroRNA miR-183 functions as an oncogene by targeting the transcription factor EGR1 and promoting tumor cell migration.** *Cancer Res* 2010, **70**:9570-9580.
- Bandres E, Cubedo E, Agirre X, Malumbres R, Zarate R, Ramirez N, Abajo A, Navarro A, Moreno I, Monzo M: **Identification by Real-time PCR of 13 mature microRNAs differentially expressed in colorectal cancer and non-tumoral tissues.** *Mol Cancer* 2006, **5**:29.
- Navon R, Wang H, Steinfeld I, Tsalenko A, Ben-Dor A, Yakhini Z: **Novel rank-based statistical methods reveal microRNAs with differential expression in multiple cancer types.** *PLoS one* 2009, **4**:e8003.
- Wang J, Chen J, Chang P, LeBlanc A, Li D, Abbruzzese JL, Frazier ML, Killy AM, Sen S: **MicroRNAs in plasma of pancreatic ductal adenocarcinoma patients as novel blood-based biomarkers of disease.** *Cancer Prev Res* 2009, **2**:807.
- Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-Delta Delta C} (T_T) method.** *Methods* 2001, **25**:402-408.
- Liu X, Zhu W, Huang Y, Ma L, Zhou S, Wang Y, Zeng F, Zhou J, Zhang Y: **High expression of serum miR-21 and tumor miR-200c associated with poor prognosis in patients with lung cancer.** *Med Oncol* 2011.
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, Sweet-Cordero A, Ebert BL, Mak RH, Ferrando AA: **MicroRNA expression profiles classify human cancers.** *Nature* 2005, **435**:834-838.
- Asaga S, Kuo C, Nguyen T, Terpenning M, Giuliano AE, Hoon DSB: **Direct serum assay for microRNA-21 concentrations in early and advanced breast cancer.** *Clin Chem* 2011, **57**:84-91.
- Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanian EL, Peterson A, Noteboom J, O'Brian KC, Allen A: **Circulating microRNAs as stable blood-based markers for cancer detection.** *Proc Natl Acad Sci* 2008, **105**:10513-10518.
- Boeri M, Verri C, Conte D, Roz L, Modena P, Facchinetti F, Calabr E, Croce CM, Pastorino U, Sozzi G: **MicroRNA signatures in tissues and plasma predict development and prognosis of computed tomography detected lung cancer.** *Proc Natl Acad Sci* 2011, **108**:3713-3718.
- Barshack I, Lithwick-Yanai G, Afek A, Rosenblatt K, Tabibian-Keissar H, Zepeniuk M, Cohen L, Dan H, Zion O, Strenov Y: **MicroRNA expression differentiates between primary lung tumors and metastases to the lung.** *Pathol Res Pract* 2010, **206**:578-584.
- Segura MF, Hanniford D, Menendez S, Reavie L, Zou X, Alvarez-Diaz S, Zakrzewski J, Blochin E, Rose A, Bogunovic D: **Aberrant miR-182 expression promotes melanoma metastasis by repressing FOXO3 and microphthalmia-associated transcription factor.** *Proc Natl Acad Sci* 2009, **106**:1814-1819.
- Guttilla IK, White BA: **Coordinate regulation of FOXO1 by miR-27a, miR-96, and miR-182 in breast cancer cells.** *J Biol Chem* 2009, **284**:23204-23216.
- Myatt SS, Wang J, Monteiro LJ, Christian M, Ho KK, Fusi L, Dina RE, Brosens JJ, Ghaem-Maghani S, Lam EWF: **Definition of microRNAs that**

- repress expression of the tumor suppressor gene FOXO1 in endometrial cancer. *Cancer Res* 2010, **70**:367-377.
33. Li J, Fu H, Xu C, Tie Y, Xing R, Zhu J, Qin Y, Sun Z, Zheng X: miR-183 inhibits TGF- β 1-induced apoptosis by downregulation of PDCD 4 expression in human hepatocellular carcinoma cells. *BMC cancer* 2010, **10**:354.
 34. Abraham D, Jackson NE, Gundara JS, Zhao J, Gill AJ, Delbridge L, Robinson B, Sidhu S: MicroRNA profiling of sporadic and hereditary medullary thyroid cancer identifies predictors of nodal metastasis, prognosis and potential therapeutic targets. *Clin Cancer Res* 2011, **17**:4772-4781.
 35. Yu S, Lu Z, Liu C, Meng Y, Ma Y, Zhao W, Liu J, Yu J, Chen J: miRNA-96 suppresses KRAS and functions as a tumor suppressor gene in pancreatic cancer. *Cancer Res* 2010, **70**:6015-6025.
 36. Zhang L, Liu T, Huang Y, Liu J: microRNA-182 inhibits the proliferation and invasion of human lung adenocarcinoma cells through its effect on human cortical actin-associated protein. *Int J Mol Med* 2011, **28**:381-383.
 37. Lowery AJ, Miller N, Dwyer RM, Kerin MJ: Dysregulated miR-183 inhibits migration in breast cancer cells. *BMC cancer* 2010, **10**:502.
 38. Hyun S, Lee JH, Jin H, Nam JW, Namkoong B, Lee G, Chung J, Kim VN: Conserved MicroRNA miR-8/miR-200 and its target USH/FOG2 control growth by regulating PI3K. *Cell* 2009, **139**:1096-1108.
 39. Hu Z, Chen X, Zhao Y, Tian T, Jin G, Shu Y, Chen Y, Xu L, Zen K, Zhang C: Serum microRNA signatures identified in a genome-wide serum microRNA expression profiling predict survival of non-small cell lung cancer. *J Clin Oncol* 2010, **28**:1721-1726.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2407/11/393/prepub>

doi:10.1186/1471-2407-11-393

Cite this article as: Zhu et al.: Overexpression of members of the microRNA-183 family is a risk factor for lung cancer: A case control study. *BMC Cancer* 2011 **11**:393.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

