

Research article

Open Access

Polymorphisms in estrogen-metabolizing and estrogen receptor genes and the risk of developing breast cancer among a cohort of women with benign breast disease

Lisa Gallicchio*^{1,2}, Sonja I Berndt¹, Meghan A McSorley¹,
Craig J Newschaffer¹, Lucy W Thuita¹, Pedram Argani³, Sandra C Hoffman¹
and Kathy J Helzlsouer^{1,2}

Address: ¹Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA, ²Prevention and Research Center, The Weinberg Center for Women's Health and Medicine, Mercy Medical Center, Baltimore, Maryland, USA and ³Department of Pathology, Johns Hopkins School of Medicine, Baltimore, Maryland, USA

Email: Lisa Gallicchio* - lgallic@mdmercy.com; Sonja I Berndt - sberndt@jhsp.h.edu;
Meghan A McSorley - mcsorley.meghan@medstudent.pitt.edu; Craig J Newschaffer - cnewscha@jhsp.h.edu; Lucy W Thuita - lthuita@jhsp.h.edu;
Pedram Argani - pargani1@jhmi.edu; Sandra C Hoffman - shoffman@jhsp.h.edu; Kathy J Helzlsouer - khelzso@mdmercy.com

* Corresponding author

Published: 29 June 2006

Received: 24 February 2006

BMC Cancer 2006, 6:173 doi:10.1186/1471-2407-6-173

Accepted: 29 June 2006

This article is available from: <http://www.biomedcentral.com/1471-2407/6/173>

© 2006 Gallicchio et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: A cohort study was conducted to examine the role of genetic polymorphisms in three estrogen metabolizing enzymes (*COMT*, *CYP1A1*, *CYP1B1*) and the two estrogen receptors (*ESR1*, *ESR2*) in the progression of benign breast disease (BBD) to breast cancer.

Methods: Among participants in an ongoing cohort study, 1438 Caucasian women had a breast biopsy for BBD and were successfully genotyped for at least one of the polymorphisms examined in this study. Genotypes were determined using DNA extracted from blood specimens collected in 1989. Incident cases of breast cancer occurring subsequent to BBD diagnosis up to 2003 were identified through cancer registries.

Results: Among all participants, the *ESR2* *5772G allele was associated with a significant decrease in the risk of breast cancer among women with BBD (Odds Ratio (OR) 0.38; 95% Confidence Interval (CI) 0.15, 0.96). Compared to the reference wild-type genotypes, marginally significant associations with the development of breast cancer were observed between carriers of the variant *ESR1* – 104062T allele (OR 0.70, 95% CI 0.45, 1.09), the variant *ESR2* *38A allele (OR 1.40; 95% CI 0.88, 2.25), and the variant *CYP1B1* 453Ser allele (OR 1.48, 95% CI 0.95, 2.32).

Conclusion: The results indicate that specific polymorphisms in the *CYP1B1*, *ESR1*, and *ESR2* genes may play a role in progression of BBD to breast cancer among Caucasian women. Although additional studies are needed to confirm or refute our findings, these results suggest that genetic markers may aid in the identification of women who are at risk for progression of BBD to cancer.

Background

Biopsy-proven benign breast disease (BBD) is a well-defined risk factor for developing invasive breast cancer [1,2]. However, little is known about which women with BBD progress to have a subsequent *in situ* or invasive cancer. A woman's level of endogenous estrogen may be important in the progression of BBD to invasive breast cancer. Abundant evidence suggests that estrogen is an important factor in the development of breast cancer among average-risk women. This evidence includes findings from recent prospective studies that showed significantly increased levels of serum estrogen in women who developed breast cancer compared to women who did not develop breast cancer [3-7]. In addition, numerous studies have reported that risk factors associated with prolonged estrogen exposure, such as early age at menarche, nulliparity, and older age at first birth, increase a woman's risk of developing breast cancer (reviewed by Mitrunen and Hirvonen [8]).

Because of the importance of estrogen in the development of breast cancer among average-risk women, it is plausible that polymorphisms in estrogen metabolizing genes may be associated with the risk of developing breast cancer among women with BBD. Estrogens undergo oxidative metabolism by several cytochrome P450 (CYPs) enzymes [9]; the major metabolic pathways are 2-hydroxylation, which is primarily catalyzed by *CYP1A1*, and 4-hydroxylation, which is dominated by *CYP1B1* [10]. The hydroxylated estrogens are then either converted to semiquinones or quinones, which can produce DNA adducts and lead to oxidative damage to lipids and DNA, or inactivated by catechol O-methyltransferase (*COMT*) [10-13]. Within the genes encoding for *CYP1A1*, *CYP1B1*, and *COMT*, a number of polymorphisms have been identified, some which have been shown in laboratory studies to be functional (8). The associations of some of these polymorphisms with the risk of developing breast cancer among average risk women have been studied, and most of the findings have been inconsistent (reviewed in Mitrunen and Hirvonen [8]). No study has examined the associations of polymorphisms in *CYP1A1*, *CYP1B1*, and *COMT* with breast cancer risk among women with BBD.

While the estrogen metabolizing enzymes may influence the levels of circulating endogenous estrogens and, therefore, contribute to breast cancer risk, the estrogen receptors determine, in part, the action of estrogens on the mammary gland. Estrogen acts on target tissues by binding to the estrogen receptors, which exist in two forms, ER- α or *ESR1* and ER- β or *ESR2* [14,15]. Like the estrogen metabolizing enzymes, a number of polymorphisms in both ER genes have been identified [16], and it has been hypothesized that these polymorphisms may increase or decrease a woman's breast cancer risk depending on the

functional consequence of the polymorphism. Several studies have investigated the associations between polymorphisms in the *ESR1* and *ESR2* genes and the risk of breast cancer among average-risk women and have reported mixed results [16-29]. To our knowledge, however, no study has examined the associations of polymorphisms in *ESR1* and *ESR2* with breast cancer risk among women with BBD.

The aim of the present study was to analyze the associations between genetic polymorphisms in three estrogen metabolizing enzymes (*COMT*, *CYP1A1*, *CYP1B1*) and the two estrogen receptors (*ESR1*, *ESR2*) and the risk of developing breast cancer among women with BBD. To address this aim, data from a cohort of women with BBD from Washington County, Maryland enrolled in the CLUE-II study were analyzed.

Methods

CLUE II cohort

In 1989, a campaign named CLUE II ("*Campaign against Cancer and Heart Disease*") was conducted in Washington County, Maryland to establish a cohort with associated blood samples [30]. At baseline, participants signed a consent form, completed a brief medical and exposure history and a food frequency questionnaire, and donated a blood sample. The blood sample was collected in heparinized tubes and centrifuged to obtain separation of the buffy coat which was then aliquotted and stored at -70°C. DNA was subsequently extracted from the buffy coat using standard phenol extraction [31].

A total of 25,081 persons (10,456 men and 14,625 women), about 30 percent of the Washington County residents, participated in the study. In addition to the information collected from this cohort at baseline in 1989, follow-up data were obtained from questionnaires administered in 1996 and approximately every 2 years thereafter. The Institutional Review Board at the Johns Hopkins Bloomberg School of Public Health approved this study.

BBD cohort

The BBD cohort is a sub-set of CLUE II. Details of the selection of CLUE II participants included in the BBD cohort are described elsewhere [32]. Briefly, women who responded on the 1996 CLUE II follow-up questionnaire that they had undergone a "breast biopsy or lumpectomy" were considered for inclusion into the BBD cohort. Women were included in the BBD cohort if either they were found to have BBD on their first located pathology report or a pathology report was not found but a breast biopsy was reported prior to or in the absence of a subsequent breast cancer diagnosis. Women who were diagnosed with breast cancer at their first biopsy, as reported

by the participant or as determined using the pathology report search, were excluded from this analysis. In total, of the 1683 women who reported that they had a breast biopsy, 1467 women (87.2%) met the criteria above and were included in the BBD cohort. Among these 1467 women, pathology report data were collected for 362 (24.7%). Seventy percent of these pathology reports indicated a diagnosis of non-proliferative BBD, 26% indicated a diagnosis of proliferative disease without atypia, and 4% reported a diagnosis of proliferative disease with atypia.

For this analysis, 17 of the 1467 women were excluded because DNA could not be obtained from their stored blood samples and polymorphisms could not be genotyped. Further, all non-Caucasians ($n = 12$) were excluded from the analysis because previous studies have shown that race is an important confounder or effect modifier in investigations of polymorphisms and disease. These exclusions accounted for less than 1% of the BBD cohort. After applying these exclusions, the final analytic cohort included 1438 Caucasian women who were successfully genotyped for at least one of the genetic polymorphisms investigated in this study.

Outcome measurement

Incident cases of *in situ* or invasive breast cancer (International Classification of Diseases, Ninth Revision, code 233, 174) were identified through linkage of the cohort participants with the Washington County Cancer Registry and, since 1992, with the Maryland State Cancer Registry. The Washington County Cancer Registry identifies cancer cases from discharge records and pathology reports from Washington County Hospital, which is the only hospital in the county, and from death certificates. Since 1992, all hospitals cancer diagnostic laboratories, and radiation therapy centers in Maryland have been required to report incident cancer cases to the Maryland Cancer Registry, and

currently, the Maryland Cancer Registry is certified as being more than 95% complete by the North American Association of Central Cancer Registries. While it is possible that cases of *in situ* or invasive breast cancer may have been missed due to migration out of the state after 1996, all women in the BBD cohort had at least 7 years of follow-up data (1989 to 1996). Further, approximately 80% of the BBD cohort responded to the 2000 follow-up questionnaire, indicating that the vast majority of the cohort remained under active surveillance in 2000.

Active case surveillance for the BBD cohort ended on April 28, 2003. In total, among all of the women included in the BBD cohort, 15 incident cases of *in situ* and 76 cases of invasive breast cancer were identified after the diagnosis of BBD.

Genotyping

Single nucleotide polymorphisms (SNPs) analyzed in the present study were selected as part of a larger study investigating the broad impact of genetic variation in candidate genes and their interactions with environmental exposures on cancer incidence and survival. For the larger study, polymorphisms that were suggested to alter function, encoded for a nonsynonymous amino acid change, or were located within the 5' or 3' untranslated region (UTR) of the gene and thus could potentially alter mRNA stability were chosen. Descriptions and dbSNP identifiers of the polymorphisms selected are shown on Table 1. Genotyping of the chosen SNPs in *COMT*, *CYP1B1*, *CYP1A1*, *ESR1* and *ESR2* was conducted at Applied Biosystems (ABI) and Celera Laboratories. All polymorphisms were genotyped using TaqMan technology (ABI, Foster City, CA, USA). The genotyping success rates of the polymorphisms selected ranged from 91.8% to 97.6%, with the exception of the *CYP1B1* Arg48Gly polymorphism for which the genotyping success rate was 84.2%.

Table 1: Description of single nucleotide polymorphisms examined in the progression of BBD to breast cancer

Gene	dbSNP rs #	snp description	Amino Acid Substitution
<i>COMT</i>	4680	Ex3-12G>A	Val158Met
<i>CYP1A1</i>	1048943	Ex7+131A>G	Ile462Val
<i>CYP1B1</i>	1800440	Ex3+315A>G	Asn453Ser
<i>CYP1B1</i>	10012	Ex2+143C>G	Arg48Gly
<i>CYP1B1</i>	1056836	Ex3+251G>C	Val432Leu
<i>ESR1</i>	1801132	Ex4-122G>C	Pro325Pro
<i>ESR1</i>	2077647	Ex1+392T>C	Ser10Ser
<i>ESR1</i>	746432	Ex1-192G>C	Ala87Ala
<i>ESR1</i>	2273206	IVS6+52G>T	
<i>ESR1</i>	851982	-104062C>T	
<i>ESR1</i>	2228480	Ex8+229G>A	Thr594Thr
<i>ESR2</i>	4986938	38 bp 3' of STP C>T	
<i>ESR2</i>	928554	5659 bp 3' of STP G>A	
<i>ESR2</i>	none	5696 bp 3' of STP A>G	
<i>ESR2</i>	8018687	5772 bp 3' of STP A>G	

The 84.2% genotyping success rate for *CYP1B1* Arg48Gly polymorphism was due to a problem with the laboratory assay that was not resolved during the study period.

Statistical analysis

The outcome variable examined in these analyses was the development of *in situ* or invasive breast cancer. Unconditional logistic regression was used to estimate age-adjusted odds ratios (ORs) and 95% confidence intervals (95% CIs) for the associations between the polymorphisms the development of breast cancer. Because previous studies indicate that the associations between several of the analyzed polymorphisms and the development of cancer may differ according to body mass index, smoking status, and alcohol intake, we conducted exploratory analyses to examine the associations between all of the polymorphisms and the development of breast cancer among strata of these variables (BMI: <25 kg/m², ≥25 kg/m²; smoking status: ever, never; alcohol intake: current, no current use). Further, these analyses were conducted for all participants and for pre- and post-menopausal participants separately. Because the number of events among the pre-menopausal women was low (n = 24) and ORs for some of the genotypes could not be calculated, only the analyses for all participants and the post-menopausal subgroup are reported. As an additional exploratory analysis, logic regression [33] was employed using the software package R 2.0.1 to examine whether any interactions between the genotypes existed.

To address the issue of multiple testing in this study, p-values for the associations between single nucleotide polymorphisms and breast cancer risk were adjusted for the false discovery rate using the method proposed by Benjamini and Hochberg [34] in the software package R 2.0.1.

Unless otherwise specified, analyses were performed using SAS Version 8.2 (Cary, NC). A p-value of less than 0.05 was considered to be statistically significant.

Results

Characteristics of the analytic cohort are reported in Table 2. The mean age of the participants was 53.5 years (standard deviation 12.1), and the majority of the women had at least a high school degree (82.2%). Approximately 88% of the women had been pregnant at least once in their lifetime, 16.2% were current smokers, and 28.8% were current drinkers. Among the 1438 women in this sample, 91 (6.3%) developed invasive or *in situ* breast cancer after an initial breast biopsy for BBD.

All of the polymorphisms examined were in Hardy-Weinberg equilibrium among women not developing breast cancer, with the exception of the *CYP1B1* Ile462Val polymorphism (p-value = 0.02). Although statistically signifi-

cant, the observed genotype counts for the *CYP1B1* Ile462Val polymorphism were similar to the expected genotype counts under Hardy-Weinberg equilibrium (observed/expected: AA-1233/1229.5; AG-80/86.9; GG-5/1.5). All *ESR2* polymorphisms were in strong linkage disequilibrium ($D' > 90$). Among the *ESR1* polymorphisms, the *Ser10Ser* and *Ala87Ala* polymorphisms were strongly linked ($D' > 90$).

The associations between polymorphisms in the estrogen metabolizing enzymes and the risk of developing invasive breast cancer among women with BBD are shown in Table 3. Among all participants and post-menopausal participants only, carriers of at least one *CYP1B1* 453Ser allele had a borderline statistically significant increase in the risk of developing breast cancer compared to women carrying the referent *Asn/Asn* genotype (all participants: OR 1.48, 95% CI 0.95, 2.32; post-menopausal participants: OR 1.61; 95% CI 0.94, 2.74). The other polymorphisms in the estrogen metabolizing enzymes were not associated with breast cancer risk among women with BBD in this study. Results were not substantially altered by BMI, alcohol intake or smoking status (results not shown). No significant interactions between the genotypes were observed. To note, after adjustment for multiple testing, none of the associations between polymorphisms in the estrogen metabolizing enzymes and the risk of developing invasive breast cancer were statistically significant.

The associations between polymorphisms in *ESR1* and *ESR2* and the risk of developing invasive breast cancer among women with BBD are shown in Tables 4 and 5. A marginally significant decrease in the risk of developing breast cancer was observed among carriers of the *ESR1* – 104062T allele compared to carriers of the referent genotype (all participants: OR 0.70, 95% CI 0.45, 1.09). Among all participants, the *ESR2* *5772G allele was associated with a statistically significant decrease in the risk of breast cancer among women with BBD (OR 0.38; 95% CI 0.15, 0.96). A marginally significant increase in the risk of developing cancer was observed among women carrying at least one *ESR2* *38A allele compared to the women homozygous for the GG genotype (all participants: OR 1.40; 95% CI 0.88, 2.25). OR estimates of the *ESR2* *5772G and the *ESR2* *38A polymorphisms were attenuated after adjustment for the presence of the other polymorphism (*ESR2* *5772G OR 0.42; 95% CI 0.17, 1.05; *ESR2* *38 OR 1.27; 95% CI 0.78, 2.04). Other *ESR1* and *ESR2* polymorphisms examined were not associated with breast cancer risk among women with BBD. Results did not differ substantially by BMI, alcohol intake or smoking status (results not shown). No significant interactions between the *ESR1* and *ESR2* genotypes were observed. After adjustment for multiple testing, the associations

Table 2: Characteristics of the study cohort (n = 1438)

Characteristic	n	%
Race		
Caucasian	1438	100.0
Age, years ^a		
<50	533	37.1
50 to 60	411	28.6
>60	494	34.3
Missing	0	0.0
Education, years ^a		
less than a high school degree	259	18.0
high school degree	714	49.7
some college	463	32.2
Missing	2	0.1
Menopausal status ^a		
Pre-menopausal	377	26.2
Peri-menopausal	35	2.4
Post-menopausal	995	69.2
Missing	31	2.2
Family history of breast cancer ^a		
No	715	49.7
Yes	275	19.1
Missing	448	31.2
Ever been pregnant (%) ^b		
No	176	12.2
Yes	1255	87.3
Missing	7	0.5
Age at first birth (years, %) ^b		
Nulliparous	176	12.2
20	347	24.1
20 to 24	539	37.5
≥25	339	23.6
Missing	37	2.6
Age at menarche (years, %) ^b		
<12	268	18.6
12 to 13	763	53.1
>13	386	26.8
Missing	21	1.5
Height, inches ^a		
<62	411	28.6
62–64	419	29.1
>64	608	42.3
Missing	0	0.0
Weight, pounds ^a		
<130	354	24.6
130–<145	318	22.1
145–<165	377	26.2
≥165	387	26.9
Missing	2	0.1
BMI, kg/m ² . ^a		
<24.9	744	51.7
25.0 – 29.9	433	30.1
≥30	259	18.0
Missing	2	0.1
Smoking status ^a		
Never	839	58.3
Former	366	25.5
Current	232	16.1
Missing	1	0.1
Current alcohol drinker ^a		
No	814	56.6

Table 2: Characteristics of the study cohort (n = 1438) (Continued)

Yes	413	28.7
Missing	211	14.7

^a data from 1989 questionnaire

^b data from 1996 questionnaire

between the *ESR1* and *ESR2* polymorphisms and breast cancer described above were not statistically significant.

Discussion

In this population-based cohort of women with BBD, there was evidence of an increased risk of breast cancer among women who carried the *CYP1B1* 453Ser allele compared to women who carried two copies of the *CYP1B1* 453Asn allele. None of the previously published studies on the *CYP1B1* Asn453Ser polymorphism and breast cancer conducted among average-risk women reported an increase in risk of breast cancer associated with this polymorphism [35-38], including a recent meta-analysis that reported an OR of 0.91 (95% CI 0.79, 1.04) associated with the heterozygous Asn/Ser genotype and an OR of 0.85 (95% CI 0.54, 1.34) associated with the homozygous Ser/Ser genotype [39]. However, functionality studies demonstrate that the *CYP1B1* polymorphic variant is associated with both higher catalytic activity in converting estrogen to 4-hydroxy estrogens and greater levels of DNA damage [39-41]. This mechanism may be more relevant for progression among the subgroup of women with BBD who progress to breast cancer, thus it is reasonable to examine this association among women biopsy proven BBD.

A marginally significant positive association was observed for the *ESR2* *38G>A. This finding is not consistent with results of published studies among either average-risk or high-risk women. Maguire et al [26] reported no increase in breast cancer risk associated with two copies of the variant *38A allele among average-risk (OR 0.81; 95% CI 0.50, 1.31) or high risk (OR 1.00; 95% CI 0.58, 1.71). In addition, Gold et al [16] observed that haplotypes containing the common *ESR2* *38G allele were associated with an increased risk of breast cancer among Ashkenazi Jews. These differences in findings may be due to the different populations under study; Maguire et al [26] examined *ESR2* polymorphisms among patients in Sweden and Gold et al [16] investigated *ESR2* and breast cancer among the Ashkenazi Jews. Neither investigated the association of *ESR2* polymorphisms among women diagnosed with BBD.

In addition, results from this study suggest that women with BBD who carry the *ESR2* *5772A>G polymorphism or the *ESR1* - 104062C>T polymorphism are at decreased risk of developing breast cancer. To our knowledge, neither of these genotypes has been examined in women

with BBD or among average risk women. The regions surrounding each polymorphism are well conserved with little recombination, therefore, it likely that, if the associations are real, the observed associations are due to linkage disequilibrium with another functional polymorphism in the region. No information on functionality is known for either *ESR2* *5772A>G or *ESR1* - 104062C>T.

There was no evidence of an association between the other polymorphisms examined in this study, including those in *COMT* and *CYP1A1*, and breast cancer risk. Although no published study has investigated the *COMT* Val158Met and *CYP1A1* Ile462Val polymorphisms in the progression of BBD to invasive breast cancer, previous studies of average-risk women have reported discrepant results regarding the association between the *COMT* Val158Met and the *CYP1A1* Ile462Val polymorphisms and the risk of breast cancer [8,42]. Both polymorphisms have been observed to be functional; the *COMT* 158Met allele is associated with three- to four-fold decreased activity compared to the wildtype allele [43] and the *CYP1A1* 462Val allele is consistent with higher enzyme activity [44]. Therefore, an association between these polymorphisms and breast cancer is biologically plausible and support the findings of studies that show an association between the polymorphisms and breast cancer risk. It may be, however, that both polymorphisms only modify the association between endogenous estrogens and the risk of breast cancer. Incorporating data on circulating estrogen concentrations may clarify the importance of carrying a specific genotype. Alternatively, it may also be that the effect of any single polymorphisms on the risk of breast cancer is small and that it is necessary to look at combinations of variants in the hormone metabolism pathway.

While this study provides novel information regarding the role of polymorphisms in estrogen metabolizing enzymes and the estrogen receptor genes in the progression of BBD to breast cancer, this study has several limitations inherent to its design. As reported in a previously published investigation on this cohort of women [32], pathology reports were not able to be obtained for all women who reported that they had undergone a surgical biopsy. This was due both to difficulties in obtaining permission for pathology review for some participants as well as to an inability to obtain records of biopsies done more than 15 to 20 years ago. Since concordance of finding a pathology report and report of biopsy among women granting permission for pathology review who reported that they had undergone

Table 3: Associations between estrogen metabolizing polymorphisms and breast cancer risk in the BBD cohort

	dbSNP rs#	Cancer (n = 91)		No Cancer (n = 1347)		All Participants	Post-menopausal Only
		n	%	n	%	Age- Adjusted OR (95% CI)	Age- Adjusted OR (95% CI)
<i>COMT</i>							
	4680						
<i>Val158Met</i>							
GG		16	17.6	272	20.2	1.00	1.00
GA		41	45.1	608	45.1	1.15 (0.63, 2.08)	0.89 (0.45, 1.77)
AA		24	26.4	371	27.5	1.10 (0.57, 2.11)	0.90 (0.43, 1.90)
GA/AA		65	71.4	979	72.7	1.13 (0.64, 1.98)	0.89 (0.47, 1.70)
Missing		10	11.0	96	7.1		
<i>CYP1A1</i>							
	1048943						
<i>Ile462Val</i>							
AA		78	85.7	1233	91.5	1.00	1.00
AG		7	7.7	80	5.9	1.38 (0.62, 3.10)	1.30 (0.45, 3.74)
GG		1	1.1	5	0.4	3.33 (0.38, 28.99)	8.33 (0.74, 93.62)
<i>p-for-trend</i>						0.20	0.20
AG/GG		8	8.8	85	6.3	1.49 (0.70, 3.19)	1.56 (0.60, 4.08)
Missing		5	5.5	29	2.2		
<i>CYP1B1</i>							
	1800440						
<i>Asn453Ser</i>							
AA		51	56.0	866	64.3	1.00	1.00
AG		31	34.1	358	26.6	1.45 (0.91, 2.31)	1.51 (0.87, 2.65)
GG		4	4.4	37	2.7	1.77 (0.61, 5.16)	2.44 (0.81, 7.36)
<i>p-for-trend</i>						0.08	0.05
AG/GG		35	38.5	395	29.3	1.48 (0.95, 2.32)	1.61 (0.94, 2.74)
Missing		5	5.5	86	6.4		
<i>CYP1B1</i>							
	10012						
<i>Arg48Gly</i>							
CC		39	42.9	600	44.5	1.00	1.00
CG		33	36.3	460	34.1	1.10 (0.68, 1.78)	0.95 (0.53, 1.69)
GG		3	3.3	76	5.6	0.60 (0.18, 2.00)	0.25 (0.03, 1.88)
CG/GG		36	39.6	536	39.8	1.02 (0.65, 1.64)	0.84 (0.48, 1.50)
Missing		16	17.6	211	15.7		
<i>CYP1B1</i>							
	1056836						
<i>Val432Leu</i>							
CC		25	27.5	365	27.1	1.00	1.00
CG		39	42.9	653	48.5	0.88 (0.53, 1.48)	1.04 (0.56, 1.98)
GG		17	18.7	270	20.0	0.93 (0.49, 1.76)	0.93 (0.41, 2.09)
CG/GG		56	61.5	923	68.5	0.90 (0.55, 1.46)	1.01 (0.56, 1.85)
Missing		10	11.0	59	4.4		

a biopsy after 1980 was high (92.5%), all women who reported a breast biopsy were included in the analyses. However, this may have led to the inclusion of a small number of women without BBD. Further, because data on the type of BBD were not collected for all participants, the

analyses conducted in this study were not stratified by type of BBD, and, therefore, the ORs reported may underestimate (or overestimate) the true associations among women with certain types of BBD.

Table 4: Associations between estrogen receptor-alpha (ESR1) polymorphisms and breast cancer risk in the BBD cohort

	dbSNP rs#	Cancer (n = 91)		No Cancer (n = 1347)		All Participants	
		n	%	n	%	Age- Adjusted OR (95% CI)	Post- menopausal Only Age-Adjusted OR (95% CI)
<i>ESR1 Pro325Pro</i>	1801132						
CC		52	57.1	794	58.9	1.00	1.00
CG		31	34.1	440	32.7	1.07 (0.67, 1.69)	1.06 (0.61, 1.87)
GG		7	7.7	64	4.8	1.65 (0.72, 3.78)	2.03 (0.81, 5.10)
CG/GG		38	41.8	504	37.4	1.14 (0.74, 1.76)	1.19 (0.70, 2.01)
Missing		1	1.1	49	3.6		
<i>ESR1 Ser10Ser</i>	2077647						
TT		26	28.6	335	24.9	1.00	1.00
TC		36	39.6	642	47.7	0.71 (0.42, 1.20)	0.90 (0.46, 1.77)
CC		27	29.7	304	22.6	1.14 (0.65, 1.99)	1.88 (0.94, 3.75)
TC/CC		63	69.2	946	68.8	0.85 (0.53, 1.37)	1.21 (0.65, 2.24)
Missing		2	2.2	66	4.8		
<i>ESR1 Ala87Ala</i>	746432						
GG		67	73.6	1077	80.0	1.00	1.00
GC		15	16.5	181	13.4	1.32 (0.74, 2.36)	1.24 (0.61, 2.51)
CC		1	1.1	7	0.5	2.30 (0.28, 18.95)	3.34 (0.38, 29.18)
<i>p-for-trend</i>						0.3	0.3
GC/CC		16	17.6	188	14.0	1.36 (0.77, 2.39)	1.31 (0.66, 2.60)
Missing		8	8.8	82	6.1		
<i>ESR1 IVS6+52</i>	2273206						
GG		69	75.8	1060	78.7	1.00	1.00
GT		13	14.3	229	17.0	0.86 (0.48, 1.59)	0.71 (0.33, 1.53)
TT		3	3.3	10	0.7	4.79 (1.29, 17.88)	5.31 (1.04, 27.28)
GT/TT		16	17.6	239	17.7	1.02 (0.58, 1.79)	0.86 (0.43, 1.73)
Missing		6	6.6	48	3.6		
<i>ESR1 Thr594Thr</i>	2228480						
AA		2	2.2	41	3.0	1.00	1.00
AG		20	22.0	358	26.6	1.14 (0.26, 5.05)	0.88 (0.19, 4.04)
GG		58	63.7	841	62.4	1.42 (0.34, 6.01)	0.86 (0.20, 3.77)
AG/GG		78	85.7	1199	89.0	1.34 (0.32, 5.62)	0.87 (0.20, 3.75)
Missing		11	12.1	107	7.9		
<i>ESR1 - 104062C>T</i>	851982						
CC		38	41.8	470	34.9	1.00	1.00
CT		33	36.3	612	45.4	0.65 (0.40, 1.05)	0.55 (0.30, 0.99)
TT		14	15.4	190	14.1	0.88 (0.47, 1.67)	0.89 (0.43, 1.85)
CT/TT		47	51.6	802	59.5	0.70 (0.45, 1.09)	0.63 (0.37, 1.08)
Missing		6	6.6	75	5.6		

Table 5: Associations between estrogen receptor-beta (ESR2) polymorphisms and breast cancer risk in the BBD cohort

	dbSNP rs#	Cancer (n = 91)		No Cancer (n = 1347)		All Participants	Post-menopausal Only
		n	%	n	%	Age- Adjusted OR (95% CI)	Age-Adjusted OR (95% CI)
<i>ESR2 38 bp 3' of STP</i>							
	4986938						
		26	28.6	470	34.9	1.00	1.00
		43	47.3	612	45.4	1.27 (0.77, 2.10)	1.46 (0.80, 2.68)
		19	20.9	190	14.1	1.83 (0.99, 3.38)	1.71 (0.78, 3.76)
	<i>p-for-trend</i>					0.06	0.1
		62	68.1	802	59.5	1.40 (0.88, 2.25)	1.52 (0.85, 2.71)
	Missing	3	3.3	75	5.6		
<i>ESR2 5659 bp 3' of STP</i>							
	928554						
		10	11.0	202	15.0	1.00	1.00
		44	48.4	632	46.9	1.41 (0.70, 2.86)	1.04 (0.48, 2.26)
		28	30.8	419	31.1	1.38 (0.66, 2.89)	0.88 (0.38, 2.05)
		72	79.1	1051	78.0	1.40 (0.71, 2.76)	0.98 (0.47, 2.06)
	Missing	9	9.9	94	7.0		
<i>ESR2 5696 bp 3' of STP</i>							
	none						
		75	82.4	1101	81.7	1.00	1.00
		12	13.2	180	13.4	0.96 (0.51, 1.81)	0.74 (0.33, 1.67)
		1	1.1	4	0.3	3.68 (0.41, 33.37)	--
		13	14.3	184	13.7	1.02 (0.56, 1.88)	0.73 (0.33, 1.64)
	Missing	3	3.3	62	4.6		
<i>ESR2 5772 bp 3' of STP</i>							
	8018687						
		86	94.5	1133	84.1	1.00	1.00
		5	5.5	169	12.5	0.39 (0.16, 0.99)	0.62 (0.24, 1.59)
		0	0.0	5	0.4	--	--
		5	5.5	174	12.9	0.38 (0.15, 0.96)	0.61 (0.24, 1.55)
	Missing	0	0.0	40	3.0		

In addition, though the base cohort is relatively large, the number of cases of breast cancer occurring limited the power of the study to make firm conclusions about the relationships between polymorphisms in *COMT*, *CYP1B1*, *CYP1A1*, *ESR1* and *ESR2* and the risk of breast cancer in Caucasian women with BBD. We had limited power to detect small associations and, therefore, were also limited in our ability to correct for multiple testing. As stated in the results section, the associations between the polymorphisms and the risk of developing breast cancer were not statistically significant after correction for multiple testing and this may have been due, in part, to the small sample size. However, while we may be able to rule out large changes in risk of clinical and public health

consequence associated with certain genotypes, we still observed, with the small number of events, statistically significant changes in the risk of breast cancer in this cohort. Taken together with functional considerations of these polymorphisms, it is important to follow-up on these observations to determine if we can better understand and perhaps intervene for those women with BBD who are most at risk to progress to cancer.

Conclusion

The findings of this study indicate that specific polymorphisms in the *CYP1B1*, *ESR1*, and *ESR2* genes may play a role in progression of BBD to breast cancer among Caucasian women. Although additional studies are needed to

confirm or refute our findings, these results suggest that genetic markers may aid in the identification of women who are at risk for progression of BBD to cancer.

Abbreviations

BBD Benign breast disease

COMT Catechol O-methyltransferase

CYPs Cytochrome P450 enzymes

ESR1 Estrogen receptor-alpha

ESR2 Estrogen receptor-beta

SNP Single nucleotide polymorphism

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

LG contributed to the statistical analysis, interpretation of the data, and the writing of the manuscript. SIB provided expertise on the more advanced analyses of the genetic data and contributed to the interpretation of the data. MAM, LWT, and SAH contributed to the data collection and statistical analysis. PA conducted reviews of the pathology report data and pathology specimens. CJN and KJN contributed to the conception and design of the study and the interpretation of the data.

Acknowledgements

This research was supported by a National Cancer Institute SPORE grant CA8843 and a National Institute on Aging grant AG18033 from the National Institutes of Health. The funding agencies had no role in the study design, collection, analysis and interpretation of the data, or in the writing of the manuscript. Dr. Gallicchio is supported by an Institutional National Research Service Award from the National Cancer Institute (T32 CA09314). The authors thank Dr. Ingo Ruzcinski for providing his biostatistics expertise in this analysis. The authors thank Judy Bolton-Hoffman and Alyce Burke at the Comstock Center for Public Health Research and Prevention for their help in carrying out this study.

References

- Wang J, Costantino JP, Tan-Chiu E, Wickerham DL, Paik S, Wolmark N: **Lower-category benign breast disease and the risk of invasive breast cancer.** *J Natl Cancer Inst* 2004, **96**:616-620.
- Hartman LC, Sellers TA, Frost MH, Lingle WL, Denghim AC, Ghosh K, Vierkant RA, Maloney SD, Pankratz VS, Hillman DW, Suman VJ, Johnson J, Blake C, Tilsty T, Vachon CM, Melton LJ, Visscher DW: **Benign breast disease and the risk of breast cancer.** *N Engl J Med* 2005, **353**:229-237.
- Missmer SA, Eliassen AH, Barbieri RL, Hankinson SE: **Endogenous estrogen, androgen, and progesterone concentrations and breast cancer risk among postmenopausal women.** *J Natl Cancer Inst* 2004, **96**:1856-1865.
- Dorgan JF, Longcope C, Stephenson HEJ, Falk RT, Miller R, Franz C, Kahle L, Campbell WS, Tangrea JA, Schatzkin A: **Serum sex hormone levels are related to breast cancer risk in postmenopausal women.** *Environ Health Perspect* 1997, **105 Suppl 3**:583-585.
- Key TJ: **Serum oestradiol and breast cancer risk.** *Endocr Relat Cancer* 1999, **6**:175-180.
- Key T, Appleby P, Barnes I, Reeves G: **Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies.** *J Natl Cancer Inst* 2002, **94**:606-616.
- Kabuto M, Akiba S, Stevens RG, Neriishi K, Land CE: **A prospective study of estradiol and breast cancer in Japanese women.** *Cancer Epidemiol Biomarkers Prev* 2000, **9**:575-579.
- Mitrunen K, Hirvonen A: **Molecular epidemiology of sporadic breast cancer. The role of polymorphic genes involved in oestrogen biosynthesis and metabolism.** *Mutat Res* 2003, **544**:9-41.
- Thompson PA, Ambrosone C: **Molecular epidemiology of genetic polymorphisms in estrogen metabolizing enzymes in human breast cancer.** *J Natl Cancer Inst Monogr* 2000, **27**:125-134.
- Zhu BT, Conney AH: **Functional role of estrogen metabolism in target cells: review and perspectives.** *Carcinogenesis* 1998, **19**:1-27.
- Liehr JG: **Is estradiol a genotoxic mutagenic carcinogen?** *Endocr Rev* 2000, **21**:40-54.
- International Agency for Research on Cancer: **Overall Evaluations of Carcinogenicity: An Updating of IARC Monographs.** 1987, volumes 1-42 (suppl. 6):.
- Yager JD: **Endogenous estrogens as carcinogens through metabolic activation.** *J Natl Cancer Inst Monogr* 2000:67-73.
- Greene GL, Gilna P, Waterfield M, Baker A, Hort Y, Shine J: **Sequence and expression of human estrogen receptor complementary DNA.** *Science* 1986, **231**:1150-1154.
- Mosselman S, Polman J, Dijkema R: **ER beta: identification and characterization of a novel human estrogen receptor.** *FEBS Lett* 1996, **392**:49-53.
- Gold B, Kalush F, Bergeron J, Scott K, Mitra N, Wilson K, Ellis N, Huang H, Chen M, Lippert R, Halldorsson BV, Woodworth B, White T, Clark AG, Parl FF, Broder S, Dean M, Offit K: **Estrogen receptor genotypes and haplotypes associated with breast cancer risk.** *Cancer Res* 2004, **64**:8891-8900.
- Roodi N, Bailey LR, Kao WY, Verrier CS, Yee CJ, Dupont WD, Parl FF: **Estrogen receptor gene analysis in estrogen receptor-positive and receptor-negative primary breast cancer.** *J Natl Cancer Inst* 1995, **87**:446-451.
- Andersen TI, Heimdal KR, Skrede M, Tveit K, Berg K, Borresen AL: **Oestrogen receptor (ESR) polymorphisms and breast cancer susceptibility.** *Hum Genet* 1994, **94**:665-670.
- Iwase H, Greenman JM, Barnes DM, Hodgson S, Bobrow L, Mathew CG: **Sequence variants of the estrogen receptor (ER) gene found in breast cancer patients with ER negative and progesterone receptor positive tumors.** *Cancer Lett* 1996, **108**:179-184.
- Wang H, Strobel HW: **Regulation of CYP3A9 gene expression by estrogen and catalytic studies using cytochrome P450 3A9 expressed in Escherichia coli.** *Arch Biochem Biophys* 1997, **344**:365-372.
- Cai Q, Shu XO, Jin F, Dai Q, Wen W, Cheng JR, Gao YT, Zheng W: **Genetic polymorphisms in the estrogen receptor alpha gene and risk of breast cancer: results from the Shanghai Breast Cancer Study.** *Cancer Epidemiol Biomarkers Prev* 2003, **12**:853-859.
- Cai Q, Gao YT, Wen W, Shu XO, Jin F, Smith JR, Zheng W: **Association of breast cancer risk with a GT dinucleotide repeat polymorphism upstream of the estrogen receptor-alpha gene.** *Cancer Res* 2003, **63**:5727-5730.
- Southey MC, Batten LE, McCredie MR, Giles GG, Dite G, Hopper JL, Venter DJ: **Estrogen receptor polymorphism at codon 325 and risk of breast cancer in women before age forty.** *J Natl Cancer Inst* 1998, **90**:532-536.
- Schubert EL, Lee MK, Newman B, King MC: **Single nucleotide polymorphisms (SNPs) in the estrogen receptor gene and breast cancer susceptibility.** *J Steroid Biochem Mol Biol* 1999, **71**:21-27.
- Kang HJ, Kim SW, Kim HJ, Ahn SJ, Bae JY, Park SK, Kang D, Hirvonen A, Choe KJ, Noh DY: **Polymorphisms in the estrogen receptor-alpha gene and breast cancer risk.** *Cancer Lett* 2002, **178**:175-180.
- Maguire P, Margolin S, Skoglund J, Sun XF, Gustafsson JA, Borresen-Dale AL, Lindblom A: **Estrogen Receptor Beta (ESR2) Polymorphisms in Familial and Sporadic Breast Cancer.** *Breast Cancer Res Treat* 2005, **94**:145-152.

27. Onland-Moret NC, van Gils CH, Roest M, Grobbee DE, Peeters PH: **The estrogen receptor alpha gene and breast cancer risk (The Netherlands).** *Cancer Causes Control* 2005, **16**:1195-1202.
28. Modugno F, Zmuda JM, Potter D, Cai C, Ziv E, Cummings SR, Stone KL, Morin PA, Greene D, Cauley JA: **Association of estrogen receptor alpha polymorphisms with breast cancer risk in older Caucasian women.** *Int J Cancer* 2005, **116**:984-991.
29. Zheng SL, Zheng W, Chang BL, Shu XO, Cai Q, Yu H, Dai Q, Xu J, Gao YT: **Joint effect of estrogen receptor beta sequence variants and endogenous estrogen exposure on breast cancer risk in Chinese women.** *Cancer Res* 2003, **63**:7624-7629.
30. Helzlsouer KJ, Alberg AJ, Huang HY, Hoffman SC, Strickland PT, Brock JW, Burse VW, Needham LL, Bell DA, Lavinge JA, Yager JD, Comstock GW: **Serum concentrations of organochlorine compounds and the subsequent development of breast cancer.** *Cancer Epidemiol Biomarkers Prev* 1999, **8**:525-532.
31. Platz EA, De Marzo AM, Erlinger TP, Rifai N, Visvanathan K, Hoffman SC, Helzlsouer KJ: **No association between pre-diagnostic plasma C-reactive protein concentration and subsequent prostate cancer.** *Prostate* 2004, **59**:393-400.
32. Gallicchio L, McSorley MA, Newschaffer CJ, Thuita LW, Huang HY, Hoffman SC, Helzlsouer KJ: **Nonsteroidal anti-inflammatory drugs, cyclooxygenase polymorphisms, and the risk of developing breast cancer among women with benign breast disease.** *Cancer* 2005, **106**(7):1443-52.
33. Ruczinski I, Kooperberg C, LeBlanc ML: **Exploring interactions in high-dimensional genomic data: an overview of Logistic Regression, with applications.** *Journal of Multivariate Analysis* 2004, **90**:178-195.
34. Benjamini Y, Hochberg Y: **Controlling the false discovery rate: a practical and powerful approach to multiple testing.** *J R Stat Soc B* 1995, **57**:289-300.
35. Rylander-Rudqvist T, Wedren S, Granath F, Humphreys K, Ahlberg S, Weiderpass E, Oscarson M, Ingelman-Sundberg M, Persson I: **Cytochrome P450 IBI gene polymorphisms and postmenopausal breast cancer risk.** *Carcinogenesis* 2003, **24**:1533-1539.
36. Bailey LR, Roodi N, Verrier CS, Yee CJ, Dupont WD, Parl FF: **Breast cancer and CYP1A1, GSTM1, and GSTT1 polymorphisms: evidence of a lack of association in Caucasians and African Americans.** *Cancer Res* 1998, **58**:65-70.
37. De VI, Hankinson SE, Li L, Colditz GA, Hunter DJ: **Association of CYP1B1 polymorphisms and breast cancer risk.** *Cancer Epidemiol Biomarkers Prev* 2002, **11**:489-492.
38. Bailey LR, Roodi N, Dupont WD, Parl FF: **Association of cytochrome P450 IBI (CYP1B1) polymorphism with steroid receptor status in breast cancer.** *Cancer Res* 1998, **58**:5038-5041.
39. Wen W, Cai Q, Shu XO, Cheng JR, Parl F, Pierce L, Gao YT, Zheng W: **Cytochrome P450 IBI and catechol-O-methyltransferase genetic polymorphisms and breast cancer risk in Chinese women: results from the shanghai breast cancer study and a meta-analysis.** *Cancer Epidemiol Biomarkers Prev* 2005, **14**:329-335.
40. Hanna IH, Dawling S, Roodi N, Guengerich FP, Parl FF: **Cytochrome P450 IBI (CYP1B1) pharmacogenetics: association of polymorphisms with functional differences in estrogen hydroxylation activity.** *Cancer Res* 2000, **60**:3440-3444.
41. Shimada T, Watanabe J, Kawajiri K, Sutter TR, Guengerich FP, Gillam EM, Inoue K: **Catalytic properties of polymorphic human cytochrome P450 IBI variants.** *Carcinogenesis* 1999, **20**:1607-1613.
42. Le ML, Donlon T, Kolonel LN, Henderson BE, Wilkens LR: **Estrogen metabolism-related genes and breast cancer risk: the multiethnic cohort study.** *Cancer Epidemiol Biomarkers Prev* 2005, **14**:1998-2003.
43. Lachman HM, Papolos DF, Saito T, Yu YM, Szumlanski CL, Weinshilboum RM: **Human catechol-O-methyltransferase pharmacogenetics: description of a functional polymorphism and its potential application to neuropsychiatric disorders.** *Pharmacogenetics* 1996, **6**:243-250.
44. Landi MT, Sinha R, Lang NP, Kadlubar FF: **Human cytochrome P450IA2**. In *IARC Scientific Publications* Lyon, IARC; 1999:173-195.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/6/173/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

