

RESEARCH ARTICLE

Open Access

Antiproliferative activity and induction of apoptosis by *Annona muricata* (Annonaceae) extract on human cancer cells

Constant Anatole Pieme^{1,2*}, Santosh Guru Kumar², Mireille Sylviane Dongmo³, Bruno Moukette Moukette¹, Fabrice Fekam Boyoum⁴, Jeanne Yonkeu Ngogang¹ and Ajit Kumar Saxena^{2*}

Abstract

Background: *Annona muricata* (*A. muricata*) is widely distributed in Asia, Africa and South America. Different parts of this plant are used to treat several diseases in Cameroon. The aim of this study is to determine the *in vitro* anti-proliferative effects and apoptotic events of *A. muricata* extracts on HL-60 cells as well as to quantify its phenols content.

Methods: The cell viability was measured by using 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) assay while the changes in morphology of HL-60 cells, membrane mitochondrial potential (MMP) and the cell cycle were used for assessment apoptosis induction.

Results: The results show that the concentration of phenols, flavonoids and flavonols in the extracts varied depending on the part of the plant. All the extracts tested inhibited the proliferation of HL-60 cells in a concentration dependent manner with IC₅₀ varied from 6–49 µg/mL. The growth inhibition of the cells by extracts was associated with the disruption of MMP, reactive oxygen species (ROS) generation and the G₀/G₁ cell arrest.

Conclusion: These findings suggest that the extracts from *A. muricata* have strong antiproliferation potential and can induce apoptosis through loss of MMP and G₀/G₁ phase cell arrest.

Keywords: Apoptosis, Membrane mitochondrial potential, Antiproliferative, Cell cycle, *A. muricata*

Background

Chemopreventive properties have long been attributed to polyphenolic compounds present in the human diet. The interest on these natural substances is increasing because of their higher potential sources of anticancer compounds. Plants and plant-based medicaments are used as the basis of many modern pharmaceuticals industries today for the treatment of our various ailments [1]. According to world health organization (WHO), more than 80% of the total world's population depends on the traditional medicines to satisfy their primary health care needs. Several phytochemical molecules from natural products capable of exerting a physiologic action on the

human body were studied and characterized. These bioactive compounds such as alkaloids, flavonoids, tannins and phenols were considered to be most important. The phytochemical research that has been done based on the ethno pharmacological information forms the effective approach in the discovery of new anti-infective agents from higher plants [2].

Annona muricata L. (*A. muricata*) commonly known as graviola, soursop or corossol, belongs to the Annonaceae family. It is a widespread small tree and has its native in Central America [3]. It is a typical tropical tree with heart shaped edible fruits and widely distributed in most of tropical countries. All parts of *A. muricata* tree are used in natural medicine in the tropic including the twigs, leaf, root, fruit and seeds. Generally, the fruit and fruit juice are taken to eliminate worms and parasites, cool fever, increase mother's milk after child birth, and as an astringent for diarrhea and dysentery [4]. The crushed seeds are used against internal and external parasites, head

* Correspondence: apieme@yahoo.fr; aksaxena@iiim.ac.in

¹Department of Physiological Sciences and Biochemistry, Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, PO Box 1364, Yaoundé, Cameroon

²Cancer Pharmacology Division, Indian Institute of Integrative Medicine, 180001, Canal Road, 18001 Jammu, India

Full list of author information is available at the end of the article

lice and warms. The twigs, leaf are considered sedative and antispasmodic [4]. A decoction of *A. muricata* leaf is used to kill bed bug and head lice to reduce fever. For the latter it can have the same effect taken orally or added to bathing water [5]. The creamy and delectable flesh of the fruit consist of 80% water, 1% protein, 18% carbohydrates and fair amount of vitamins B, B2 and C, potassium and dietary fiber [6]. The leaf are lanceolate with glossy and dark green in color had been traditionally used to treat headaches, hypertension, cough, asthma and used as antispasmodic, sedative and nervine for heart condition [7,8]. Previous reports have demonstrated that the leaf, twigs, root, stem, and fruit seed extracts of *A. muricata* have several biological activities such as anti-bacterial [9], antifungal [10] and anti-malarial [11]. Its leaf extract were also found to possess antioxidant [12] and molluscicidal properties [13]. Recently, it has also been reported to exhibit anti-inflammatory and analgesic effects [14], cytotoxicity and apoptosis inducing activities on T47D breast cancer [15], antiviral activity [16] and antidiabetic activity. Phytochemical investigation of the leaf of *A. muricata* showed the presence of alkaloids [17], essential oils [18] and acetogenins [19]. These acetogenins demonstrated to be selectively toxic against various types of the cancerous cells without harming healthy cells [20]. Acetogenin 1 was reported to exhibit cytotoxic activities against the human pancreatic tumor cell line (PACA-2), human prostate adenocarcinoma (PC-3) and human lung carcinoma (A-549), while Acetogenin 2 was reported to exhibit cytotoxicity against human hepatoma carcinoma cell line (Hep G2) [21]. Seven isoquinoline alkaloids including reticuline, coclaurine, coreximine, atherosperminine, stepharine, anomurine and anomuricine have been isolated from the leaves, root and stem barks of *A. muricata* [22]. The essential oil of the fresh fruit pulp of *A. muricata* contains 2-hexenoic acid methyl ester (23.9%), 2-hexenoic acid ethyl ester (8.6%), 2-octenoic acid methyl ester (5.4%), 2-butenic acid methyl ester (2.4%), β -caryophyllene (12.7%), 1,8-cineole (9.9%), linalool (7.8%), α -terpineol (2.8%), lialyl propionate (2.2%) and calarence (2.2%) [23]. Therefore, we attempted to investigate the growth-inhibitory and apoptotic effects of extracts from leaf, twigs and roots from *A. muricata* against Human promyelocytic leukemia (HL-60 cells).

Methods

Preparation of extracts

The plant material including leaf, twigs and roots of were collected in Yaounde, capital city of Cameroon and authenticated by Mr NANA, a botanist of the National Herbarium of Cameroon in comparison to the voucher specimens under the reference number of 3289/HNC. They were then shade dried and grounded using a blender.

The obtained powder was cold macerated with ethanol 95°C with occasional stirring for 3 days. After 3 days, the suspension was filtered through filtered paper and the filtrate was evaporated to dryness at low temperature, using a rotary evaporator. The procedure was repeated 3 times and until total decoloration of the mixture observed. The obtained extracts were stored and used for further analysis.

Determination of phenolic composition of the extracts

Total phenol determination

The total phenol was determined by the Folin–Ciocalteu method, the reaction mixture contains: 200 μ L of diluted spice extract, 800 μ L of freshly prepared diluted Folin Ciocalteu reagent and 2 mL of 7.5% sodium carbonate. The final mixture will be diluted to 7 mL with deionized water. Mixture was kept in the dark at ambient conditions for 2 h to complete the reaction. The absorbance at 765 nm will be measured. Garlic acid was used as standard and the results were expressed as mg garlic acid (GAE)/g of dried material.

Determination of total flavonoid content

Total flavonoid content was determined using aluminium chloride (AlCl_3) according to a known method using quercetin as a standard. The spice extract (0.1 mL) was added to 0.3 mL distilled water followed by 5% NaNO_2 (0.03 mL). After 5 min at 25°C, AlCl_3 (0.03 mL, 10%) was added. After further 5 min, the reaction mixture was treated with 0.2 mL of 1 mM NaOH. Finally, the reaction mixture was diluted to 1 mL with water and the absorbance was measured at 510 nm. The results will be expressed as mg of quercetin (QE)/g of dried material

Determination of total flavonols

Total flavonols in the plant extracts were estimated using the method of Kumaran and Karunakaran [24]. To 2.0 mL of sample (standard), 2.0 mL of 2% AlCl_3 ethanol and 3.0 mL (50 g/L) sodium acetate solutions were added. The absorption at 440 nm was read after 2.5 h at 20°C. Extract samples were evaluated at a final concentration of 0.1 mg/mL. Total flavonoid content was calculated as mg of quercetin (mg/g) using the following equation $y = 5.3911x + 0.0313$, $R^2 = 0.9967$ based on the calibration curve, where x was the absorbance and y was the concentration of quercetin (mg/g).

Antitumor activity of the extracts

Cell culture

The Human promyelocytic leukemia (HL-60 cells) was obtained from European Collection of Cells Culture (ECCC), Sigma–Aldrich, India. They were grown in RPMI-1640 medium containing 10% Fetal bovine serum

(FBS), penicillin (100 IU/mL) and streptomycin (100 µg/mL medium). The cells suspension was kept in the incubator (Thermo Electron Corporation, USA) at 37°C, 5% CO₂; 95% humidity. Cells were used for different assays during logarithmic growth phase while the untreated control cultures received only the vehicle (dimethyl sulfoxide [DMSO] < 0.1%).

Cells viability and treatments

HL-60 cells were seeded in 96 different well plates containing 15×10^3 µL/well, respectively. The cultured cells were then treated (triplicate wells per condition) by adding of 100 µL of serial dilutions of the three extracts in DMSO to give a final concentration of 100, 30, 10 and 1 µg/mL. In addition, the DMSO alone was added to another set of cells as the solvent control (DMSO < 0.1%). The cells were then incubated for another 48 h prior to the addition of 20 µL of 2.5 mg/mL solution of 3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide (MTT) into each well. The incubation was continued for 3 h before the media was removed. A mixture of DMSO (150 µL) was added to each well and mixed to ensure dissolving of the crystal formazan before the absorbance at 570 nm was measured. Three replications of each experiment were performed and fifty percent of inhibitory concentration (IC₅₀) of each extract was calculated.

Hoechst 33258 staining of cells for nuclear morphology

HL-60 cells (2×10^6 cells/3 mL/well) were treated with different extracts at different concentrations of extract for 24 h. They were collected, centrifuged at 400 g and washed once with PBS. A solution of Hoechst (Hoechst, 10 µg/mL; citric 10 mM; Na₂HPO₄ 0.45 M; Tween-20 0.05%) was added in each tube and kept in the dark at room temperature for 30 min. The mixture was washed with PBS and the pellet suspended in 100 µL of PBS/glycerol (1:1). The solution (10 µL) was poured into the slide and nuclear morphology alterations observed under fluorescence microscope (Olympus X 70, magnification 20 X) [25].

Reactive oxygen species (ROS) assay

ROS production was monitored by flow cytometry using 2', 7'- dichlorodihydrofluorescein diacetate (DCFH₂-DA). This dye is a stable non polar compound that readily diffuses into cells and is hydrolyzed by intracellular esterase to yield 2',7' dichlorodihydrofluorescein (DCFH₂), which is trapped within the cells. Hydrogen peroxide or low molecular weight peroxides produced by the cells oxidizes DCFH₂ to a highly fluorescent compound 2',7'-dichlorofluorescein (DCF). Thus, the fluorescence intensity was proportional to the amount of hydrogen peroxide produced by the cells. Briefly, HL-60 cells (1×10^6

cells/2 mL/well) were treated with extracts of leaf, roots and twigs of *A. muricata* at different concentrations for 24 h. Thirty minutes before the end of the experiment, the cell culture was treated with DCFH₂-DA (50 µM) and kept in the dark. Cells were then collected, centrifuged (200 g; 4°C; 5 min) and the pellet was washed with 1 mL of PBS and centrifuged as mentioned earlier. The pellet was suspended in 500 µL of PBS and the fluorescence was assessed by comparing two fluorescence emission 480 nm/530 nm using a flow-cytometer (BD-LSR).

Mitochondrial membrane potential (MMP) assay

HL-60 cells (1×10^6 cells/2 mL/well) were treated with the three extracts at different concentrations for 24 h. Thirty minutes before the end of the experiment, the cell culture was treated with Rhodamine-123 (200nM) and kept in the dark for 30 mn. Cells were collected, centrifuged (400 g; 4°C; 5 min), the pellet was washed with 1 mL of PBS and centrifuged as mentioned earlier. The depolarization of mitochondrial membrane was examined by measuring the fluorescence emission shift (red to green) of the Δψ_m sensitive cationic Rh-123 dye. The fluorescence intensity of 10,000 events was analyzed in FL-1 channel on BD FACS Calibur (Becton Dickinson, USA) flow cytometer. The decrease in fluorescence intensity because of MMP loss was analyzed in FL-1 channel and the change of potential membrane (Δψ_m) was assessed by comparing fluorescence.

DNA content and cell cycle phase distribution

HL-60 cells (1×10^6 cells/2 mL/well) were treated with extracts at 20, 50, 100 µg/mL for 24 h. They were harvested and washed with 1 mL of PBS, then centrifuged at 400 g for 5 min at 4°C. The pellet was suspended in 100 µL of PBS and 900 µL of hypertonic buffer (PI-25 µg/mL, RNAase- 40 µg/mL, sodium citrate-0.1% and Triton-100X-0.03%) and incubated at 37°C in dark for 20 min. Finally, cells were analyzed immediately on flow cytometer FACS Calibur (Becton Dickinson, USA). The data were collected in list mode on 10,000 events and illustrated in a histogram, where the number of cells (counts) is plotted against the relative fluorescence intensity of PI (FL-2; λ_{em}: 585 nm; red fluorescence). The resulting DNA distributions were analyzed by Modfit (Verity Software House Inc., Topsham, ME) for the proportions of cells in G₀/G₁, S phase, and G₂/M phases of the cell cycle [26].

Statistical analysis

The viability experiments were done in triplicates and each data point represents the average of at least 3 independent experiments. The distributions of the data are abnormal. The data was expressed as mean ± SD. In

order to carry out statistical analysis, the data was analyzed using SPSS (Version 11.5; SPSS Inc.) and M.S. Office, Excel software. One way analysis of variance technique was applied to observe the significance between the groups. The post hoc test Duncan's multiple range test was performed to know the significant difference among the groups. Entire statistical analysis was carried out at $p < 0.05$.

Results

Phenolic contents of *A. muricata*

The results of this study showed that the level of polyphenols, flavonoids and flavonols varied depending on the part of the plant (Table 1). The concentration of these three groups of molecules is higher in the leaves compare other parts of plant extract. The lowest concentration of flavonoids and flavonols was found on the stem barks of *A. muricata* while its roots show the lower phenols content.

In vitro anticancer activity

Effects of A. muricata extracts on the proliferation of HL-60

In this study we used a microculture assay based on metabolic reduction of MTT to evaluate the cytotoxic effect extracts of *A. muricata* on HL-60 cells. This technique permitted us to evaluate dose-dependent effect, by linear regression analysis showing acceptable R^2 values and correlation coefficients. As shown in the Figure 1, the addition of extracts at different concentration to the cultured cells inhibited dramatically and significantly the proliferation of the cells in a dose-dependent manner. The values of IC_{50} of the extract after 48 h was between 6–12 $\mu\text{g/mL}$ which is lower than 20 $\mu\text{g/mL}$ (Table 1).

Morphological changes of apoptotic treated HL-60 cells with A. muricata extracts

To investigate of *A. muricata* extracts on the nuclear modification on HL-60 cells, the Hoechst 33258 staining test was performed at different concentrations (20, 50 and 100 $\mu\text{g/mL}$) after 24 h of treatment. Hoechst 33258 reagent is a membrane-permeable blue fluorescent dye which stained cell nucleus. As observed in Figure 2, the control or untreated cells present the characteristics of healthy cells. They appeared to be intact oval shape and the uniform nuclei were stained with a less bright blue fluorescence. Cells treated with tested extracts exhibited

a bright blue color when the concentration of extract increases. These results demonstrated typical features of apoptosis such as cell shrinkage, chromatin condensation, and fragmentation to multiple aggregate of apoptotic bodies and cell decrement (Figure 2). The apoptotic nuclei clearly showed highly condensed or fragmented chromatin. At 100 $\mu\text{g/mL}$, most of the cells undergo apoptosis and the number of apoptotic bodies increases.

A. muricata extracts induce apoptosis by generation of ROS

To investigate whether extracts of *A. muricata* inhibit the HL-60 cells by through the generation of ROS, we monitored the redox status of the HL-60 treated cells using the oxidation sensitive fluorescent dye DCFDA. As shown in Figure 3A, B and C, the ROS levels generated after 24 h treatment of HL-60 cells varied with the concentration of extract and the part of plant used in the study. However, the higher ROS levels were found at the concentration of 20 $\mu\text{g/mL}$ (9.99% for twigs), 50 $\mu\text{g/mL}$ (4.08% for Leaf) and 100 $\mu\text{g/mL}$ (3.07% for the twigs) (Figure 3D). At the higher concentration (100 $\mu\text{g/mL}$), the ROS produced by cells with the extract of the roots and leaves were lower than that of untreated control cells (Figure 3D).

A. muricata extracts disrupt mitochondrial membrane potential in HL-60 cells

After treatment of cells with extracts at different concentration, we observed an increase of fluorescence intensity indicating the mitochondrial membrane depolarization as Figure 4A, B & C. The depolarization of mitochondrial membrane varied with concentration and the extracts. A dose-dependent increase of fluorescence observed is ranging from 17.34% to 98.91% (Figure 4B). Among the extract tested, the twigs extract demonstrated a higher depolarization of mitochondrial membrane (21.75%) at 20 $\mu\text{g/mL}$ while those of leaf and roots showed a maximum depolarization at 100 $\mu\text{g/mL}$ (98.29 and 98.91% respectively). All the extracts at 50 $\mu\text{g/mL}$ exhibited more than 50% of depolarization of mitochondrial membrane (Figure 4D). These results show that extracts induced apoptosis on HL-60 cells after 24 h through the disruption of mitochondrial membrane.

Table 1 Phenolic composition and fifty percent inhibition of extracts of *A. muricata*

Parts of plant used	HL-60 cells IC_{50} ($\mu\text{g/mL}$)	Total polyphenol (mg/g dried material)	Flavonoids (mg/g dried material)	Flavonols (mg/g dried material)
T	49 ± 3.2^b	19.88 ± 1.52^b	6.03 ± 0.92^b	2.40 ± 0.90^b
R	9 ± 0.8^a	13.01 ± 0.58^c	6.82 ± 1.35^b	3.63 ± 2.81^b
L	14 ± 2.4^b	23.07 ± 1.56^a	9.96 ± 1.53^a	6.57 ± 1.42^a

Values are represented as mean \pm SD, n = 3; L: Leaf; R: Roots; T: twigs, Values affected with different letters are significantly different ($p < 0.05$).




Figure 1 Viability of HL-60 cells after 48 h treatment with extracts of *A. muricata*; (n = 3); (A) HL-60 cells; L (Leaf); R (Roots); T (twigs).

A. muricata extracts induce a G0/G1 cell cycle arrest in HL-60 cells

The changes in the cell cycle distribution were shown in Figure 5A and the apoptotic cells were counted based on G0 DNA contents. The results show no changes in the cell cycle distribution of the control group, however, the accumulation of cells was found in apoptotic (G0) with significant modification of G2/M and S phases when the concentration of all the tested extracts increase (Figure 5A, B & C). The cell population of G0

phase significantly increased from 2.51 – 96% (leaf), 2.83 – 95% (roots) and 5.01 – 98% (twigs) and in the meantime the proportion G0/G1 drastically increased (Figure 5D). The results demonstrated that all the tested extracts induced apoptosis on HL-60 cells through G0/G1 phase cell cycle arrest.

Discussion

Natural polyphenols are secondary metabolites produced by plants for their defense against different types of






Figure 2 Effect of extracts *A. muricata* on nuclear morphological changes of HL-60 cells. After 24 h of treatment cells staining with Hoechst 33258 incubated for 30 min, and observed under fluorescence microscope. Olympus, Tokyo, Japan; magnification 200x. Marked morphological changes of cell apoptosis such as condensation of chromatin and nuclear fragmentations were found clearly. Apoptotic cells gradually increased in a dose-dependent manner; (A): Leaf; (B): Roots; (C): twigs.



stress, e.g. ultraviolet radiation, aggression of pathogens, low soil fertility, changes of environmental temperature, severe drought, and grazing pressure [27]. The interest on plant phenols is increasing in the recent decade because of their health promoting potential. It is widely known that diets containing an abundance of phenols have protective effects against a variety of diseases, particularly cardiovascular disease and cancer. Phenols from herbal extract are raising great interest as powerful and safe anticancer strategy for their broad range targeting capability and low side effects. Depending on the chemical structure, several beneficial effects of polyphenols and their implications in the human health have been identified including in cancer [28], neuroprotection [6,29], cardiovascular system dysfunction and damage, the metabolic syndrome, diabetes, aging, and different inflammation-related pathologies [27,30,31]. Chemotherapy drugs from polyphenols could improve the survival of cancer patients, with low side effects. Thus, it is urgent to develop novel drugs which are more effective

[32]. Novel strategies for determination of natural products with biological activity require the implementation of large-scale screening programs.

The antiproliferative activities of *A. muricata* have been carried out on several cancer cells with significant positive results [15,21]. Plant extracts with IC₅₀ values ≤ 30 µg/mL are considered pharmaceutically active [33]. Our MTT results (Table 1) indicated that extracts of *A. muricata* inhibited significantly the HL-60 cells *in vitro* and can be considered as active according to the suggestions from the National Cancer Institute (NCI) which stated that the IC₅₀ lower or equal to 20 µg/mL can be used as a benchmark for suitable screening cancer drugs from plants and herbs [34]. A dose-dependent inhibitory effect of extracts was also observed in HL-60 treated cells (Figure 1A). Among these extracts the roots exhibited the higher cytotoxic effects than other extracts (Table 1). Our results demonstrated that *A. muricata* extracts have significant cytotoxic potential on HL-60 cells.



Apoptosis is a crucial mode of programmed cell death, which is an active physiological process to eliminate selectively unnecessary cells [35]. Induction of cell apoptosis in tumor tissue is the best stage for cancer therapy [35]. Apoptosis is a common mode of action of chemotherapeutic agents, including the natural product-derived drugs. Furthermore, induction of apoptosis is recognized as an efficient strategy for cancer chemotherapy and a useful indicator for cancer treatment and prevention. Hence several researchers nowadays have performed apoptotic screening of natural products from herbal extracts in Cameroon [36,37]. Apoptosis involves specific morphological and biochemical changes such as chromatin condensation, membrane blebbing, cell shrinkage, DNA fragmentation, etc. Induction of apoptosis is the key to success of plant products as anticancer agents [38,39]. After 24 h of treatment with *A. muricata* extracts, the characteristics of apoptotic cells, including the increase of bright blue color of Hoechst 33258 staining was observed (Figure 2A, B & C) as well as evident DNA fragmentations at 100 µg/mL, which are the important hallmarks of

apoptosis [40]. The results indicated that all the extracts from *A. muricata* induce apoptosis of HL-60 cells.

Mitochondrion is an integral part of apoptotic machinery and events such as loss of mitochondrial membrane potential is classical evidence for apoptosis [35]. Usually, the decrease in MMP occurs during the early stage of apoptosis before the cell morphology changes. The sharp decrease in the membrane potential indicates the irreversible occurrence of early apoptosis due to an increase in the permeability of the mitochondrial membrane follow by the release of apoptotic factors, including cytochrome c [41]. Based on the current research, we propose that the following mechanism is responsible for extracts-induced HL-60 cell apoptosis. Briefly, extracts of *A. muricata* acts through the disruption of membrane mitochondrial to arrest cells in the G0/G1 phase and inhibit cell proliferation. Cell cycle checkpoints are control mechanisms that ensure the proper progression of cell cycle events [42]. *In vitro* apoptosis assay study presented here (Figure 5A, B, C), showed that extracts from *A. muricata* induced apoptosis in HL-60 tumor cells in a




Figure 5 Cell cycle analysis of extracts of *A. muricata* on HL-60 cells. After 24 h, treated cells were incubated with RNase (40 µg/mL), stained with propidium iodide (25 µg/mL), and analyzed by BD-FACS Caliber flow-cytometer; **(A)**: Leaf; **(B)**: Roots; **(C)**: twigs; Cells distribution after treatment with extracts of *A. muricata*; values are expressed as means ± standard error (n = 3). Values affected with different letters are significantly different (p < 0.05) from the control. L: Leaf; R: Roots; T: twigs; Ctl: control; app: apoptotic phase.

dose-dependent manner, compared to untreated cells. In this experiment, the extracts of *A. muricata* induced G0/G1 cell cycle arrest in HL-60 cells at different concentration after 24 h of treatment (Figure 5A, B, C). Therefore, we can suggest that the anticancer effects of *A. muricata* extract is associated with G0/G1 cell cycle arrest and cell differentiation. The cell cycle analysis revealed that all the extracts of *A. muricata* can markedly induce a G0/G1 phase arrest in HL-60 cells, but they have low effects on the G2/M phase. Although the induction mechanism of cell differentiation by *A. muricata* extract is not clear, a block of cell cycle progression at the G0/G1 phase may be closely related to the differentiation.

Most chemopreventive agents known today from plant extracts are subdivided into two categories: (i) blocking agents which inhibit the initiation step by preventing

carcinogen activation and (ii) suppressing agents, which inhibit malignant cell proliferation during promotion and progression steps of carcinogenesis [43]. Natural products, including medicinal plants, herbs and spices provide rich resources for anticancer drug discovery [44]. Phytochemical analysis of the extracts of *A. muricata* was performed in order to identify the chemical nature of the active principles. Quantitative analysis of the extracts revealed the presence of phenolics, flavonoids and trace amounts of flavonols (Table 1). Previous studies, demonstrated the presence of a number of phytochemicals, including phenols as flavonoids, tannins, anthraquinones and steroids in the leaf extracts of *A. muricata* [45]. Cinnamic acid derivative, Coumaric acid hexose, 5-Caffeoylquinic acid, Dihydrokaempferol-hexoside, p-Coumaric acid, Caffeic acid derivative and Dicafeoylquinic acid represented phenolics

acids group were isolated in its fruits [46]. Intensive chemical investigations of the leaves and seeds of *A. muricata* have resulted in the isolation and identification of a great number of acetogenins with interesting biological or pharmacological activities, such as antitumoral, cytotoxicity and apoptosis on HCT-116 and HT-29 cancer cells, [16,47]. Hence, previous studies revealed that several acetogenins act as a DNA topoisomerase I poison, arrested cancer cells at the G1 phase and induced apoptotic cell death in a Bax- and caspase-3- related pathways, and inhibited NADH-ubiquinone oxidoreductase (complex I) in mitochondria [21]. It can be suggested that the synergistic effects of phytochemicals present in this plant extracts including acetogenins may be the underlying principle behind the chemotherapy potential agents of *A. muricata*. These groups of molecules found in a large quantity in *A. muricata* extracts may act in synergy and might be responsible for their antiproliferation activity through the significant decrease in the mitochondrial membrane potential induction of apoptosis.

Conclusions

In conclusion, *A. muricata* exhibit antiproliferative effects of HL-60 cells by inducing loss of cell viability, morphology changes, loss of membrane mitochondrial potential and G0/G1 phase cell arrest. Our data confirm the potential of *A. muricata* as an agent of chemotherapeutic and cytostatic activity in HL-60 cells. These data suggest that these extracts have potential for cancer chemotherapy. However, the complete mechanisms underlying the therapeutic effects of the extracts need to be investigated as well as the identification of the active molecules present in the different parts of the plant.

Competing interests

The authors declared no potential conflicts of interest with respect to the research authorship and/or publication of this article.

Authors' contributions

PCA, GSK and BMMM have conducted the study; MSDN has collected the plant and carried out the extraction, FFB helped in building and correcting the manuscript, KSA provided reagents and supervised the research with JYG. All the authors read and approved the final manuscript.

Acknowledgements

The authors wish to thank the Federation of Indian Chambers of Commerce and Industry (FICCI), who provided scholarship to Dr. PIEME C.A. through CV Raman program to conduct this research in the Indian Institute of Integrative Medicine (CSIR).

Author details

¹Department of Physiological Sciences and Biochemistry, Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, PO Box 1364, Yaoundé, Cameroon. ²Cancer Pharmacology Division, Indian Institute of Integrative Medicine, 180001, Canal Road, 18001 Jammu, India. ³Department of Biochemistry and Molecular Biology, University of Buea, Buea, Cameroon. ⁴Department of Biochemistry, Faculty of Sciences, University of Yaoundé I, PO Box 812, Yaoundé, Cameroon.

Received: 3 June 2014 Accepted: 9 December 2014

Published: 24 December 2014

References

1. Abraham Z: *Glimpses of Indian Ethno botany*. New Delhi: Oxford and Publishing Co; 1981:308–320.
2. Duraipandiyar V, Ayyanar M, Ignacimuthu S: **Antimicrobial activity of some ethnomedical plants used by paliyar tribe from Tamil Nadu, India.** *BMC Compl Alternat Med* 2006. doi:10.1186/1472-6882-6-35. 635.
3. Alassane W, Yanjun Z, Caux C, Brouard JP, Pousset JL, Bodo B: **Annomicatin C, a novel cyclohexapeptide from the seeds of *Annona muricata*.** *C R Chimie* 2004, **7**:981–988.
4. George D, Pamplona R: **Encyclopedia of medical plants.** *Editorial Safelize Spain* 1999, **1**:381.
5. Hann J: *A history of the Tumucua Indians and mission.* Florida: University Press; 1996:13–21.
6. Pier O: *Pacific island ecosystem at risk, "Result set for: Annonaceae *Annona muricata*".* United States Geological Survey and United States Forest Service: Pier species lists; 2008:8.
7. Taylor L: *Technical data report for graviola, *Annona muricata*,* Volume 10. Austin: Sage Press; 2002:1–6.
8. Lans CA: **Ethnomedicines used in trinidad and tobago for urinary problems and diabetes mellitus.** *J Ethnobiol Ethnomed* 2006, **2**:45–55.
9. Vieira GHF, Mourão JA, Ângelo ÂM, Costa RA, Vieira SDF: **Antibacterial effect (in vitro) of *Moringa oleifera* and *Annona muricata* against gram positive and gram negative bacteria.** *Rev Inst Med Trop Sao Paulo* 2010, **52**(3):129–132.
10. Heinrich M, Kuhnt M, Wright CW, Rimpler H, Phillipson JD, Schandelmaier A, Warhurst DC: **Parasitological and microbiological evaluation of mixe Indian medicinal plants (Mexico).** *J Ethnopharmacol* 1992, **36**:81–85.
11. Antoun MD, Gerena L, Milhus WK: **Screening of the flora of Puerto rico for potential antimalarial bioactives.** *Int J Pharmacol* 1993, **31**:255–258.
12. Baskar R, Rajeswari V, Kumar TS: **In vitro antioxidant studies in leaves of annona species.** *Indian J Exp Biol* 2007, **4**:480–485.
13. Luna Jde S, De Carvalho JM, De Lima MR, Bieber LW, Bento Ede S, Franck X, Sant'ana AE: **Acetogenins in *Annona muricata* L. (annonaceae) leaves are potent molluscicides.** *Nat Prod Res* 2006, **20**:253–257.
14. Roslida AH, Tay CE, Zuraini A, Chan PF: **Anti-inflammatory and anti-nociceptive activities of the ethanolic extract of *Annona muricata* leaf.** *J Nat Remedies* 2010, **10**:97–104.
15. Eka prasasti NR, Tuti SS, Retno WA: **The breast of anticancer from leaf extract of *Annona muricata* againts cell line in t47d.** *Int J Applied Sci Technol* 2012, **2**(1):157–164.
16. Gajalakshmi S, Vijayalakshmi S, Devi Rajeswari V: **Phytochemical and pharmacological properties of *Annona muricata*: a review.** *Int J Pharm Pharm Sci* 2012, **4**(2):3–6.
17. Leboeuf M, Cavé A, Bhaumik PK, Mukherjee B, Mukherjee R: **The phytochemistry of the annonaceae.** *Phytochem* 1982, **21**:2783–2813.
18. Kossouoh C, Moudachirou M, Adjakidje V, Chalchat JC, Figueirédo G: **Essential oil chemical composition of *Annona muricata* L. leaves from Benin.** *J Ess Oil Res* 2007, **19**:307–309.
19. Chang FR, Liaw CC, Lin CY, Chou CJ, Chiu HF, Wu YC: **New adjacent bis-tetrahydrofuran annonaceous acetogenins from *Annona muricata*.** *Planta Med* 2003, **69**:241–246.
20. Ragasa CY, Soriano G, Torres OB, Don MJ, Shen CC: **Acetogenins from *Annona muricata*.** *Phcog J* 2012, **32**(4):32–37.
21. Rieser MJ, Kozlowski JF, Wood KV, McLaughlin J: **Muricatacin: a simple biologically active acetogenin derivative from the seeds of *Annona muricata* (Annonaceae).** *Tetrahedron Lett* 1991, **32**:1137–1140.
22. Sulaiman H, Roslida AH, Fezah O, Tan KL, Tor YS, Tan Cl: **Chemopreventive potential of *Annona Muricata* L. Leaves on chemically-induced skin papillomagenesis in mice.** *Asian Pac J Cancer Prev* 2012, **13**:2532–2533.
23. Jirovetz L, Buchbauer G, Ngassoum MB: **Essential oil compounds of the *Annona muricata* fresh fruit pulp from Cameroon.** *J Agric Food Chem* 1998, **46**:3719–3720.
24. Kumaran A, Karunakaran RJ: **In vitro antioxidant activities of methanol extracts of *Phyllanthus* species from India.** *Lebens-Wiss Technol* 2007, **40**:344–352.
25. Bhushan S, Kumar A, Malik F, Andotra SS, Sethi VK, Kaur IP, Taneja SC, Qazi GN, Singh J: **A triterpenediol from *Boswellia serrata* induces apoptosis through both the intrinsic and extrinsic apoptotic pathways in human leukemia HL-60 cells.** *Apoptosis* 2007, **12**:1911–1926.
26. Bhushan S, Kakkar V, Pal HC, Guru SK, Kumar A, Mondhe DM, Sharma PR, Taneja SC, Kaur IP, Singh J, Saxena AK: **Enhanced anticancer potential of**

- encapsulated solid lipid nanoparticles of TPD: a novel triterpenediol from *Boswellia serrata*. *Mol Pharmacol* 2013, **10**:225–235.
27. Rodríguez ML, Estrela JM, Ortega ÁL: **Natural polyphenols and apoptosis induction in cancer therapy.** *J Carcinog Mutag* 2013, **S6**:004. doi:10.4172/2157-2518.S6-004.
28. Korkina LG, De Luca C, Kostyuk VA, Pastore S: **Plant polyphenols and tumors: from mechanisms to therapies, prevention, and protection against toxicity of anti-cancer treatments.** *Curr Med Chem* 2009, **16**:3943–3965.
29. Gutierrez-Merino C, Lopez-Sanchez C, Lagoa R, Samhan-Arias AK, Bueno C, Garcia-Martinez V: **Neuroprotective actions of flavonoids.** *Curr Med Chem* 2011, **18**:1195–1212.
30. Queen BL, Tollefsbol TO: **Polyphenols and aging.** *Curr Aging Sci* 2010, **3**:34–42.
31. Accomando S, Pellitteri V, Corsello G: **Natural polyphenols as anti-inflammatory agents.** *Front Biosci* 2010, **2**:318–331.
32. Bengui Y, Jun L, Zu L, Jinrong Y, Ting N, Shu W: **Anti-tumor activity and relative mechanism of ethanolic extract of *Marsdenia tenacissima* (Asclepiadaceae) against human hematologic neoplasm *in vitro* and *in vivo*.** *J Ethnopharmacol* 2014, **153**:258–267.
33. Suffness M, Pezzuto JM: **Assays related to cancer drug discovery.** In *Methods in plant biochemistry: assays for bioactivity*. Edited by Hostettmann K. London: 6 Academic Press; 1990:71–133.
34. Pieme CA, Santosh KG, Ambassa P, Suresh K, Ngameni B, Ngogang YJ, Bhushan S, Saxena AK: **Induction of mitochondrial dependent apoptosis and cell cycle arrest in human promyelocytic leukemia HL-60 cells by an extract from *Dorstenia psilurus*: a spice from Cameroon.** *BMC Compl Alternat Med* 2013, **13**:1–9.
35. Xiaoxia C, Yaping X, Xiao Bing W, Pan W, Hongxia L, Han Y, Quanhong L: **Anti-tumor and pro-apoptotic activity of ethanolic extract and its various fractions from *Polytrichum commune* L.ex Hedwin L1210 cells.** *J Ethnopharmacol* 2012, **143**:49–56.
36. Choumssi AT, Loureiro R, Silva AM, Moreira AC, Pieme CA, Asonganyi T, Oliveira PJ, Penlap BV: **Toxicity evaluation of some traditional African spices on breast cancer cells and isolated rat hepatic mitochondria.** *Food Chem Toxicol* 2012, **50**:4199–4208.
37. Pieme CA, Santosh KG, Ngamegni B, Ngogang J, Shashi B, Tetsuya M, Saxena AK: **Ethyl acetate fraction of *Garcinia epunctata* induces apoptosis in human promyelocytic cells (HL-60) through the ROS generation and G0/G1 cell cycle arrest: a bioassay-guided approach.** *Environ Toxicol Pharmacol* 2013, **3**(6):865–874.
38. Hsiao WL, Liu L: **The role of traditional Chinese herbal medicines in cancer therapy-from TCM theory to mechanistic insights.** *Planta Med* 2010, **76**:1118–1131.
39. Sreelatha S, Jeyachitra A, Padma PR: **Antiproliferation and induction of apoptosis by *Moringa oleifera* leaf extract on human cancer cells.** *Food Chem Toxicol* 2011, **49**:1270–1275.
40. Hu W, Lee SK, Jung MJ, Heo SI, Hur JH, Wang MH: **Induction of cell cycle arrest and apoptosis by the ethyl acetate fraction of *Kalopanax pictus* leaves in human colon cancer cells.** *Bioresour Technol* 2010, **101**:9366–9372.
41. Ying W, Qingshuang Z, Junxia Z, Xiaojing Z, Jinxiu Z, Li-an W: **Iso-suillin from the mushroom *Suillus flavus* induces cell cycle arrest and apoptosis in K562 cell line.** *Food Chem Toxicol* 2014, **67**:17–25.
42. Myung-Ja Y, Jin-Kyung K, Seong-yeol P, Yunha K, Channy P, Eun Sook K, Kie-In P, Hong Seob S, Raekil P: **Potential anticancer properties of the water extract of *Inontus obliquus* by induction of apoptosis in melanoma B16-F10 cells.** *J Ethnopharmacol* 2009, **121**:221–228.
43. Duvoix A, Blasius R, Delhalle S, Schnekenburger M, Morceau F, Henry E, Dicato M, Diederich M: **Chemopreventive and therapeutic effects of curcumin.** *Cancer Lett* 2005, **223**:181–190.
44. Schwartzman G, Ratain MJ, Cragg GM, Wong JE, Saijo N, Parkinson DR, Fujiwara Y, Pazdur R, Newman DJ, Dagher, Di Leone L: **Anticancer drug discovery and development throughout the world.** *Am J Clin Oncol* 2002, **20**:475–595.
45. Solomon-Wisdom GO, Ugoh SC, Mohammed B: **Phytochemical screening and antimicrobial activities of *Annona muricata* (L) leaf extract.** *Am J Biol Chem Pharm Sci* 2014, **2**(1):1–7.
46. Jiménez VM, Maike G, Schweiggert RM, Reinhold C, Esquivel P: **Identification of phenolic compounds in soursop (*Annona muricata*) pulp by high-performance liquid chromatography with diode array and electrospray ionization mass spectrometric detection.** *Food Res Int* 2014, **65**:42–46.
47. Zorofchian Moghadamtousi S, Hamed K, Elham R, Mohammadjavad P, Mehran Fadaeinasab H, Abdul K: ***Annona muricata* leaves induce G1 cell cycle arrest and apoptosis through mitochondria-mediated pathway in human HCT-116 and HT-29 colon cancer cells.** *J Ethnopharmacol* 2014, **156**:277–289.

doi:10.1186/1472-6882-14-516

Cite this article as: Pieme et al.: Antiproliferative activity and induction of apoptosis by *Annona muricata* (Annonaceae) extract on human cancer cells. *BMC Complementary and Alternative Medicine* 2014 **14**:516.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

