

Research

Open Access

## Identification of anthranilate and benzoate metabolic operons of *Pseudomonas fluorescens* and functional characterization of their promoter regions

Diane M Retallack\*, Tracey C Thomas, Ying Shao, Keith L Haney, Sol M Resnick, Vincent D Lee and Charles H Squires

Address: The Dow Chemical Company, Biotechnology Research and Development, 5501 Oberlin Dr. San Diego, CA 92121, USA

Email: Diane M Retallack\* - dmretallack@dow.com; Tracey C Thomas - tcthomas@dow.com; Ying Shao - yshao@dow.com; Keith L Haney - klhaney@dow.com; Sol M Resnick - smresnick@dow.com; Vincent D Lee - vdlee@dow.com; Charles H Squires - csquires@dow.com

\* Corresponding author

Published: 05 January 2006

Received: 17 November 2005

*Microbial Cell Factories* 2006, 5:1 doi:10.1186/1475-2859-5-1

Accepted: 05 January 2006

This article is available from: <http://www.microbialcellfactories.com/content/5/1/1>

© 2006 Retallack et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** In an effort to identify alternate recombinant gene expression systems in *Pseudomonas fluorescens*, we identified genes encoding two native metabolic pathways that were inducible with inexpensive compounds: the anthranilate operon (*antABC*) and the benzoate operon (*benABCD*).

**Results:** The *antABC* and *benABCD* operons were identified by homology to the *Acinetobacter* sp. anthranilate operon and *Pseudomonas putida* benzoate operon, and were confirmed to be regulated by anthranilate or benzoate, respectively. Fusions of the putative promoter regions to the *E. coli lacZ* gene were constructed to confirm inducible gene expression. Each operon was found to be controlled by an AraC family transcriptional activator, located immediately upstream of the first structural gene in each respective operon (*antR* or *benR*).

**Conclusion:** We have found the anthranilate and benzoate promoters to be useful for tightly controlling recombinant gene expression at both small (< 1 L) and large (20 L) fermentation scales.

### Background

Ideally, to facilitate control of gene expression for production of proteins in an organism like *P. fluorescens*, which has been developed as a robust recombinant protein expression system [1,2], it is desirable to have a collection of expression cassettes. These cassettes would contain a variety of promoters that are tightly regulated, of differing strengths, induced under different growth conditions, and/or by different chemicals. These expression cassettes can then be linked to various genes of interest to achieve total control of those genes under typical fermentation

conditions. High levels of gene expression are often obtained in the *P. fluorescens* system using the *E. coli lacUV5* and *tac* promoters [1]. Several bacterial promoters have been previously shown to be effective at driving transgene expression in pseudomonads including the bacteriophage  $\lambda$  P<sub>R</sub> and P<sub>L</sub> promoters [3,4], which are regulated by a temperature sensitive repressor protein, and the *Pseudomonas* P<sub>m</sub>, P<sub>u</sub>, and P<sub>sal</sub> promoters [4,5], which are regulated by alkyl- or halotoluenes (P<sub>m</sub> and P<sub>u</sub>) or salicylates (P<sub>sal</sub>) and the T7 early promoter [6], regulated by isopropyl-thiogalactopyranoside (IPTG).

**Table 1: Bacterial strains used in this study**

Genus, Species	Strain Name	Relevant Genotype
<i>Escherichia coli</i>	JM109	<i>lacZ</i> <sup>-</sup>
<i>Pseudomonas fluorescens</i>	MB101	<i>lacZ</i> <sup>-</sup>
<i>Pseudomonas fluorescens</i>	MB214	<i>lacZ</i> <sup>+</sup>
<i>Pseudomonas fluorescens</i>	DC164	<i>lacZ</i> <sup>-</sup> $\Delta$ <i>pyrF</i>
<i>Pseudomonas fluorescens</i>	DC253	<i>lacZ</i> <sup>-</sup> $\Delta$ <i>benAB</i>
<i>Pseudomonas fluorescens</i>	DC284	<i>lacZ</i> <sup>-</sup> <i>benR</i> ::pCR2.1

Pseudomonads are capable of metabolizing a wide variety of aromatic hydrocarbons, including benzoate and anthranilate [7-11], which are inexpensive and non-toxic. Benzoate and anthranilate are converted to catechol by benzoate 1,2-dioxygenase together with 2-hydro-1,2-dihydroxybenzoate dehydrogenase, and anthranilate 1,2-dioxygenase respectively. These enzymes are encoded by the *benABCD* and *antABC* operons, which have been identified in several organisms [9-16], and are often regulated by transcriptional activators belonging to the AraC/XylS family of transcriptional regulators. Transcriptional activators BenR and BenM have been described, which activate transcription of the *benABCD* operon [7,17,18]. Recently transcriptional activators of two different anthranilate operons have been described [10,13].

We describe in this report the identification of the *P. fluorescens* strain MB214 *benABCD* and *antABC* operons, along with the genes coding for the transcriptional regulatory proteins BenR and AntR, respectively. The promoter regions of the *benABCD* and *antABC* operons have been defined, and regulation by BenR and AntR examined. A range of potential inducing compounds, in addition to benzoate and anthranilate, were tested including *o*-toluate, *m*-toluate, *p*-toluate and chlorinated anthranilates. Promoter elements are defined which are capable of driving heterologous gene expression at both small (1 L shake flask) and large (20 L fermentor) scales.

## Results

### ***P. fluorescens* MB214 can utilize anthranilate and benzoate as sole carbon source**

To characterize the functional catabolic and regulatory potential of *Pseudomonas fluorescens* strain MB214 (Table 1), the strain was tested for the ability to utilize a variety of aromatic and aliphatic hydrocarbons or aromatic acids as sole sources of carbon for growth. Carbon substrates were supplied at 5–10 mM as sole source of carbon and energy in liquid minimal medium (M9 containing 1 mM MgSO<sub>4</sub> and 5 mg/ml trace metals solution), unless stated otherwise. Growth in liquid media was measured spectrophotometrically at 600 nm. In most cases, growth was also examined on solid media of identical composition but containing 1.5% (w/v) agar. Naphthalene and toluene


were supplied in the vapor phase to agar plates containing carbon-free M9 medium. Growth of strain MB214 in media containing test substrates was compared to that observed in control media lacking a carbon test substrate. Results of the phenotypic screens showed that benzoate, anthranilate, 1-decanol, and 1-dodecanol were able to support growth of *P. fluorescens*. However, salicylate, phenylacetate, phthalate, phenol, naphthalene, *n*-octane, *n*-decane, *n*-dodecane, *n*-hexadecane and toluene were not able to support growth of *P. fluorescens*.

Subsequent experiments conducted in M9 media with glucose, glycerol, citrate, or succinate as primary growth substrates indicated that benzoate and anthranilate catabolism in strain MB214 was inducible by addition of the individual aromatic acid substrates. The functional evidence for catabolic pathways inducible by benzoate and anthranilate, or their metabolic intermediates, led to a systematic isolation and evaluation of the corresponding regulatory switches for the development of gene expression systems.

### **Characterization of the anthranilate and benzoate metabolic operons**

Sequence analysis and annotation of the *P. fluorescens* MB214 genome in comparison to public databases revealed putative anthranilate and benzoate operons similarly structured to the genes encoding the dioxygenase and reductase subunits clustered in one operon downstream of an AraC/XylS type transcriptional activator, likely driven by its own promoter (Figure 1). A second putative transcriptional activator, RXF04732, was identified downstream of the putative *benABCD* operon as well. As described later in this report, it is the upstream transcriptional activator that acts on the Pben promoter. The genes encoding the putative transcriptional activators, *antR* and *benR*, are orientated differently with respect to each dioxygenase operon (see Figure 1). Similar to other AraC/XylS type regulators and the recently described *Burkholderia cepacia* DBO1 *andR* gene [13], *antR* is oriented divergently from the putative *antABC* operon. However, *benR* is oriented in the same direction as *benABCD*, similar to the *benR* gene of *Pseudomonas putida* [7].

The proteins coded by the anthranilate and benzoate operons were compared to the UniProt protein database (01/05) using the BLASTP program [19]. A comparison of the AntA, AntB and AntC proteins to homologues of *P. aeruginosa*, *P. putida* and *P. resinovorans* is shown in Table 2. The *P. fluorescens* MB214 AntA, AntB, AntC and AntR proteins, were most similar to those *P. resinovorans*. Analysis of *P. fluorescens* BenA, BenB, BenC, BenD and BenR protein to homologues of *P. aeruginosa* and *P. putida* is shown in Table 3. These proteins were found to be homologous to those coded by the *P. aeruginosa* toluate 1,2-diox-



**Figure 1**

**Anthranilate and benzoate operons.** Above are depicted schematic representations (not drawn to scale) of the *P. fluorescens* MB214 anthranilate (*antR antABC* Genbank accession number [DQ172833](#)) and benzoate (*benR benABCD* Genbank accession number [DQ172832](#)) metabolic operons. Open reading frames for each gene are indicated by arrows, with the direction of the arrow corresponding to the direction of the open reading frame. Gene names are indicated below each arrow.

ygenase and *P. putida* benzoate catabolic operons. The similarity of these proteins to those coded for by previously identified catabolic operons, together with the functional characterization of anthranilate and benzoate metabolism, indicate that these operons likely code for the anthranilate and benzoate operons of *P. fluorescens*.

#### Isolation of the *Pant* and *Pben* promoters

Initially, approximately 500–700 bp regions upstream of the *antA* and *benA* genes (Figure 2) were amplified and cloned upstream of the *lacZ* reporter gene in pDOW1017. As shown in Figure 3A and 3B the resulting plasmids, pDOW1019 and pDOW1029 are found to contain benzoate and anthranilate inducible promoters, respectively. Uninduced cultures showed low level  $\beta$ -galactosidase activity in the presence of the *Pben::lacZ* fusion. However, no  $\beta$ -galactosidase activity was detected in uninduced cultures carrying the *Pant::lacZ* fusion. The *Pben* and *Pant* promoter containing fragments were next truncated to include only the region between the putative translation start of the first gene of the structural operon (*benA* and *antA* respectively) and the upstream putative transcriptional activator open reading frame (ORF). The truncated benzoate promoter fragment, *Pben278*, was amplified and cloned upstream of the *lacZ* reporter gene in pDOW1017. The *P. fluorescens* strain MB101 (Table 1) was transformed with the resulting construct, pDOW1028, and induced with benzoate to assess activity.

Figure 3A shows that *Pben278* exhibits benzoate inducible activity similar to *Pben509*, indicating that the 275 bp truncated promoter contains all elements necessary for benzoate-activated transcription. Moreover, the *Pben* fragment was further truncated at the 3' end to just 87 bp based on primer extension mapping of the transcriptional start site as described below. Similar to *Pben278*, *Pben87* was also found to be as active as the original *Pben509* fragment (data not shown). Refined annotation of the 5' region of the *benA* gene showed the translation start to be 25 base pairs downstream of the 3' end of *Pben87* (see Figure 2). The truncated anthranilate promoter fragment, *Pant311*, was amplified and cloned upstream of the *lacZ* reporter gene of pDOW1017. Little activity was detected from the truncated promoter when fused to the *lacZ* reporter gene (data not shown).

#### Overexpression of transcriptional activator increases *Pant* activity

In an attempt to improve the activity of the anthranilate inducible promoter, the promoter fragment was amplified to include the putative transcriptional activator ORF directly upstream of *antA*. This PCR fragment was cloned upstream of the *lacZ* reporter gene of pDOW1017. MB101 was transformed with the resulting plasmid, pDOW1035, and activity upon induction with anthranilate was assessed. The addition of the transcriptional activator in multicopy demonstrated faster and greater induction compared to *Pant713* (Figure 3B), which does not contain the entire transcriptional activator ORF. The activity of the culture harbouring pDOW1035 peaks early, then declines between 6 and 24 hours. This is likely due to the rapid decline in anthranilate concentration, compared to the culture carrying pDOW1029 (data not shown). Similar experiments were performed with a fragment containing *benR* and the *Pben509* fragment, however no difference in promoter activity were observed (data not shown).

#### Inactivation of the *benABCD* operon transcriptional activator

There are two putative transcriptional regulator coding sequences that flank the *benABCD*. To determine if the ORF *benR*, upstream of *benABCD*, acts as the benzoate transcriptional activator, we insertionally inactivated *benR* as described in Materials and Methods. Following inactivation of the putative benzoate transcriptional activator, the resultant strain DC284 was transformed with the *Pben::lacZ* fusion, pDOW1019, and tested for benzoate activated *lacZ* expression. As shown in Figure 4A, inactivation of *benR* results in an inability to activate *Pben::lacZ* expression. Moreover, benzoate is not metabolized as in the wild-type MB101 strain (Figure 4B) indicating that the chromosomal benzoate dioxygenase operon (*benABCD*) also fails to be induced in the transcriptional activator knockout strain.

**Table 2: Anthranilate operon homologues. Percent amino acid identity/similarity shown below. AntA: *P. aeruginosa* (UNI\_TREMBL:Q910X0); *P. putida* (UNI\_TREMBL:Q93SR3), and *P. resinovorans* (GenPept: NP\_758565). AntB: *P. aeruginosa* (UNI\_TREMBL:Q910SR8); *P. putida* (UNI\_TREMBL:Q90W9); *P. resinovorans* (GenPept: NP\_758547). AntC: *P. aeruginosa* (UNI\_TREMBL:Q90W8); *P. putida* (UNI\_TREMBL:Q93SR4), *P. resinovorans* (GenPept: NP\_758546). Anthranilate dioxygenase transcriptional activator AntR: *P. resinovorans* (GenPept: NP\_758551).**

	<i>P. aeruginosa</i> homologue (%identity/similarity)	<i>P. putida</i> homologue (identity/ similarity)	<i>P. resinovorans</i> homologue (%identity/similarity)
<i>P. fluorescens</i> AntA	80/90	45/66	91/95
<i>P. fluorescens</i> AntB	71/81	71/82	74/85
<i>P. fluorescens</i> AntC	63/77	63/77	74/84
<i>P. fluorescens</i> AntR			74/86

### Alternative inducers of Pant and Pben promoters

*Ortho*-, *meta*- and *para*-toluate were tested as inducers of Pant and Pben. Similar to other systems [11,16], *m*-toluate was found to be an inducer of the *P. fluorescens* benzoate operon promoter. Addition of 5 mM *m*-toluate induced Pben278 (pDOW1028) to significant levels, ~13 fold over background (Figure 5). However, addition of 5 mM *o*-toluate and *p*-toluate induced Pben278 only slightly. The *antR*-Pant construct (pDOW1035) was induced by 5 mM *o*-toluate approximately 19- fold over background, although to levels still lower than *m*-toluate induction of Pben (Figure 5). *P. fluorescens* growth on solid medium consisting of 1XM9 salts supplemented with either *m*-, *o*-, or *p*-toluate (5 mM) as a sole carbon source was tested in comparison to growth on similar solid medium supplemented with 0.5% glucose as a carbon source. While *P. fluorescens* grew on medium supplemented with glucose, growth on *m*-, *o*-, or *p*-toluate as carbon source was not observed even after 14 days of incubation (data not shown). These results indicated that although *m*-, *o*- and *p*-toluate could not be utilized as carbon sources, certain forms of toluate can act as significant inducers of the Pben278 and *antR*-Pant promoter fusions.

Analysis of anthranilate catabolism in strain MB214 revealed that several chloro-anthranilates acted as induc-

ers of the anthranilate catabolic pathway (data not shown). The pDOW1035 construct containing *antR*-Pant, which showed higher expression levels than pDOW1029 (Pant713), was used to compare Pant induction by 2 mM 6-chloro-anthranilate to that by 2 mM anthranilate. As shown in Figure 6A, the level induction of the *antR*-Pant::lacZ fusion by 6-chloroanthranilate was similar to that measured with anthranilate. Induction with 6-chloroanthranilate showed a steady increase in promoter activity over time, which appeared to level between 6 and 8 hours. Induction with anthranilate showed a rapid increase in promoter activity between 4 and 6 hours, which then declined from 6 to 8 hours. The decline in activity was likely to due to metabolism of the inducer. Figure 6B shows concentration of the inducing compounds throughout the time course of induction. While the concentration of 6-chloroanthranilate decreased slightly during the time course, the concentration of anthranilate decreased significantly. These results indicate not only that 6-chloroanthranilate can act as an inducer of Pant, but that it may act as a gratuitous inducer, as it does not appear to be metabolized significantly by *P. fluorescens*.

**Table 3: Benzoate operon homologues. Percent amino acid identity/similarity shown below. BenR: XylS of *P. aeruginosa* (UNI\_TREMBL:Q910W3) and *P. putida* (UNI\_SPROT:XY53\_PSEPU). BenA: *P. aeruginosa* toluate 1,2 dioxygenase alpha subunit (UNI\_TREMBL:Q910W4); *P. putida* BenA (UNI\_TREMBL:Q88140). BenB: *P. aeruginosa* toluate 1, 2-dioxygenase small subunit (UNI\_TREMBL:Q910W5); *P. putida* BenB (UNI\_TREMBL:Q88139). BenC: *P. aeruginosa* toluate 1,2 dioxygenase subunit (UNI\_TREMBL:Q9WWV0); *P. putida* BenC (UNI\_TREMBL:Q88138). BenD: *P. aeruginosa* cis 1,2-dihydroxycyclohexa-3,4-diene carboxylate dehydrogenase (UNI\_TREMBL:Q910W7); *P. putida* BenD (UNI\_TREMBL:Q88137).**

	<i>P. aeruginosa</i> homologue (% identity/similarity)	<i>P. putida</i> homologue (% identity/similarity)
<i>P. fluorescens</i> BenA	64/79	64/79
<i>P. fluorescens</i> BenB	66/78	70/79
<i>P. fluorescens</i> BenC	53/68	53/68
<i>P. fluorescens</i> BenD	77/84	75/82
<i>P. fluorescens</i> BenR	58/73	54/70

**A**

```

CTCGCTCATC GCGCACGGG TGATTACCAC CTGCAACTTG CGGCAGTCAC CGGAAATCGC 60
CAGCACCTGA TGCTCATGG GCGAAATGAT CACGCCTTGG TCGCGGTTGG AACTGAGACG 120
TTCACCGTTC TTGCTCAGCT CTTGCTCGCC CACCAGTGGC AGGCTCAAGC TGTAGCTGCT 180
GAAGTGCTCG GCGTCTTGA TGTCGATGGT CACATCAGTG CCGTACTCGA TCACGCCCGAG 240
GGTGGTGGCG CGGGATTGA ACACGTTGGC GCTGTGGTGA AAGCGCAGGC GCTCGGGGGT 300
TGCCGTGCGC AGGCGATGGG GCCCGCAGAT GCCGGACATC CAGCTGCGCG CGCCTTCCAG 360
GTCAAGCGT TGAATATGAA TATCGGCTGT CTGACTAGTC ATCAGGGTGC ACCCGCGCCG 420
GTTAGCGGTT TGCGCGCTGT GACGGCGCGT GGTGAACTT CGACAGCAAG TTCGAGGCCA 480
CGCCAGTGGC GTTCTCACTG GGTGGATAGC AAGGCTCGAC TATGTGGATA AAGCCGAGAG 540
TTTTGCGACC ATCGCCCGCC ATCAGCATAG CCGATGCGCT CACCGCGCGC CACCGTCATG 600
GGTATTGGCC GCCCAACTTT GCGGCCTACG TTCGCCCAIT AAGCGGATAG CCCGCCACCG 660
CATCGCAGCC GCTTAATGCG TCAACGTTTA GCATGATCA AAGGTGCGCT CCC...713

```

**B.**

```

CCAACGTCAG TGAACGTTTC CTGTACAGCC TGTTTGAGCG CCAGGTGGGG CTGTGCGCCG 60
GCGATTACGT ACGCCGCTGC AAGCTCGAAC GCGTACATGC ACGCTTGCAA CTAAGCAGCA 120
CGCGCAGCGT GACCGAGGTG GCTTTGGACC ATGGGTTTCAT GCACCTAGGG CCGTTTTCCG 180
AAGCCTATCG CAAACGCTTC GCGCAACTGC CGTGCAGAC CTGGAACCGC CATCGTTAAG 240
CGACGTGCGC CTGCGCGATA GCGATGTGCA GGCAGCGGAT ATTGACGGGC AGGGCGAGCA 300
CGTACGGTGA GGGCGCCTGA TACAAGAACA ACGGAGGGCC CGCCCATGA TCAGTACACT 360
CGACCGACTC GCCTGCAAT TGCGCGAGTC CGTACAGGAA GACCCCGCCA CTGGGGTGT 420
CCGCTGCGCG CCGCAGATCT TCACCGACC CGACCTGTTT GCCCTGGAGA TGAACACAT 480
CTTCGAAGGC GGGTGGATCT ACCTGGCCC 509

```

**Figure 2**  
**Anthranilate (A) and benzoate (B) promoter regions.**  
 The largest promoter region tested for each is shown. Pant311 = bases 413–713 (A, dotted underline), Pben278 = bases 236–509 (B), Pben87 = bases 236–323 (B, dotted underline). Underlined with solid lines are the putative -35 and -10 sites. A 9 bp direct repeat is shown in italics (A). Putative XylS- type activator binding sites are shown in italics (B). The transcriptional start for each promoter is shown in bold. Double underlined in B is 5' end of the *benA* coding region.

### Mapping transcriptional start sites of Pben and Pant promoters

Total RNA was isolated from benzoate induced cultures of MB101 carrying either pDOW1019 (Pben509) or pDOW1028 (Pben278). Total RNA from each was subjected to primer extension analysis using the lacZPE primer. Both constructs show the transcriptional start at a cystine nucleotide 196 bp upstream of the 3' end of the Pben278 clone (Figure 7A). This is consistent with the location of the putative -35 and -10 promoter sequences shown in italics and underlined in Figure 7C.

Total RNA was isolated from an anthranilate induced culture of MB101 carrying pDOW1029 (Pant713). Primer extension analysis was performed using the lacZPE2 primer. An extension product was observed corresponding to the adenosine nucleotide 31 bases upstream of the 3' end of the Pant713 clone (Figure 7B). This start site is consistent with the spacing of the putative -10 and -35 promoter sequences shown in italics and underlined in Figure 7D.


### Analysis of Pben and Pant activity at the 20 L fermentation scale

*P. fluorescens* MB101 carrying the plasmid pDOW1028 (Pben278 promoter) was grown in a standard 20 L fermentation, as described in Materials and Methods, and induced with 6 mM sodium benzoate. Promoter activity was monitored as a function of *lacZ* expression over the time course of the fermentation. However, under standard conditions using glucose as a carbon source, no promoter activity was observed. This was not consistent with results using glucose as a carbon source shown at the shake flask scale (Figure 3). Additional shake flask experiments revealed that the Pben278 promoter was active when either citrate or glycerol was used as a carbon source as well (data not shown). When glycerol was used in place of glucose as a carbon source for 20 L fermentation, Pben278 activity was observed following induction with benzoate, and increased over the course of the fermentation (Figure 8A). The level of induction was comparable to that observed at the shake flask scale.

The *antR*-Pant construct pDOW1035 was also tested at the 20 L scale. MB101 carrying pDOW1035 was grown under standard fermentation condition, with glucose as a carbon source. Following induction with either 5 mM or 10 mM anthranilate, promoter activity was observed (Figure 8B). The culture induced with 10 mM anthranilate showed a higher level of activity than that induced with 5 mM anthranilate, as expected. Activity was not detected until very late in the fermentation unlike Pben278, which showed an increase in activity between 6 and 24 hours (Figure 8A).

### Benzoate induces Pben509

To determine whether benzoate is in fact the inducer of the Pben509 promoter and not a downstream metabolite, we sought to block the metabolism of benzoate at the first step by deleting the *benA* and *benB* genes, which code for the large and small subunits of benzoate 1, 2-dioxygenase respectively. A deletion of *benAB* was constructed in the  $\Delta$ *proC* $\Delta$ *pyrF* strain DC164 as described in Materials and Methods. The resultant strain, DC253, was tested for the ability to metabolize benzoate and for the ability to support benzoate induced  $\beta$ -galactosidase activity from the Pben509::*lacZ* construct. To test Pben activity, DC253 was transformed with pDOW1019 and the resultant strain was induced with 5 mM benzoate. As shown in Figure 9A, Pben509::*lacZ* remains active in the *benAB* knockout strain, *i.e.*  $\beta$ -galactosidase activity is detected. The concentration of benzoate remaining in the cell-free broth as measured by HPLC showed that the *benAB* deletion mutants were unable to metabolize benzoate, while the parent strain DC164 did metabolize benzoate efficiently (Figure 9B). These results indicate that benzoate does indeed act as an inducer Pben509, and not necessarily a




**Figure 3**  
**Anthranilate and benzoate promoter activity.** Cultures were grown in M9 medium supplemented with 1% glucose. Induction of Pben promoter constructs for 24 hours is shown in panel A. Induction of Pant promoter constructs for 6 (grey bars) or 24 (black bars) hours is shown in panel B.  $\beta$ -galactosidase activity for each is depicted on the Y-axes. Promoter construct tested is indicated on the X-axis. White or hatched bars indicate uninduced samples and grey or black bars indicate induced samples. Standard deviation of triplicate wells is depicted by error bars. A representative experiment is shown.

downstream metabolite. This conclusion is consistent with the observation that Pben509 was not induced by cis-cis-muconate (data not shown).


**Discussion**

The work presented here describes the anthranilate and benzoate catabolic operons of *P. fluorescens* strain MB101. The anthranilate dioxygenase large and small subunits (AntA and AntB) and reductase (AntC) are most closely related to the anthranilate dioxygenase and reductase recently described in *P. resinovorans*. Although the transcriptional activator, AntR, is also closely related to the



**Figure 4**  
**Effect of benR inactivation on Pben278 activity.** Above is shown  $\beta$ -galactosidase activity (A) of either wild type (MB101) or *benR* inactivated (MB101 *benR::kan*) carrying pDOW1019. Cultures were grown in M9 medium supplemented with 1% glucose and induced with 2 mM benzoate for 24 hours. Standard deviation of triplicate wells is depicted by error bars. A representative experiment is shown. Benzoate metabolism over the course of the induction is shown in B, with benzoate concentration shown on the Y-axis and time post induction shown on the X-axis.


transcriptional activator of *P. resinovorans*, the chromosomal arrangement of *P. fluorescens antR* gene in relation to the *antABC* is different than that of *P. resinovorans* [10]. The *P. fluorescens antR* gene is divergently transcribed from the *antABC* operon, similar to the arrangement of the anthranilate dioxygenase operon of *B. cepacia* [13]. The *P. fluorescens AntR* represents further evidence of an AraC/XylS type transcriptional activator regulating anthranilate dioxygenase expression. Transcriptional activators for anthranilate operons have only recently been described [10,13]. Studies of *antABC* promoter activation presented here indicate that AntR may be the limiting factor in expression from the Pant promoter. Addition of *antR* in multicopy results in significant improvement of Pant activity (Figure 3) indicating that the presence of presumably higher levels of inducer alone was not sufficient to promote high levels of gene expression from the Pant promoter. It is possible that either AntR is produced at low



**Figure 5**  
**Induction of *antR*-Pant and Pben278 by toluate.**  
 Shown on the X-axis is  $\beta$ -galactosidase activity of Pben278 and *antR*-Pant constructs (pDOW1028 and pDOW1035 respectively) 8 hours post-induction. Cultures were grown in M9 medium supplemented with 1% glucose. The inducer (5 mM) used in each case is as described in the legend to the right of the bar graph. Standard deviation of triplicate wells is depicted by error bars. A representative experiment is shown.


levels, or that the protein binds weakly to the promoter region. In either case, increasing the overall concentration of the protein would result in improved promoter activity. The rapid metabolism of anthranilate in the pDOW1035 harbouring strain supports the observation that overexpression of *antR* results in improved Pant activity, since increased activity of the chromosomally located Pant promoter would result in increased expression of the anthranilate dioxygenase, and subsequently, increased anthranilate metabolism. Although AntR shows high homology to other AraC/XylS transcriptional activators, we were unable to detect a consensus XylS type DNA binding site (TGCA-N6-GGNTA) [20] upstream of the Pant core promoter. Further investigation of AntR expression and elucidation of its binding site in the Pant promoter may help to improve promoter activity. The Pben promoter fragment, however, does appear to have a XylS type DNA binding site as shown in italics in Figure 2B. The sequences TGCC-N6-GGATA and TGCA-N6-GGATA, separated by 5 bp, are located immediately upstream of the putative -35 signal. Further investigation is required to determine whether these sequences are important for Pben activation. Unlike the *antR*-Pant system, increasing the copy number of *benR* did not improve Pben activity (data not shown).

Anthranilate and benzoate have been shown to be inducers of the Pant and Pben promoters, respectively, as expected. Not unexpectedly, several other compounds were found to induce the promoters. *m*-toluate, which has



**Figure 6**  
**Induction of *antR*-Pant by 6-chloro-anthranilate.**  
 Above is shown a time course of Pant induction (MB101 carrying pDOW1035) performed in M9 media supplemented with 1% glucose. Cultures, induced with either 2 mM anthranilate (diamonds) or 2 mM 6-chloro-anthranilate (squares), were monitored for  $\beta$ -galactosidase activity (A) and inducer concentration (B) as shown on the Y-axes. Time post induction in hours is shown on the X-axes. Standard deviation of triplicate wells is depicted by error bars. A representative experiment is shown.


been shown to induce other benzoate operons, including those of *Rhodococcus* sp. strain 19070 [16] and *P. putida* mt-2 [11], was found to induce the Pben promoter 13-fold over background. *o*-toluate, which has not been previously reported to induce either benzoate or anthranilate operon promoters, was found to induce the Pant promoter 19-fold. However, *o*-toluate induction of the Pben promoter, along with *p*-toluate induction, was only 2.5 to 3-fold over background. The anthranilate promoter was also inducible with 6-chloro-anthranilate. The fact that 6-chloro-anthranilate was not metabolized significantly by *P. fluorescens* strain MB101 indicates that 6-chloro-anthranilate acts as a gratuitous inducer, and that anthranilate itself, and not necessarily a metabolite of anthranilate, may act as an inducer. Benzoate was shown to be the inducer of the Pben promoter by deleting the genes cod-



**Figure 7**  
**Mapping transcriptional start sites of Pben and Pant promoters.** Shown above (A) are primer extension products of 30 µg RNA isolate from MB101 carrying pDOW1028 (Pben278) induced with benzoate run next to the sequence ladder generated with the same primer (GATC). Shown in (B) is the primer extension product of 10 or 30 µg RNA isolated from MB101 carrying pDOW1029 (Pant713) induced with anthranilate is run to the right of a sequence ladder (ACGT) generated with the same primer. The DNA sequence of the Pben278 promoter is shown in (C), with the bold, underlined C indicating the transcriptional start site. The putative -35 and -10 regions are shown in italics and underlined and the 5' portion of the *benA* coding sequence is shown in bold italics. The 3' portion of the Pant promoter fragment is shown in (D), with the transcriptional start site indicated in bold and underlined and the translational start in bold italics. The putative -35 and -10 regions are shown in italics and underlined.

ing for the benzoate dioxygenase large and small subunits. As shown in Figure 9, the Pben promoter remains active in the absence of benzoate metabolism.

For Pben and Pant to be used as part of an expression system, it is important to identify inducing compounds that are not metabolized by the host organism. We have shown that the Pben and *antR*-Pant promoters can drive heterologous gene expression at the 20 L fermentation scale. Although the Pben promoter appears to be repressed under the glucose feed conditions used at the 20 L fermentations scale, the *antR*-Pant promoter construct was found to be active. However, a change in carbon source from glucose to glycerol alleviated the observed Pben repression at the 20 L scale. Additional fermentation optimization may result in improved promoter activity and heterologous gene expression. Further strain development, such as chromosomal deletion of the anthranilate




**Figure 8**  
**Promoter activity at the 20 L scale.** Induction of Pben278:*lacZ* with ~6 mM benzoate (A) and induction of *antR*-Pant:*lacZ* with 5 or 10 mM anthranilate (B) are shown above. β-galactosidase activity is shown on the Y-axes, and time post induction in hours is shown on the X-axes. Standard deviation of triplicate wells is depicted by error bars. A representative experiment is shown.

and benzoate dioxygenases should also improve expression during the course of induction and simplify the induction process, in that the inducer will not need to be added more than once during the course of induction to maintain expression levels.

**Conclusion**

It is important to identify and develop several recombinant expression systems to allow for flexibility and reliability in protein production. The availability of more than one inducible expression system will allow for differential expression of proteins during production. Differential expression of chaperones, foldases or disulfide bond isomerases from these alternate expression systems may aid in overall yield of active target protein. We have shown that the benzoate and anthranilate promoters examined in this work can be used to drive recombinant protein expression in *P. fluorescens* at both small (shake flask) and





**Figure 9**  
**Analysis of Pben509 activity in a  $\Delta$ benAB strain.** Shown is  $\beta$ -galactosidase activity of DC253 carrying pDOW1019 (Pben509::lacZ) uninduced (0 mM benzoate) or induced with 5 mM sodium benzoate (A). Standard deviation is depicted by error bars. Cultures were grown in LB medium supplemented with uracil and proline. Benzoate metabolism in the parental strain (DC164,  $\Delta$ pyrF) compared to the  $\Delta$ benAB strain (DC253,  $\Delta$ pyrF $\Delta$ benAB) is shown in B.

large (20 L) scales. Further development of these promoter systems will aid in expression of complex proteins in *P. fluorescens*.

## Methods

### Bacterial strains and growth conditions

Bacterial strains and plasmids used are listed in Tables 1 and 4. All plasmid construction and propagation was performed using JM109 (Promega, Madison, WI) as a host cell, functional assays were performed in *P. fluorescens* MB101 or derivatives of MB101 as described. The *P. fluorescens* strain MB214 is a derivative of MB101, which contains the *E. coli lacZYA* operon integrated into the genome. Tetracycline was used at a concentration of 15  $\mu$ g/ml in both *E. coli* and *P. fluorescens* hosts; kanamycin was used at a concentration of 50  $\mu$ g/ml in both *E. coli* and *P. fluorescens* hosts; ampicillin was used at a concentration of 100  $\mu$ g/ml in *E. coli* hosts. Cells were grown on LB

medium (Difco, Becton Dickinson, Sparks, MD), or when indicated minimal medium consisting of 1XM9 salts (Difco, Becton Dickinson, Sparks, MD), trace elements and either glucose, citrate or glycerol as a carbon source as indicated. *E. coli* strains were grown at 37°C, while *P. fluorescens* strains were grown at 30°C.

### 20 L fermentation

*P. fluorescens* fermentations were conducted in standard, aerated 20 L research fermentors with a mineral salts medium derived from Riesenberg, *et al.* [21]. Cultures were grown at 32°C and pH 6.5–7.0 was maintained through the addition of aqueous ammonia. Agitation and sparged airflow rates were gradually increased during the growth phase of the fermentation to control dissolved oxygen at a positive level (15% of saturation) but were fixed at maximum levels when these were reached. The cultures were operated as either a glucose or glycerol fed-batch or as a glycerol batch (glycerol concentration <200 g/L). The fermentation process was divided into an initial cell growth phase (typically 24–30 hours) and a gene expression (induction) phase in which inducer was added (anthranilate or sodium benzoate solution delivered as a bolus or fed) to initiate recombinant gene expression.

### General cloning methods

Restriction digestion, ligation and polymerase chain reaction were performed essentially as described in Sambrook and Russell. [22]. Restriction enzymes and modifying enzymes were purchased from New England Biolabs (Beverly, MA), and Taq polymerase was purchased from either Promega or Invitrogen (Carlsbad, CA). Genomic DNA was prepared using the Easy DNA isolation kit (Invitrogen) according to the manufacturer's protocol. Plasmid DNA was prepared using the Nucleospin or Nucleobond plasmid DNA kit (Clontech, Valencia, CA) according to the manufacturer's protocol. Primers used in this work are described in Table 5. *P. fluorescens* plasmids used in this study are derived from RSF1010 [23].

### Construction of benzoate promoter fusions

The Pben509 promoter fragment was amplified using the ben40PL and ben40PU primers with MB214 genomic DNA as template. The resulting fragment was digested with Pac I and BamH I and cloned into pDOW1017 to produce the Pben509::lacZ fusion pDOW1019. The Pben278 promoter fragment was amplified using primers benL278 and ben40PU, using pDOW1026 as template. PCR product was purified on a Microcon YM-100 column (Millipore, Billerica, MA). The PCR product was digested with Pac I and BamH I and cloned into the same sites of pNEB193 to produce pDOW1022. The BamH I-Pac I fragment of pDOW1022 containing the Pben278 fragment was then cloned into pDOW1017 to produce pDOW1028, a Pben278::lacZ fusion.

**Table 4: Plasmids Used in This Study**

Plasmid	Relevant Feature	Source
PCR2.1 TOPO	Cloning vector	Invitrogen
pNEB193	Cloning vector	New England Biolabs
pGEM-T Easy	Cloning vector	Promega
pDOW1017	Promoterless <i>lacZ</i>	This work
pDOW1019	Pben509:: <i>lacZ</i>	This work
pDOW1020	<i>antR</i> -Pant	This work
pDOW1021	<i>benR</i>	This work
pDOW1022	Pben278	This work
pDOW1026	Pben509	This work
pDOW1028	Pben278:: <i>lacZ</i>	This work
pDOW1029	Pant713:: <i>lacZ</i>	This work
pDOW1033	Promoterless <i>phoA</i>	This work
pDOW1035	<i>antR</i> Pant:: <i>lacZ</i>	This work
pDOW1041	Pben278:: <i>phoA</i>	This work
pDOW1055	Pant311:: <i>phoA</i>	This work
pDOW1081	Pben88:: <i>phoA</i>	This work
pDOW1101	Pant311:: <i>lacZ</i>	This work
pDOW1131	400 bp <i>benR</i> fragment	This work
pDOW1139	3' <i>benR</i> + 5' <i>benC</i>	This work
pDOW1261-2	<i>tetAR pyrF</i>	This work

#### Construction of anthranilate promoter fusions

The Pant713 Promoter was amplified using primers antPU and antPL and MB214 genomic DNA as the template. The resulting fragment was cloned into pDOW1017 to produce the Pant713::*lacZ* fusion pDOW1029. The Pant311 promoter was amplified using primers ant311 and antPL+SD using MB214 genomic DNA as the template. The PCR product was purified with a Microcon YM-100 column (Millipore) and the volume brought up to 50  $\mu$ l with 5 mM Tris-Cl, pH8.0 (~100 ng/ $\mu$ l). The PCR product was cloned into the pGEM-T Easy vector (Promega) and sequenced using M13F and M13R primers. The pGEM clone was digested with BamH I and Pac I then cloned into the same sites of pDOW1033 creating a Pant311::*phoA* fusion, pDOW1055.

#### Construction of promoter fusions with transcriptional activators

The putative benzoate operon transcriptional activator gene that lies upstream of *benA* was amplified using the primers benact5 and benact3, using MB214 genomic DNA as template. An ~1.2 kb fragment was isolated using the Prep-a-Gene kit (Biorad, Hercules, CA), digested with Pme I and Hind III and cloned into the same sites to pNEB193 (New England Biolabs) to produce pDOW1021. The Pme I - Hind III fragment of pDOW1021 was then cloned into the same sites of pDOW1019, upstream of the Pben509::*lacZ* fusion.

The anthranilate operon transcriptional activator that lies upstream of *antA* was amplified using the 3'antactiv primer and antPL+SD primer using MB214 genomic DNA

as template. An ~1.3 kb fragment was purified using the Prep-a-Gene kit (Biorad), digested with Hind III and BamH I and cloned into the same sites to pNEB193 to produce pDOW1020. The BamH I -Hind III fragment of pDOW1020 was then cloned into the same site of pDOW1017 to produce pDOW1035.

#### $\beta$ -galactosidase assay

Strains of interest were grown overnight (30°C shaking at 250 rpm) in 1 $\times$  M9 supplemented with 1% (w/v) glucose, 1 mM MgSO<sub>4</sub> and trace elements. Strains were then sub-cultured and induced with indicated concentrations of anthranilate or benzoate.  $\beta$ -galactosidase activity of samples was analyzed in a 96-well format. For each sample well, buffer was prepared as follows: 152  $\mu$ l Z buffer (0.06 M Na<sub>2</sub>HPO<sub>4</sub>-7H<sub>2</sub>O, 0.04 M NaH<sub>2</sub>PO<sub>4</sub>-H<sub>2</sub>O, 0.01 M KCl, 0.001 M MgSO<sub>4</sub>-7H<sub>2</sub>O) was mixed with 8  $\mu$ l 1 M  $\beta$ -mercaptoethanol. Buffer was prepared in bulk quantities: to each 900  $\mu$ l of this mix were added 1 drop 0.1%SDS and 2 drops CHCl<sub>3</sub>, and vortexed to mix. In each well, 144  $\mu$ l of the above reaction mix and 16  $\mu$ l of cells were combined. The microtiter plate was sealed and vortexed for 10 seconds, then equilibrated at room temperature for 5 minutes before addition of 50  $\mu$ l 4 mg/ml 2-nitrophenyl  $\beta$ -D-galactopyranoside (ONPG). When a significant yellow color developed, 90  $\mu$ l stop solution (1 M Na<sub>2</sub>CO<sub>3</sub>) was added and time was recorded. Reactions were read at A<sub>420</sub> and A<sub>550</sub>, soon after stopping the reaction; cell density was read at A<sub>600</sub>. Miller Units were calculated as follows:  $1000 * ((A_{420} - (1.75 * A_{550})) / (\text{time (in minutes)} * 0.1 * A_{600}))$ . An average of triplicate wells is reported for each sample.

**Table 5: Primers Used in This Study**

Primer Name	Sequence
phoA5bam	CGGGATCCGGGGCCTCCTTGCGGGT
phoA3Xho	CCGCTCGAGTTATTTAGCCCCAGAGC
ant3II	CCTTAATTAACAGGGTGCACCCACG
antPL+SD	CGGGATCCGGGAGGCACCTTTTGATCAT
antPL	CGGGATCCCTTTTGATCATGGCTAAC
antPU	GCTCTAGACTCGCTCATCGCCGACG
AntAKO5	GGAATTCTTCGTGACGATGCG
AntAKO3	CGGGATCCGCTCGCGATGCTGC
lacZPE	GGATGTGCTGCAAGGC
lacZPE2	GTAACCATGGTCATCGC
ben40PU	CGGGATCCGGGGCCAGGTAGATCCAC
ben40PL	CCTTAATTAACCAACGTCAGTGAACGTT
benactko-for	CGCGACACATTGCTGCCAG
benactko-rev	AGTATCAGCCATCGCACCTT
benL278	CCTTAATTAAGTTAAGCGACGTGCGC
3' antactiv	CCCAAGCTTCTATCGAGGCAAGCCAG
benact5'	AGCTTTGTTTAAACGCATGACGTTGTTGATTC
benact3'	CCCAAGCTTCCCGTCAATATCCGCTG
H3_5'benAKOclean	CCCAAGCTTGCCATGAGGCGGAAAACGCTGC
H3_3'benBKOclean	CCCAAGCTTCGGTGATCGCCACGCTGTGCGC
benKOmega	CATACGTCATGGCCCTCCGTTGTTC
invbenKOmega	GAACAACGGAGGGCCATGACGTATG
5'benA_seq	CTGCTGGAAAACGCCTGCCTGGAG
seq_3'benB	GAGCACTTCAAGCATCGACAGGAAC
I261-8378F	CTTCAGATCCAGACTCACCAG
I261-103R	GACCATGATTACGCCAAGCGC
M13F	GTAAAACGACGGCCAGT
M13R	CACACAGGAAACAGCTATGAC

**Analysis of anthranilate and benzoate concentrations**

HPLC analyses of culture supernatant was carried out on a Hewlett-Packard Series 1100 HPLC using an isocratic method capable of separating anthranilate, benzoate, catechol, 6-chloroanthranilate and *o*-toluate. The separation column was a ZORBAX Eclipse XDB-C8 (4.6 × 150 mm, 5 μm; Agilent P/N 993967-906) equipped with a Supelguard Discovery C18 guard column (4 × 20 mm; Supelco P/N 505129). The mobile phase contained 25% acetonitrile in 25 mM sodium dihydrogen phosphate (NaH<sub>2</sub>PO<sub>4</sub>), pH 2.5, and the flow rate was 1.0 mL/min. The method used a standard sample injection volume of 2.5 μL and compounds were detected by monitoring absorbance at 254 nm. Under these conditions, compound retention times were as follows: catechol (2.20 min), 6-chloroanthranilate (3.01 min), anthranilate (3.46 min), benzoate (4.98 min), and *o*-toluate (6.53 min). Area responses were linear (R<sup>2</sup> 0.99) for calibration standards ranging from 0.5 to 5.0 mM.

**RNA Isolation**

An overnight culture of MB101 carrying the appropriate plasmid was grown in 1 × M9 supplemented with 1% glucose (w/v), 1 mM MgSO<sub>4</sub> and trace elements. The culture was induced with 5 mM benzoate or anthranilate as

appropriate for 8 or 24 hours. Cells were pelleted and total RNA isolated using RNeasy maxi kit (Qiagen). The RNA was resuspended to a final volume of 200 μl and treated with 10 units DNaseI (RNase free) (Ambion) according to manufacturer's protocol. Following DNaseI treatment, the RNA was purified using an RNeasy midi or mini column (Qiagen). RNA concentration was determined using Ribogreen (Molecular Probes).

**Primer extension**

Primer Labeling: 1 μl 10 μM primer (either lacZPE or lacZPE2), 1 μl 10 × T4 kinase buffer, 5 μl <sup>32</sup>P γ ATP (50 uCi, Amersham Biosciences, Piscataway, NJ), 1 μl T4 kinase, 2 μl ddH<sub>2</sub>O, was incubated at 37 °C for 30–60 minutes. 5 μl of the reaction was reserved to use for sequencing ladder. 20 μl TE was added to the remaining 5 μl and purified using a G25 sephadex column (Amersham-Pharmacia) to remove unincorporated nucleotides, yielding a final concentration of 0.2 μM labelled primer.

Sequencing ladder: The Promega fmol kit, along with 1 μM labelled primer was used to generate a sequencing ladder. Plasmid template used corresponds with that contained in the strain from which RNA was isolated for the extension reaction.

Primer Extension reaction: 10–20 µg of total RNA was mixed with 0.2 pmol primer to yield a final volume of 12 µl, and incubated at 70°C for 10 min. To this was added 4 µl 5× Superscript II buffer, 2 µl 1 M DTT, 1 µl 10 mM dNTPs, 1 µl Superscript II or Thermoscript Reverse transcriptase (Invitrogen) and incubated 42°C (Superscript) or 55°C (Thermoscript) for 1 hour. 4 µl of sequencing stop solution was added to the reaction. All reactions were heated at 70°C for 2 minutes immediately before being loaded onto a 6% Long Ranger (BioWhittaker, Rockland, ME)/8 M urea/1.2× TBE gel next to the sequencing ladder. The gel was run in 0.6× TBE buffer, then dried, exposed to phosphor screen, and imaged on the Typhoon phosphorimager (Molecular Dynamics).

#### Inactivation of benzoate transcriptional activator

A DNA fragment containing a portion of the open reading frame (ORF) upstream of the *benA* gene was amplified by PCR using MB214 genomic DNA as template with the *benactKO-for* and *benactKO-rev* primers. JM109 was transformed with the resulting product cloned into the pCR2.1 TOPO vector (Invitrogen). Transformants were screened for insert by colony PCR using M13F and M13R primers, and the positive clones were further confirmed by DNA sequencing. The resulting plasmid, pDOW1131, was used to insertionally inactivate the corresponding ORF. *P. fluorescens* strain MB101 was transformed with pDOW1131, selecting on LB agar supplemented with tetracycline. Primers *benactKO-for* and either M13F or M13R were used to confirm insertion of the plasmid into the desired ORF by colony PCR. The resulting strain was named DC284.

#### Construction of *P. fluorescens* Δ*benAB*Δ*proC*Δ*pyrF*

Following the method previously described for construction of gene deletions in *P. fluorescens* [24], the plasmid pDOW1139 was constructed to facilitate deletion of the *benAB* genes as follows. The 3' portion of the *benR* gene and the 5' portion of the *benC* gene were amplified using MB214 genomic DNA as template. The *benR* region was amplified using primers H3\_5'*benAKOclean* and *benKOmega*. The *benC* region was amplified using primers H3\_3'*benBKOclean* and *invbenKOmega*. The *benR* and *benC* fragments were fused using primers H3\_5'*benAKOclean* and H3\_3'*benBKOclean* with both fragments as template. The expected 1.1 kb fragment was gel-purified using Qiaex II (Qiagen) and cloned into Srf I digested pDOW1261-2[24]. *P. fluorescens* MB101 derivative DC164 [24] was transformed with the resulting plasmid, pDOW1139. Transformants were selected by plating on LB-proline-uracil with tetracycline for selection. Since the plasmid could not replicate in *P. fluorescens*, tetracycline resistant colonies isolated following transformation resulted from the plasmid integration into the chromosome. The site of plasmid integration was analyzed by

PCR. To obtain strains that have lost the integrated plasmid by recombination between the homologous regions, single colonies of the transformants were inoculated into liquid LB supplemented with 250 µg/mL each proline and uracil (LB-proline-uracil), grown overnight, and then plated onto LB-proline-uracil and 500 µg/mL 5'-fluoro-orotic acid (FOA) to counter select for the loss of the plasmid [25]. Isolates having the phenotype expected (i.e., tetracycline sensitive, uracil auxotrophic, and FOA resistant) were selected. DNA from the resulting strain (DC253) was analyzed by PCR to confirm removal of the *benAB* region using primers 5'*benA\_seq*, *seq\_3'**benB*, M13R21, 1261-8378F and 1261-103R.

#### Authors' contributions

DMR participated in design and coordination of experiments, performed sequence alignments, identified promoter regions and participated in promoter analyses, transcriptional start site analyses, 20 L fermentation analyses and in drafting the manuscript. YS and TCT participated in promoter analyses, and in drafting portions of the manuscript. TCT also participated in anthranilate promoter analysis. YS also constructed and analyzed *benR* deletion mutant. KLH participated in 20 L fermentation analyses and in drafting portions of the manuscript. SMR examined carbon source utilization, developed HPLC methods and drafted a portion of the manuscript. VDL constructed and analyzed the *benAB* deletion mutant and contributed to a portion of the manuscript draft. CHS participated in design of experiments. All authors have read and approved the manuscript.

#### Acknowledgements

The authors would like to thank John D. Stankowski, Angel Salazar, Miracle Foy and Joe Illeman for excellent technical assistance. We also thank Lawrence Chew, Carrie Schneider, Tom Ramseier and Mani Subramanian for helpful discussions. This work was supported by the Dow Chemical Company.

#### References

1. Squires CH, Retallack DM, Chew LC, Ramseier TM, Schneider JC, Talbot HW: **Heterologous Protein Production in *P. fluorescens***. *BioProcess International* 2004, **December 2004**:54-58.
2. Chew LC, Ramseier TM, Retallack DM, Schneider JC, Squires CH, Talbot HW: ***Pseudomonas fluorescens***. In *Production of Recombinant Protein* Edited by: Gellissen G, Weinheim, Wiley-VCH; 2005.
3. Sanchez-Romero JM, De Lorenzo V: **Genetic Engineering of Non-pathogenic *Pseudomonas* strains as Biocatalysts for Industrial and Environmental Processes**. In *Manual of Industrial Microbiology and Biotechnology* Second edition. Edited by: Demain AL, Davies JE. Washington, D.C., ASM Press; 1999:460-474.
4. Schweizer HP: **Vectors to express foreign genes and techniques to monitor gene expression in *Pseudomonads***. *Curr Opin Biotechnol* 2001, **12(5)**:439-445.
5. Slater RJ, Williams R: **The Expression of Foreign DNA in Bacteria**. In *Molecular Biology and Biotechnology* Fourth edition. Edited by: Walker JM, Rapley R. Cambridge, UK, The Royal Society of Chemistry; 2000:125-154.
6. Suh SJ, Silo-Suh LA, Ohman DE: **Development of tools for the genetic manipulation of *Pseudomonas aeruginosa***. *Journal of Microbiological Methods* 2004, **58(2)**:203-212.

7. Cowles CE, Nichols NN, Harwood CS: **BenR, a XylS homologue, regulates three different pathways of aromatic acid degradation in *Pseudomonas putida*.** *J Bacteriol* 2000, **182(22)**:6339-6346.
8. Duetz WA, Marques S, de Jong C, Ramos JL, van Andel JG: **Inducibility of the TOL catabolic pathway in *Pseudomonas putida* (pWW0) growing on succinate in continuous culture: evidence of carbon catabolite repression control.** *J Bacteriol* 1994, **176(8)**:2354-2361.
9. Kessler B, Marques S, Kohler T, Ramos JL, Timmis KN, de Lorenzo V: **Cross talk between catabolic pathways in *Pseudomonas putida*: XylS-dependent and -independent activation of the TOL meta operon requires the same cis-acting sequences within the Pm promoter.** *J Bacteriol* 1994, **176(17)**:5578-5582.
10. Urata M, Miyakoshi M, Kai S, Maeda K, Habe H, Omori T, Yamane H, Nojiri H: **Transcriptional Regulation of the ant Operon, Encoding Two-Component Anthranilate 1,2-Dioxygenase, on the Carbazole-Degradative Plasmid pCARI of *Pseudomonas resinovorans* Strain CA10.** *J Bacteriol* 2004, **186(20)**:6815-6823.
11. Williams PA, Murray K: **Metabolism of benzoate and the methylbenzoates by *Pseudomonas putida* (arvilla) mt-2: evidence for the existence of a TOL plasmid.** *J Bacteriol* 1974, **120**:416-423.
12. Bundy BM, Campbell AL, Neidle EL: **Similarities between the antABC-encoded anthranilate dioxygenase and the benABC-encoded benzoate dioxygenase of *Acinetobacter* sp. strain ADPI.** *J Bacteriol* 1998, **180(17)**:4466-4474.
13. Chang HK, Mohseni P, Zylstra GJ: **Characterization and Regulation of the Genes for a Novel Anthranilate 1,2-Dioxygenase from *Burkholderia cepacia* DBO1.** *J Bacteriol* 2003, **185(19)**:5871-5881.
14. Jeffrey WH, Cuskey SM, Chapman PJ, Resnick S, Olsen RH: **Characterization of *Pseudomonas putida* mutants unable to catabolize benzoate: cloning and characterization of *Pseudomonas* genes involved in benzoate catabolism and isolation of a chromosomal DNA fragment able to substitute for xylS in activation of the TOL lower-pathway promoter.** *J Bacteriol* 1992, **174(15)**:4986-4996.
15. Neidle EL, Hartnett C, Ornston LN, Bairoch A, Reikik M, Harayama S: **Nucleotide sequences of the *Acinetobacter calcoaceticus* benABC genes for benzoate 1,2-dioxygenase reveal evolutionary relationships among multicomponent oxygenases.** *J Bacteriol* 1991, **173(17)**:5385-5395.
16. Haddad S, Eby DM, Neidle EL: **Cloning and expression of the benzoate dioxygenase genes from *Rhodococcus* sp. strain 19070.** *Appl Environ Microbiol* 2001, **67(6)**:2507-2514.
17. Collier LS, Gaines GL, Neidle EL: **Regulation of benzoate degradation in *Acinetobacter* sp. strain ADPI by BenM, a LysR-type transcriptional activator.** *J Bacteriol* 1998, **180(9)**:2493-2501.
18. Clark TJ, Phillips RS, Bundy BM, Momany C, Neidle EL: **Benzoate Decreases the Binding of cis,cis-Muconate to the BenM Regulator despite the Synergistic Effect of Both Compounds on Transcriptional Activation.** *J Bacteriol* 2004, **186(4)**:1200-1204.
19. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search tool.** *J Mol Biol* 1990, **215(3)**:403-410.
20. Gonzalez-Perez MM, Ramos JL, Gallegos MT, Marques S: **Critical Nucleotides in the Upstream Region of the XylS-dependent TOL meta-Cleavage Pathway Operon Promoter as Deduced from Analysis of Mutants.** *J Biol Chem* 1999, **274(4)**:2286-2290.
21. Riesenberger D, Horn U, Strittmatter W: **Method for improved recombinant protein expression in bacteria by monitoring and modulating protein folding using a degP promoter-based reporter system and coexpression of the chaperone Skp.** Application: WO WO , (Merck Patent G.m.b.H., Germany; Riesenberger, Ulrike); 2003:27 pp.
22. Sambrook J, Russell DW: **Molecular Cloning A Laboratory Manual.** third edition. Cold Spring Harbor , Cold Spring Harbor Press; 2001.
23. Scholz P, Haring V, Wittmann-Liebold B, Ashman K, Bagdasarian M, Scherzinger E: **Complete nucleotide sequence and gene organization of the broad-host-range plasmid RSF1010.** *Gene* 1989, **75(2)**:271-288.
24. Schneider JC, Jenings AF, Mun DM, McGovern PM, Chew LC: **Auxotrophic markers pyrF and proC can replace antibiotic markers on protein production plasmids in high-cell-density *Pseudomonas fluorescens* fermentation.** *Biotech Prog* 2005, **21(2)**:343-348.
25. Toyn JH, Gunyuzlu PL, White WH, Thompson LA, Hollis GF: **A counterselection for the tryptophan pathway in yeast: 5-fluoroanthranilic acid resistance.** *Yeast* 2000, **16(6)**:553-560.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

