

RESEARCH

Open Access

Asymptomatic *Plasmodium falciparum* infection is associated with anaemia in pregnancy and can be more cost-effectively detected by rapid diagnostic test than by microscopy in Kinshasa, Democratic Republic of the Congo

Junior R Matangila^{1,3*}, Jean Lufuluabo², Axel L Ibalanky¹, Raquel A Inocêncio da Luz³, Pascal Lutumba¹ and Jean-Pierre Van Geertruyden³

Abstract

Background: In areas of high malaria transmission, *Plasmodium falciparum* infection during pregnancy is characterized by malaria-related anaemia, placental malaria and does not always result in clinical symptoms. This situation is associated with poor pregnancy outcomes. The aim of this study was to determine the extent of asymptomatic *P. falciparum* infection, its relation with anaemia as well as the most cost-effective technique for its diagnosis in healthy pregnant women living in Kinshasa, Democratic Republic of the Congo.

Methods: In a cross-sectional study design, information on socio-demographic characteristics and cost data were collected in healthy pregnant women attending antenatal care consultations. *Plasmodium falciparum* infection was diagnosed using rapid diagnostic test (RDT), microscopy and polymerase chain reaction (PCR). Haemoglobin concentration was also determined.

Results: In total, 332 pregnant women were enrolled. RDT and microscopy data were available for all the blood samples and 166 samples were analysed by PCR. The prevalence of asymptomatic *P. falciparum* infection using microscopy, RDTs and PCR, were respectively 21.6%, 27.4% and 29.5%. Taking PCR as a reference, RDTs had a sensitivity of 81.6% and a specificity of 94.9% to diagnose asymptomatic *P. falciparum* infection. The corresponding values for microscopy were 67.3% and 97.4%. The prevalence of anaemia was 61.1% and asymptomatic malaria increased five times the odds ($p < 0.001$) of having anaemia. RDTs were more cost-effective compared to microscopy. Incremental cost-effectiveness ratio was US\$ 63.47 per microscopy adequately diagnosed case.

Conclusion: These alarming results emphasize the need to actively diagnose and treat asymptomatic malaria infection during all antenatal care visits. Moreover, in DRC, malaria and anaemia control efforts should be strengthened by promoting the use of insecticide-treated nets, intermittent preventive treatment with sulphadoxine-pyrimethamine and iron and folic acid supplements.

Keywords: Asymptomatic *P. falciparum* infection, Anaemia, Pregnancy, Cost-effectiveness, Democratic Republic of the Congo

* Correspondence: matangilaj@yahoo.fr

¹Département de Médecine Tropicale, Université de Kinshasa, B.P. 747, Kinshasa, XI, République Démocratique du Congo

³International Health Unit, Department of Epidemiology, University of Antwerp, Campus DrieEiken, Universiteitsplein 1, Wilrijk 2610, Belgium

Full list of author information is available at the end of the article

Background

Malaria infection in pregnant women is a major problem in sub-Saharan Africa (SSA) with significant risks for the mother and their offspring. Malaria in pregnancy is characterized by a secondary anaemia and the presence of parasites in the placenta known as placental malaria, leading to low birth weight, source of perinatal morbidity and mortality. In areas of high *P. falciparum* transmission, where a high proportion of the population is semi-immune, most pregnant women with malaria are asymptomatic [1-4].

The sequestration of infected erythrocytes in the intervillous spaces induces a local inflammation and a massive infiltration of immune cells (macrophages, monocytes and lymphocytes) and is often observed and described as 'inflammatory placental malaria' [2,5]. This condition disturbs exchanges between the mother and the foetus and leads to poor pregnancy outcomes, such as abortion, stillbirths, low birth weight and infant mortality [6]. Annually, up to 200,000 infant deaths are attributed to malaria during pregnancy [7,8]. Moreover, untreated asymptomatic malaria evolves in a chronic infection characterized by marked dyserythropoietic changes in the red cell precursors and increased erythrophagocytosis. It has been suggested that these changes may be mediated, at least in part, by high levels of tumour necrosis factor (TNF) [9-11]. Maternal anaemia consequently increases the incidence of maternal death during pregnancy or post-partum, and is associated with increased foetal and infant mortality, prematurity and low birth weight across many populations [12]. In malaria-endemic areas, it is estimated that 25% of pregnant women are infected with malaria parasites, with the greatest risk of infection and morbidity in primiparous adolescents, and those co-infected with HIV [13].

Diagnosis of asymptomatic *P. falciparum* infection during pregnancy is, therefore, essential to prevent poor pregnancy outcomes, but presents a great challenge. The absence of clinical signs decreases the chances of a clinical diagnosis and a management guided by a decisional algorithm. Moreover, microscopic diagnosis is often challenged by fluctuating and very low parasite densities, due to placental parasite sequestration, the most common form of asymptomatic malaria during pregnancy [14]. Previous studies showed that RDTs detecting *P. falciparum* histidine-rich protein 2 (HRP2) are more sensitive than microscopy and appear to be reliable predictors of adverse outcomes of malaria in pregnancy [15]. However, more recent studies have brought conflicting results [16]. While RDTs have been extensively evaluated in malaria among the non-pregnant population, there are few published data on the performance and cost-effectiveness of RDT for diagnosing asymptomatic *P. falciparum* infection during pregnancy in malaria-endemic areas.

In the Democratic Republic of the Congo (DRC), previous studies have evaluated the prevalence of malaria and placental infection [17], the practice of intermittent preventive treatment (IPTp) [18] and the use of insecticide-treated net (ITN) [19] in pregnant women. However, there is very little data on asymptomatic *P. falciparum* infection and its contribution to anaemia among pregnant women. The latest studies were conducted in 1988 and 2003 and reported a prevalence of malaria infection among women attending antenatal clinics (ANCs) to be respectively 22% and 49% [20,21]. These prevalence estimates in endemic settings were done by microscopy and may have been different if other methods, such as RDT (Rapid Diagnostic Test) and PCR (Polymerase Chain Reaction), had been used [22].

Therefore, this study determined through microscopy, RDTs and PCR the prevalence of asymptomatic *P. falciparum* infection during pregnancy and its relation to anaemia. In addition, their cost and cost-effectiveness to diagnose asymptomatic *P. falciparum* infection during pregnancy, in semi-rural setting in Kinshasa, were assessed.

Methods

This study was conducted at the Centre Hospitalier de Kingasani II (CHK), commonly known as Maternité des Soeurs. With twenty to thirty deliveries daily, it is the most frequented Centre in Kinshasa and countrywide. Located in the heart of the semi-rural areas of the south-eastern suburbs of Kinshasa, the maternity provides invaluable health care at an accessible rate.

In a cross-sectional survey, the prevalence of asymptomatic *P. falciparum* infection was determined in apparently healthy pregnant women, going to the CHK for their first antenatal care (ANC) visit. Data and blood samples were collected prior to any routine administration of intermittent preventive treatment (IPT). Participants were recruited from July 27th, 2012 to August 27th, 2012. Women not providing written informed consent or presenting fever, muscle aches or other symptoms suggestive for malaria were excluded. Women who received anti-malarial treatment within the past two weeks were also excluded from the study. A structured questionnaire was used to obtain information on age, parity, and gestational age, level of education, previous or current use of anti-malarial drugs and the use of bed nets.

Blood sample collection

Blood samples were collected from a finger prick for laboratory analysis which included: thick blood smears for microscopy, RDT performance, determination of haemoglobin concentration and molecular analysis. For molecular analysis, blood was collected on a filter paper (Whatmann 3 MM), dried thoroughly, put in individual zip lock plastic

bags containing desiccant and stored at room temperature (< 25°C) until completion of the study and then transported to the Tropical Diseases Research Centre (TDRC) in Zambia for molecular analyses.

Laboratory analysis

Microscopy, RDTs and PCR were used to detect *P. falciparum* infection. Giemsa-stained thick blood smears (TBS) were used for microscopy. Blood slides were examined using light microscopy at 1,000 × magnification. Hundred microscopic fields were examined in the thick smear before concluding that a blood slide was negative. All slides were read twice by experienced microscopists. If the discrepancy was greater than 15%, a third reader was used to confirm diagnosis. The parasite density per microlitre of blood was computed using the following formula: (Number of trophozoites × 6,000)/Number of leucocytes. Besides microscopy, rapid diagnostic tests (RDTs) were also performed. In this study the SD Bioline Malaria Ag Pf[®] detecting HRP2 was used.

Plasmodium species-specific diagnostic PCR assay

As microscopy in pregnancy has a lower sensitivity due to placental sequestering of parasites and RDTs have internal validity problems, species specific PCR was conducted on all microscopy negative samples but RDT positive samples. In addition, 166 randomly chosen samples were analysed using nested Plasmodium species diagnostic PCR assay. Molecular analysis was performed at TDRC Ndola, Zambia.

The blood spot samples of 5 mm diameter size were soaked in 0.5% saponin in phosphate-buffered saline (PBS), incubated for 10 minutes at room temperature in a 1.5-ml tube, and centrifuged at 14000 rpm. The supernatant was discarded and blood spots were washed in 1 ml of PBS. One hundred Fifty microlitres of a 2% Chelex-100 resin work solution (Bio-Rad Richmond, CA) and 50 µl of water (pH 9.5) were added to the sample in a 1.5-ml tube and incubated at 100°C for 10 minutes. After centrifugation at 10,000 g for 1 min, the supernatant was collected and stored at -20°C, prior to the PCR assay.

Polymerase chain reaction investigation

Nested PCR assay was performed as a two-step procedure. Firstly, amplification of Plasmodium genus specific fragment was carried out as follow: PCR mixture containing buffer, dNTPs, MgCl₂, primers, Taq polymerases and sterile water. All PCR reactions were carried out in a total volume of 20 µl. One µl of the purified template DNA was used for the first reaction, in which the fragment spanned by rPLU5 (5' CCTGTT GTTGCCCTTAAACTTC 3') and rPLU6 (5' TTAAAATGTGTT GCAGTTAAAACG 3') was amplified. Secondly, a 1 µl aliquot from the product of the first PCR reaction was subsequently used as a template for

Plasmodium falciparum-specific fragment amplification using FAL1 (5' TTAAACTGGTTTGGGAAAACCAAAT ATATT3') and FAL2 (5' ACACAATGAACTCAA TCAT GACTACCCGTC 3') specific primers [23]. Positive and negative controls were always included in the assays. A negative control without DNA template and a positive control with appropriate template (3D7) were always included.

PCR products were detected by running 20 µl of DNA product on a 3% agarose gel, small fragments (Eurogentec[®]), which was subsequently stained with a 0.5 µg/ml ethidium bromide solution and visualized under ultraviolet transillumination. The specific size of the PCR product (second amplification) was 205 bp for *P. falciparum*.

Haemoglobin concentration

Determining the concentration of haemoglobin was performed with a portable HemoControl[®] device. Anaemia was defined by a haemoglobin concentration <11 g/dL. Anaemia was classified as severe anaemia Hb <7 g/dl, moderate anaemia Hb: 7–9.9 g/dl and mild anaemia Hb: 10–10.9 g/dl. [24].

Statistical analysis

The population of pregnant women in the health zone of Kingasani was estimated at 7,187 [25]. With an expected malaria prevalence of 30%, a desired accuracy set at 5% and a confidence interval of 95%, the minimum sample size was calculated at 309 pregnant women.

Data was entered and stored in Epi info[™]7. Descriptive statistics were employed for the analysis of socio-demographic data. Frequencies were used to assess the prevalence of asymptomatic malaria and anaemia in pregnant women. The χ^2 test was used to investigate associations between categorical variables. Odds ratios (ORs), 95% confidence intervals (CIs) were calculated and $p < 0.05$ values were considered to be statistically significant. Multivariate logistic regression models were constructed to identify factors associated with asymptomatic malaria or anaemia during pregnancy. Based on a priori knowledge and a P value less than 0.05 considered as significant, the following variables: Age, sex, parity, marital status, ownership of bed net, presence of lattice on windows, malaria infection and geophagia were included in the model. Backward regression technique was used to construct the model. Statistical analyses were performed using SPSS statistical program, version 17 (SPSS, Chicago, IL, USA).

The sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of microscopy and RDT (SD Bioline malaria Ag Pf[®]) were determined with PCR as gold standard, for the 166 samples (50%) of which PCR results were available.

Collection of data to estimate the cost of *P. falciparum* infection diagnosis

Financial data were obtained by conducting interviews with Hospital managers and laboratory staff in Kingasani health Zone. Costs were collected in Congolese Francs (CDF) and converted to US dollars (exchange rate US\$1 = CDF 920, July 2012). The cost for one thick blood smear includes: a glass slide, cotton, lancet, Giemsa and other stains, immersion oil, alcohol and gloves. The personnel cost for the service provided corresponded to the time spent from drawing blood samples, preparing the thick and thin smears, staining, reading and reporting the results. At CHK II as in all public maternities in Kinshasa, RDT kits are donated and the personnel service is free of charge. In the present study, for RDT diagnostic strategy, the effective contact time with those seeking care, which accounted for the personnel cost, and the cost of RDT kit were the main input parameters to figure out the amount of money a pregnant women would pay. The effective contact time was comprised of: drawing blood samples from patients, applying samples onto the test, test reading and reporting of results. The time was recorded on a "time sheet" by laboratory personnel for every RDT and microscopy service provided. Laboratory staff work eight hours from Monday to Friday and 5 hours on Saturday, thus 68 hours (4,080 minutes) weekly. Assuming a month of four weeks; laboratory staff work 272 hours (16,320 minutes) monthly. Staff salary data were provided by the staff manager.

Ethical considerations

This study was approved by the Ethical Committee board of the University of Kinshasa. Test results of the thick blood smear were revealed to all participants three days after sample collection. All women were given mebendazole, iron and folic acid on a routine basis and were treated with SP regardless of the laboratory test results.

Results

Socio-demographic characteristics of the pregnant women

In total, 332 pregnant women from 18 to 39 years old agreed to participate in the study. The median age was 27 years (interquartile range (IQR) = 22-33), and 26% were primigravidae, the median gravidity was 3 (IQR = 1-5). Eighty one percent of the subjects were married and 66.2% had not completed secondary school. Thirty-seven percent (37%) of the pregnant women had started their antenatal care in the third trimester. The mean gestational age in primigravidae at the time of the first prenatal visit was 21.5 weeks (SD \pm 4.2) vs. 23.6 weeks (SD \pm 5.8) in multi-gravida ($p < 0.01$) (Table 1).

Prevalence of asymptomatic *P. falciparum* infection in pregnant women at the first prenatal care (PNC) visit

RDT and microscopy data were available for 332 blood samples and of these, 166 samples were analysed by PCR. The prevalence of asymptomatic *P. falciparum* infection using microscopy, RDTs and PCR, were respectively 21.6% (95% CI:17.4-26.6%), 27.4% (95% CI:22.5-32.6%) and 29.5% (95% CI:22.7-37.1%). *Plasmodium malariae* was diagnosed in five women by microscopy, but was not confirmed by PCR. The median parasite density was 126/ μ l (IQR:105-162). In bivariate analyses, age, gravidity, marital status and the use of bed nets demonstrated a significant association with asymptomatic malaria. The regression model shows that being younger than 20 years old was an independent risk factor associated with asymptomatic *P. falciparum* infection (AOR 3.2;95% CI:1.2-8.2, $p = 0.02$), while being married (AOR 0.3;95% CI:0.2-0.8, $p = 0.01$) and spending the night under a bed net (AOR 0.4; 95% CI:0.2-0.7, $p < 0.01$) was associated with low prevalence of asymptomatic *P. falciparum* infection (Table 1).

Prevalence of anaemia

The mean haemoglobin concentration was 10.5 ± 1.4 g/dL and the prevalence of anaemia was 61.1% (95% CI:55.7-66.4). Of these anaemic pregnant women, 70 (21.1%), 132 (39.7%), 1 (0.3%), had respectively, mild, moderate and severe anaemia. In multivariate analyses, asymptomatic malaria increased five times the likelihood of having anaemia (AOR 5 95% CI:2.3-10.1; $p < 0.001$) (Tables 1 and 2).

Possession and use of insecticide-treated nets

Of all participants, 43.1% (95% CI:37.7-48.6%) possessed a bed net. Of these, 69.8% stated that their nets were treated with insecticide and 81.8% reported having slept under the bed net the night before the interview (Table 1).

PCR confirmation of the RDT positive but microscopy negative cases

Out of 332 samples, 92 were RDT positive and 72 were microscopy positive. All 72 microscopy positive samples were confirmed by RDT. Analysis of the 20 samples that were microscopy negative but RDT positive revealed that 65% (13/20) were positive by PCR (95% CI: 40.8- 84.6) (Figure 1).

Validity of microscopy and RDT using species specific PCR as golden standard

PCR was performed on 166 random chosen samples. Using PCR as the gold standard, microscopy and RDTs had a sensitivity of respectively 67.3% (95% CI:52.5-80.1) and 81.6% (95% CI:68.0-91.2) to diagnose asymptomatic *P. falciparum* infection. Specificity was 97.4% (95% CI:92.7-99.5) and 94.9% (95% CI:89.2-98.1) for microscopy

Table 1 Predictors for asymptomatic *P. falciparum* infection and anaemia in asymptomatic pregnant women in semi-rural suburbs of Kinshasa, 2012

Socio-demographic characteristic	N	Asymptomatic <i>P. falciparum</i>						Anaemia				
		Positive TBS	OR	P value	AOR 95% CI	P value	Anaemia	OR	P value	AOR	P value	
Age	<20	47(14.2)	24(51.1)	5.2	<0.001	3.2 (1.2-8.2)	0.02*	34(72.3)	1.8	0.09	0.9 (0.3-2.4)	0.8
	20-49	285(85.8)	48(16.9)	1				169(59.3)	1			
Gravidity	Primigravidae	86(25.9)	27(31.4)	2.0	0.01	0.8 (0.3-1.7)	0.5	59(68.6)	1.5	0.1	1.5 (0.8-2.9)	0.2
	Multigravidae	246(74.1)	45(18.3)	1				144(58.5)	1			
Gestational age	1 st et 2 ^{ème} trimester	209(63.0)	42(20.1)	0.8	0.4	0.7 (0.4-1.3)	0.3	134(61.8)	1.1	0.7	1.1 (0.6-1.8)	0.7
	3 rd trimester	123(37.0)	30(24.4)	1				69(60.0)	1			
Marital Status ^o	Married	259(80.9)	41(15.8)	0.22	<0.001	0.3 (0.2-0.8)	0.01*	153(59.1)	0.7	0.2	1.3 (0.6-2.9)	0.6
	Single	61(19.1)	28(45.9)	1				41(67.2)	1			
Ownership and use of bed net	Yes and used	117(35.2)	13(11.1)	0.4	0.002	0.4 (0.2-0.7)	0.005*	72(61.5)	1			
	Yes not used	26(7.8)	10(38.5)	1.8	0.2	1.9 (0.8-4.9)	0.2	17(65.4)	1	0.8	0.9 (0.3-2.4)	0.8
	No	189(56.9)	49(25.9)	1				114(60.3)	1.2	0.7	0.7 (0.5-1.2)	0.2
Presence of lattice on windows	Yes	55(16.6)	13(23.6)	1.1	0.7	1.1 (0.5-2.5)	0.7	36(65.5)	1.2	0.5	1.6 (0.5-4.4)	0.4
	No	277(83.4)	59(21.3)	1				164(60.3)	1			
Malaria	Yes	72(21.7)	NA					60 (83.3)	4.1	<0.001*	5 (2.3-10.1)	<0.001*
	No	260(78.3)	NA					143(55.0)	1			
Geophagia	Yes	180 (54.2)	NA					103(57.2)	0.7	0.1	0.7 (0.5-1.2)	0.2
	No	152 (45.8)	NA					100(65.8)	1			

NA: Not applicable *Significant at $p < 0.05$, OR- odds ratio, AOR- adjusted odds ratio, C.I- confidence interval. ^oN = 320.

Table 2 Prevalence of anaemia in pregnant women in semi-rural suburbs of Kinshasa, 2012 (N = 332)

			Percentage	Median (Parasite density)
Haemoglobin concentration	Severe anaemia <7 g/dl	1	0.3%	128 (IQR: 128-128)
	Moderate anaemia 7-9.9 g/dl	135	40.7%	100 (IQR: 78-132)
	Slight anaemia 10-10.9 g/dl	67	20.2%	127 (IQR: 103-159)
	Non-anaemic ≥ 11 g/dl	129	38.9%	120 (IQR: 83-135)

and RDTs, respectively. RDTs demonstrated a positive predictive value (PPV) of 87.0% (95% CI:73.7-95.1%) and a negative predictive value (NPV) of 92.5% (95% CI:86.2-96.5%). The corresponding values for microscopy were 91.7% (95% CI:77.5-98.2%) and 87.7% (95% CI:80.8-92.8%). Microscopy combined to RDT demonstrated a sensitivity of 81.6% (95% CI:68.0-91.2), a specificity of 94.9% (95% CI:89.2-98.1), a PPV of 87.0% (95% CI:73.7-95.1%) and NPV of 92.5% (95% CI:86.2-96.5%). The differences were not significant for any comparison (Table 3).

Cost-effectiveness of RDT and microscopy

On the basis of the 166 samples for which PCR results were available, malaria RDT was the most effective with the number of cases correctly diagnosed being 40 (90.9%) compared to 33(88.6%) for microscopy (Table 4). The corresponding proportion of patients correctly diagnosed was 90.9% and 88.6%, respectively. Routine use of RDT would result into an additional 2.3% of patients correctly diagnosed in comparison to microscopy. Thus, incremental cost-effectiveness ratio was US\$ 63.47 per adequately diagnosed case for microscopy when compared to RDT strategy (Table 5).

Discussion

In the most frequented maternity in Kinshasa, located in the heart of the semi-rural areas Kinshasa, the prevalence of asymptomatic malaria at the first antenatal visit was 27% and 21%, using respectively RDT and microscopy. PCR showed that almost one third and one fifth were missed by microscopy and RDT respectively. Microscopy results (21%) were similar to the 22% value reported in 1988 [20].

These values are paradoxically lower than the 42% found in Lubumbashi, an area of low malaria transmission [21]. Asymptomatic infection is a characteristic in areas of high malaria transmission; therefore, one would expect a higher prevalence in Kinshasa. It can be argued that the prevalence of asymptomatic *P. falciparum* infection may have been somewhat lower, because the study was carried out during the dry season, period subjected to a mild/low malaria transmission. Therefore, another survey during the rainy season may be informative.

Parasite density was also found to be very low in all infected pregnant women. *P. falciparum*-infected but asymptomatic individuals tend to have low and submicroscopic parasite densities [26,27]. Moreover, during pregnancy placental sequestration of parasitized red blood cells may lead to the absence or low grade of peripheral parasitaemia [28].

Other studies also found a protective effect of bed nets against malaria in pregnancy [29]. Other studies have shown that younger women may be more susceptible than older women to malaria because they are still in the process of acquiring natural immunity to pregnancy related malaria [28]. Single women were more likely to have asymptomatic *P. falciparum* infection compared to married women. In addition, owning a mosquito net was associated with being married, as previously reported by Pettifor *et al.* in 2008 [19]. This indicates that marriage or living with a partner, may offer women an ideal setting to promote maternal health. Gravidity did not show any association with asymptomatic *P. falciparum* infection. This indicates that malaria endemicity may be lower than expected impairing the development of immunity against pregnancy-related malaria




Figure 1 Flow diagram for malaria diagnosis and subsequent PCR analysis in semi-rural suburbs of Kinshasa, 2012.

Table 3 Comparison of RDT and microscopy performance using PCR as golden standard in semi-rural suburbs of Kinshasa, 2012 (N = 166)

		PCR results			Performances			
		Positive	Negative	Total	Sensitivity	Specificity	PPV	NPV
Microscopy	+	33	3	36	67.3% (52.5-80.1)	97.4% (92.7-99.5)	91.7% (77.5-98.2)	87.7% (80.8-92.8)
	-	16	114	130				
RDT	+	40	6	46	81.6% (68.0-91.2)	94.9% (89.2-98.1)	87.0% (73.7-95.1)	92.5% (86.2-96.5)
	-	9	111	120				
Microscopy and RDT	+	40	6	46	81.6% (68.0-91.2)	94.9% (89.2-98.1)	87.0% (73.7-95.1)	92.5% (86.2-96.5)
	-	9	111	120				

during successive pregnancies. Similar observations were made in other low malaria-endemic areas [30-33], stressing the need for further studies to assess the endemicity in the DRC. An individual predictor impairing acquisition of immunity is HIV infection [34]. It is, however, very unlikely that this would have affected the present results, as HIV prevalence in ANC in Kinshasa is estimated at 3.9% [35].

Pregnant women were coming for their first antenatal visit at the time of enrolment, thus prior to any administration of IPTp with SP. However *P. falciparum* infection in early pregnancy has been associated with adverse outcomes such as low birth weight LBW [33]. The use of bed nets would, therefore, be a suitable option to protect pregnant women during this period. Almost half of the surveyed women 43% declared to own a bed net. These results are higher than the 26%, 33% and 7% previously reported by UNDP in 2009 [35], Pettifor *et al.* in 2008 [19] and the Programme National de Lutte contre le Paludisme (PNLP) in 2007 [36]. However, only 35% of all pregnant women slept under a bed net, which may suggest a slight awareness of malaria during pregnancy in this specific group [19].

High prevalence of anaemia was observed and strongly correlated with asymptomatic *P. falciparum* infection. This prevalence is similar to those reported in Kisangani in 2000 (65%) and Lubumbashi (in average 65%) in 2003

[37,21]. However, Kinshasa, Kisangani and Lubumbashi belong to three different epidemiological malaria transmission patterns. These findings and the fact that a high proportion of non-malaria infected women were anaemic suggest the presence of other risk factors for anaemia in pregnant women, although the contribution of malaria seems to be significant. The aim of the study was not to assess the risk factors for anaemia in pregnant women; a dedicated study would be more appropriate to disclose this useful information. There was no significant difference in the density of parasitaemia in those with mild, moderate and severe anaemia (Table 2). A similar observation was also reported from Nigeria in 2009 [38]. A limitation of this study was that HIV infection, which is also known to cause anaemia [39], was not documented. Considering HIV when constructing the regression model could have provided more accurate information on the contribution of asymptomatic malaria.

In the present study, no significant difference was found between microscopy and RDT using PCR as the gold standard. This observation was also made elsewhere [16,40]. Combination results of RDTs and microscopy did not improve the sensitivity, specificity, PPV as well as NPV to diagnose asymptomatic *P. falciparum* infection. In addition, all the RDT positive but microscopy negative samples were analysed by PCR and it showed that 65% were in fact sub-microscopic infections. The PCR negative samples could be either explained by the persistence of HRP-2 circulation in the blood more than two weeks even after successful clearance of infected erythrocytes in the bloodstream or by the sequestration of the parasites in the placenta while HRP-2

Table 4 Comparison of effectiveness of the two diagnostic strategies (N = 166)

Tests results	Diagnostic strategy	
	RDT	Microscopy
True positive	40	33
False positive	6	3
True negative	111	114
False negative	9	16
Total sample analysed	166	166
Number correctly diagnosed	151	147
Proportion correctly diagnosed	90.9	88.6

Table 5 Results of the cost-effectiveness analysis of the microscopy compared to RDT, Kingasani health zone, Kinshasa, 2013

Diagnostic strategy	Cost (US\$)	Additional cost (US\$)	Effectiveness	Additional effectiveness	ICER (US\$)
Microscopy	2.62	1.46	0.886		63.47
RDT	1.16		0.909	0.023	

ICER: Incremental Cost-effectiveness Ratio.

Table 6 Cost components and unit costs considered for monthly asymptomatic *P. falciparum* diagnosis, at CHK II, in Kingasani Health Zone, Kinshasa in 2013

Items	Unit cost (US\$)	Microscopy strategy (US\$) (n = 332)	RDT strategy (US\$) (n = 332)
Exams and supplies			
Thick blood smear ^a	0.72	239.04	-
SD Bioline Malaria Antigen Pf® - one test	0.79	-	262.28
Salary			
Laboratory technician salary	304.34 (monthly)	588.18	123.82
Equipment			
Microscope- one unit – value	520.82 (annual)	43.4 (monthly)	-
Training*			
Microscopy - one annual course	-	-	-
RDT-one annual course	-	-	-
Total		870.62	386.1

^aCorresponds to the individual cost of an examination, which includes a glass slide, Giemsa and other stains (all components of stains), oil immersion, lancet, cotton, alcohol and gloves.

*Training costs were not available.

circulate [41]. PCR showed that almost one third and one fifth were missed by microscopy and RDT respectively. These findings highlight the necessity of strengthening IPTp in pregnancy.

This study demonstrated that RDT technique was more cost-effective than microscopy. This finding is supported by other studies carried out in other settings [42,43]. In malaria microscopy, the major cost input was the personnel salary and laboratory equipment. The major parameter that determined the cost of RDT-based strategy was cost of the test. The study did not consider the capital costs of setting up the infrastructure for microscopy diagnosis, which is known to be costly, as well as personnel training from which data were not available (Table 6). If considered, microscopy costs would be higher. Studies have shown that preventive efforts (IPT with SP and insecticide-treated bed net) still leave a large proportion of women with peripheral parasitaemia which is associated with anaemia and impaired delivery outcomes [44]. This highlights the necessity of assessing other strategies such as prompt screening and treatment of women when they present to antenatal care. Therefore, RDT use may be the best available diagnostic alternative for asymptomatic malaria screening in remote or semi-rural endemic areas where infrastructure for microscopy diagnosis is not in place, or where high accuracy of microscopy cannot be assured like in Kinshasa [45].

Conclusion

This study reports a high prevalence of anaemia among pregnant women in the health zone of Kingasani, Kinshasa and anaemia is strongly correlated to asymptomatic *P. falciparum* infection. These results are quite alarming and emphasize the need to actively diagnose and treat asymptomatic malaria infection using appropriate techniques during ANC visits in areas of high malaria transmission,

regardless of IPTp. On the other hand, though the use of ITNs in DRC is slightly increasing, the promotion of ITN use should be further encouraged. The present study suggests that RDTs may replace blood smears screening of asymptomatic *P. falciparum* infection in pregnancy in some malaria-endemic settings by virtue of their ease-to-use, ability to detect sub-microscopic infection and cost-effectiveness compared with microscopy.

Abbreviations

AOR: Adjusted Odds-ratio; CI: Confidence Interval; DRC: Democratic Republic of the Congo; IQR: Inter Quartile Range; OR: Odds-ratio; PCR: Polymerase Chain Reaction; PNLP: Programme National de Lutte contre le Paludisme; PRDT: Rapid Diagnostic Test; SSA: Sub-Saharan Africa; TBS: Thick Blood Smear.

Competing interests

The authors declare that they have no competing interest.

Author's contributions

JMR participated in the conception and design of the study protocol, conducted the study, drafted the manuscript and participated in molecular analysis. AI and L participated in the design of the study protocol and conducted the study. RIL, PL, JPV reviewed the manuscript and provided critical inputs. All authors read and approved the final manuscript.

Acknowledgements

The authors thank all pregnant women who participated in this study and the staff of Centre Hospitalier de Kingasani II (Centre des Soeurs). We also thank Rose Tujibikila, Bruno Nsilulu, and Celestin Atero for assisting in sample collection and analysis. Eric Njunju is thanked for his technical assistance in the setting up of the PCR assays. The team thanks David Mwakazanga for assisting in statistical analysis and Professor Paul Gwakisa for participating in the revision of the manuscript. This study was sponsored by Southern African Centre of Infectious Diseases (SACIDS) and VLIR UOS project (ZRDC2012 70447).

Author details

¹Département de Médecine Tropicale, Université de Kinshasa, B.P. 747, Kinshasa, XI, République Démocratique du Congo. ²Institut Supérieur de Techniques Médicales, Kinshasa RDC, Kinshasa, République Démocratique du Congo. ³International Health Unit, Department of Epidemiology, University of Antwerp, Campus DrieEiken, Universiteitsplein 1, Wilrijk 2610, Belgium.

Received: 24 December 2013 Accepted: 28 March 2014
 Published: 2 April 2014

References

1. Steketee RW, Wirima JJ, Campbell CC: **Developing effective strategies for malaria prevention programs for pregnant African women.** *Am J Trop Med Hyg* 1996, **55**(Suppl 1):95–100.
2. Nosten F, Rogerson SJ, Beeson JG, McGready R, Mutabingwa TK, Brabin B: **Malaria in pregnancy and the endemicity spectrum: what can we learn?** *Trends Parasitol* 2004, **20**:425–432.
3. Sullivan AD, Nyirenda T, Cullinan T, Taylor T, Harlow SD, James SA, Meshnick SR: **Malaria infection during pregnancy: intrauterine growth retardation and preterm delivery in Malawi.** *J Infect Dis* 1999, **179**:1580–1583.
4. Verhoeff FH, Brabin BJ, Chimsuku L, Kazembe P, Broadhead RL: **Malaria in pregnancy and its consequences for the infant in rural Malawi.** *Ann Trop Med Parasitol* 1999, **93**(Suppl 1):S25–S33.
5. Ordi J, Menendez C, Ismail MR, Ventura PJ, Palacin A, Kahigwa E, Ferrer B, Cardesa A, Alonso PL: **Placental malaria is associated with cell-mediated inflammatory responses with selective absence of natural killer cells.** *J Infect Dis* 2001, **183**:1100–1107.
6. Menendez C, Ordi J, Ismail MR, Ventura PJ, Aponte JJ, Kahigwa E, Font F, Alonso PL: **The impact of placental malaria on gestational age and birth weight.** *J Infect Dis* 2000, **181**:1740–1745.
7. Ismail MR, Ordi J, Menendez C, Ventura PJ, Aponte JJ, Kahigwa E, Hirt R, Cardesa A, Alonso PL: **Placental pathology in malaria: a histological, immunohistochemical, and quantitative study.** *Hum Pathol* 2000, **31**:85–93.
8. van Geertruyden JP, Thomas F, Erhart A, D'Alessandro U: **The contribution of malaria in pregnancy to perinatal mortality.** *Am J Trop Med Hyg* 2004, **71**(Suppl 2):35–40.
9. Steketee RW, Nahlen BL, Parise ME, Menendez C: **The burden of malaria in pregnancy in malaria-endemic areas.** *Am J Trop Med Hyg* 2001, **64**(Suppl 1-2):28–35.
10. Faquin WC, Schneider TJ, Goldberg MA: **Effect of inflammatory cytokines on hypoxia-induced erythropoietin production.** *Blood* 1992, **79**:1987–1994.
11. Jelkmann W, Pagel H, Wolff M, Fandrey J: **Monokines inhibiting erythropoietin production in human hepatoma cultures and in isolated perfused rat kidneys.** *Life Sci* 1992, **50**:301–308.
12. Leopardi O, Naughten W, Salvia L, Colecchia M, Matteelli A, Zucchi A, Shein A, Muchi JA, Carosi G, Ghione M: **Malaric placentas. A quantitative study and clinic-pathological correlations.** *Pathol Res Pract* 1996, **192**:892–898.
13. Schantz-Dunn J, Nour NM: **Malaria and pregnancy: a global health perspective.** *Rev Obstet Gynecol* 2009, **2**:186–192.
14. Brabin BJ, Romagosa C, Abdelgalil S, Menendez C, Verhoeff FH, McGready R, Fletcher KA, Owens S, D'Alessandro U, Nosten F, Fischer PR, Ordi J: **The sick placenta: the role of malaria.** *Placenta* 2004, **25**:359–378.
15. Minja DT, Schmiegelow C, Oesterholt M, Magistrado PA, Bostrom S, John D, Pehrson C, ndersen D, Deloron P, Salanti A, Lemnge M, Luty AJ, Alifrangis M, Theander T, Lusingu JP: **Reliability of rapid diagnostic tests in diagnosing pregnancy-associated malaria in north-eastern Tanzania.** *Malar J* 2012, **11**:211.
16. Kyabayinze DJ, Tibenderana JK, Nassali M, Tumwine LK, Riches C, Montague M, Counihan H, Hamade P, van Geertruyden JP, Meek S: **Placental Plasmodium falciparum malaria infection: operational accuracy of HRP2 rapid diagnostic tests in a malaria endemic setting.** *Malar J* 2011, **10**:306.
17. Lukuka KA, Fumie OS, Mulumbu MR, Lokombe BJ, Muyembe TJ: **[Malaria prevalence at delivery in four maternity hospitals of Kinshasa City, Democratic Republic of Congo]. (in French).** *Bull Soc Pathol Exot* 2006, **99**:200–201.
18. Likwela JL, D'Alessandro U, Lokwa BL, Meuris S, Dramaix MW: **Sulfadoxine-pyrimethamine resistance and intermittent preventive treatment during pregnancy: a retrospective analysis of birth weight data in the Democratic Republic of Congo (DRC).** *Trop Med Int Health* 2012, **17**:322–329.
19. Pettifor A, Taylor E, Nku D, Duvall S, Tabala M, Meshnick S, Behets F: **Bed net ownership, use and perceptions among women seeking antenatal care in Kinshasa, Democratic Republic of the Congo (DRC): opportunities for improved maternal and child health.** *BMC Public Health* 2008, **8**:331.
20. Mbanzulu PN, Leng JJ, Kaba S, Mputu L, Ngimbi NP, Makengo N, Ngbege: **[Malaria and pregnancy. Epidemiological situation in Kinshasa (Zaire)] (in French).** *Rev Fr Gynecol Obstet* 1988, **83**:99–103.
21. Kalenga MK, Nyembo MK, Nshimba M, Foidart JM: **[Anemia associated with malaria and intestinal helminthiasis in Lubumbashi](in French).** *Sante Publ* 2003, **15**:413–421.
22. Rantala AM, Taylor SM, Trottman PA, Luntamo M, Mbewe B, Maleta K, Kulmala T, Ashorn P, Meshnick SR: **Comparison of real-time PCR and microscopy for malaria parasite detection in Malawian pregnant women.** *Malar J* 2010, **9**:269.
23. Snounou G, Viriyakosol S, Zhu XP, Jarra W, Pinheiro L, do Rosario VE, Thaithong S, Brown KN: **High sensitivity of detection of human malaria parasites by the use of nested polymerase chain reaction.** *Mol Biochem Parasitol* 1993, **61**:315–320.
24. WHO: **Iron deficiency anaemia assessment, prevention and control: a guide for programme managers.** Geneva, Switzerland: World Health Organization; 2001.
25. Ministère de la Santé: **Présentation de la Zone de Santé de Kingasani.** ; 2008.
26. Trape JF, Peelman P, Morault-Peelman B: **Criteria for diagnosing clinical malaria among a semi-immune population exposed to intense and perennial transmission.** *Trans R Soc Trop Med Hyg* 1985, **79**:435–442.
27. Roucher C, Rogier C, Dieye-Ba F, Sokhna C, Tall A, Trape JF: **Changing malaria epidemiology and diagnostic criteria for Plasmodium falciparum clinical malaria.** *PLoS One* 2012, **7**:e46188.
28. Brabin BJ: **An analysis of malaria in pregnancy in Africa.** *Bull World Health Organ* 1983, **61**:1005–10016.
29. Mathanga DP, Campbell CH, Taylor TE, Barlow R, Wilson ML: **Socially marketed insecticide-treated nets effectively reduce Plasmodium infection and anaemia among children in urban Malawi.** *Trop Med Int Health* 2006, **11**:1367–1374.
30. Agan T, Ekabua J, Udoh A, Ekanem E, Efiok E, Mgbekem M: **Prevalence of anemia in women with asymptomatic malaria parasitemia at first antenatal care visit at the University of Calabar Teaching Hospital, Calabar, Nigeria.** *Int J Womens Health* 2010, **2**:229–233.
31. Anchang-Kimbi JK, Achidi EA, Nkegoum B, Sverre-remark-Ekstrom E, Troye-Blomberg M: **Diagnostic comparison of malaria infection in peripheral blood, placental blood and placental biopsies in Cameroonian parturient women.** *Malar J* 2009, **8**:126.
32. Douamba Z, Bisseye C, Djigma FW, Compaore TR, Bazie VJ, Pietra V, Nikiema JB, Simpore J: **Asymptomatic malaria correlates with anaemia in pregnant women at Ouagadougou, Burkina Faso.** *J Biomed Biotechnol* 2012, **2012**:198317.
33. Valea I, Tinto H, Drabo MK, Huybregts L, Sorgho H, Ouedraogo JB, Guiguemde RT, van Geertruyden JP, Kolsteren P, D'Alessandro U: **An analysis of timing and frequency of malaria infection during pregnancy in relation to the risk of low birth weight, anaemia and perinatal mortality in Burkina Faso.** *Malar J* 2012, **11**:71.
34. Mount AM, Mwapasa V, Elliott SR, Beeson JG, Tadesse E, Lema VM, Molyneux ME, Meshnick SR, Rogerson SJ: **Impairment of humoral immunity to Plasmodium falciparum malaria in pregnancy by HIV infection.** *Lancet* 2004, **363**:1860–1867.
35. UNDP-DRC: **Kinshasa, profile summary: Poverty and living conditions of households.** ; 2009.
36. President's Malaria Initiative: **Malaria Operational Plan Fiscal Year 2012.** Democratic Republic of Congo; 2011. http://www.pmi.gov/countries/mops/fy12/drc_mop_fy12.pdf.
37. Kanku T, Labama L, Wembodinga U: **Anémie de la femme enceinte à Kisangani : Prévalence et risque liés à quelques indications de risque.** *Congo Med* 2000, **13**:915–920.
38. Nwagha UI, Ugwu VO, Nwagha TU, Anyaehie BU: **Asymptomatic Plasmodium parasitaemia in pregnant Nigerian women: almost a decade after Roll Back Malaria.** *Trans R Soc Trop Med Hyg* 2009, **103**:16–20.
39. Nkhoma ET, Kalilani-Phiri L, Mwapasa V, Rogerson SJ, Meshnick SR: **Effect of HIV infection and Plasmodium falciparum parasitemia on pregnancy outcomes in Malawi.** *Am J Trop Med Hyg* 2012, **87**:29–34.
40. Dhorda M, Piola P, Nyehangane D, Tumwezeze B, Nalusaji A, Nabasumba C, Turyakira E, McGready R, Ashley E, Guerin PJ, Snounou G: **Performance of a histidine-rich protein 2 rapid diagnostic test, Paracheck Pf(R), for detection of malaria infections in Ugandan pregnant women.** *Am J Trop Med Hyg* 2012, **86**:93–95.
41. Humar A, Ohrt C, Harrington MA, Pillai D, Kain KC: **Parasight F test compared with the polymerase chain reaction and microscopy for the diagnosis of Plasmodium falciparum malaria in travelers.** *Am J Trop Med Hyg* 1997, **56**:44–48.
42. Lubell Y, Reyburn H, Mbakiliwa H, Mwangi R, Chonya K, Whitty CJ, Mills A: **The cost-effectiveness of parasitologic diagnosis for malaria-suspected patients in an era of combination therapy.** *Am J Trop Med Hyg* 2007, **77**(Suppl 6):128–132.
43. Uzochukwu BS, Obikeze EN, Onwujekwe OE, Onoka CA, Griffiths UK: **Cost-effectiveness analysis of rapid diagnostic test, microscopy and**

syndromic approach in the diagnosis of malaria in Nigeria: implications for scaling-up deployment of ACT. *Malar J* 2009, **8**:265.

44. De Beudrap P, Turyakira E, White LJ, Nabasumba C, Tumwebaze B, Muehlenbachs A, Guérin PJ, Boum Y, McGready R, Piola P: **Impact of malaria during pregnancy on pregnancy outcomes in a Ugandan prospective cohort with intensive malaria screening and prompt treatment.** *Malar J* 2013, **12**:139.
45. Muhindo HM, Ilombe G, Meya R, Mitashi PM, Kutekemeni A, Gasigwa D, Lutumba P, van Geertruyden JP: **Accuracy of malaria rapid diagnosis test Optimal-IT((R)) in Kinshasa, the Democratic Republic of Congo.** *Malar J* 2012, **11**:224.

doi:10.1186/1475-2875-13-132

Cite this article as: Matangila *et al.*: Asymptomatic *Plasmodium falciparum* infection is associated with anaemia in pregnancy and can be more cost-effectively detected by rapid diagnostic test than by microscopy in Kinshasa, Democratic Republic of the Congo. *Malaria Journal* 2014 **13**:132.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

