

RESEARCH

Open Access

# Cytochrome P450 single nucleotide polymorphisms in an indigenous Tanzanian population: a concern about the metabolism of artemisinin-based combinations

Karol J Marwa<sup>1\*</sup>, Theresa Schmidt<sup>2</sup>, Maria Sjögren<sup>2</sup>, Omary MS Minzi<sup>3</sup>, Erasmus Kamugisha<sup>4</sup> and Göte Swedberg<sup>2</sup>

## Abstract

**Background:** Artemisinin-based combinations currently recommended for treatment of uncomplicated *Plasmodium falciparum* malaria in many countries of sub-Saharan Africa are substrates of CYP enzymes. The cytochrome enzyme system is responsible for metabolism of about 80-90% of clinically used drugs. It is, therefore, important to obtain the pharmacogenetics of the population in the region with respect to these combinations and thereby enable practitioners to predict treatment outcomes. The aim of this study was to detect and determine allelic frequencies of CYP2C8\*2, CYP2C8\*3, CYP3A4\*1B, CYP3A5\*3 and CYP2B6\*6 variant alleles in a Tanzanian indigenous population.

**Methods:** Genomic DNA extraction from blood obtained from 256 participants who escorted patients at Karume Health Centre in Mwanza Tanzania, was carried out using the Gene JET™ Genomic DNA purification kit (Thermo Scientific). Genotyping for the cytochrome P450 variant alleles was performed using predesigned primers. Amplification was done by PCR while differentiation between alleles was done by restriction fragment length polymorphism (PCR-RFLP) (for CYP2C8\*2, CYP2C8\*3) and sequencing (for CYP2B6\*6, CYP3A5\*3 and CYP3A4\*1B).

**Results:** CYP2C8\*2, CYP2C8\*3, CYP3A5\*3, CYP3A4\*1B and CYP2B6\*6 variant allelic frequencies were found to be 19,10,16,78 and 36% respectively.

**Conclusion:** Prevalence of CYP2C8\*2, CYP3A5\*3, CYP3A4\*1B and CYP2B6\*6 mutations in a Tanzanian population/ subjects are common. The impact of these point mutations on the metabolism of anti-malarial drugs, particularly artemisinin-based combinations, and their potential drug-drug interactions (DDIs) needs to be further evaluated.

**Keywords:** Cytochrome P450, Artemisinin-based combination therapy, Poor metabolizers, Tanzania, *Plasmodium falciparum* malaria

## Background

Prevention and treatment of malaria is mainly based on the use of drugs [1]. Treatment response in *Plasmodium falciparum* malaria depends on various factors such as parasite resistance, host natural immunity, drug quality and the pharmacokinetics of the administered drug [2]. Various studies have been carried out on molecular mechanisms for resistance in malaria chemotherapy [3]. However limited efforts have been made to determine

genetic polymorphisms in cytochrome P450 enzymes, which may be associated with treatment failure (in extensive metabolizers) as a result of subtherapeutic drug concentrations. Cytochrome P450 (CYP) polymorphisms may also cause toxicity (in slow metabolizers) due to high drug concentrations and emergence or spread of drug resistance (in extensive metabolizers) as a result of subtherapeutic drug concentrations. Recent studies have also associated drug resistant selection with poor metabolizer (PM) phenotype.

The increasing use of artemisinin-based combinations as an effective treatment of resistant malaria demands determination of genetic polymorphisms in the metabolism of

\* Correspondence: carol\_maro@yahoo.com

<sup>1</sup>Department of Pharmacology, Catholic University of Health and Allied Sciences, Mwanza, Tanzania

Full list of author information is available at the end of the article

these drugs and its influence on pharmacokinetic profiles between individuals, adverse drug reactions and clinical outcome.

Polymorphisms in gene encoding for drug-metabolizing enzymes and transporters are associated with individual variation in drug response [4]. The frequency of variant alleles encoding for CYP families varies among populations according to race and ethnic background [5].

Cytochrome P450 metabolize about 80-90% of clinically used drugs [5-8]. About 40% of cytochrome P450 dependent-drug metabolism is due to polymorphic enzymes [9]. Polymorphisms occur in all members of the CYP2C subfamily, which are CYP2C8, CYP2C9, CYP2C18 and CYP2C19 [10,11]. The genes for these subfamilies are located on chromosome 10q24 [12]. The role of CYP genes in metabolism varies significantly. For example, CYP2C has been shown to be responsible for metabolism of about 20% of clinically used drugs and endogenous substances [10].

CYP2C8 has been proven to make up 7% of the total CYP content of the liver [13] and plays a major role in the metabolism of many clinically available drugs [14-16] accounting for about 5% of the drugs used clinically [17]. The CYP2C8\*2 allele has been reported to be common in Africans, but rare in Asians and Caucasians, whereas CYP2C8\*3 has been reported to be common in Caucasians, but rare in Africans or Asians. The latter allele (CYP2C8\*3) has been associated with a marked reduction in amodiaquine metabolism in Caucasian population [14].

CYP2B6 is located within the CYP2B gene cluster on chromosome 19 [18] and is one of the most polymorphic genes in humans [19]. This gene contributes 2-10% of the total CYP content [20]. CYP2B6\*6 is more frequent in African descent than Caucasoids [21,22] and is associated with raised plasma concentrations of anti-retroviral (efavirenz, nevirapine) [19] and anti-malarial drugs (artemisinin and arteether) [1,19].

Cytochrome P450 subfamily CYP3A is the most abundant CYP in the human liver and small intestine [23]. CYP3A4, which is part of CYP3A gene cluster, is involved in the metabolism of approximately 50% of clinically used drugs [5,24].

CYP3A4 is widely considered to be the dominant CYP3A isoform, but this has been questioned lately when data have shown that the amount of CYP3A5 is much larger than previously assumed in individuals expressing this enzyme [25]. The most common single nucleotide polymorphism (SNP) within the CYP3A4 family is CYP3A4\*1B [26], an A to G transition at 392 of the proximal promoter region of the gene. The CYP3A4\*1B variant allele is shown to be associated with higher expression of CYP3A4 than the wild type allele, which could be caused by a repressor element in the wild

type promoter [27,28]. CYP3A4\*1B is associated with poor metabolism of artemether, lumefantrine [29,30] and quinine [31].

The expression of CYP3A5 in the human liver differs vastly between ethnic groups and within the groups. CYP3A5\*3 is the most frequently recognized non-functional allele and one of the most frequent polymorphism among cytochrome P450 genes being recognized [32]. In Caucasians, the frequency of CYP3A5\*3 has been showed to be  $\geq 90\%$  [33] whilst the occurrence among black Africans differs between 11 and 78% over the African continent [34].

This study aimed at determining presence of CYP 450 variant alleles (CYP2C8\*2, CYP2C8\*3, CYP2B6\*6, CYP3A4\*1B and CYP3A5\*3) responsible for the metabolism of artemisinin-based combinations employed in the treatment of uncomplicated malaria and complicated malaria (injectable artesunate) in Tanzania where malaria is endemic.

## Methods

### Study area and subjects

This was a cross-sectional study conducted from April 2013 to April 2014 at Karume Health Centre in Igombe, a semi-urban and malaria meso-endemic area in Ilemela district, Mwanza region, Tanzania, with a population of about 40,000.

### Sampling procedure

Serial sampling method was employed where every third and sixth individual was selected until the sample size was reached.

### Blood sample collection and DNA extraction

Venous blood (1-2 ml) was collected into sterile EDTA-containing tubes from 256 healthy volunteers after obtaining informed consent from each of the participants. The collected blood was stored at  $-40^{\circ}\text{C}$  until genomic DNA extraction was done. Genomic DNA extraction was performed with Gene JET Genomic DNA Purification Kit (Thermo Scientific) according to the manufacturer's instructions.

### Genotyping

#### PCR conditions

The PCR master mix contained 2  $\mu\text{l}$  1X Dream Taq Green Buffer (Fermentas), 2  $\mu\text{l}$  1 mM dNTP Mix (Thermo Scientific), 10  $\mu\text{l}$  0.5  $\mu\text{M}$  reverse primer (Eurofins), 10  $\mu\text{l}$  0.5  $\mu\text{M}$  forward primer (Eurofins), 0.25  $\mu\text{l}$  1.25 U Dream Taq (Thermo Scientific).

PCR amplification was done using primers shown in Table 1 and conditions indicated in Table 2: 2.5% agarose gel electrophoresis was performed to confirm successful PCR amplification for all samples.

**Table 1 Primers used**

SNP	Primers		Reference
	Forward	Reverse	
CYP3A4*1B	5'-ATGGCCAAGTCTGGGATGAG-3'	5'-CTCACCTCTGTT CAGGGAAAC-3'	[29,35]
CYP2B6*6	5'-TGAGTGATGGCAGACAATCACA-3'	5'-CAAGTTGAGCATCTTCAGGAACT-3'	[29,35]
CYP3A5*3	5'-TGGAGAGTGGCATAGGAGATAC-3'	5'CCATACCCCTAGTTGTACGACACA-3'	[29,35]
CYP2C8*2	5'-GAACCAAGCATCACTGGA-3'	GAAATCAAATACTGATCTGTTGC-3'	[36]
CYP2C8*3	5'-CTAAAGGACTTGGTAGGTGCA-3'	5'-CAGGATGCGCAATGAAGACC-3'	[29,35]

**Restriction fragment length polymorphism (PCR-RFLP) for CYP2C8\*2 and CYP2C8\*3**

The PCR product (10 µL) was digested with ten units of **BclI** (for CYP2C8\*2) in Buffer G (Thermo Scientific) at 55°C for three hours or five units of **BseRI** (for CYP2C8\*3) in CutSmart Buffer (New England Biolabs) at 37°C for 15 min. Digestion was followed by thermal inactivation of BclI at 80°C for 20 min. Five µL of inactivated digested product mixed with a drop of loading dye was analysed by electrophoresis on a 2.5% agarose gel containing ethidium bromide and visualized under UV light illuminator. The mutant variant of CYP2C8\*2 was undigested and observed as a band of 107 bp, while wild type DNA was observed as bands of 57 and 50 bp after BclI digestion. Results with bands at 107, 57 and 50 bp were recorded as heterozygotes. For CYP2C8\*3, wild type DNA was observed as bands of 187 and 282 bp after BseRI digestion. Mutants were undigested with one band at 439 bp and heterozygotes yielded bands of 157,282 and 439 bp.

**CYP3A4\*1B, CYP3A5\*3 and CYP2B6\*6 sequencing**

The PCR products were purified from excess dNTP and primers. To every PCR sample of 15 µl, 7.5 µl of ExoSAP Mix was added consisting of 0.019 µl Exonuclease I (20 U/µl), 0.190 µl SAP (Shrimp Alkaline Phosphate 1 U/µl) and 7.290 µl of sterile water. The samples were incubated at 37°C for 30 min and then at 90°C in a thermocycler for 5 min. In order to prepare for sequencing 2 µl of the PCR-product was mixed with 0.4 pmol of the primer and 15.6 µl of sterile water. The samples were

then sent for sequencing at Uppsala Genome Centre after DNA quantification.

**Ethical clearance**

Ethical and study approval was granted by the joint Catholic University of Health and Allied Sciences (CUHAS)/Bugando Medical Centre (BMC) Institutional Review Board. All participants signed a written informed consent.

**Statistical analysis**

Allelic frequencies were analysed using SPSS 17, SNPs analysed were evaluated using the online tool Hardy-Weinberg equilibrium calculator including analysis for ascertainment bias [37].

**Results**

A total of 256 subjects were studied for the frequency of SNPs in this study. The median age of the subjects was 28 (IQR 22-35) years and 80.5% (206/256) were females. All study subjects were natives. Successfully amplified samples were 239 for CYP2C8\*2 out of 256, 92 for CYP3A4\*1B out of 100, 97 for CYP3A5\*3 out of 100, 100 for CYP2C8\*3 out of 100 and 95 for CYP2B6\*6 out of 100. For CYP3A4, the A392G mutation known as CYP3A4\*1B was investigated. The allele prevalence was found to be 78% among the 92 subjects included in the study, where 57 (62%) of them were homozygote for the mutant variant. The allele prevalence of the A6986G mutation in CYP3A5, named CYP3A5\*3 was 16% among 97 subjects. Out of the 97 subjects, only two (2%) were homozygous for the mutation.

**Table 2 PCR conditions employed for various SNPs**

SNP	PCR conditions					
	Initial denaturation	Denaturation	Annealing	Extension	Final extension	Hold temp
CYP3A4*1B	94°C for 2 min	94°C for 30 s (30 cycles)	60°C for 30 s (30 cycles)	72°C for 90 s (30 cycles)	72°C for 5 min	4°C
CYP2B6*6	96°C for 3 min	96°C for 30 s (40 cycles)	64°C for 1 min 30 s (40 cycles)	72°C for 1 min 30 s (40 cycles)	72°C for 10 min	4°C
CYP3A5*3	96°C for 3 min	96°C for 30 s (40 cycles)	59.3°C for 30 s (40 cycles)	72°C for 30 s (40 cycles)	72°C for 10 min	4°C
CYP2C8*2	96°C for 3 min	96°C for 30 s (40 cycles)	56°C for 1 min (40 cycles)	72°C for 1 min (40 cycles)	72°C for 10 min	4°C
CYP2C8*3	96°C for 3 min	96°C for 30 s (30 cycles)	64°C 1 min 30 s (30 cycles)	72°C 1 min 30 s (30 cycles)	72°C for 10 min	4°C

For CYP2C8 the A805T mutation known as CYP2C8\*2 was investigated. The allele prevalence was found to be 19% among the 239 subjects included in the study, where 11 (4.6%) of them were homozygote for the mutant variant. The G416A mutation (CYP2C8\*3) was also analysed, the allele prevalence was identified to be 10% among the 100 subjects included in the study, where 0 (0%) were homozygote for the mutant variant. For CYP2B6, the G15631T mutation (CYP2B6\*6) was investigated. The allele prevalence was found to be 36% among the 95 subjects included in the study, where 11 (11.6%) were identified as homozygote for the mutant variant. The allele frequencies were all at Hardy Weinberg equilibrium. These results are in agreement with several earlier studies from other African countries (Tables 3 and 4).

## Discussion

The present study was designed to establish the frequencies of common CYP polymorphisms in Mwanza, Tanzania, as a background for future studies on drug metabolism in clinical trials with anti-malarial drugs. Since artemisinin-based combination therapy (ACT) is the recommended treatment, the study focused on the markers that have been shown to be mostly related with metabolism of anti-malarial drugs, taking into account that some of these mutations have been shown to affect therapeutic response and safety during treatment.

In general, there were no big differences in allele frequencies in comparison with other studies done on African populations (Tables 3 and 4). Despite there being limited information in Tanzania, a few studies on CYP markers done have been performed in Zanzibar, where the population history is different from mainland Tanzania, and Mwanza is geographically distant from Zanzibar. It cannot thus be taken for granted that genetic polymorphisms should be the same in these locations.

**Table 3 Allele frequency of CYP3A4\*1B and CYP3A5\*3 reported from previous studies in African populations**

Country	Number of subjects	CYP3A4*1B (%)	CYP3A5*3 (%)
Senegal	178	79 [38]	-
Cameroon	72	78.0 [39]	-
East Africa	-	-	36 [40]
South Africa	980,987	73.0-76.0 [41,42]	15.0 [41]
Ghana	129-204	72.0-81 [38,43]	15.0 [43]
Mozambique	86	73.8 [44]	-
Sao Tomé and	78	67.7 [44]	-
Tanzania (Zanzibar)	99, 92	69.2 [45]	15.8 [45]
Tanzania (Mainland)	165	74 [29]	18 [29]
Guinea Bissau	-	74 [46]	-
Zimbabwe	100	-	77.6 [47]

**Table 4 Allele frequencies of CYP2C8\*2, CYP2C8\*3 and CYP2B6\*6 reported from previous studies in African populations**

Country number	CYP2C8*2 (%)	CYP2C8*3 (%)	CYP2B6*6 (%)
Burundi 60	-	-	50.0 [48]
Burkina Faso 275	15.5 [49]	0.3 [49]	-
Ghana 200	17-17.9 [50-52]	0 [50]	46.9 [22]
Zimbabwe 74	-	-	49 [53]
Mozambique 115	16.0 [54]	-	-
Senegal 88	22.0 [36]	-	-
Tanzania 165 (Zanzibar)	13.9 [55]	2.1 [55]	-
Tanzania 165 (Mainland)	-	-	34-41.8 [29,56]
South Africa 182 (Xhosa)	-	-	32.14 [57]
Madagascar 153	15.0 [36]	-	-

This study identified similar findings to Zanzibar in the CYP3A5\*3 variant allele: 16 vs 15.8% [45], and for CYP2C8\*3 where none of the patients was homozygous mutant [55]. However the study recorded 10% heterozygous CYP2C8\*3 which differs from the 0% recorded from earlier studies in Zanzibar [55] and a slightly higher CYP2C8\*2 (19%) compared to 13.9% recorded in Zanzibar [55].

This study identified that 4.6% of participants were homozygotes for the slow metabolizer allele (CYP2C8\*2) representing a significant population at high risk to amodiaquine toxicity due to amodiaquine accumulation, taking into account artesunate-amodiaquine is an alternative to artemether-lumefantrine in the treatment of uncomplicated malaria in Tanzania and other sub-Saharan African countries [1,58].

The mutation A6986G leads to splicing defects in intron 3, introducing a stop codon causing severely decreased enzyme activity [25]. The homozygous mutants of CYP3A5\*3 was detected in 2% of the study subjects suggesting poor metabolism to anti-malarial drugs such as quinine, quinidine, artemether, chloroquine, halofantrine, lumefantrine, mefloquine, primaquine, artemisinin, arteether, arterlinic acid and proguanil [1], which suggests that a significant proportion of the population is exposed to toxicity of these anti-malarial drugs, including ACT, such as artemether-lumefantrine and artesunate-mefloquine during treatment taking into account that there is a high diversity in CYP3A5 in Africa.

CYP3A5 diversity is higher in East Africa [6,40], which has higher frequencies than other parts of sub-Saharan Africa [6]. The diversity explains a difference between CYP3A5\*3 data from this study and that observed from other areas of East Africa [40], but is similar to data from other parts of Tanzania mainland [29] and South Africa [41]. A high frequency of CYP2B6\*6 homozygous mutant allele from this study indicates that a significant

**Table 5 Allele and genotype frequencies of CYP3A4\*1B and CYP3A5\*3 in Igombe, Tanzania, and expected frequencies according to Hardy Weinberg equilibrium**

Gene	SNP	Allele frequency	Genotype frequency			Gene allele
			wt/wt	wt/mut	mut/mut	
CYP3A4	A392G	G, 78% A, 22%	5.4% (5/92)	32.6% (30/92)	62% (57/92)	CYP3A4*1B wild type
Expected			4	31	56	
CYP3A5	A6986G	G, 16% A, 84%	70.1 (68/97)	27.9% (27/97)	2% (2/97)	CYP3A5*3 wild type
Expected			68	26	2	
CYP2C8	G416A	A, 10% G, 90%	80% (80/100)	20% (20/100)	0% (0/100)	CYP2C8*3 wild type
Expected			81	18	1	
CYP2C8	A805T	T, 19% A, 81%	66.1 (158/239)	29.3% (70/239)	4.6% (11/239)	CYP2C8*2 wild type
Expected			159	74	9	
CYP2B6	G15631T	T, 36% G, 64%	39% (37/95)	49.4% (47/95)	11.6% (11/95)	CYP2B6*6 wild type
Expected			39	44	12	

Note: wt/wt means homozygous wild type, wt/mut means heterozygous mutant and mut/mut means homozygous mutant.

proportion of the population are poor metabolizers of artemisinin and arteether [1] resulting in an increased risk to toxicity of the above anti-malarial drugs.

For CYP3A4\*1B, a very high proportion of the poor metabolizer allele (PM) in the population indicates high risk to artemisinin, artemether and lumefantrine toxicity due to a less efficient metabolism as discussed with CYP3A5\*3 and CYP2C8\*2 above. The high frequency recorded in this study demonstrates the previous documentations on a higher occurrence of this variant allele in Africans than Caucasians [59]. The high allele frequency observed from the study population is similar to other African populations, Ghana being one example [43]. Population variability for CYP3A4 is also exceptionally high, more than 100-fold [60]. The frequency of the CYP3A4\*1B mutation differs between populations. Among Caucasians and Hispanics the frequency is 3-5%, among Asians it seems to be absent [60] while the variant is more common among Africans as shown by findings from this study and previous studies (Tables 3, 4 and 5).

CYP2C8\*3 has been shown to affect metabolism to a higher extent than CYP2C8\*2. This study did not identify any homozygous patients who are PM of amodiaquine. This finding is not surprising since this variant allele has been shown to be rare/uncommon among indigenous Africans and predominantly concentrated in Caucasians [14]. CYP2C8\*3 frequencies recorded in this study are similar to Uganda [61] and other African countries [50].

Findings from this study demonstrate the existence of diverse genetics of the above CYP 450 s in Tanzania,

thus calling for an evaluation of the impact of CYP 450 polymorphisms on metabolism of artemisinin-based combinations and its impact on clinical outcome and toxicity among *P. falciparum* malaria-infected patients in African populations.

## Conclusion

Prevalence of CYP2C8\*2, CYP3A5\*3, CYP3A4\*1B and CYP2B6\*6 mutations are common while CYP2C8\*3 mutations are rare in a Tanzanian population. The frequencies recorded in this study are comparable to data obtained from previous studies on African populations. Pharmacogenomics data, such as that presented in this paper, provides a basis for further studies on impact of polymorphism in ACT safety and efficacy.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

KJM participated in proposal development, sample collection, carrying out molecular genetic analysis, data analysis and manuscript drafting. TS and MS carried out molecular genetic analysis and data analysis. OMS, EK and GS participated in proposal development, supervision of the research group and revising and approving the manuscript for publication. All authors read and approved the final manuscript.

## Acknowledgements

We sincerely thank the participants who donated blood and appreciate the contribution of the nurses, laboratory technicians and clinicians at Karume Health Centre, Igombe Mwanza, Tanzania. This work was funded by the Swedish Research Council, Department of Medical Biochemistry and Microbiology, Uppsala University and the Catholic University of Health and Allied Sciences.

#### Author details

<sup>1</sup>Department of Pharmacology, Catholic University of Health and Allied Sciences, Mwanza, Tanzania. <sup>2</sup>Institute of Medical Biochemistry and Microbiology, Uppsala University, Uppsala, Sweden. <sup>3</sup>Unit of Pharmacology and Therapeutics, Muhimbili University of Health and Allied Health Sciences, Dar Es Salaam, Tanzania. <sup>4</sup>Department of Biochemistry, Catholic University of Health and Allied Sciences, Mwanza, Tanzania.

Received: 23 August 2014 Accepted: 25 October 2014

Published: 3 November 2014

#### References

1. Kerb R, Fux R, Morike K, Krensner PG, Gil JP, Gleiter CH, Schwab M: **Pharmacogenetics of antimalarial drugs: effect on metabolism and transport.** *Lancet Infect Dis* 2009, **9**:760–774.
2. Obua C, Hellgren U, Ntale M, Gustafsson LL, Ogwal-Okeng JW, Gordi T, Jerling M: **Population pharmacokinetics of chloroquine and sulfadoxine and treatment response in children with malaria: suggestions for an improved dose regimen.** *Br J Clin Pharmacol* 2008, **65**:493–501.
3. Arav-Boger R, Shapiro TA: **Molecular mechanisms of resistance in antimalarial chemotherapy: the unmet challenge.** *Annu Rev Pharmacol Toxicol* 2005, **45**:565–585.
4. Shah RR: **Pharmacogenetics in drug regulation: promise, potential and pitfalls.** *Philos Trans R Soc Lond B Biol Sci* 2005, **360**:1617–1638.
5. Serijpa D, Pilotto A, Panza F, Matera MG: **Pharmacogenetics of cytochrome P450 (CYP) in the elderly.** *Ageing Res Rev* 2010, **9**:457–474.
6. Bains RK: **African variation at cytochrome P450 genes evolutionary aspects and the implications for the treatment of infectious diseases.** *Evol Med Public Health* 2013, **2013**:118–134.
7. Guengerich FP: **Cytochrome P450s and other enzymes in drug metabolism and toxicity.** *Aaps J* 2006, **8**:E101–E111.
8. Lynch T, Price A: **The effect of cytochrome P450 metabolism on drug response, interactions, and adverse effects.** *Am Fam Physician* 2007, **76**:391–396.
9. Ingelman-Sundberg M, Oscarson M, McLellan RA: **Polymorphic human cytochrome P450 enzymes: an opportunity for individualized drug treatment.** *Trends Pharmacol Sci* 1999, **20**:342–349.
10. Goldstein JA: **Clinical relevance of genetic polymorphisms in the human CYP2C subfamily.** *Br J Clin Pharmacol* 2001, **52**:349–355.
11. Miners JO, Birkett DJ: **Cytochrome P4502C9: an enzyme of major importance in human drug metabolism.** *Br J Clin Pharmacol* 1998, **45**:525–538.
12. Finta C, Zaphiropoulos PG: **The human CYP2C locus: a prototype for intergenic and exon repetition splicing events.** *Genomics* 2000, **63**:433–438.
13. Totah RA, Rettie AE: **Cytochrome P450 2C8: substrates, inhibitors, pharmacogenetics, and clinical relevance.** *Clin Pharmacol Ther* 2005, **77**:341–352.
14. Daily EB, Aquilante CL: **Cytochrome P450 2C8 pharmacogenetics: a review of clinical studies.** *Pharmacogenomics* 2009, **10**:1489–1510.
15. Schoch GA, Yano JK, Wester MR, Griffin KJ, Stout CD, Johnson EF: **Structure of human microsomal cytochrome P450 2C8: evidence for a peripheral fatty acid binding site.** *J Biol Chem* 2004, **279**:9497–9503.
16. Schoch GA, Yano JK, Sansen S, Dansette PM, Stout CD, Johnson EF: **Determinants of cytochrome P450 2C8 substrate binding: structures of complexes with montelukast, troglitazone, felodipine, and 9-cis-retinoic acid.** *J Biol Chem* 2008, **283**:17227–17237.
17. Naraharsetti SB, Lin YS, Rieder MJ, Marcianti KD, Psaty BM, Thummel KE, Totah RA: **Human liver expression of CYP2C8: gender, age, and genotype effects.** *Drug Metab Dispos* 2010, **38**:889–893.
18. Hoffman SM, Nelson DR, Keeney DS: **Organization, structure and evolution of the CYP2 gene cluster on human chromosome 19.** *Pharmacogenetics* 2001, **11**:687–698.
19. Zanger UM, Klein K, Saussele T, Blievnick J, Hofmann MH, Schwab M: **Polymorphic CYP2B6: molecular mechanisms and emerging clinical significance.** *Pharmacogenomics* 2007, **8**:743–759.
20. Wang H, Tompkins LM: **CYP2B6: new insights into a historically overlooked cytochrome P450 isozyme.** *Curr Drug Metab* 2008, **9**:598.
21. King J, Aberg JA: **Clinical impact of patient population differences and genetic variation in efavirenz therapy.** *Aids* 2008, **22**:1709–1717.
22. Klein K, Lang T, Saussele T, Barbosa-Sicard E, Schunck WH, Eichelbaum M, Schwab M, Zanger UM: **Genetic variability of CYP2B6 in populations of African and Asian origin: allele frequencies, novel functional variants, and possible implications for anti-HIV therapy with efavirenz.** *Pharmacogenet Genomics* 2005, **15**:861–873.
23. Shimada T, Yamazaki H, Mimura M, Inui Y, Guengerich FP: **Interindividual variations in human liver cytochrome P-450 enzymes involved in the oxidation of drugs, carcinogens and toxic chemicals: studies with liver microsomes of 30 Japanese and 30 Caucasians.** *J Pharmacol Exp Ther* 1994, **270**:414–423.
24. Thummel KE, Wilkinson GR: **In vitro and in vivo drug interactions involving human CYP3A.** *Annu Rev Pharmacol Toxicol* 1998, **38**:389–430.
25. Kuehl P, Zhang J, Lin Y, Lamba J, Assem M, Schuetz J, Watkins PB, Daly A, Wrighton SA, Hall SD, Maurel P, Relling M, Brimer C, Yasuda K, Venkataramanan R, Strom S, Thummel K, Boguski MS, Schuetz E: **Sequence diversity in CYP3A promoters and characterization of the genetic basis of polymorphic CYP3A5 expression.** *Nat Genet* 2001, **27**:383–391.
26. Lamba JK, Lin YS, Schuetz EG, Thummel KE: **Genetic contribution to variable human CYP3A-mediated metabolism.** *Adv Drug Deliv Rev* 2002, **54**:1271–1294.
27. Amirimani B, Ning B, Deitz AC, Weber BL, Kadlubar FF, Rebbeck TR: **Increased transcriptional activity of the CYP3A4\*1B promoter variant.** *Environ Mol Mutagen* 2003, **42**:299–305.
28. Hashimoto H, Toide K, Kitamura R, Fujita M, Tagawa S, Itoh S, Kamataki T: **Gene structure of CYP3A4, an adult-specific form of cytochrome P450 in human livers, and its transcriptional control.** *Eur J Biochem* 1993, **218**:585–595.
29. Staehli Hodel EM, Csajka C, Ariey F, Guidi M, Kabanyanyi AM, Duong S, Decosterd LA, Olliaro P, Beck HP, Genton B: **Effect of single nucleotide polymorphisms in cytochrome P450 isoenzyme and N-acetyltransferase 2 genes on the metabolism of artemisinin-based combination therapies in malaria patients from Cambodia and Tanzania.** *Antimicrob Agents Chemother* 2013, **57**:950–958.
30. Piedade R, Gil JP: **The pharmacogenetics of antimalaria artemisinin combination therapy.** *Expert Opin Drug Metab Toxicol* 2011, **7**:1185–1200.
31. Rodriguez-Antona C, Sayi JG, Gustafsson LL, Bertilsson L, Ingelman-Sundberg M: **Phenotype-genotype variability in the human CYP3A locus as assessed by the probe drug quinine and analyses of variant CYP3A4 alleles.** *Biochem Biophys Res Commun* 2005, **338**:299–305.
32. Temesvari M, Kobori L, Paulik J, Sarvary E, Belic A, Monostory K: **Estimation of drug-metabolizing capacity by cytochrome P450 genotyping and expression.** *J Pharmacol Exp Ther* 2012, **341**:294–305.
33. Zhou SF, Liu JP, Chowbay B: **Polymorphism of human cytochrome P450 enzymes and its clinical impact.** *Drug Metab Rev* 2009, **41**:89–295.
34. Alessandrini M, Asfaha S, Dodgen TM, Warnich L, Pepper MS: **Cytochrome P450 pharmacogenetics in African populations.** *Drug Metab Rev* 2013, **45**:253–275.
35. Hodel EM, Ley SD, Qi W, Ariey F, Genton B, Beck HP: **A microarray-based system for the simultaneous analysis of single nucleotide polymorphisms in human genes involved in the metabolism of anti-malarial drugs.** *Malar J* 2009, **8**:285.
36. Paganotti GM, Gramolelli S, Tabacchi F, Russo G, Modiano D, Coluzzi M, Romano R: **Distribution of human CYP2C8\*2 allele in three different African populations.** *Malar J* 2012, **11**:125.
37. Rodriguez S, Gaunt TR, Day IN: **Hardy-Weinberg equilibrium testing of biological ascertainment for Mendelian randomization studies.** *Am J Epidemiol* 2009, **169**:505–514.
38. Zeigler-Johnson C, Walker A, Mancke B, Spangler E, Jalloh M, McBride S, Deitz A, Malkowicz S, Ofori-Adjei D, Gueye S: **Ethnic differences in the frequency of prostate cancer susceptibility alleles at SRD5A2 and CYP3A4.** *Hum Hered* 2002, **54**:13–21.
39. Swart M, Skelton M, Wonkam A, Kannemeyer L, Dandara C: **CYP1A2, CYP2A6, CYP2B6, CYP3A4 and CYP3A5 polymorphisms in two Bantu-speaking populations from Cameroon and South Africa: implications for global pharmacogenetics.** *Curr Pharmacogenomics Personalized Med* 2012, **10**:43–53.
40. Bains RK, Kovacevic M, Plaster CA, Tarekegn A, Bekele E, Bradman NN, Thomas MG: **Molecular diversity and population structure at the cytochrome P450 3A5 gene in Africa.** *BMC Genet* 2013, **14**:1–18.
41. Dandara C, Lombard Z, Du Plooy I, McLellan T, Norris SA, Ramsay M: **Genetic variants in CYP (-1A2, -2C9, -2C19, -3A4 and -3A5), VKORC1 and**

- ABCB1 genes in a black South African population: a window into diversity. *Pharmacogenomics* 2011, **12**:1663–1670.
42. Drogemoller B, Plummer M, Korkie L, Agenbag G, Dunaiski A, Niehaus D, Koen L, Gebhardt S, Schneider N, Olckers A, Wright G, Warnich L: **Characterization of the genetic variation present in CYP3A4 in three South African populations.** *Front Genet* 2013, **4**:17.
43. Kudzi W, Dodoo AN, Mills JJ: **Genetic polymorphisms in MDR1, CYP3A4 and CYP3A5 genes in a Ghanaian population: a plausible explanation for altered metabolism of ivermectin in humans?** *BMC Med Genet* 2010, **11**:111.
44. Oliveira E, Pereira R, Amorim A, McLeod H, Prata MJ: **Patterns of pharmacogenetic diversity in African populations: role of ancient and recent history.** *Pharmacogenomics* 2009, **10**:1413–1422.
45. Ferreira PE, Veiga MI, Cavaco I, Martins JP, Andersson B, Mushin S, Ali AS, Bhattarai A, Ribeiro V, Bjorkman A, Gil JP: **Polymorphism of antimalaria drug metabolizing, nuclear receptor, and drug transport genes among malaria patients in Zanzibar, East Africa.** *Ther Drug Monit* 2008, **30**:10–15.
46. Cavaco I, Reis R, Gil JP, Ribeiro V: **CYP3A4\*1B and NAT2\*14 alleles in a native African population.** *Clin Chem Lab Med* 2003, **41**:606–609.
47. Roy J-N, Lajoie J, Zijenah LS, Barama A, Poirier C, Ward BJ, Roger M: **CYP3A5 genetic polymorphisms in different ethnic populations.** *Drug Metab Dispos* 2005, **33**:884–887.
48. Calcagno A, D'Avolio A, Simiele M, Cusato J, Rostagno R, Libanore V, Baietto L, Siccardi M, Bonora S, Di Perri G: **Influence of CYP2B6 and ABCB1 SNPs on nevirapine plasma concentrations in Burundese HIV-positive patients using dried sample spot devices.** *Br J Clin Pharmacol* 2012, **74**:134–140.
49. Parikh S, Ouedraogo JB, Goldstein JA, Rosenthal PJ, Kroetz DL: **Amodiaquine metabolism is impaired by common polymorphisms in CYP2C8: implications for malaria treatment in Africa.** *Clin Pharmacol Ther* 2007, **82**:197–203.
50. Kudzi W, Dodoo AN, Mills JJ: **Characterisation of CYP2C8, CYP2C9 and CYP2C19 polymorphisms in a Ghanaian population.** *BMC Med Genet* 2009, **10**:124.
51. Adjei GO, Kristensen K, Goka BQ, Hoegberg LC, Alifrangis M, Rodrigues OP, Kurtzhals JA: **Effect of concomitant artesunate administration and cytochrome P4502C8 polymorphisms on the pharmacokinetics of amodiaquine in Ghanaian children with uncomplicated malaria.** *Antimicrob Agents Chemother* 2008, **52**:4400–4406.
52. Rower S, Bienzle U, Weise A, Lambertz U, Forst T, Otchwemah RN, Pfutzner A, Mockenhaupt FP: **High prevalence of the cytochrome P450 2C8\*2 mutation in Northern Ghana.** *Trop Med Int Health* 2005, **10**:1271–1273.
53. Nyakutira C, Roshammar D, Chigutsa E, Chonzi P, Ashton M, Nhachi C, Masimirembwa C: **High prevalence of the CYP2B6 516G→T(\*6) variant and effect on the population pharmacokinetics of efavirenz in HIV/AIDS outpatients in Zimbabwe.** *Eur J Clin Pharmacol* 2008, **64**:357–365.
54. Arnaldo P, Thompson RE, Lopes MQ, Suffys PN, Santos AR: **Frequencies of cytochrome P450 2B6 and 2C8 allelic variants in the Mozambican population.** *Malays J Med Sci* 2013, **20**:13–23.
55. Cavaco I, Stromberg-Norklit J, Kaneko A, Msellem MI, Dahoma M, Ribeiro VL, Bjorkman A, Gil JP: **CYP2C8 polymorphism frequencies among malaria patients in Zanzibar.** *Eur J Clin Pharmacol* 2005, **61**:15–18.
56. Ngaimisi E, Habtewold A, Minzi O, Makonnen E, Mugusi S, Amogne W, Yimer G, Riedel KD, Janabi M, Aderaye G, Mugusi F, Bertilsson L, Akillu E, Burhenne J: **Importance of ethnicity, CYP2B6 and ABCB1 genotype for efavirenz pharmacokinetics and treatment outcomes: a parallel-group prospective cohort study in two sub-Saharan Africa populations.** *PLoS One* 2013, **8**:e67946.
57. Parathyras J, Gebhardt S, Hillermann-Rebello R, Grobbelaar N, Venter M, Warnich L: **A pharmacogenetic study of CD4 recovery in response to HIV antiretroviral therapy in two South African population groups.** *J Hum Genet* 2009, **54**:261–265.
58. Maiteki-Sebuguzi C, Jagannathan P, Yau VM, Clark TD, Njama-Meya D, Nzarubara B, Talisuna AO, Kanya MR, Rosenthal PJ, Dorsey G, Staedke SG: **Safety and tolerability of combination antimalarial therapies for uncomplicated falciparum malaria in Ugandan children.** *Malar J* 2008, **7**:106.
59. Tayeb MT, Clark C, Ameyaw MM, Haites NE, Evans DA, Tariq M, Mobarek A, Ofori-Adjei D, McLeod HL: **CYP3A4 promoter variant in Saudi, Ghanaian and Scottish Caucasian populations.** *Pharmacogenetics* 2000, **10**:753–756.
60. Zanger UM, Schwab M: **Cytochrome P450 enzymes in drug metabolism: regulation of gene expression, enzyme activities, and impact of genetic variation.** *Pharmacol Ther* 2013, **138**:103–141.
61. Paganotti GM, Cosentino V, Russo G, Tabacchi F, Gramolelli S, Coluzzi M, Romano R: **Absence of the human CYP2C8\*3 allele in Ugandan children exposed to Plasmodium falciparum malaria.** *Infect Genet Evol* 2014, **27**:432–435.

doi:10.1186/1475-2875-13-420

**Cite this article as:** Marwa et al.: Cytochrome P450 single nucleotide polymorphisms in an indigenous Tanzanian population: a concern about the metabolism of artemisinin-based combinations. *Malaria Journal* 2014 **13**:420.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

