

Research

Open Access

Basal shuttle of NF- κ B/I κ B α in resting T lymphocytes regulates HIV-1 LTR dependent expression

Mayte Coiras^{†1}, María Rosa López-Huertas^{†1}, Joaquín Rullas¹,
Maria Mittelbrunn² and José Alcamí^{*1}

Address: ¹AIDS Immunopathology Unit, National Center of Microbiology, Instituto de Salud Carlos III, Majadahonda, Madrid, Spain and ²Immunology Service, Hospital de La Princesa, Universidad Autonoma de Madrid, Madrid, Spain

Email: Mayte Coiras - mcoiras@isciii.es; María Rosa López-Huertas - mrlhuertas@isciii.es; Joaquín Rullas - joaquin.m.rullas@gsk.com; Maria Mittelbrunn - mmittelbrun.hlpr@salud.madrid.org; José Alcamí* - ppalcami@isciii.es

* Corresponding author †Equal contributors

Published: 8 August 2007

Received: 17 May 2007

Retrovirology 2007, 4:56 doi:10.1186/1742-4690-4-56

Accepted: 8 August 2007

This article is available from: <http://www.retrovirology.com/content/4/1/56>

© 2007 Coiras et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: In HIV-infected T lymphocytes, NF- κ B/Rel transcription factors are major elements involved in the activation of LTR-dependent transcription from latency. Most NF- κ B heterodimer p65/p50 is sequestered as an inactive form in the cytoplasm of resting T lymphocytes via its interaction with I κ B inhibitors. In these cells, both absolute HIV latency and low level ongoing HIV replication have been described. These situations could be related to differences in the balance between NF- κ B and I κ B α ratio. Actually, control of I κ B α by cellular factors such as Murr-1 plays a critical role in maintaining HIV latency in unstimulated T lymphocytes. Formerly, our group demonstrated the presence of nuclear I κ B α in T cells after PMA activation. Now we attempt to determine the dynamics of NF- κ B/I κ B α nucleocytoplasmic transport in absence of activation as a mechanism to explain both the maintenance of latency and the existence of low level ongoing HIV replication in resting CD₄⁺ T lymphocytes.

Results and conclusion: We show that the inhibition of the nuclear export by leptomycin B in resting CD₄⁺ T cells resulted in nuclear accumulation of both I κ B α and p65/RelA, as well as formation of NF- κ B/I κ B α complexes. This proves the existence of a rapid shuttling of I κ B α between nucleus and cytosol even in absence of cellular activation. The nuclear accumulation of I κ B α in resting CD₄⁺ T lymphocytes results in inhibition of HIV-LTR dependent transcription as well as restrains HIV replication in CD₄⁺ T lymphocytes. On the other hand, basal NF- κ B activity detected in resting CD₄⁺ T lymphocytes was related to low level HIV replication in these cells.

Background

The nuclear factor κ B (NF- κ B) family of proteins are inducible transcription factors that play a central role in regulating the expression of a wide variety of genes associated with cell proliferation, immune response, inflammation, cell survival, and oncogenesis [1,2]. Functionally competent NF- κ B is mainly composed by heterodimers of

p65/RelA or c-Rel proteins complexed to p50/NF- κ B1. NF- κ B activity is regulated partially at subcellular level because active NF- κ B heterodimers are normally sequestered in the cytoplasm via its non-covalent interaction with a family of inhibitory proteins termed I κ Bs, being I κ B α the major NF- κ B inhibitor protein. NF- κ B activation is initiated by a variety of stimuli such as cytokines and

growth factors, which lead to activation of I κ B kinase complex (IKK). IKK in turn phosphorylates I κ B α , resulting in its degradation via the ubiquitin-mediated proteolytic pathway. This permits NF- κ B translocation into the nucleus, where engages cognate κ B enhancer elements and modulates gene expression [1,2].

Control over NF- κ B activity is not only accomplished through association with I κ B α in the cytosol, but a role for nuclear I κ B α in the control of NF- κ B-driven transcription has been proposed [3,4]. In this model, newly synthesized I κ B α would be able to shuttle actively between the cytoplasm and the nucleus, and then remove NF- κ B from the κ B consensus sequences. Thus, nuclear I κ B α would promote the return of NF- κ B to the cytoplasm and the termination of its transcriptional response. The shuttle of NF- κ B and I κ B α between nucleus and cytosol in tumor cell lines has been described previously [3-5] as well as its influence on κ B dependent gene expression. However, in normal human CD $_4^+$ T lymphocytes in a resting state, NF- κ B binding activity is low and consists predominantly of inactive p50/p50 homodimers. In these cells, functional p50/p65 complexes are induced by cell activation [6]. Our group described previously that I κ B α can translocate to the nucleus in T lymphocytes activated with phorbol-12-myristate-13-acetate (PMA) [7], but little is known about the existence of a NF- κ B/I κ B α shuttling in resting blood T cells.

The NF- κ B pathway provides an attractive target to viral pathogens. Activation of NF- κ B is a rapid, immediate early event that occurs within minutes after exposure to a stimulus, does not require *de novo* protein synthesis, and produces a strong transcriptional activation of several viral genes [6]. As a result, NF- κ B is essential in the regulation of the HIV-1 long terminal repeat (LTR) promoter [8-10]. The promoter-proximal (enhancer) region of the HIV LTR contains two adjacent NF- κ B binding sites (-109 to -79) that play a central role in mediating inducible HIV gene expression. These NF- κ B responsive elements are major elements in triggering HIV LTR-transcription in blood CD $_4^+$ T cells [6,9-11]. Accordingly, HIV production in T cells is mainly associated with the activation induced by different stimuli, whereas resting or unstimulated CD $_4^+$ T lymphocytes offer a cellular environment for latency due to low permissiveness to HIV LTR activity [6]. However, the existence of a low-level ongoing replication in resting CD $_4^+$ T lymphocytes has been described [12-14].

To reconcile these contradictory data, the hypothesis that the existence of a basal NF- κ B activity could contribute to the low viral replication detected in HIV-infected CD $_4^+$ T lymphocytes in a resting state is proposed. To this aim, the molecular mechanisms involved in the NF- κ B/I κ B α traffic between cytoplasm and nucleus of resting T lymphocytes

from human blood have been analyzed. When resting CD $_4^+$ T lymphocytes were cultured in presence of leptomycin B (LMB), a nuclear export inhibitor [15], both p65/RelA and I κ B α were accumulated and associated in the nucleus, suggesting a rapid shuttling of both proteins in unstimulated T cells. In fact, HIV LTR-driven transactivation and HIV replication can be blocked in resting as well as activated T cells by I κ B α over-expression. Our findings suggest that the balance between NF- κ B and I κ B α at nuclear level would be a key mechanism involved in both the maintenance of HIV latency and the induction of low-level HIV replication in resting CD $_4^+$ T lymphocytes.

Results

Analysis of I κ B α and p65/RelA subcellular distribution in resting CD $_4^+$ T lymphocytes

Resting non-activated CD $_4^+$ T lymphocytes were negatively isolated from human PBMCs by depletion of B cells, NK cells, monocytes, CD $_8^+$ T cells and activated lymphocytes. Analysis by flow cytometry revealed they were CD $_4^+$ CD $_25^-$ CD $_69^-$ HLA-DR $^-$ with a purity >95%.

I κ B α and p65/RelA shuttling between nucleus and cytosol was analyzed in resting blood CD $_4^+$ T cells by using LMB, a specific inhibitor of the nuclear protein export. The subcellular distribution of I κ B α and p65/RelA was first analyzed by immunofluorescence assays. Both I κ B α and p65/RelA were localized in the cytosol of unstimulated CD $_4^+$ T cells (Fig. 1), but after treatment with LMB, both I κ B α and p65/RelA were retained in the nucleus. This nuclear translocation was observed in the absence of any stimulus and was not due to serum activation since similar results were observed in serum deprivation conditions (data not shown).

These results were confirmed using chimeric proteins formed by the enhanced yellow fluorescent protein (EYFP) fused to I κ B α or p65/RelA. Resting CD $_4^+$ T cells were transiently transfected with plasmids pEYFP-p65 and pEYFP-I κ B α separately. Analysis was performed 24 hours after transfection by confocal microscopy. There was low quantity of both I κ B α and p65/RelA in the nucleus of the resting T cells before LMB treatment (Fig. 2a) but after exposure to LMB, both EYFP-I κ B α and EYFP-p65 fusion proteins were retained in the nucleus. Plasmid pEYFP-C1 containing the EYFP under the control of CMV promoter was used as control of non-specific intracellular distribution.

To exclude that nucleoporation could induce NF- κ B activity, electrophoretic mobility shift assays (EMSA) were performed in nuclear extracts from CD $_4^+$ T lymphocytes transfected with a control plasmid (pcDNA3.1) by two different methods: the Amaxa Nucleofector system and




Figure 1

Subcellular localization of IκBα and p65/RelA in CD₄⁺ T lymphocytes. Cells were treated or not with 20 nM LMB and then fixed, permeabilized and stained with specific antibodies against IκBα and p65/RelA. A secondary antibody conjugated with Texas Red (Molecular Probes) was used. Images were taken by confocal microscopy.

classical electroporation using an Equibio electroporator (Figure 2b).

In order to determine the dynamics of IκBα shuttling, resting CD₄⁺ T cells were transiently transfected with EYFP-IκBα vector, attached to fibronectine-coated slides and filmed *in vivo* by time-lapse confocal microscopy during treatment with LMB. Photographs were taken each minute after adding LMB and it was determined that less than 6 minutes were enough to saturate the nucleus with IκBα (Fig. 3 and additional file 1).

LMB toxicity was assessed by propidium iodide staining and flow cytometry in resting CD₄⁺ T cells treated up to 24 hours. Mortality due to LMB treatment (20 nM) was increased only 10% above controls after the longest incubation time (data not shown).

Analysis of nuclear protein-protein interactions

Because more than 10⁸ blood T lymphocytes for each experimental point were required to perform these experiments, T cells were expanded according to a protocol previously developed in our laboratory. PBMCs were cultured for 3 days with 5 μg/ml PHA and for the consecutive 9 days with 300 U/ml IL-2. These long-term cultures of PHA-treated T lymphocytes were maintained without supplemental IL-2 18 hours before the experiment to assure they were in a resting state concerning NF-κB activity. Following this protocol, it was proved that basal and induced NF-κB was similar as in resting T lymphocytes [7] (see Additional file 2).

Consequently, association between IκBα and p65/RelA was determined in the nucleus of long-term cultures of PHA-treated T lymphocytes. For this purpose, nuclear and




Figure 2

Subcellular localization of EYFP-IκBα and EYFP-p65 fusion proteins in CD₄⁺ T lymphocytes. (a) Cells were transiently transfected with 1 μg of either EYFP-IκBα or EYFP-p65 expression vectors per million of cells. LMB was added immediately after transfection. After 18–24 hours of incubation, cells were analyzed by confocal microscopy. pEYFP-C1 vector was used as control of unspecific distribution. (b) Resting purified CD₄⁺ T lymphocytes were transiently transfected with the control plasmid pcDNA3.1 by using an Amaxa nucleofector and a classical electroporator (Equibio). As occurs in untransfected resting T cells (lane 1), NF-κB was not induced in resting CD₄⁺ T lymphocytes after electroporation (lanes 3 and 4). As a positive control, NF-κB (p50/p65) binding was induced in these cells by PMA activation (lane 2).

cytosolic protein extracts were analyzed by immunoblotting assays. As previously shown for resting CD₄⁺ T lymphocytes (Fig. 1), both nuclear IκBα and p65/RelA levels increased in cells treated with LMB (Fig. 4a, Nucleus, lane 2). The accumulation of cytosolic proteins in the nucleus after LMB treatment has been ruled out by immunoblotting of cytosolic and nuclear extracts from PHA-treated T cells by using an antibody against both p105 and p50/NF-κB1 proteins (Fig. 4b). The p105 protein is the precursor of the p50 subunit and it presents exclusively a cytosolic location.

Immunoprecipitation assays with an antibody against p65/RelA showed the presence of NF-κB/IκBα complexes in the nucleus of T cells treated or not with LMB (Fig. 4a,

Immunoprecipitation, Nucleus, lanes 1 and 2), whereas no association between p65/RelA and IκBα was observed in cells activated with PMA (Fig. 4a, Immunoprecipitation, Nucleus, lane 3).

NF-κB DNA-binding activity in unstimulated T lymphocytes

Once it was confirmed that both p65/RelA and IκBα were able to shuttle between nucleus and cytosol in unstimulated T cells, NF-κB DNA binding activity was analyzed by EMSA. Despite the presence of p65/RelA in the nucleus, no binding was detected in unstimulated T cells treated or not with LMB (Fig. 4c, lanes 1 and 2). This correlated with the detection of NF-κB/IκBα complexes in the nucleus of these cells (Fig. 4a, Immunoprecipitation, Nucleus, lanes




Figure 3
Kinetic analysis of nuclear IκBα translocation. One CD₄⁺ T lymphocyte transfected with EYFP-IκBα vector was photographed before and after treatment with LMB up to 30 minutes. Photographs were taken *in vivo* by confocal microscopy every minute after adding LMB.

1 and 2). As expected, NF-κB kept the binding activity to -κB motif in PMA-activated T cells (Fig. 4c, lane 3), due to the absence of NF-κB/IκBα complexes in the nucleus of these cells (Fig. 4a, Immunoprecipitation, Nucleus, lane 3).

Analysis of IκBα resynthesis in resting T cells

Unstimulated T cells were incubated with CHX for 30 minutes before adding other stimulus in order to stop *de novo* protein synthesis. Then, LMB or PMA were added to the culture medium. Immunoblotting assays showed a decrease of IκBα levels in the nucleus of T cells incubated with both CHX and LMB (Fig. 4d, lane 2) or with CHX, LMB and PMA (Fig. 4d, lane 3), but not in those cells only incubated with LMB (Fig. 4d, lane 1). These data not only confirm previous results showing that nuclear translocation of IκBα is dependent on protein resynthesis [3] but also asserts that this *de novo* protein synthesis is carried out even in unstimulated T cells.

Basal NF-κB activity can activate HIV-LTR promoter in CD₄⁺ T lymphocytes

Resting blood CD₄⁺ T cells were transfected with a LTR-LUC vector alone or together with a Tat expression vector under the control of the CMV promoter in order to assess NF-κB-dependent transcriptional activity in these cells by measurement of luciferase activity (Fig. 5a and 5b). As expected, both Tat over-expression and PMA activation enhanced LTR-dependent transcription, as previously described [11,16]. However, when nuclear levels of IκBα were increased by LMB (Fig. 5a) or transient transfection of CMV-IκBα vector (Fig. 5b), a dramatic decrease in luciferase activity was observed, both in PMA-activated T cells and cells in which Tat was over-expressed. Interestingly, a basal NF-κB activity able to induce a low LTR transactiva-

tion was detected in unstimulated CD₄⁺ T cells. This low LTR transactivation was annulled when IκBα was over-expressed by both LMB or CMV-IκBα transfection, thus proving this basal LTR transactivation was due to a residual NF-κB activity in resting CD₄⁺ T cells.

Progression of HIV replication in resting CD₄⁺ T lymphocytes

To assess the role of basal NF-κB activity and IκBα over-expression on a model of HIV production in resting and activated CD₄⁺ T cells, highly purified CD₄⁺ CD₂₅⁻ CD₆₉⁻ DR⁻ T lymphocytes obtained from blood of different healthy donors were transfected with a full-length infectious HIV clone (NL4.3) together with a CMV-IκBα expression vector or pcDNA3 as negative control. Cells were maintained in culture up to 7 days either in the absence of activation or activated with two different stimuli, PHA and CD3 antibodies. HIV p24-gag was quantified 5 and 7 days after transfection. An intense HIV replication was detected in activated CD₄⁺ T cells after 7 days in culture (Fig. 6b). Besides, a discrete but significant HIV p24-gag production was assessed in resting CD₄⁺ T cells after 5 days of transfection (Fig. 6a). When IκBα was over-expressed in these cells, p24-gag production decreased as compared to cells transfected with a control plasmid and this difference was significant ($p < 0.05$) for resting and anti-CD₃ activated T cells. Although more than five-fold decrease was observed at day 7 for PHA-activated lymphocytes when IκBα was over-expressed, this result did not reach statistical significance ($p = 0.081$).

Discussion

Initiation of HIV transcription from a quiescent state is regulated through the concerted action of different cellular factors acting at LTR sequences [17,18]. Among them, NF-κB proteins are the most important inducible elements involved in initiation of HIV transcription in normal T cells [6,11,19-21]. As a result, a strong control of nuclear NF-κB translocation would be required to maintain HIV latency.

Nuclear translocation and activity of NF-κB is regulated through different mechanisms including association with its main inhibitor IκBα as a cytosolic inactive form. An additional mechanism of NF-κB control is the nuclear location of IκBα that act as a terminator of -κB dependent transactivation [4,5]. In fact, a dynamic shuttling of NF-κB has been described in established cell lines by balancing fluxes into and out of the nucleus [22-24] as well as the capacity of IκBα to enter the nucleus of T cells activated with PMA [7]. However, the nucleocytoplasmic shuttling of both NF-κB and IκBα in T cells in a resting state and its potential role in the maintenance of latency or the initiation of HIV transcription has not been determined so far. This is a very important issue, because resting CD₄⁺ T cells




Figure 4

Analysis of nuclear NF-κB/IκBα complexes in CD₄⁺ T cells and IκBα pool dependence on *de novo* protein synthesis. (a) Analysis of subcellular distribution of p65/RelA and IκBα in CD₄⁺ T cells and presence of NF-κB/IκBα complexes in the nucleus after treatment with LMB or PMA. Ten micrograms of cytosolic and nuclear extracts from CD₄⁺ T cells treated with either PMA or LMB during 4 and 6 hours respectively were analyzed by Western Blot using antibodies against p65/RelA and IκBα. Immunoprecipitation assays were performed using 100 μg of these cytosolic and nuclear extracts, which were incubated with 5 μg of an antibody against p65/RelA conjugated with agarose. IκBα and p65/RelA complexes were characterized by immunoblotting. (b) Contamination with cytosolic proteins during nuclear protein extraction or accumulation of cytosolic proteins in the nucleus after treatment with LMB was assessed by Western Blot using an antibody against both p105 and p50/NF-κB1 proteins. (c) Analysis of NF-κB DNA-binding activity in CD₄⁺ T cells treated with either PMA or LMB. Three micrograms of nuclear extract were incubated with an oligonucleotide containing the double consensus motif κB present in the HIV LTR labeled with [α-³²P]-dCTP. Protein extracts were obtained from CD₄⁺ T cells after treatment with either LMB or PMA for 6 and 4 hours respectively. (d) Analysis of IκBα pool dependence on *de novo* protein synthesis. Ten micrograms of nuclear extracts from CD₄⁺ T cells incubated with 20 nM LMB for 4 hours and 10 μg/ml CHX and/or 25 ng/ml PMA for 4 hours, 30 min and 2 hours, respectively, were analyzed by Western Blot.

containing integrated HIV provirus constitute one of the long-lived cellular reservoirs of HIV *in vivo* [25,26] and represent a main obstacle to the eradication of the virus [27,28]. This HIV reservoir had been thought to be quiescent with regard to virus replication based on the principle that HIV production in T cells is linked to cellular activation. However, HIV production may occur in T cells that

have not undergone classic T cell activation [29] and even in CD₄⁺ T lymphocytes lacking any activation markers [13].

These observations raise the question of whether NF-κB would be able to initiate the transcription of its target genes in resting T cells. In normal human CD₄⁺ T cells, NF-




Figure 5

Influence of IκBα over-expression on HIV-LTR transactivation. Resting CD₄⁺ T cells were transfected with LTR-LUC vector together with (a) pcDNA3.1 and/or CMV-Tat expression vectors or (b) pcDNA3.1 and/or CMV-Tat and/or CMV-IκBα expression vectors, as indicated. Cells were treated with LMB immediately after transfection and/or with PMA two hours after transfection, as indicated. Luciferase activity was measured 18 hours after transfection. Numbers on the top of the bars represent fold transcriptional activity relative to unstimulated T cells transfected with pcDNA3.1.

κB binding activity is low and consists predominantly of p50/p50 complexes, but not p50/p65. T-cell activation results in the formation of p50/p65 complexes and the induction of HIV-LTR transactivation. According to this hypothesis, both p65/RelA and IκBα showed a predominant cytosolic distribution in resting CD₄⁺ T cells (Fig. 1). However, a sharp increase in both nuclear IκBα and p65/RelA was found when nuclear export was inhibited by LMB, even in the absence of activation (Fig. 1 and 2). Moreover, *in vivo* kinetic studies determined that IκBα completely filled the nucleus of resting CD₄⁺ T cells in less than 6 minutes after adding LMB to the culture medium (Fig. 3 and additional file 1). NF-κB was associated to IκBα in the nucleus of resting T cells (Fig. 4a) and only p50/p50 heterodimers were able to bind DNA (Fig. 4c). In contrast, in PMA-activated T cells no association between

IκBα and p65/RelA was found despite the presence of both proteins in the nuclear compartment (Fig. 4a), and consequently p50/p65 heterodimers could bind DNA (Fig. 4c). These results suggest the existence of post-translational modifications in p65/RelA and/or IκBα in PMA-activated T lymphocytes that would decrease the affinity between both proteins allowing DNA binding of active NF-κB. On the other hand, it has been described that only newly synthesized IκBα can enter the nucleus [4]. Accordingly, a sharp decrease in nuclear IκBα levels was observed in resting T cells when *de novo* protein synthesis was inhibited, whereas p65/RelA exhibited a longer half-life due to the existence of a pre-synthesized pool or a less active degradation (Fig. 4d). Therefore, a rapid degradation of IκBα occurs in T cells in the absence of activation and continuous synthesis is required to maintain a cytosolic pool of

dissociated I κ B α , able to translocate to the nucleus and capture NF- κ B. These data proved not only the existence of a nucleocytoplasmic shuttling of I κ B α and NF- κ B in resting T lymphocytes but also that it is an extremely dynamic process detected exclusively when nuclear export is inhibited.

It has been described that HIV replication may occur within CD₄⁺T cells activated below the threshold required for proliferation [12,13]. Indeed, it has been proposed that basal nuclear NF- κ B translocation is required for the activation of genes involved in cell survival and these small discharges of nuclear NF- κ B could be the cause of the low level replication observed in resting HIV-infected T cells. In support of this hypothesis, when CD₄⁺T cells were transfected with luciferase expression vectors under the control of the HIV-LTR, low but consistent transcriptional activity, which was enhanced by Tat expression, was detected (Fig. 5a). In order to confirm that NF- κ B was responsible for this low level LTR activity in resting T cells, nuclear levels of I κ B α were increased by LMB (Fig. 5a) or transient transfection of CMV-I κ B α vector (Fig. 5b). In both cases, LTR transcriptional activation decreased, even when Tat was also over-expressed. Moreover, despite the observation that in PMA-activated lymphocytes NF- κ B was not bound to I κ B α in the nucleus (Fig. 4a), I κ B α over-expression resulted in strong decrease in HIV-LTR transactivation. It has been previously shown [3,4] that I κ B α can bind p65/RelA and transport it back to the cytosol. When this pathway is blocked by LMB, I κ B α cumulates in the nucleus at higher concentrations than during normal trafficking. We hypothesize that in these conditions NF- κ B activity could be inhibited by high I κ B α concentrations (Fig. 5). This observation supports that mechanisms involved in post-translational modifications of p65/RelA and/or I κ B α induced by PMA, which block the formation of NF- κ B/I κ B α complexes, can be overcome by I κ B α over-expression. Besides, low LTR transactivation detected in resting CD₄⁺T cells was also annulled by I κ B α over-expression, proving this basal LTR transactivation was due to a residual NF- κ B activity in these cells.

To confirm the role of I κ B α in an infectious model, a full-length proviral clone (NL4.3) was transfected in non-stimulated CD₄⁺T cells together with a CMV-I κ B α expression vector or pcDNA3.1 as negative control. This transfection method was used because the main goal was to analyze the role of I κ B α over-expression on HIV replication in both resting and activated lymphocytes and classical infection models require previous T cell activation. In this system, low transfection rates of T lymphocytes are usually achieved but they were enough to induce full HIV replication after stimulation with PHA or anti-CD₃. One open question in this model is whether p24-gag production derives from plasmid driven transient virus produc-

tion and not yet full viral replication. Because T cell activation induces both HIV integration and further proviral transcription, full viral replication was achieved in PHA and anti-CD₃-activated T lymphocytes. Moreover, increasing concentrations of p24-gag were detected throughout culture time, thereby suggesting several cycles of infection (Fig. 6). In this experimental system, inhibition of HIV replication by I κ B α over-expression is probably produced during the first cycle of replication, because in subsequent replication cycles I κ B α will not be over-expressed in non-transfected lymphocytes. Actually, a delay in HIV spread in culture due to partial inhibition of the first replication cycle in CMV-I κ B α -transfected cells was observed (Fig. 6). Moreover, decrease in p24-gag production in CMV-I κ B α -transfected cells was significant ($p < 0.05$) for resting and anti-CD₃ activated T cells. Although for PHA-activated lymphocytes this difference was not significant, a five-fold decrease was observed at day 7 and a trend towards statistical significance was found ($p = 0.081$). On the other hand, it is difficult to precise if the mechanism involved in p24-gag production in non-activated T lymphocytes is due to plasmid-driven transient virus production and not yet to viral replication. However, our results showed a decrease in LTR transactivation (Fig. 5) and p24-gag production (Fig. 6) in resting CD₄⁺T lymphocytes when I κ B α is over-expressed. It suggests that increasing I κ B α levels in naturally HIV-infected CD₄⁺T lymphocytes carrying an integrated provirus could contribute to NF- κ B inhibition and subsequent low-level viral production or absolute latency, as described in resting CD₄⁺T lymphocytes *in vivo* [12-14,30,31].

On the other hand, it has been described that HIV can integrate into the genomes of *in vitro*-inoculated resting CD₄⁺T cells that have not received activating stimuli [32]. Accordingly, HIV replication can also start in these cells although it cannot further progress unless these CD₄⁺T cells were subsequently activated and NF- κ B activity were maintained.

Overall, these data suggest that LTR transcriptional activation can be initiated by basal NF- κ B activity in resting CD₄⁺T cells in the absence of previous stimuli. Alternatively, the presence of high levels of nuclear I κ B α would result in NF- κ B control and viral latency. These data are supported by the existence of transdominant mutants of I κ B α that block NF- κ B induction and inhibit *de novo* HIV infection in T cells by interfering with viral transcription [20,33]. Besides, control of I κ B α by other cellular factors such as Murr1, have been also involved in the maintenance of HIV latency in resting CD₄⁺T lymphocytes [34].

Conclusion

The maintenance of HIV latency should be considered an active cellular process. In resting CD₄⁺T cells, both I κ B α




Figure 6

HIV replication in resting or activated CD₄⁺ T cells transfected with an infectious molecular HIV-1 clone. Highly purified CD₄⁺ CD₂₅⁻ CD₆₉⁻ DR⁻ T cells were transfected with the NL4.3 infectious molecular HIV-1 clone together with CMV-IkBa or pcDNA3.1 as negative control, and then activated with anti-CD₃ and IL-2, PHA and IL-2, or maintained in the absence of activation. Viral replication was determined by quantification of HIV p24-gag antigen in culture supernatants (a) after 5 days of transfection or (b) after 7 days of transfection. Numbers on the top of the bars represent fold HIV-replication relative to unstimulated T cells transfected with pcDNA3.1. Differences in p24-gag production were significant for resting and anti-CD₃-activated T cells ($p < 0.05$) and a trend towards statistical significance was found in PHA-activated T cells ($p = 0.081$).

and NF- κ B are continuously shuttling between cytosol and nucleus, as well as continuously associating and dissociating to permit a low transcriptional activity necessary for the activation of genes involved in cell survival. In resting HIV-infected T cells, the balance between free NF- κ B and NF- κ B/I κ B α complexes in the nucleus could directly participate in the maintenance of HIV-latency when I κ B α predominates as well as in the low ongoing HIV replication when NF- κ B escapes I κ B α control. Both phenomena have been characterized *in vivo* and constitute major pathogenic mechanisms in the persistence of long-lived cellular reservoirs of HIV [12-14,30,31]. Increased understanding of the control of NF- κ B activation and repression would permit not only the development of new strategies to stop active HIV replication but also alternative treatments aimed at reactivation of latent HIV reservoirs in order to reduce them and contribute to viral eradication.

Methods

Cells

Peripheral blood mononuclear cells (PBMCs) were isolated from blood of healthy donors by centrifugation through a Ficoll-Hypaque gradient (Pharmacia Corporation, North Peapack, NJ). Cells were collected in supplemented RPMI and maintained at a concentration of 2×10^6 cells/ml. PHA-treated T lymphocytes were obtained from PBMCs incubated for 3 days with 5 μ g/ml phytohemagglutinin (PHA) (Sigma-Aldrich, St. Louis, MO) and for the consecutive 9 days with 300 U/ml IL-2 (Chiron, Emeryville, CA). These long-term cultures of PHA-treated T lymphocytes were maintained without supplemental IL-2 18 hours before the experiment. These PHA-treated T lymphocytes remained at a pre-activated status and expressed activation markers [35] although NF- κ B did not show DNA-binding activity (Additional file 2).

Resting CD $_4^+$ T lymphocytes were isolated from PBMCs by negative selection with CD $_4$ Negative Isolation Kit (T helper/inducer cells) (DynaL Biotech, Oslo, Norway), according to the manufacturer's instructions. Subsequently, isolated CD $_4^+$ T cells were depleted of CD $_{25}^+$ by positive selection with Dynabeads CD $_{25}$ (DynaL Biotech). Purity of isolated CD $_4^+$ CD $_{25}^-$ T cells was analyzed by flow cytometry with a FACScalibur flow cytometer (BD Biosciences, Erembodegem, Belgium). Cells were stained with monoclonal antibodies (mAb) against CD $_4$ and HLA-DR conjugated with fluorescein isothiocyanate (FITC), and anti-CD $_{25}$, -CD $_{69}$, and -CD $_3$ conjugated with phycoerythrin (PE), all provided by BD Biosciences. Analysis by flow cytometry revealed that the phenotype of isolated T lymphocytes was CD $_4^+$ CD $_{25}^-$ CD $_{69}^-$ HLA-DR $^-$ with a purity >95%.

Reagents and antibodies

Cells were incubated with 25 ng/ml of 5-phorbol 12-myristate 13-acetate (PMA) (Sigma-Aldrich) for 30 min-18 hours. Leptomycin B (LMB) was used at 20 nM (Sigma-Aldrich). Cells treated with 10 μ g/ml of cycloheximide (CHX) (Sigma-Aldrich) were incubated with this reagent 30 minutes before adding other stimuli. Primary antibodies against p65/RelA, p105/p50 and I κ B α were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Secondary antibodies conjugated to horseradish peroxidase were purchased from GE Healthcare (Uppsala, Sweden). Secondary antibodies conjugated to Alexa 488 or Texas Red were purchased from Molecular Probes (Eugene, OR).

Vectors

Luciferase reporter gene under the control of the U3+R regions of the HIV-long terminal repeat (LTR) (LAI strain) was previously reported [36]. pSV- β -Galactosidase vector (Promega, Madison, WI) was used to cotransfect the cells as an internal control reporter. I κ B α gene cloned in pcDNA3.1(+) vector under the control of CMV promoter (CMV-I κ B α) was described previously [37]. Viral Tat gene under the control of the CMV promoter (CMV-Tat) was also described previously [38]. pcDNA3.1(+) vector was used as negative control (Invitrogen, Carlsbad, CA). The vector pNL4.3 that contained the HIV complete genome and induced an infectious progeny after transfection in several cell lines was kindly provided by Dr M.A. Martin [39; National Institute of health AIDS Research and Reference Reagent Program #3418]. Dr. Johannes Schmid kindly provided the constructions of p65/RelA and I κ B α genes in the enhanced yellow fluorescent protein vector (pEYFP-p65 and pEYFP-I κ B α , respectively) [40,41]. Expression vector pEYFP-C1 (Clontech, BD Biosciences) that contains the yellow fluorescent protein gene under CMV promoter control was used as negative control. All plasmids were purified using Qiagen Plasmid Maxi Kit (Qiagen, CA), following the manufacturer's instructions.

Transfection assays

CD $_4^+$ T cells (5×10^6) were transiently transfected with 2 μ g of plasmid DNA under U-14 electroporation program conditions by nucleoporation with an Amaxa Nucleofector (Amaxa, Cologne, Germany) according to the manufacturer's instructions. Alternatively, CD $_4^+$ T cells were also transfected by electroporation with an Easyjet Plus Electroporator (Equibio, Middlesex, UK). In brief, 10×10^6 cells were resuspended in 350 μ l of RPMI without supplements and mixed with 1 μ g of plasmid DNA per 10^6 cells in a 4 mm electroporation cuvette (Equibio). Cells were transfected at 320 V, 1500 μ F and maximum resistance. After transfection, cells were incubated in supplemented RPMI at 37°C for 24 hours before analysis. Luciferase and β -Galactosidase activities were assayed using Luciferase Assay System and β -Galactosidase Enzyme Assay System,

respectively, according to manufacture's instructions (Promega).

Western blot assays

Cytosolic and nuclear protein extracts were obtained as described previously [7]. Ten micrograms of nuclear extracts were fractionated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto Hybond-ECL nitrocellulose paper (GE Healthcare). After blocking and incubation with primary and secondary antibodies, proteins were detected with SuperSignal West Pico Chemiluminescent Substrate (Pierce, Rockford, IL).

Immunoprecipitation assays

Cytosolic and nuclear protein extracts were subjected to immunoprecipitation with agarose-conjugated antibody against p65/RelA (Santa Cruz Biotechnology). In brief, nuclear or cytosolic proteins (100 µg) were incubated overnight at 4°C with 10 µg of specific agarose-conjugated antibody in RIPA buffer (PBS 1×, 0.1% SDS, 1% NP-40) and 0.5% sodium deoxycholate (DOC). Immunoprecipitate was collected by centrifugation at 4°C, 2,500 rpm for 5 minutes and washed four times with RIPA/DOC buffer. Finally, the agarose pellet was denatured at 95°C for 2 minutes and analyzed by SDS-PAGE, followed by immunoblotting with the specific antibodies.

Electrophoretic mobility shift assays (EMSA)

Nuclear protein extracts (3 µg) were analyzed using the [α -³²P]-dCTP-labeled double-stranded synthetic wild-type HIV enhancer oligonucleotide 5'-AGCTTACAAGGGACTTCCGCTGGGGACTTCCAGGGA-3' containing both κB consensus motifs. The nucleoprotein-oligonucleotide complexes were analyzed by electrophoresis on a non-denaturing 6% polyacrylamide gel.

HIV replication assay

Resting CD₄⁺ T cells were transfected with pNL4.3 vector alone or with either CMV-IκBα or pcDNA3.1(+) vectors and further activated with 1 µg/ml anti-CD₃ (BD Biosciences) plus 300 IU/ml IL-2. Viral replication was assessed by quantification of HIV p24 gag antigen in culture supernatants every 48 hours using an enzyme-like immunoassay (Innotest™ HIV Ag mAb, Innogenetics, Barcelona, Spain).

Confocal microscopy

For immunofluorescence assays, cells were immobilized in PolyPrep slides (Sigma-Aldrich) for 15 minutes and then fixed with 2% paraformaldehyde-0.025% glutaraldehyde in PBS for 10 minutes at room temperature. After washing twice with 0.1% glycine/PBS, cells were permeabilized with 0.1% Triton X-100/PBS for 10 minutes. After washing, cells were treated with 1 mg/ml NaBH₄ for 10

minutes. Incubation for 1 hour at room temperature with each primary and secondary antibodies and subsequent washes were performed with PBS/2% bovine serum albumin (BSA)/0.05% saponine buffer. Coverslips were immobilized with 70% glycerol/PBS. Images were obtained with a Radiance 2100 confocal microscope (BioRad, Hercules, CA). For time-lapse fluorescence confocal microscopy, coverslips were coated with fibronectin (20 µg/ml) for 2 h at 37°C and blocked with PBS containing BSA 0.1%. Then, coverslips were washed with 1× Hanks Balanced Salt Solution (HBSS) and mounted in Attofluor open chambers (Molecular Probes). Cells were allowed to adhere on these chambers for 30 min. Confocal images were acquired using a Leica TCS-SP Confocal Laser Scanning Unit (Leica, Heidelberg, Germany) equipped with Ar and He-Ne laser beams and attached to a Leica DMIRBE Inverted Epi-Fluorescence Microscope. Images were processed and assembled into movies using Leica confocal software.

Statistical analysis

Differences in HIV replication in the presence of IκBα over-expression were assessed by Mann-Whitney test using Statistical Product and Service Solutions (SPSS) software v14 (Addlink Software Científico, Madrid, Spain).

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

MT carried out all the molecular biology studies and drafted the manuscript.

MRLH carried out the CD₄⁺ T cell isolation and performed the HIV replication assays.

JR and MM participated in the analyses by confocal microscopy.

JA conceived of the study, and participated in its design and coordination and helped to draft the manuscript.

All authors read and approved the final manuscript.

Additional material

Additional file 1

Kinetic analysis of nuclear I κ B α translocation. Movie of one CD4⁺T lymphocyte transfected with EYFP-I κ B α vector photographed before and after treatment with LMB up to 30 minutes. Photographs were taken in vivo by confocal microscopy every minute after adding LMB. This movie corresponds to Figure 3, which has been constructed with some pictures taken at different moments before and after adding LMB. It is an .avi file that can be viewed with Windows Media Player.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1742-4690-4-56-S1.avi>]

Additional file 2

Absence of NF- κ B binding activity in nuclear protein extracts from unstimulated PHA-treated T cells. (a) Binding of NF- κ B in nuclear extracts from PHA-treated T cells to its cognate DNA sequence was analyzed. PBMCs were cultured for 3 days with 5 μ g/ml PHA and for the consecutive 9 days with 300 U/ml IL-2. These long-term cultures of PHA-treated T lymphocytes were maintained without supplemental IL-2 18 hours. Three micrograms of nuclear extracts from IL-2 depleted T cells (lane 1) and activated with PMA for 30 min or 5 hours (lanes 2 and 3, respectively) were incubated with an oligonucleotide containing double - κ B consensus motif from HIV LTR labeled with [α -³²P]-dCTP. (b) Analysis of the NF- κ B complexes composition by supershift assay. Three micrograms of nuclear extracts from PHA-treated T cells activated with PMA for 2 hours were incubated with antibodies against p50/NF- κ B1 (lane 3), p65/RelA (lane 4) or c-Rel (lane 5) before the incubation with an oligonucleotide containing double - κ B consensus motif from HIV LTR labeled with [α -³²P]-dCTP. Lane 2 shows the specificity of binding of the NF- κ B complexes using excess (100 \times) of unlabelled - κ B-motif oligonucleotide as competitor.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1742-4690-4-56-S2.ppt>]

Acknowledgements

We would like to thank Belén García-Fernández and Elena Mateos for excellent technical assistance and Olga Palao for secretarial assistance. We also thank Dr. Johannes Schmid (Department of Vascular Biology and Thrombosis Research, University of Vienna, Austria) for the gift of EYFP-I κ B α and EYFP-p65 constructions, Dr. Fernando Arenzana-Seisdedos (Institut Pasteur, Paris, France) for the gift of the CMV-I κ B α vector, and Dr. Carles Suñé (National Centre for Biotechnology, Madrid, Spain) for the gift of CMV-Tat construction. We also thank to the Center of Blood Transfusions (Comunidad de Madrid) for kindly providing blood of healthy donors. This work was supported by the following projects: ISCIII-RETIC-RD06/0006/0037, SAF 2004-04258, SAF 2000-00/0028, FIPSE 36453/03, VIRHORST Network from Comunidad de Madrid, Spain.

References

- Ghosh S, Karin M: **Missing pieces in the NF κ B puzzle.** *Cell Suppl* 2002, **109**:S81-S96.
- Li Q, Verma IM: **NF- κ B regulation in the immune system.** *Nat Rev Immunol* 2002, **2**:725-734.
- Arenzana-Seisdedos F, Thompson J, Rodríguez MS, Bachelier F, Thomas D, Hay RT: **Inducible nuclear expression of newly synthesized I κ B alpha negatively regulates DNA-binding and transcriptional activities of NF- κ B.** *Mol Cell Biol* 1995, **15**:2689-2696.
- Arenzana-Seisdedos F, Turpin P, Rodríguez M, Thomas D, Hay RT, Virelizier JL, Dargemont C: **Nuclear localization of I κ B**

- alpha promotes active transport of NF- κ B from the nucleus to the cytoplasm.** *J Cell Sci* 1997, **110**:369-378.
- Turpin P, Hay RT, Dargemont C: **Characterization of I κ B α nuclear import pathway.** *J Biol Chem* 1999, **274**:6804-6812.
- Hiscott J, Kwon H, Genin P: **Hostile takeovers: viral appropriation of the NF- κ B pathway.** *J Clin Invest* 2001, **107**:143-151.
- Lain de Lera T, Folgueira L, Martín AG, Dargemont C, Pedraza MA, Bermejo M, Bonay P, Fresno M, Alcami J: **Expression of I κ B α in the nucleus of human peripheral blood T lymphocytes.** *Oncogene* 1999, **18**:1581-1588.
- Nabel G, Baltimore D: **An inducible transcription factor activates expression of human immunodeficiency virus in T cells.** *Nature* 1987, **326**:711-713.
- Roulston A, Lin R, Beauparlant P, Wainberg MA, Hiscott J: **Regulation of HIV-1 and cytokine gene expression in myeloid cells by NF- κ B/Rel transcription factors.** *Microbiol Rev* 1995, **59**:481-505.
- Baldwin AS Jr: **The NF- κ B and I κ B proteins: new discoveries and insights.** *Annu Rev Immunol* 1996, **14**:649-681.
- Alcami J, Lain de Lera T, Folgueira L, Pedraza MA, Jacqué JM, Bachelier F, Noriega AR, Hay RT, Harrich D, Gaynor RB, Virelizier JL, Arenzana-Seisdedos F: **Absolute dependence on κ B responsive elements for initiation and Tat-mediated amplification of HIV transcription in blood CD4 T lymphocytes.** *EMBO J* 1995, **14**:1552-1560.
- Zhang Z, Schuler T, Zupancic M, Wietgreffe S, Staskus KA, Reimann KA, Reinhart TA, Rogan M, Cavert W, Miller CJ, Veazey RS, Notermans D, Little S, Danner SA, Richman DD, Havlir D, Wong J, Jordan HL, Schacker TW, Racz P, Tenner-Racz K, Letvin NL, Wolinsky S, Haase AT: **Sexual transmission and propagation of SIV and HIV in resting and activated CD4⁺ T cells.** *Science* 1999, **286**:1353-1357.
- Eckstein DA, Penn ML, Korin YD, Scripture-Adams DD, Zack JA, Kreisberg JF, Roederer M, Sherman MP, Chin PS, Goldsmith MA: **HIV-1 actively replicates in naive CD4(+) T cells residing within human lymphoid tissues.** *Immunity* 2001, **15**:671-682.
- Pomerantz RJ: **Residual HIV-1 disease in the era of highly active antiretroviral therapy.** *N Engl J Med* 1999, **340**:1672-1674.
- Nishi K, Yoshida M, Fujiwara D, Nishikawa M, Horinouchi S, Beppu T: **Leptomycin B targets a regulatory cascade of crm1, a fission yeast nuclear protein, involved in control of higher order chromosome structure and gene expression.** *J Biol Chem* 1994, **269**:6320-6324.
- Beauparlant P, Kwon H, Clarke M, Lin R, Sonenberg N, Wainberg M, Hiscott J: **Transdominant mutants of I κ B alpha block Tat-tumor necrosis factor synergistic activation of human immunodeficiency virus type I gene expression and virus multiplication.** *J Virol* 1996, **70**:5777-5785.
- Persaud D, Zhou Y, Siliciano JM, Siliciano RF: **Latency in human immunodeficiency virus type I infection: no easy answers.** *J Virol* 2003, **77**:1659-1665.
- Rohr O, Marban C, Aunis D, Schaeffer E: **Regulation of HIV-1 gene transcription: from lymphocytes to microglial cells.** *J Leukoc Biol* 2003, **74**:736-749.
- Palmieri C, Trimboli F, Puca A, Fiume G, Scala G, Quinto I: **Inhibition of HIV-1 replication in primary human monocytes by the I κ B α -alphaS32/36A repressor of NF- κ B.** *Retrovirology* 2004, **1**:45.
- Quinto I, Mallardo M, Baldassarre F, Scala G, Englund G, Jeang KT: **Potent and stable attenuation of live-HIV-1 by gain of a proteolysis-resistant inhibitor of NF- κ B (I κ B α -alphaS32/36A) and the implications for vaccine development.** *J Biol Chem* 1999, **274**:17567-17572.
- Quinto I, Puca A, Greenhouse J, Silvera P, Yalley-Ogunro J, Lewis MG, Palmieri C, Trimboli F, Byrum R, Adelsberger J, Venzon D, Chen X, Scala G: **High attenuation and immunogenicity of a simian immunodeficiency virus expressing a proteolysis-resistant inhibitor of NF- κ B.** *J Biol Chem* 2004, **279**:1720-1728.
- Johnson C, Van Antwerp D, Hope TJ: **An N-terminal nuclear export signal is required for the nucleocytoplasmic shuttling of I κ B α .** *EMBO J* 1999, **18**:6682-6693.
- Carloti F, Dower SK, Qwarnstrom EE: **Dynamic shuttling of nuclear factor κ B between the nucleus and cytoplasm as a consequence of inhibitor dissociation.** *J Biol Chem* 2000, **275**:41028-41034.
- Huang TT, Kudo N, Yoshida M, Miyamoto SA: **Nuclear export signal in the N-terminal regulatory domain of I κ B α controls cytoplasmic localization of inactive NF- κ B/I κ B α complexes.** *Proc Natl Acad Sci USA* 2000, **97**:1014-1019.

25. Chun TW, Finzi D, Margolick J, Chadwick K, Schwartz D, Siliciano RF: **In vivo fate of HIV-1-infected T cells: quantitative analysis of the transition to stable latency.** *Nat Med* 1995, **1**:1284-1290.
26. Pierson T, McArthur J, Siliciano RF: **Reservoirs for HIV-1: mechanisms for viral persistence in the presence of antiviral immune responses and antiretroviral therapy.** *Annu Rev Immunol* 2000, **18**:665-708.
27. Finzi D, Hermankova M, Pierson T, Carruth LM, Buck C, Chaisson RE, Quinn TC, Chadwick K, Margolick J, Brookmeyer R, Gallant J, Markowitz M, Ho DD, Richman DD, Siliciano RF: **Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy.** *Science* 1997, **278**:1295-1300.
28. Wong JK, Hezareh M, Gunthard HF, Havlir DV, Ignacio CC, Spina CA, Richman DD: **Recovery of replication-competent HIV despite prolonged suppression of plasma viremia.** *Science* 1997, **278**:1291-1295.
29. Kinter AL, Umscheid CA, Arthos J, Cicala C, Lin Y, Jackson R, Donoghue E, Ehler L, Adelsberger J, Rabin RL, Fauci AS: **HIV envelope induces virus expression from resting CD4+ T cells isolated from HIV-infected individuals in the absence of markers of cellular activation or apoptosis.** *J Immunol* 2003, **170**:2449-2455.
30. Lassen KG, Bailey JR, Siliciano RF: **Analysis of human immunodeficiency virus type 1 transcriptional elongation in resting CD4+ T cells in vivo.** *J Virol* 2004, **78**:9105-9114.
31. Zhang ZQ, Wietgrefe SW, Li Q, Shore MD, Duan L, Reilly C, Lifson JD, Haase AT: **Roles of substrate availability and infection of resting and activated CD4+ T cells in transmission and acute simian immunodeficiency virus infection.** *Proc Natl Acad Sci USA* 2004, **101**:5640-5645.
32. Swiggard WJ, Baytop C, Yu JJ, Dai J, Li C, Schretzenmair R, Theodosopoulos T, O'Doherty U: **Human immunodeficiency virus type 1 can establish latent infection in resting CD4+ T cells in the absence of activating stimuli.** *J Virol* 2005, **79**:14179-14188.
33. Kwon H, Pelletier N, DeLuca C, Genin P, Cisternas S, Lin R, Wainberg MA, Hiscott J: **Inducible expression of I κ B α repressor mutants interferes with NF- κ B activity and HIV-1 replication in Jurkat T cells.** *J Biol Chem* 1998, **273**:7431-7440.
34. Ganesh L, Burstein E, Guha-Niyogi A, Louder MK, Mascola JR, Klomp LW, Wijmenga C, Duckett CS, Nabel GJ: **The gene product Murr1 restricts HIV-1 replication in resting CD4+ lymphocytes.** *Nature* 2003, **426**:853-857.
35. Li ZW, Rickert RC, Karin M: **Genetic dissection of antigen receptor induced-NF-kappaB activation.** *Mol Immunol* 2004, **41**:701-714.
36. Bachelier F, Alcamí J, Arenzana-Seisdedos F, Virelizier JL: **HIV enhancer activity perpetuated by NF-kappaB induction on infection of monocytes.** *Nature* 1991, **350**:709-712.
37. Rodríguez MS, Michalopoulos I, Arenzana-Seisdedos F, Hay RT: **Inducible degradation of I κ B α in vitro and in vivo requires the acidic C-terminal domain of the protein.** *Mol Cell Biol* 1995, **15**:2413-2419.
38. Malim MH, Hauber J, Fenrick R, Cullen BR: **Immunodeficiency virus rev trans-activator modulates the expression of the viral regulatory genes.** *Nature* 1988, **335**:181-183.
39. Adachi A, Gendelman HE, Koenig S, Folks T, Willey R, Rabson A, Martin MA: **Production of acquired immunodeficiency syndrome-associated retrovirus in human and non-human cells transfected with an infectious molecular clone.** *J Virol* 1988, **59**:284-291.
40. Schmid JA, Birbach A, Hofer-Warbinek R, Peng M, Burner U, Furtmüller PG, Binder BR, de Martin R: **Dynamics of NF kappa B and I kappa B alpha studied with green fluorescent protein (GFP) fusion proteins. Investigation of GFP-p65 binding to DNA by fluorescence resonance energy transfer.** *J Biol Chem* 2000, **275**:17035-17042.
41. Birbach A, Gold P, Binder BR, Hofer E, de Martin R, Schmid JA: **Signaling molecules of the NF-kappa B pathway shuttle constitutively between cytoplasm and nucleus.** *J Biol Chem* 2002, **277**:10842-10851.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

