

RESEARCH ARTICLE

Open Access

Conditional BDNF release under pathological conditions improves Huntington's disease pathology by delaying neuronal dysfunction

Albert Giralt^{1,2,3}, Olga Carretón^{1,2,3}, Cristina Lao-Peregrin⁴, Eduardo D Martín⁴ and Jordi Alberch^{1,2,3*}

Abstract

Background: Brain-Derived Neurotrophic Factor (BDNF) is the main candidate for neuroprotective therapy for Huntington's disease (HD), but its conditional administration is one of its most challenging problems.

Results: Here we used transgenic mice that over-express BDNF under the control of the Glial Fibrillary Acidic Protein (GFAP) promoter (pGFAP-BDNF mice) to test whether up-regulation and release of BDNF, dependent on astrogliosis, could be protective in HD. Thus, we cross-mated pGFAP-BDNF mice with R6/2 mice to generate a double-mutant mouse with mutant huntingtin protein and with a conditional over-expression of BDNF, only under pathological conditions. In these R6/2;pGFAP-BDNF animals, the decrease in striatal BDNF levels induced by mutant huntingtin was prevented in comparison to R6/2 animals at 12 weeks of age. The recovery of the neurotrophin levels in R6/2;pGFAP-BDNF mice correlated with an improvement in several motor coordination tasks and with a significant delay in anxiety and clasping alterations. Therefore, we next examined a possible improvement in cortico-striatal connectivity in R6/2;pGFAP-BDNF mice. Interestingly, we found that the over-expression of BDNF prevented the decrease of cortico-striatal presynaptic (VGLUT1) and postsynaptic (PSD-95) markers in the R6/2:pGFAP-BDNF striatum. Electrophysiological studies also showed that basal synaptic transmission and synaptic fatigue both improved in R6/2;pGFAP-BDNF mice.

Conclusions: These results indicate that the conditional administration of BDNF under the GFAP promoter could become a therapeutic strategy for HD due to its positive effects on synaptic plasticity.

Keywords: Huntingtin, synaptic plasticity, therapy, astrocytes, learning, Long-term potentiation

Background

Huntington's disease (HD) is an inherited neurodegenerative disorder, which results from the expansion of a CAG trinucleotide repeat in the *huntingtin* (*htt*) gene [1,2]. Mutant *htt* (*mhtt*) causes cortical atrophy and the preferential death of medium-sized spiny neurons in the striatum. The demise of these neurons causes motor and cognitive dysfunction [3]. In exon-1 and full-length mouse models of HD, a progressive decline in cortico-striatal synaptic function that correlates with the pathology has been described [4,5]. Thus, molecules regulating

cortico-striatal connectivity would be extremely relevant to HD therapy perspectives.

Brain-Derived Neurotrophic Factor (BDNF) is an important neuroprotective factor for the preservation of medium-sized spiny neurons in the striatum [6]. However, the trophic support of striatal neurons is provided by cortical afferents [7,8]. A reduction in BDNF levels has been described in HD patients [9,10] and similar results have been found in several animal models [11-17]. Furthermore, the transport and function of BDNF are also disrupted in HD [18,19]. Thus, as one pathogenic mechanism leading to dysfunction and loss of striatal neurons in HD might be the reduction in BDNF levels, its administration should improve these alterations [20-22]. In fact, BDNF is a strong candidate for neuroprotective therapies, as it has been tested in

* Correspondence: alberch@ub.edu

¹Departament de Biologia Cel·lular, Immunologia i Neurociències, Facultat de Medicina, Universitat de Barcelona, Barcelona, Spain

Full list of author information is available at the end of the article

acute [23,24] and in transgenic mouse models [12,25] of HD. However, the main challenge is to find the way to administer BDNF chronically and conditionally to the target neurons [22,26-28].

We previously demonstrated in an excitotoxic model of HD that transgenic astrocytes engineered to over-express BDNF under the control of the GFAP promoter, when grafted in wt mice, release higher levels of BDNF than control astrocytes [23]. This enhanced release of BDNF protects striatal neurons. Here we tested whether conditional and pathology-dependent delivery of BDNF could be neuroprotective in a transgenic mouse model of HD. We cross-mated R6/2 [29] with pGFAP-BDNF mice [23] to obtain double-mutant animals: R6/2:pGFAP-BDNF. Our results show that BDNF over-expression from striatal astrocytes improved HD phenotype by preventing cortico-striatal synaptic dysfunction.

Results


GFAP up-regulation increases BDNF levels in the striatum of R6/2:pGFAP-BDNF mice

In this study, we crossed mice that over-express BDNF under the control of the GFAP promoter (pGFAP-BDNF) [23] with the HD mouse model R6/2 [29]. With this procedure we obtained double-mutant R6/2:pGFAP-BDNF mice, which would express BDNF depending on the degree of astrogliosis.

First, we checked whether 12-week-old R6/2 and R6/2:pGFAP-BDNF mice displayed astrogliosis by performing immunohistochemistry against GFAP. The results showed clear astroglial reactivity in the striatum of HD mouse models (Figure 1A). Accordingly, quantification by Western blot showed a significantly equal increase of striatal GFAP in R6/2 and R6/2:pGFAP-BDNF mice (Figure 1B). As previously observed in the excitotoxic model [23], this neuropathological hallmark induced an up-regulation of the pGFAP-BDNF transgene activity only in R6/2:pGFAP-BDNF mice, which correlates with significantly higher striatal BDNF levels than in R6/2 mice (Figure 1C). These results validated our model because although both, R6/2 and R6/2:pGFAP-BDNF mice displayed equal levels of striatal GFAP, the presence of the pGFAP-BDNF transgene only in the R6/2:pGFAP-BDNF mice prompted an increase and recovery of striatal BDNF levels in these mice relative to R6/2 mice who lacked the pGFAP-BDNF transgene.

R6/2:pGFAP-BDNF mice show improvements in motor coordination

We next evaluated motor coordination in wt, pGFAP-BDNF, R6/2 and R6/2:pGFAP-BDNF mice using the rotarod test. Disturbed motor coordination was observed in R6/2, compared to wt and pGFAP-BDNF mice at 9-10 weeks of age, at 16 rpm (Figure 2A) and 24 rpm



(Figure 2B) and steadily worsened up to 12 weeks of age (Figure 2A-B). However, motor coordination deficits in R6/2:pGFAP-BDNF mice advanced with significantly less severity than in R6/2 mice (Figure 2A-B).

To further confirm better motor coordination in R6/2:pGFAP-BDNF than in R6/2 mice, we also evaluated their performance on the balance beam. The test with the balance beam was performed at 7, 10 and 12 weeks of age. The distance covered was the parameter used to check motor coordination. The results showed that performance was strongly affected in both R6/2 and R6/2:




Figure 2 Preservation of striatal BDNF levels in R6/2;pGFAP-BDNF mice is associated with an improvement of their motor performance over R6/2 mice. Motor coordination and balance were analyzed by performing the rotarod task at 16 (A) and 24 (B) rpm in 9-, 10-, 11- and 12-week-old wt, pGFAP-BDNF, R6/2 and R6/2;pGFAP-BDNF mice. Motor coordination and balance were further assessed by the balance beam test at 7, 10 and 12 weeks of age (C). Ataxia and gait abnormalities were assessed by the footprint test, analyzing front paw length (D) and paw overlap (E) at 7, 10 and 12 weeks of age in wt, pGFAP-BDNF, R6/2 and R6/2;pGFAP-BDNF mice. Values are expressed as mean \pm s.e.m. Data were analyzed by one-way ANOVA at different stages of the disease's progression in the case of the footprint and the balance beam tests because different cohorts of animals were used. In the case of the rotarod task, two-way ANOVA with repeated measures was performed. For clarity, in A and B only statistical comparisons between R6/2 and R6/2;pGFAP-BDNF mice are depicted. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$, as compared to wt mice; \$ $p < 0.05$, \$\$ $p < 0.01$ and \$\$\$ $p < 0.001$, as compared to pGFAP-BDNF mice; and # $p < 0.05$ and ## $p < 0.01$, as compared to R6/2 mice (n = 10-12/genotype).

pGFAP-BDNF mice from 7 weeks of age (Figure 2C), before any overt phenotype is detectable. Interestingly, these alterations worsened in R6/2 but not in R6/2:pGFAP-BDNF mice, resulting in a significantly higher ability to cross the beam of R6/2:pGFAP-BDNF mice than of R6/2 mice at 12 weeks of age (Figure 2C).


Ataxia and gait abnormalities were evaluated by the footprint test. We found that both R6/2 and R6/2:pGFAP-BDNF mice displayed significant abnormalities in their front paw length (Figure 2D) and paw overlap (Figure 2E), when compared with their controls. Front paw length was significantly shorter in both R6/2 and R6/2:pGFAP-BDNF mice from 7 weeks of age, when loss of body weight was not yet evident (Figure 3A). No significant differences were observed between R6/2 and R6/2:pGFAP-BDNF mice (Figure 2D). On the other hand, paw overlap was significantly higher at 12 weeks of age in both R6/2 and R6/2:pGFAP-BDNF mice than in wt and pGFAP-BDNF mice (Figure 2E). In conclusion, the footprint test showed that gait and ataxia abnormalities were disturbed equally in R6/2 and R6/2:pGFAP-BDNF mice.

The onset of the clasping reflex is ameliorated in R6/2:pGFAP-BDNF mice

Next, we studied additional phenotype alterations characteristic of R6/2 mice, such as loss of body weight and muscular strength and the development of the clasping reflex. Both R6/2 and R6/2:pGFAP-BDNF mice developed a normal body weight increase, compared with wt and pGFAP-BDNF mice, up to 10 weeks of age (Figure 3A). However, at this age R6/2 and R6/2:pGFAP-BDNF mice stopped gaining body weight; and at 11 weeks of age, both began to lose weight, with no significant differences between them (Figure 3A). Muscular strength was evaluated by the hiring wire test. This test showed that the muscular strength of R6/2 and R6/2:pGFAP-BDNF mice is normal, compared with wt and pGFAP-BDNF mice, up to 12 weeks of age, when both groups showed disorders in this parameter, as shown by a significant increase in falls from the grid (Figure 3B). On the other hand, both R6/2 and R6/2:pGFAP-BDNF mice steadily increased their clasping behavior (Figure 3C). However, R6/2 mice showed an early onset (8 weeks of age) of clasping behavior, whereas R6/2:pGFAP-BDNF mice started at 11 weeks of age (Figure 3C), suggesting significant delay of the associated neuronal dysfunction.


Anxiety alterations are delayed in R6/2:pGFAP-BDNF mice

To further characterize the behavioral phenotype improvements in R6/2:pGFAP-BDNF mice, compared to R6/2 mice, we also used the plus maze and the open field paradigms, which mainly check anxiety levels and spontaneous locomotor activity, respectively (see



Methods). The results obtained in the plus maze indicated that at all ages analyzed (7, 10 and 12 weeks of age) R6/2 mice spent significantly more time in the open arms than wt and pGFAP-BDNF mice did (Figure 4A). This result suggests aberrant, decreased anxiety levels, as described elsewhere for these mice [30]. On the other hand, R6/2:pGFAP-BDNF mice showed a significant delay in the appearance of these alterations, which was not significant relative to wt mice until 12 weeks of age (Figure 4A). Thus, R6/2:pGFAP-BDNF mice showed an improvement in anxiety-like disorders.

To study the locomotor activity and exploration levels, we also performed the open field test. As previously described in HD mouse models [31,32], R6/2 and R6/2:pGFAP-BDNF mice showed a progressive decrease in spontaneous locomotor and exploratory activity in the open field, which reach significance relative to wt and pGFAP-BDNF mice at 12 weeks of age (Figure 4B). However, this significant trend to hypoactivity was equally developed in both R6/2 and R6/2:pGFAP-BDNF mice. These results suggest that less anxiety-like behavior in the plus maze shown by HD transgenic animals



is unlikely to be caused by higher exploratory activity. Furthermore, the results indicate that the normalization of anxiety levels observed in R6/2:pGFAP-BDNF mice is not prompted by simple recovery of spontaneous locomotor activity.

Loss of striatal volume and soma size of calbindin neurons and nuclear inclusions formation are ameliorated in R6/2:pGFAP-BDNF mice

As R6/2:pGFAP-BDNF mice showed greater preservation of their striatal-dependent motor coordination than R6/2 mice did, we next evaluated whether striatal neuropathology correlated with these behavioral results. First, although the pGFAP-BDNF transgene was expressed in the whole brain in R6/2:pGFAP-BDNF mice, this expression pattern was not enough to recover the reduction of total brain weight induced by mutant huntingtin (in grams; wt: 0.463 ± 0.013 ; pGFAP-BDNF: 0.455 ± 0.005 ; R6/2: 0.396 ± 0.014 ; R6/2:pGFAP-BDNF: 0.397 ± 0.020 ; p value < 0.05 for R6/2 and R6/2:pGFAP-BDNF mice compared with wt and pGFAP-BDNF mice). In contrast, stereological estimation of the striatal volume revealed that R6/2:pGFAP-BDNF mice have larger striata than R6/2 mice did (Figure 5A). Next we checked whether the striatal volume preservation found in R6/2:pGFAP-BDNF mice was due to greater survival of striatal projection neurons. Stereological counting of the calbindin-positive projection neurons in all groups indicated that both R6/2 and R6/2:pGFAP-BDNF mice have a similar significant decrease in the number of these neurons, in comparison with wt and pGFAP-BDNF mice (Figure 5B). However, when we measured the soma volume of calbindin-positive neurons, we found a significant volume improvement in R6/2:pGFAP-BDNF over R6/2 mice (Figure 5C). Finally, we wanted to check the number of Nuclear Inclusions (NIIs) in the striatum from our mouse models (Figure 5D). We observed that density of NIIs were lower in R6/2:pGFAP-BDNF relative to R6/2 mice striata (Figure 5D).

Effect of BDNF over-expression on levels of DARPP-32 and enkephalin in the R6/2:pGFAP-BDNF striatum

Enkephalin protein levels fall strongly in human HD brains and in mouse models [3,25,33,34]. Therefore, we used a specific antibody for Western blot that recognizes two enkephalin-positive bands at ~30 kDa [35] as previously described [36]. Our results show that R6/2:pGFAP-BDNF mice showed significant improvement in striatal enkephalin protein levels, compared with R6/2 mice (Figure 6A). To analyze the biochemical improvements in the striatum of R6/2:pGFAP-BDNF mice further, DARPP-32 protein levels were also analyzed in the striata from all genotypes (Figure 6B). DARPP-32

protein levels were significantly higher in R6/2:pGFAP-BDNF mice than in R6/2 ones (Figure 6B).

Cortico-striatal synapses are preserved in R6/2:pGFAP-BDNF mice

As we found a preservation of the volume of striatal projection neurons in R6/2:pGFAP-BDNF mice, but not of their number, in comparison with R6/2 animals, we hypothesized that mainly subtle micro-structural changes were involved in R6/2:pGFAP-BDNF mouse phenotype improvements. Thus, we additionally tested the state of cortico-striatal connectivity by counting VGLUT1- and PSD-95-positive particle densities in the striatum of wt, pGFAP-BDNF, R6/2 and R6/2:pGFAP-BDNF 12-week-old mice.

Confocal image analysis of coronal sections at 12 weeks of age showed that only R6/2 mice had a significant decrease in striatal VGLUT1- (Figure 7A) and PSD-95-positive (Figure 7B) particle density, in comparison with wt mice. However, R6/2:pGFAP-BDNF mice displayed highly intact striatal VGLUT1- and PSD-95-positive particle density, compared with wt mice (Figure 7). These results suggest that both excitatory terminals (VGLUT1) and dendritic spines (PSD-95) are preserved in R6/2:pGFAP-BDNF mice. Moreover, these results are in line with the volume preservation of the whole striatum and striatal calbindin-positive neurons in R6/2:pGFAP-BDNF mice, compared with R6/2 ones.

Intact cortico-striatal synaptic transmission and resistance to synaptic fatigue in R6/2:pGFAP-BDNF mice

Due to the improvement in the number of synaptic particles in R6/2:pGFAP-BDNF mice over R6/2 mice, we next examined glutamatergic synaptic transmission in the different groups. We made extracellular recordings of field potentials in acute cortico-striatal brain slices. As described in previous studies, single-pulse stimulation of striatal slices elicited a characteristic response with different components [37-39]. One of these components, the population spike (PS), reflects excitatory monosynaptic transmission in a population of striatal neurons due to endogenous glutamate release upon electrical stimulation of glutamate-containing fibers. This component is mediated by activation of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) glutamate receptors [38]. First, we analyzed basal synaptic transmission by applying isolated stimuli of increasing intensity. Extracellular field recordings showed that R6/2 mice tended to exhibit decreased PS responses, which were statistically significant at high stimulus strengths (Figure 8A; $V = 10$ V, $p < 0.01$). Interestingly, the amplitude of R6/2:pGFAP-BDNF PS responses was not significantly different from PS response in wt or pGFAP-BDNF mice (Figure 8A), indicating that the





Figure 5 Striatal neuropathology is improved in R6/2;pGFAP-BDNF mice compared to R6/2 mice. Striatal volume (A) was stereologically measured in 12-week-old wt, pGFAP-BDNF, R6/2 and R6/2;pGFAP-BDNF mice. A representative microphotograph of coronal sections immunostained for calbindin is depicted (A). The number of calbindin-positive projecting neurons (B) was stereologically counted in all groups. The soma volume of calbindin-positive projecting neurons (C) was stereologically calculated in all genotypes by the nucleator method. Density of EM48-positive Nuclear Inclusions (NIIs) were stereologically counted in R6/2 and R6/2;pGFAP-BDNF mice (D). Values are expressed as mean \pm s.e.m. Data were analyzed by one-way ANOVA or unpaired t-test. * $p < 0.05$ and ** $p < 0.01$, as compared to wt mice; # $p < 0.05$, as compared to R6/2 mice ($n = 5-6$ /genotype). Scale bar (A) 1 mm and (D) 10 μ m.



efficacy of excitatory synaptic inputs is greater in R6/2: pGFAP-BDNF mice. Since mhtt has been shown to impair synaptic transmission in the cortico-striatal pathway [40,41], we next examined synaptic fatigue by means of PS recordings evoked by a four-stimulus train (100 Hz). After the second stimulus in the train, the PS response was significantly smaller in R6/2 slices than in wt slices (Figure 8B, $p < 0.05$). Interestingly, the PS response in pGFAP-BDNF and R6/2:pGFAP-BDNF mice was significantly greater than in wt and R6/2 mice (Figure 8B), indicating that R6/2:pGFAP-BDNF mice were more resistant to synaptic fatigue.

We next investigated cortico-striatal synaptic plasticity by inducing LTP by high-frequency conditioning tetanus (see Methods). Baseline responses were monitored for 10-30 min before conditioning and were found to be stable. Tetanus conditioning revealed a marked difference in the ability of transgenic HD mice to support LTP, with strengthening of PS significantly low in R6/2 and R6/2:pGFAP-BDNF mice, whereas wt and pGFAP-BDNF mice had a significant PS response (Figure 8C, $p < 0.001$).

Discussion

Here we demonstrate that conditional BDNF production and release from astrocytes is a safe and neuroprotective method to improve the R6/2 mice phenotype. Double-mutant R6/2:pGFAP-BDNF mice show improvements in striatal-dependent behavior, anxiety alterations and clasping levels, compared with R6/2 mice. In support of this, R6/2:pGFAP-BDNF mice show associated significant preservation of neuronal architecture, which correlates with improved cortico-striatal connectivity.

The neurotrophin BDNF has been widely put forward as a possible therapeutic molecule for HD treatment [20,22]. However, the key challenge in the field of growth factor therapy is drug delivery to the central nervous system. Another important consideration in the design of growth factor therapies for neurodegenerative disorders is the need for extended periods of treatment [42]. To date, there are no successful systems for delivering BDNF, because it does not cross the blood-brain barrier *via* peripheral administration [43,44]. On the other hand, approaches such as infection/lipotransfection and cell therapy are still inaccessible because of their collateral toxicity effects and the likelihood of their inducing aberrant proliferations, respectively. In addition, they are invasive procedures [27,45].

Both R6/2 mice [46-48] and HD human patients [49] show a striatal increase in astrogliosis reactivity with an associated up-regulation of GFAP levels. In the present study, we took advantage of striatal astrogliosis and the





Figure 7 Preservation of cortico-striatal connectivity in R6/2;pGFAP-BDNF mice. Immunohistochemical staining against the pre-synaptic marker VGLUT1 (A) and the post-synaptic marker PSD-95 (B) was performed in 12-week-old wt, pGFAP-BDNF, R6/2 and R6/2;pGFAP-BDNF mice. The images were taken by confocal microscopy. VGLUT1- (A) and PSD-95- (B) positive particles were counted by the ImageJ software and represented in a graph as density of particles/field. Values are expressed as mean \pm s.e.m. Data were analyzed by one-way ANOVA. * $p < 0.05$ and ** $p < 0.01$ as compared to wt mice; \$ $p < 0.05$ and \$\$ $p < 0.01$ as compared to pGFAP-BDNF mice; # $p < 0.05$ as compared to R6/2 mice (n = 4/group). Scale bar 15 μ m.



consequent up-regulation of GFAP promoter activity in R6/2 mice to produce and release BDNF physiologically. We accomplished this aim by crossing R6/2 mice with pGFAP-BDNF mice and obtaining double-mutant mice (R6/2:pGFAP-BDNF). The transgene works in a conditional fashion because it depends on the presence of astrogliosis, since pGFAP-BDNF mice did not show any increase in striatal BDNF levels compared with wt mice. Therefore, the increase in striatal BDNF in R6/2:pGFAP-BDNF mice, in comparison with R6/2 mice, was due to a conditional release, with auto-regulated mechanisms and only when it is pathologically required.

In this model, astrogliosis enables striatal BDNF expression to be significantly restored to close to control levels. Previous studies employed a *Bdnf* transgene driven by the promoter for the alpha subunit of Ca^{2+} /calmodulin-dependent kinase II to over-express BDNF in the forebrain of R6/1 [12] and YAC128 [34] mice, showing high over-expression of BDNF that slowed the progression of the disease. Intriguingly, the beneficial effects observed in the present study are very similar to those described in these studies. Therefore, in an HD context, a small increase in BDNF levels is enough to provide a significant symptomatic improvement, which demonstrates the high relevance of this neurotrophin in the pathophysiology of the disease. This emphasizes the need to control the amount of BDNF delivered, since excessive chronic production and delivery of the neurotrophin to the brain may induce neuronal dysfunction and toxicity [50,51].

The behavioral characterization of R6/2:pGFAP-BDNF mice showed that the dysfunctions in motor coordination, evaluated by the rotarod and the balance beam tests, were delayed and improved, suggesting that the preservation of striatal BDNF levels had a strong positive effect on these mice. These results provide further support for the importance of BDNF in motor coordination function [52] and specifically of the striatum-dependent component [7]. Intriguingly, we did not find any changes in gait and locomotor deficits when mice performed the footprint and open field tests, which indicated that the improved striatal BDNF levels in R6/2:pGFAP-BDNF mice were not enough to ameliorate these symptoms. On the other hand, not only was motor coordination enhanced in R6/2:pGFAP-BDNF, but other neurological alterations typical of R6/2 mice were also improved, such as anxiety disturbances and the clasp reflex [29,30]. The improvement in anxiety alterations corroborated previous studies showing that one molecular mechanism able to modulate these processes was BDNF [53,54]. On the other hand, these results do not rule out possible functional improvements

in other brain regions involved in the regulation of anxiety, such as the limbic-medial prefrontal circuit or the amygdala [55], in R6/2:pGFAP-BDNF mice and point to the need for further studies on this possibility.

Our data support the idea that neural dysfunction is a critical and early point in the pathophysiology of HD [56,57]. We found that the loss of soma volume in R6/2 projection neurons [40,58] improved in R6/2:pGFAP-BDNF mice. Interestingly, we also detected an amelioration of striatal NII density in R6/2:pGFAP-BDNF mice which is in accordance with their improved cellular phenotype. Moreover, the density of glutamatergic cortical terminals (VGLUT1-positive) and dendritic spines (PSD-95-positive) in the striatum were both down-regulated in R6/2 striata, but preserved in R6/2:pGFAP-BDNF striata. These positive changes in synaptic markers and neuronal soma volume correlated with an absence of basal synaptic transmission dysfunction and a greater resistance to synaptic fatigue in R6/2:pGFAP-BDNF mice than in R6/2 mice. Intriguingly, pGFAP-BDNF mice showed a significant increase in synaptic fatigue relative to wt mice. This result suggests a subtle increase on BDNF levels in those mice which we were unable to detect. On the other hand, our results corroborate previous studies showing that BDNF application increases glutamatergic currents [59] and the number of dendritic spines [15,60]. However, cortico-striatal LTP alterations did not improve in R6/2:pGFAP-BDNF mice. Although striatal BDNF levels were higher in R6/2:pGFAP-BDNF mice than in R6/2, they did not reach wt levels. This result could explain the lack of improvement in LTP expression in cortico-striatal slices from R6/2:pGFAP-BDNF mice. In fact, Simmons and others showed that there was a reversal of the impairment in hippocampal LTP in knock-in mice when levels of BDNF were completely recovered [61]. These findings confirm the need for complete integrity in the BDNF pathway in order to regulate cortico-striatal LTP [62].

The finding that BDNF released from transgenic astrocytes in R6/2:pGFAP-BDNF mice improves and regulates their striatal-dependent behavior and synaptic transmission corroborates the recent hypothesis that astrocytes are involved in neuro-plasticity phenomena such as the regulation and strengthening of neurosignals and connectivity of neuronal networks, some of which are likely to be involved in cognitive functions at least [63-65]. Thus, it is important to highlight that our results suggest that astrocytes could modulate some of these plasticity processes *via* production and delivery of BDNF, probably into the excitatory synapse. This is possible since astrocytes are capable, though at low levels, of producing and releasing BDNF *per se* [23,66,67] and send very fine projections to the excitatory synapses in order to strengthen them [68]. Thus, our results suggest

a possible role for BDNF released from astrocytes in the regulation of basal synaptic transmission and synaptic fatigue in excitatory cortico-striatal synapses.

In the present study we wanted to administer BDNF non-invasively using a genetic approach. However, other non-invasive strategies have been reported to increase BDNF in the HD brains such as drug administration [61,69,70]. Thus, combination therapy with the pGFAP-BDNF construct and administration of a drug such as ampakines [69] could be crucial to fully recover the HD phenotype.

Finally, it should also be noted that other neurodegenerative diseases are associated with reactive astrogliosis and the consequent up-regulation of GFAP gene activity. Along these lines, pathologies such as Alzheimer's disease [71,72], Parkinson's disease [73,74] and amyotrophic lateral sclerosis [75] show increased astrogliosis reactivity with a consequent up-regulation of GFAP levels. It would be highly relevant to test whether the pGFAP-BDNF transgene might benefit these diseases. This statement takes relevance since BDNF delivery in a transgenic AD mouse model prevents loss of glutamatergic synapses and improves cognition [76]. In conclusion, conditional BDNF delivery regulated by the GFAP promoter in astrocytes could well be a therapeutic strategy for treatment of HD and other neurodegenerative diseases.

Methods

Animal handling and care

To obtain double-mutant animals (R6/2:pGFAP-BDNF mice) we cross-mated male R6/2 mice [29] with female pGFAP-BDNF mice [23]. Both R6/2 and pGFAP-BDNF mice were obtained from distinct colonies bred in a B6CBA background strain. Wild type (wt) littermate animals were used as the control group. Mice were housed together in numerical birth order in groups of mixed genotypes with access to food and water *ad libitum* in a colony room kept at constant temperature (19-22°C) and humidity (40-50%) on a 12 h light/dark cycle. All experiments used male mice, blind-coded for genotype. Data were recorded for analysis by microchip mouse number. All animal-related procedures were in accordance with the European Community guidelines for the care and use of laboratory animals (89/609//EEC) and were approved by the local animal care committee of the University of Barcelona (99/01) and Generalitat de Catalunya (99/1094).

Rotarod

Motor coordination and balance were evaluated on the rotarod apparatus at distinct rotations per minute (rpm), as described elsewhere [25]. In brief, animals at 5 weeks of age were trained at constant speed (24 rpm) for 60

sec. We performed two trials per day for three consecutive days; and the latency to fall and the number of falls during 60 sec was recorded. No differences between groups were detected at this period. After training and starting at 7 weeks of age, animals were evaluated once a week at 16 and 24 rpm, and the number of falls in a total of 60 sec was recorded. The animals were put on the rotarod several times until the addition of the latency to fall off reached a total of 60 sec. The curves representing the behavioral pattern were compared and the percentage of motor coordination function impairment was calculated as described elsewhere [77].

Balance beam

The motor coordination and balance of mice were also assessed by measuring their ability to traverse a narrow beam [25]. The beam consisted of a long steel cylinder (50 cm) with a 50 mm-square cross-section and a 15 mm-round diameter. The beam was placed horizontally, 50 cm above the bench surface, with each end mounted on a narrow support. Animals were allowed to walk for 1 min along the beam, while their latency to fall, number of falls and distance covered were measured.

Footprint

The footprint test was performed as described elsewhere [25]. Mice were trained to walk in a corridor that was 50 cm long and 7 cm wide. The forefeet and hindfeet of the mice were painted with non-toxic red and blue ink, respectively, and then given one run. The footprint pattern was analyzed for the number of steps on the white paper, the stride length was measured as the average distance of forward movement between each stride, and the forebase and hindbase widths were measured as the perpendicular distance between the left and right footprints of a given step.

Hanging wire

Neuromuscular abnormalities were analyzed by the hanging wire test as described elsewhere [25,78] in 7-, 10- and 12-week old wt, pGFAP-BDNF, R6/2 and R6/2: pGFAP-BDNF mice. A standard wire cage lid was used. To test balance and grip strength, mice were placed on top of a wire cage lid. The lid was shaken slightly to cause the mouse to grip the wires and then turned upside down for 60 sec. The number of falls of each mouse was recorded.

Clasping

Clasping was measured by suspending mice from their tails at least 1 foot above a surface for 1 min. A clasping event was defined by the retraction of either or both hindlimbs into the body and toward the midline. Mice

were scored according to the following criteria: 0 = no clasping, 1 = clasping 2 paws, and 2 = clasping all paws.

Plus Maze

To analyze mouse anxiety, we used the plus maze paradigm. The apparatus used was similar to that described elsewhere [30]. Briefly, the raised plus maze was made of wood and consisted of two opposing 30 × 8 cm open arms, and two opposing 30 × 8 cm arms enclosed by 15 cm-high walls. The maze was raised 50 cm from the floor and lit by dim light. Each mouse was placed in the central square of the raised plus maze, facing an open arm and its behavior was scored for 5 min. At the end of each trial, any defecation was removed and the apparatus was wiped with 70% alcohol. We recorded the time spent in the open arms, which normally correlates with low levels of anxiety. Animals were tracked and recorded with SMART junior software (Panlab, Spain).

Open Field

To check spontaneous locomotor activity we used the open field [79]. Briefly, the apparatus consisted of a white circular arena measuring 40 cm in diameter and 40 cm in height. Dimly light intensity was 40 lux throughout the arena. Animals were placed on the arena center and allowed to explore freely for 10 min. Spontaneous locomotor activity was measured. At the end of each trial, any defecation was removed and the apparatus was wiped with 70% ethanol. Animals were tracked and recorded with SMART junior software (Panlab, Spain).

Immunohistochemistry

Animals were deeply anesthetized with pentobarbital (60 mg/kg) and intracardially perfused with a 4% paraformaldehyde solution in 0.1 M sodium phosphate, pH 7.2. Brains were removed and post-fixed for 2 h in the same solution, cryoprotected with 30% sucrose in phosphate buffered saline (PBS) with 0.02% sodium azide and frozen in dry-ice cooled isopentane. Serial coronal sections (40 μm) obtained with a cryostat were processed for free-floating immunohistochemistry.

The sections were washed three times in PBS, permeabilized and blocked for 15 min by shaking at room temperature with PBS containing 0.3% Triton X-100 and 3% normal goat serum (Pierce Biotechnology). After three washes, brain slices were incubated overnight by shaking at 4°C with the corresponding primary antibodies in PBS with 0.02% sodium azide buffer. Antibodies used were the anti-PSD-95 1:500 (Affinity BioReagents) and anti-VGLUT1 1:500 (Synaptic systems). After primary antibody incubation, slices were washed three times and then incubated for 2 h shaking at room temperature

with subtype-specific fluorescent secondary antibodies: Cy3 goat anti-rabbit and Cy2 goat anti-mouse (both 1:100; from Jackson ImmunoResearch). No signal was detected in controls incubated in the absence of the primary antibody. Slight modifications were performed for VGLUT1 immunohistochemistry [80].

For diaminobenzidine immunohistochemistry experiments, endogenous peroxidases were blocked for 30-45 min in PBS containing 10% methanol and 3% H₂O₂. Then, non-specific protein interactions were blocked with normal serum or bovine serum albumin. Tissue was incubated overnight at 4°C with the following primary antibodies: anti-GFAP (1:500; Dako A/S), anti-calbindin (1:10000; Sigma Chemical Co.) and anti-EM48 (1:500; Chemicon). Sections were washed three times in PBS and incubated with a biotinylated secondary antibody (1:200; Pierce) at room temperature for 2 h. The immunohistochemical reaction was developed using the ABC kit (Pierce) and made visible with diaminobenzidine. No signal was detected in controls in which the primary antibodies had been omitted.

Stereology

Striatal volume estimations were performed as described elsewhere [25]. Unbiased blinded to genotype counting for genotype and condition was performed with Computer-Assisted Stereology Toolbox (CAST) software (Olympus Danmark A/S). For estimating mean cellular/perikaryal volumes of neurons (so-called local volumes) with design-based stereology, the “nucleator” method was used, as described elsewhere [81]. To determine neuronal and NIIs sub-population densities (neurons or NIIs per mm³) in the striatum, we used the dissector counting procedure in coronal sections spaced 240 µm apart, as described elsewhere [82].

Confocal analysis and dendritic spine-like structures counting

Fluorescently stained coronal sections were examined blinded to genotype by confocal microscopy, using a Leica TCS SL laser scanning confocal spectral microscope with argon and helium-neon lasers. Images were taken with a 63× numerical aperture lens with 4× digital zoom and standard (one Airy disc) pinhole. For each mouse, at least 4 slices of 40 µm containing striatal tissue were analyzed. Between 4 and 6 representative images, from dorsomedial striatum, were obtained from each slice. For each image, the entire three-dimensional stack of images was obtained by the use of the Z drive present in the Leica TCS SL microscope, and the size of the optical image was 0.5 µm, with a separation of 4 µm between each one. The number of dendritic spine-like structures was counted by the freeware NIH ImageJ version 1.33 by Wayne Rasband (National Institutes of

Health, Bethesda, MD), similar to described elsewhere [15].

Western blotting

Mice (12 weeks old; n: 7-14 per genotype) were deeply anesthetized in a CO₂ chamber. The brain was quickly removed and the striatum was dissected, frozen in dry ice and stored at -80°C until use. Briefly, striatal tissue was sonicated in 250 µl lysis buffer (PBS, 1% Nonidet P40, 0.1% SDS, 0.5% sodium deoxycholate, 1 mM PMSF, 10 mg/ml aprotinin, 1 mg/ml leupeptin, 2 mg/ml sodium orthovanadate) and centrifuged at 12,000 rpm for 20 min. Striatal proteins (15 µg) were analyzed by 7.5% SDS-polyacrylamide gel electrophoresis and transferred to Immobilon-P membranes (Millipore). Gel blots were blocked in TBS-T (150 mM NaCl, 20 mM Tris-HCl (pH 7.5), 0.05% Tween 20) with 5% non-fat dry milk and 5% bovine serum albumin. Immunoblots were probed with anti-DARPP-32 (1:1000; BD Biosciences), anti-GFAP (1:1000; Dako A/S), anti-BDNF (1:1000; Santa Cruz Biotechnology) and an affinity-purified rabbit anti-enkephalin applicable for Western blotting [35]. All blots were incubated with the primary antibody overnight at 4°C in PBS 0.02% sodium azide buffer with shaking. After several washes in TBS-T, blots were incubated with IgG HRP-conjugated anti-mouse or anti-rabbit antibodies (Promega Biotechnology) and developed using the ECL Western blotting analysis system (Santa Cruz). Incubation with a monoclonal anti-β-tubulin antibody (1:50,000; Sigma Chemical Co.) was used as a loading control.

Electrophysiology

Transverse brain slices (450 µm thickness) were prepared from mice, using conventional methods [83], and incubated for more than 1 h at room temperature (21-24°C) in artificial cerebrospinal fluid (aCSF). The aCSF contained (in mM) 124 NaCl, 2.69 KCl, 1.25 KH₂PO₄, 2 MgSO₄, 26 NaHCO₃, 2 CaCl₂ and 10 glucose, and was gassed with 95% O₂ and 5% CO₂. Brain slices containing both striatum and cortex were transferred to an immersion recording chamber and superfused (2.5 ml/min) with gassed aCSF warmed to 32-34°C. After 1 h of equilibration, extracellular field potentials were recorded in the dorsomedial striatum by a glass microelectrode filled with 1 M NaCl solution on stimulation of the white matter between the cortex and the striatum with a bipolar tungsten electrode *via* a 2100 isolated pulse stimulator (A-M Systems, Inc.). Long-term potentiation (LTP) was induced by applying four 1 s, 100 Hz trains delivered every 10 s, and potentiation was measured for 1 h after LTP induction at 0.1 Hz. For each experiment, population spike (PS) amplitude was expressed as a percentage of average pre-tetanus baseline amplitude

values. Synaptic fatigue was induced by a train at 100 Hz. Data were filtered (highpass, 0.1 Hz; lowpass 3 kHz) and digitized using a PowerLab 4/26 acquisition system (AD Instruments). The software Scope (AD Instruments) was used to display PS and measurements of the amplitude of PS.

Statistical analysis

All data are expressed as mean \pm s.e.m. Different statistical analyses were performed as appropriate and are indicated in the figure legends. Values of $p < 0.05$ were considered statistically significant.

Acknowledgements

We thank Maria Teresa Muñoz and Ana López for technical assistance, and Dra. Teresa Rodrigo Calduch and the staff of the animal care facility (Facultat de Psicologia, Universitat de Barcelona) for their help. We also thank Dra. Maria Calvo and Anna Bosch from the Confocal Microscopy Facility (Campus Casanova) of the Centre Científic i Tecnològic (Universitat de Barcelona) for their support and advice with all confocal techniques. We also thank Dr. Eric W. Dickson (Department of Emergency Medicine, University of Iowa) for kindly providing the anti-ENK antibody. The current work was supported by grants from Ministerio de Ciencia e Innovación (SAF2008-04360, J.A.; BFU2008-04196/BFI, E.D.M.), INCReCyT project (JCCM and PCYTA; E.D.M.), and Instituto de Salud Carlos III: CIBERNED and Red de Terapia Celular (RD06/0010/0006). A.G. and O.C. were a fellow of Ministerio de Ciencia e Innovación, Spain.

Author details

¹Departament de Biologia Cel·lular, Immunologia i Neurociències, Facultat de Medicina, Universitat de Barcelona, Barcelona, Spain. ²Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Casanova 143, Barcelona, Spain. ³Centro de Investigación Biomédica en Red sobre Enfermedades Neurodegenerativas (CIBERNED), Spain. ⁴Laboratory of Neurophysiology and Synaptic Plasticity, Albacete Science and Technology Park (PCYTA). Institute for Research in Neurological Disabilities (IDINE), University of Castilla-La Mancha, Albacete, Spain.

Authors' contributions

AG and OC carried out the behavioral tests. AG performed morphological and biochemical studies. CLP and EDM carried out the electrophysiological experiments. JA designed the experiments and wrote the paper with assistance from all other authors. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 19 July 2011 Accepted: 10 October 2011

Published: 10 October 2011

References

- Andrew SE, Goldberg YP, Kremer B, Telenius H, Theilmann J, Adam S, Starr E, Squitieri F, Lin B, Kalchman MA: **The relationship between trinucleotide (CAG) repeat length and clinical features of Huntington's disease.** *Nat Genet* 1993, **4**:398-403.
- Benitez J, Fernandez E, Garcia RP, Robledo M, Ayuso C, Garcia YJ: **[The gene responsible for Huntington's disease in Spanish families: its diagnostic value and the relation between trinucleotide expansion and the clinical characteristics].** *Rev Clin Esp* 1994, **194**:591-593.
- Reiner A, Albin RL, Anderson KD, D'Amato CJ, Penney JB, Young AB: **Differential loss of striatal projection neurons in Huntington disease.** *Proc Natl Acad Sci USA* 1988, **85**:5733-5737.
- Cepeda C, Hurst RS, Calvert CR, Hernandez-Echeagaray E, Nguyen OK, Jocoy E, Christian LJ, Ariano MA, Levine MS: **Transient and progressive electrophysiological alterations in the corticostriatal pathway in a mouse model of Huntington's disease.** *J Neurosci* 2003, **23**:961-969.
- Joshi PR, Wu NP, Andre VM, Cummings DM, Cepeda C, Joyce JA, Carroll JB, Leavitt BR, Hayden MR, Levine MS, Bamford NS: **Age-dependent alterations of corticostriatal activity in the YAC128 mouse model of Huntington disease.** *J Neurosci* 2009, **29**:2414-2427.
- Gavalda N, Perez-Navarro E, Gratacos E, Comella JX, Alberch J: **Differential involvement of phosphatidylinositol 3-kinase and p42/p44 mitogen activated protein kinase pathways in brain-derived neurotrophic factor-induced trophic effects on cultured striatal neurons.** *Mol Cell Neurosci* 2004, **25**:460-468.
- Baquet ZC, Gorski JA, Jones KR: **Early striatal dendrite deficits followed by neuron loss with advanced age in the absence of anterograde cortical brain-derived neurotrophic factor.** *J Neurosci* 2004, **24**:4250-4258.
- Strand AD, Baquet ZC, Aragaki AK, Holmans P, Yang L, Cleren C, Beal MF, Jones L, Kooperberg C, Olson JM, Jones KR: **Expression profiling of Huntington's disease models suggests that brain-derived neurotrophic factor depletion plays a major role in striatal degeneration.** *J Neurosci* 2007, **27**:11758-11768.
- Ferrer I, Goutan E, Marin C, Rey MJ, Ribalta T: **Brain-derived neurotrophic factor in Huntington disease.** *Brain Res* 2000, **866**:257-261.
- Zuccato C, Ciammola A, Rigamonti D, Leavitt BR, Goffredo D, Conti L, MacDonald ME, Friedlander RM, Silani V, Hayden MR, et al: **Loss of huntingtin-mediated BDNF gene transcription in Huntington's disease.** *Science* 2001, **293**:493-498.
- Diekmann H, Anichtchik O, Fleming A, Futter M, Goldsmith P, Roach A, Rubinsztein DC: **Decreased BDNF levels are a major contributor to the embryonic phenotype of huntingtin knockdown zebrafish.** *J Neurosci* 2009, **29**:1343-1349.
- Gharami K, Xie Y, An JJ, Tonegawa S, Xu B: **Brain-derived neurotrophic factor over-expression in the forebrain ameliorates Huntington's disease phenotypes in mice.** *J Neurochem* 2008, **2**:369-379.
- Gines S, Seong IS, Fossale E, Ivanova E, Trettel F, Gusella JF, Wheeler VC, Persichetti F, MacDonald ME: **Specific progressive cAMP reduction implicates energy deficit in presymptomatic Huntington's disease knock-in mice.** *Hum Mol Genet* 2003, **12**:497-508.
- Giralt A, Rodrigo T, Martin ED, Gonzalez JR, Mila M, Cena V, Dierssen M, Canals JM, Alberch J: **Brain-derived neurotrophic factor modulates the severity of cognitive alterations induced by mutant huntingtin: Involvement of phospholipase Cgamma activity and glutamate receptor expression.** *Neuroscience* 2009, **158**:1234-1250.
- Lynch G, Kramar EA, Rex CS, Jia Y, Chappas D, Gall CM, Simmons DA: **Brain-derived neurotrophic factor restores synaptic plasticity in a knock-in mouse model of Huntington's disease.** *J Neurosci* 2007, **27**:4424-4434.
- Simmons DA, Mehta RA, Lauterborn JC, Gall CM, Lynch G: **Brief ampakine treatments slow the progression of Huntington's disease phenotypes in R6/2 mice.** *Neurobiol Dis* 2011, **41**:436-444.
- Spies TL, Grote HE, Varshney NK, Cordery PM, van DA, Blakemore C, Hannan AJ: **Environmental enrichment rescues protein deficits in a mouse model of Huntington's disease, indicating a possible disease mechanism.** *J Neurosci* 2004, **24**:2270-2276.
- del Toro Daniel, Canals JM, Gines S, Kojima M, Egea G, Alberch J: **Mutant huntingtin impairs the post-Golgi trafficking of brain-derived neurotrophic factor but not its Val66Met polymorphism.** *J Neurosci* 2006, **26**:12748-12757.
- Gauthier LR, Charrin BC, Borrell-Pages M, Dompierre JP, Rangone H, Cordelieres FP, De MJ, MacDonald ME, Lessmann V, Humbert S, Saudou F: **Huntingtin controls neurotrophic support and survival of neurons by enhancing BDNF vesicular transport along microtubules.** *Cell* 2004, **118**:127-138.
- Alberch J, Perez-Navarro E, Canals JM: **Neurotrophic factors in Huntington's disease.** *Prog Brain Res* 2004, **146**:195-229.
- Cho SR, Benraiss A, Chmielnicki E, Samdani A, Economides A, Goldman SA: **Induction of neostriatal neurogenesis slows disease progression in a transgenic murine model of Huntington disease.** *J Clin Invest* 2007, **117**:2889-2902.
- Zuccato C, Cattaneo E: **Brain-derived neurotrophic factor in neurodegenerative diseases.** *Nat Rev Neurol* 2009, **5**:311-322.
- Giralt A, Friedman HC, Caneda-Ferron B, Urban N, Moreno E, Rubio N, Blanco J, Peterson A, Canals JM, Alberch J: **BDNF regulation under GFAP**

- promoter provides engineered astrocytes as a new approach for long-term protection in Huntington's disease. *Gene Ther* 2010, **17**:1294-1308.
24. Perez-Navarro E, Canudas AM, Akerund P, Alberch J, Arenas E: **Brain-derived neurotrophic factor, neurotrophin-3, and neurotrophin-4/5 prevent the death of striatal projection neurons in a rodent model of Huntington's disease.** *J Neurochem* 2000, **75**:2190-2199.
 25. Canals JM, Pineda JR, Torres-Peraza JF, Bosch M, Martin-Ibanez R, Munoz MT, Mengod G, Enfors P, Alberch J: **Brain-derived neurotrophic factor regulates the onset and severity of motor dysfunction associated with enkephalinergic neuronal degeneration in Huntington's disease.** *J Neurosci* 2004, **24**:7727-7739.
 26. Lindvall O, Bjorklund A: **First step towards cell therapy for Huntington's disease.** *Lancet* 2000, **356**:1945-1946.
 27. Lindvall O, Kokaia Z, Martinez-Serrano A: **Stem cell therapy for human neurodegenerative disorders-how to make it work.** *Nat Med* 2004, **10**(Suppl):S42-S50.
 28. Zuccato C, Cattaneo E: **Role of brain-derived neurotrophic factor in Huntington's disease.** *Prog Neurobiol* 2007, **81**:294-330.
 29. Mangiarini L, Sathasivam K, Seller M, Cozens B, Harper A, Hetherington C, Lawton M, Trotter Y, Lehrach H, Davies SW, Bates GP: **Exon 1 of the HD gene with an expanded CAG repeat is sufficient to cause a progressive neurological phenotype in transgenic mice.** *Cell* 1996, **87**:493-506.
 30. File SE, Mahal A, Mangiarini L, Bates GP: **Striking changes in anxiety in Huntington's disease transgenic mice.** *Brain Res* 1998, **805**:234-240.
 31. Luesse HG, Schiefer J, Spruenken A, Puls C, Block F, Kosinski CM: **Evaluation of R6/2 HD transgenic mice for therapeutic studies in Huntington's disease: behavioral testing and impact of diabetes mellitus.** *Behav Brain Res* 2001, **126**:185-195.
 32. Menalled LB: **Knock-in mouse models of Huntington's disease.** *NeuroRx* 2005, **2**:465-470.
 33. Albin RL, Reiner A, Anderson KD, Dure LS, Handelin B, Balfour R, Whetsell WO, Penney JB, Young AB: **Preferential loss of striato-external pallidal projection neurons in presymptomatic Huntington's disease.** *Ann Neurol* 1992, **31**:425-430.
 34. Xie Y, Hayden MR, Xu B: **BDNF overexpression in the forebrain rescues Huntington's disease phenotypes in YAC128 mice.** *J Neurosci* 2010, **30**:14708-14718.
 35. Denning GM, Ackermann LW, Barna TJ, Armstrong JG, Stoll LL, Weintraub NL, Dickson EW: **Proenkephalin expression and enkephalin release are widely observed in non-neuronal tissues.** *Peptides* 2008, **29**:83-92.
 36. Watson JD, Varley JG, Tomlin SJ, Medbak S, Rees LH, Hinds CJ: **Biochemical characterization of circulating Met-enkephalins in canine endotoxin shock.** *J Endocrinol* 1986, **111**:329-334.
 37. Flaggmeyer I, Haas HL, Stevens DR: **Adenosine A1 receptor-mediated depression of corticostriatal and thalamostriatal glutamatergic synaptic potentials in vitro.** *Brain Res* 1997, **778**:178-185.
 38. Schotanus SM, Fredholm BB, Chergui K: **NMDA depresses glutamatergic synaptic transmission in the striatum through the activation of adenosine A1 receptors: evidence from knockout mice.** *Neuropharmacology* 2006, **51**:272-282.
 39. Yin HH, Park BS, Adermark L, Lovinger DM: **Ethanol reverses the direction of long-term synaptic plasticity in the dorsomedial striatum.** *Eur J Neurosci* 2007, **25**:3226-3232.
 40. Klapstein GJ, Fisher RS, Zanjani H, Cepeda C, Jokel ES, Chesselet MF, Levine MS: **Electrophysiological and morphological changes in striatal spiny neurons in R6/2 Huntington's disease transgenic mice.** *J Neurophysiol* 2001, **86**:2667-2677.
 41. Laforet GA, Sapp E, Chase K, McIntyre C, Boyce FM, Campbell M, Cadigan BA, Warzecki L, Tagle DA, Reddy PH, et al: **Changes in cortical and striatal neurons predict behavioral and electrophysiological abnormalities in a transgenic murine model of Huntington's disease.** *J Neurosci* 2001, **21**:9112-9123.
 42. Nagahara AH, Tuszynski MH: **Potential therapeutic uses of BDNF in neurological and psychiatric disorders.** *Nat Rev Drug Discov* 2011, **10**:209-219.
 43. Zhang Y, Pardridge WM: **Neuroprotection in transient focal brain ischemia after delayed intravenous administration of brain-derived neurotrophic factor conjugated to a blood-brain barrier drug targeting system.** *Stroke* 2001, **32**:1378-1384.
 44. Zhang Y, Pardridge WM: **Blood-brain barrier targeting of BDNF improves motor function in rats with middle cerebral artery occlusion.** *Brain Res* 2006, **1111**:227-229.
 45. Hoffman R, Tong J, Brandt J, Traycoff C, Bruno E, McGuire BW, Gordon MS, McNiece I, Srour EF: **The in vitro and in vivo effects of stem cell factor on human hematopoiesis.** *Stem Cells* 1993, **11**(Suppl 2):76-82.
 46. Kusakabe M, Mangiarini L, Laywell ED, Bates GP, Yoshiki A, Hiraiwa N, Inoue J, Steindler DA: **Loss of cortical and thalamic neuronal tenascin-C expression in a transgenic mouse expressing exon 1 of the human Huntington disease gene.** *J Comp Neurol* 2001, **430**:485-500.
 47. Stack EC, Kubilus JK, Smith K, Cormier K, Del Signore SJ, Guelin E, Ryu H, Hersch SM, Ferrante RJ: **Chronology of behavioral symptoms and neuropathological sequela in R6/2 Huntington's disease transgenic mice.** *J Comp Neurol* 2005, **490**:354-370.
 48. Yu ZX, Li SH, Evans J, Pillarisetti A, Li H, Li XJ: **Mutant huntingtin causes context-dependent neurodegeneration in mice with Huntington's disease.** *J Neurosci* 2003, **23**:2193-2202.
 49. Faideau M, Kim J, Cormier K, Gilmore R, Welch M, Auregan G, Dufour N, Guillemier M, Brouillet E, Hantraye P, et al: **In vivo expression of polyglutamine-expanded huntingtin by mouse striatal astrocytes impairs glutamate transport: a correlation with Huntington's disease subjects.** *Hum Mol Genet* 2010, **19**:3053-3067.
 50. Croll SD, Suri C, Compton DL, Simmons MV, Yancopoulos GD, Lindsay RM, Wiegand SJ, Rudge JS, Scharfman HE: **Brain-derived neurotrophic factor transgenic mice exhibit passive avoidance deficits, increased seizure severity and in vitro hyperexcitability in the hippocampus and entorhinal cortex.** *Neuroscience* 1999, **93**:1491-1506.
 51. Kells AP, Henry RA, Connor B: **AAV-BDNF mediated attenuation of quinolinic acid-induced neuropathology and motor function impairment.** *Gene Ther* 2008, **15**:966-977.
 52. Fritsch B, Reis J, Martinowich K, Schambra HM, Ji Y, Cohen LG, Lu B: **Direct current stimulation promotes BDNF-dependent synaptic plasticity: potential implications for motor learning.** *Neuron* 2010, **66**:198-204.
 53. Castren E: **Neurotrophins as mediators of drug effects on mood, addiction, and neuroprotection.** *Mol Neurobiol* 2004, **29**:289-302.
 54. Pardon MC: **Role of neurotrophic factors in behavioral processes: implications for the treatment of psychiatric and neurodegenerative disorders.** *Vitam Horm* 2010, **82**:185-200.
 55. Canteras NS, Resstel LB, Bertoglio LJ, Carobrez AP, Guimaraes FS: **Neuroanatomy of anxiety.** *Curr Top Behav Neurosci* 2010, **2**:77-96.
 56. Cepeda C, Wu N, Andre VM, Cummings DM, Levine MS: **The corticostriatal pathway in Huntington's disease.** *Prog Neurobiol* 2007, **81**:253-271.
 57. Milnerwood AJ, Raymond LA: **Early synaptic pathophysiology in neurodegeneration: insights from Huntington's disease.** *Trends Neurosci* 2010, **33**:513-523.
 58. Levine MS, Klapstein GJ, Koppel A, Gruen E, Cepeda C, Vargas ME, Jokel ES, Carpenter EM, Zanjani H, Hurst RS, et al: **Enhanced sensitivity to N-methyl-D-aspartate receptor activation in transgenic and knockin mouse models of Huntington's disease.** *J Neurosci Res* 1999, **58**:515-532.
 59. Li YX, Zhang Y, Lester HA, Schuman EM, Davidson N: **Enhancement of neurotransmitter release induced by brain-derived neurotrophic factor in cultured hippocampal neurons.** *J Neurosci* 1998, **18**:10231-10240.
 60. Rex CS, Lin CY, Kramar EA, Chen LY, Gall CM, Lynch G: **Brain-derived neurotrophic factor promotes long-term potentiation-related cytoskeletal changes in adult hippocampus.** *J Neurosci* 2007, **27**:3017-3029.
 61. Simmons DA, Rex CS, Palmer L, Pandeyarajan V, Fedulov V, Gall CM, Lynch G: **Up-regulating BDNF with an ampakine rescues synaptic plasticity and memory in Huntington's disease knockin mice.** *Proc Natl Acad Sci USA* 2009, **106**:4906-4911.
 62. Jia Y, Gall CM, Lynch G: **Presynaptic BDNF promotes postsynaptic long-term potentiation in the dorsal striatum.** *J Neurosci* 2010, **30**:14440-14445.
 63. Haydon PG: **GLIA: listening and talking to the synapse.** *Nat Rev Neurosci* 2001, **2**:185-193.
 64. Santello M, Volterra A: **Neuroscience: Astrocytes as aide-memoires.** *Nature* 2010, **463**:169-170.
 65. Suzuki A, Stern SA, Bozdagi O, Huntley GW, Walker RH, Magistretti PJ, Alberini CM: **Astrocyte-neuron lactate transport is required for long-term memory formation.** *Cell* 2011, **144**:810-823.

66. Dougherty KD, Dreyfus CF, Black IB: **Brain-derived neurotrophic factor in astrocytes, oligodendrocytes, and microglia/macrophages after spinal cord injury.** *Neurobiol Dis* 2000, **7**:574-585.
67. Riley CP, Cope TC, Buck CR: **CNS neurotrophins are biologically active and expressed by multiple cell types.** *J Mol Histol* 2004, **35**:771-783.
68. Fellin T: **Communication between neurons and astrocytes: relevance to the modulation of synaptic and network activity.** *J Neurochem* 2009, **108**:533-544.
69. Simmons DA, Mehta RA, Lauterborn JC, Gall CM, Lynch G: **Brief ampakine treatments slow the progression of Huntington's disease phenotypes in R6/2 mice.** *Neurobiol Dis* 2011, **41**:436-444.
70. Squitieri F, Orobello S, Cannella M, Martino T, Romanelli P, Giovacchini G, Frati L, Mansi L, Ciarmiello A: **Riluzole protects Huntington disease patients from brain glucose hypometabolism and grey matter volume loss and increases production of neurotrophins.** *Eur J Nucl Med Mol Imaging* 2009, **36**:1113-1120.
71. Ingelsson M, Fukumoto H, Newell KL, Growdon JH, Hedley-Whyte ET, Frosch MP, Albert MS, Hyman BT, Irizarry MC: **Early Abeta accumulation and progressive synaptic loss, gliosis, and tangle formation in AD brain.** *Neurology* 2004, **62**:925-931.
72. Verkhratsky A, Olabarria M, Noristani HN, Yeh CY, Rodriguez JJ: **Astrocytes in Alzheimer's disease.** *Neurotherapeutics* 2010, **7**:399-412.
73. Braak H, Sastre M, Del TK: **Development of alpha-synuclein immunoreactive astrocytes in the forebrain parallels stages of intraneuronal pathology in sporadic Parkinson's disease.** *Acta Neuropathol* 2007, **114**:231-241.
74. Terada S, Kuroda S, Ishizu H: **[Parkinson's disease].** *Ryoikibetsu Shokogun Shirizu* 2003, **257**:260.
75. Vargas MR, Johnson JA: **Astroglialosis in amyotrophic lateral sclerosis: role and therapeutic potential of astrocytes.** *Neurotherapeutics* 2010, **7**:471-481.
76. Blurton-Jones M, Kitazawa M, Martinez-Coria H, Castello NA, Muller FJ, Loring JF, Yamasaki TR, Poon WW, Green KN, LaFerla FM: **Neural stem cells improve cognition via BDNF in a transgenic model of Alzheimer disease.** *Proc Natl Acad Sci USA* 2009, **106**:13594-13599.
77. Ferrante RJ, Andreassen OA, Dedeoglu A, Ferrante KL, Jenkins BG, Hersch SM, Beal MF: **Therapeutic effects of coenzyme Q10 and remacemide in transgenic mouse models of Huntington's disease.** *J Neurosci* 2002, **22**:1592-1599.
78. Gimenez-Llort L, Fernandez-Teruel A, Escorihuela RM, Fredholm BB, Tobena A, Pekny M, Johansson B: **Mice lacking the adenosine A1 receptor are anxious and aggressive, but are normal learners with reduced muscle strength and survival rate.** *Eur J Neurosci* 2002, **16**:547-550.
79. Glynn D, Bortnick RA, Morton AJ: **Complexin II is essential for normal neurological function in mice.** *Hum Mol Genet* 2003, **12**:2431-2448.
80. Martin-Ibanez R, Jenstad M, Berghuis P, Edwards RH, Hioki H, Kaneko T, Mulder J, Canals JM, Ernfors P, Chaudhry FA, Harkany T: **Vesicular glutamate transporter 3 (VGLUT3) identifies spatially segregated excitatory terminals in the rat substantia nigra.** *Eur J Neurosci* 2006, **23**:1063-1070.
81. Gundersen HJ: **The nucleator.** *J Microsc* 1988, **151**:3-21.
82. Torres-Peraza J, Pezzi S, Canals JM, Gavalda N, Garcia-Martinez JM, Perez-Navarro E, Alberch J: **Mice heterozygous for neurotrophin-3 display enhanced vulnerability to excitotoxicity in the striatum through increased expression of N-methyl-D-aspartate receptors.** *Neuroscience* 2007, **144**:462-471.
83. Martin ED, Buno W: **Stabilizing effects of extracellular ATP on synaptic efficacy and plasticity in hippocampal pyramidal neurons.** *Eur J Neurosci* 2005, **21**:936-944.

doi:10.1186/1750-1326-6-71

Cite this article as: Giralt *et al.*: Conditional BDNF release under pathological conditions improves Huntington's disease pathology by delaying neuronal dysfunction. *Molecular Neurodegeneration* 2011 **6**:71.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

