

Research article

Open Access

Mutations on the Switch III region and the $\alpha 3$ helix of $G\alpha_{16}$ differentially affect receptor coupling and regulation of downstream effectors

May YM Yu, Maurice KC Ho, Andrew MF Liu and Yung H Wong*

Address: Department of Biochemistry, Molecular Neuroscience Center and Biotechnology Research Institute, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong, PR China

Email: May YM Yu - mayyu@ust.hk; Maurice KC Ho - bokcmho@ust.hk; Andrew MF Liu - andyliu@ust.hk; Yung H Wong* - boyung@ust.hk

* Corresponding author

Published: 22 November 2008

Received: 13 August 2008

Journal of Molecular Signaling 2008, 3:17 doi:10.1186/1750-2187-3-17

Accepted: 22 November 2008

This article is available from: <http://www.jmolecularsignaling.com/content/3/1/17>

© 2008 Yu et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: $G\alpha_{16}$ can activate phospholipase C β (PLC β) directly like $G\alpha_q$. It also couples to tetrapeptide repeat I (TPRI) which is linked to Ras activation. It is unknown whether PLC β and TPRI interact with the same regions on $G\alpha_{16}$. Previous studies on $G\alpha_q$ have defined two minimal clusters of amino acids that are essential for the coupling to PLC β . Cognate residues in $G\alpha_{16}$ might also be essential for interacting with PLC β , and possibly contribute to TPRI interaction and other signaling events.

Results: Alanine mutations were introduced to the two amino acid clusters (246–248 and 259–260) in the switch III region and $\alpha 3$ helix of $G\alpha_{16}$. Regulations of PLC β and STAT3 were partially weakened by each cluster mutant. A mutant harboring mutations at both clusters generally produced stronger suppressions. Activation of Jun N-terminal kinase (JNK) by $G\alpha_{16}$ was completely abolished by mutating either clusters. Contrastingly, phosphorylations of extracellular signal-regulated kinase (ERK) and nuclear factor κ B (NF- κ B) were not significantly affected by these mutations. The interactions between the mutants and PLC $\beta 2$ and TPRI were also reduced in co-immunoprecipitation assays. Coupling between G_{16} and different categories of receptors was impaired by the mutations, with the effect of switch III mutations being more pronounced than those in the $\alpha 3$ helix. Mutations of both clusters almost completely abolished the receptor coupling and prevent receptor-induced $G\beta\gamma$ release.

Conclusion: The integrity of the switch III region and $\alpha 3$ helix of $G\alpha_{16}$ is critical for the activation of PLC β , STAT3, and JNK but not ERK or NF- κ B. Binding of $G\alpha_{16}$ to PLC $\beta 2$ or TPRI was reduced by the mutations of either cluster. The same region could also differentially affect the effectiveness of receptor coupling to G_{16} . The studied region was shown to bear multiple functionally important roles of G_{16} .

Background

As the major group of cell-surface detectors for hormones and neurotransmitters, G protein-coupled receptors

(GPCRs) employ a variety of signal transduction pathways to regulate cellular functions. One of the primary signaling routes initiated upon activation of GPCRs is

through the stimulation of PLC β by members of the G α_q subfamily. PLC β activity can in turn regulate many downstream kinases and transcription factors, thereby modulating cellular processes such as growth and differentiation. The interactions between PLC β and G α_q subfamily members have been examined by mutagenesis studies. Alanine scanning mutagenesis of G α_q has identified a stretch of amino acids (Ile²¹⁷-Lys²⁷⁶) that may be responsible for PLC β interaction. Within this region, two groups of amino acids (Asp²⁴³, Asn²⁴⁴, Glu²⁴⁵ and Arg²⁵⁶, Thr²⁵⁷; Figure 1A and 1B) have been suggested to be crucial for PLC β interaction [1]. These two clusters of amino acids are located in the $\alpha 3$ helix and $\beta 4$ - $\alpha 3$ loop (Figure 1A) which exhibits dramatic conformational changes during G protein activation [2,3].

G α_{16} is a member of G α_q subfamily which can activate PLC β [4], and its unique promiscuity for GPCRs [4] highlights its importance in cellular signaling, especially in hematopoietic cells where it is restrictively expressed [5]. Recent studies have revealed that G α_{16} possesses additional signaling properties which may be independent of PLC β activity. It has been demonstrated early on that interleukin-2 and interleukin-8 induce G α_{16} -mediated activation of ERK [6]. The use of a constitutively active mutant of G α_{16} (G α_{16} QL) confirmed that it can indeed stimulate the activities of ERK [7] and JNK [8,9] in various cell types. Presumably these stimulatory signals proceed via PLC β which triggers the cleavage of phosphatidylinositol bisphosphate to form IP₃ and DAG, and the latter can modulate numerous signaling cascades through the activation of protein kinase C (PKC). The ability of G α_{16} QL to activate transcription factors such as STAT3 [7,10] and NF- κ B [11,12] also requires PLC β activity. The discovery of a novel binding partner of G α_{16} , tetratricopeptide repeat 1 (TPR1) [13] opens up new possibilities for the regulation of ERK and its downstream effectors. Since TPR1 prefers to bind to active Ras, its association with G α_{16} may facilitate signaling along the Ras/Raf-1/MEK/ERK axis. However, no study has yet addressed the relative contributions of the PLC β and TPR1 on the G α_{16} -mediated signaling events.

Although G α_{16} shares only 55% identity to G α_q in the amino acid sequence, the two clusters of putative PLC β -interacting residues can be found in the homologous regions of G α_{16} . Cluster 1 includes Asn²⁴⁶, Gln²⁴⁷ and Glu²⁴⁸, while cluster 2 consists of Gly²⁵⁹ and Thr²⁶⁰. It is noteworthy that both the glutamate and threonine at the end of each cluster are highly conserved among the G α_q subfamily members (Figure 1A). Mutations of these residues may impair PLC β activation, and affect other downstream effectors which are dependent on PLC β activity. Signals channeled through the TPR1/Ras route are unlikely to be affected by such mutations unless TPR1 and

PLC β interact with similar regions on G α_{16} . To date, the TPR1-interacting domain(s) of G α_{16} has not been defined. Indeed, there are precedents for multifunctional domains in G α subunits. The receptor- and effector-interacting regions (e.g., $\alpha 2$ - $\beta 4$, $\alpha 3$ - $\beta 5$, and $\alpha 4$ - $\beta 6$ loops) in G α_s and G α_i subunits have substantial overlaps ([14-20], also see Figure 1B). Therefore, it would be necessary to examine the possible impacts of these five residues on the ability of G α_{16} to interact with different molecular partners. Furthermore, both the switch III region and $\alpha 3$ helix show substantial spatial rearrangements during the activation of a G α subunit, mutations in these two regions may also perturb the receptor-mediated G protein activation.

In this study, the functional impacts of the two clusters of amino acids were investigated for receptor-dependent and independent regulation of different effectors of G α_{16} . The results suggested that the two regions of interest were important for the activation of PLC β , STAT3, JNK but not ERK or NF- κ B. Besides, interactions of G α_{16} with TPR1 and PLC β were also reduced in a similar fashion. The same regions could also differentially affect the effectiveness of receptor coupling to G α_{16} .

Results

Design and expression of the G α_{16} mutants

A previous study has shown that the residues lying on the switch III region and the $\alpha 3$ helix of G α_q are required for interaction with PLC β [1]. The corresponding two clusters of amino acids on G α_{16} are Asn²⁴⁵-Glu²⁴⁷ and Gly²⁵⁹-Thr²⁶⁰ (Figure 1A). Mutations of these two clusters of residues into alanine created 3 mutants (Figure 2A) denoted as 3A (Asn²⁴⁵-Glu²⁴⁷ \rightarrow Ala), 2A (Gly²⁵⁹-Thr²⁶⁰ \rightarrow Ala) and 5A (all 5 residues \rightarrow Ala). The Gln²¹² \rightarrow Leu (QL) mutation [8,21] was also introduced to individual alanine mutants to generate a constitutively active phenotype for studying effector interactions in a receptor-independent fashion. Previous studies have already confirmed the capability of G α_{16} QL to activate various downstream effectors such as ERK, NF- κ B, STAT3 and JNK [7,12].

Since COS-7 cells do not express G α_{16} endogenously, they are an excellent platform for studying G α_{16} -regulated signaling pathways by over-expressing G α_{16} or its mutants. A tailor-designed antiserum [22] which recognizes the N-terminus (amino acid residues 13-27) of G α_{16} was used for detecting the expression level of G α_{16} and its mutants. Upon expression in COS-7 cells, a single band at around 43 kDa was detected by the anti-G α_{16} antiserum for each construct (Figure 2B, lower part). Plasma membrane proteins prepared from non-transfected COS-7 cells served as a negative control (Figure 2B, leftmost lane). All of the mutants were expressed in COS-7 cells in a comparable level as G α_{16} and G α_{16} QL, indicating that the mutations

A

<<<< SW3 >>>

\\ \\ \alpha3 \\ \\

$G\alpha_q$	(235-261)	YDQVLVESDNENRMEE SKALFRTIITY
$G\alpha_{16}$	(238-264)	YDQCLEENNQENRMKE SLALFGTILEL
$G\alpha_{i1}$	(230-256)	YDLVLAEDEEMNRMHE SMKLFDS ICNN
$G\alpha_z$	(231-257)	YDLKLYEDNQT SRMAE SLRLFDS ICNN
$G\alpha_{t1}$	(226-252)	YDMVLVEDDEVNRMHE SLHLFNS ICNH
$G\alpha_{s1}$	(253-279)	YNMVIREDNQTNRLQEALNLFKSIWNN
$G\alpha_{12}$	(255-281)	YDQVLMEDRRTNRLVESMNI FETIVNN

* : : * . * : * : : * : *




Figure 1
Sequence alignment and molecular model of $G\alpha_{16}$. (A) The sequences corresponding to the switch III region and $\alpha 3$ helix of various $G\alpha$'s were aligned. The consensus sequences are indicated as asterisks, colons and dots for strictly conserved, closely related and barely related residues among the candidates. The regions corresponding to the two clusters of putative PLC-interacting residues of $G\alpha_q$ are highlighted in orange. (B) A stereogram of the constructed molecular model of $G\alpha_{16}$ is shown. Portions of the molecular surface were colored as blue, grey and cyan for the regions interacting with receptor, effector, or both, respectively, based on the studies of different G proteins. The side chains of the residues studied here are shown in spheres as indicated (except for Gly²⁵⁹ which is devoid of any side chain).




Figure 2
Effects of 2A, 3A and 5A mutations on Gα₁₆-mediated PLCβ activation. (A) Positions of the alanine mutations on the corresponding Gα₁₆ mutants were shown as an alignment with the Gα₁₆ sequence. Identical residues were simplified with dots. (B) *Top*: COS-7 cells were transiently transfected with 0.25 μg/ml cDNAs encoding the wild type or QL mutants of Gα₁₆, 2A, 3A and 5A. Transfectants were labeled with [³H]myo-inositol and assayed for IP accumulation. * IP accumulation stimulated by constitutively active mutants was significantly higher than that obtained with their wild type counterparts; ‡ Constitutive activity was significantly lower than that obtained with Gα₁₆QL; Tukey-Kramer's test, p < 0.05. *Bottom*: Transfected COS-7 cells were harvested and membrane proteins were extracted for immunodetection. A Gα₁₆-specific custom antiserum was used for recognition of Gα₁₆ and its mutants. Fluorographs were visualized with the ECL chemiluminescence detection kit. Untransfected COS-7 cells served as the negative control. Two separate sets of transfected cells yielded similar results. (C) COS-7 cells were transfected with increasing amounts of cDNA encoding Gα₁₆QL, 2A-QL, 3A-QL or 5A-QL. Empty vector pcDNA3 was added to balance the amount of cDNA used in the transfection for each sample. Gα₁₆-transfected cells served as the negative control (hollow square). *Top*: IP production increased dose-dependently with increasing expression levels of the constitutively active form of alanine mutants and Gα₁₆. *Bottom*: Expression level of constitutively active counterparts of Gα₁₆ and its mutants were determined by Western blotting.

introduced did not affect the apparent expression or stability of Gα₁₆.

Impairment of PLCβ Regulation

The same sets of transfected cells examined in expression study were tested for their ability to stimulate PLCβ. Expression of Gα₁₆QL in COS-7 cells significantly increased IP formation by ~7 fold as compared to cells expressing wild-type (WT) Gα₁₆. Expression of constitu-

tively active counterparts of the mutants (2A-QL, 3A-QL and 5A-QL) in COS-7 cells significantly increased IP accumulation as compared to the responses of their corresponding non-QL counterparts. Responses induced by QL forms of the mutants were significantly reduced as compared with the response elicited by Gα₁₆QL (Figure 2B). 2A-QL and 3A-QL retained 42% and 73% of the IP production of Gα₁₆QL, respectively, whereas 5A-QL preserved about 61% of the IP accumulation of Gα₁₆QL.

However, the differences between the mutants were not statistically significant. Apparently, the results indicated that the mutations partially impaired the ability of $G\alpha_{16}$ to activate PLC β .

Further experiments were carried out to verify the observed variations of the induced IP formation between different constitutively active mutants. Increasing amounts of individual QL mutant cDNA were transfected into COS-7 cells and the IP accumulation in the transfected cells were measured. For all of the QL mutants, increasing levels of IP accumulation were observed as the cDNA amounts were increased (Figure 2C, upper part). The expression levels of individual QL mutants were correspondingly increased as examined by Western blotting (Figure 2C, lower part). There were obvious differences in the PLC β responses induced by the different mutants. $G\alpha_{16}$ QL and 2A-QL induced maximal stimulation of IP production at ~ 0.1 $\mu\text{g/ml}$, while maximal responses for 3A-QL and 5A-QL were only obtained at cDNA concentrations of 0.5 $\mu\text{g/ml}$ or higher. Despite similar levels of expression, the maximal response stimulated by 2A-QL was only about half of that elicited by $G\alpha_{16}$ QL. The estimated EC_{50} values of IP accumulation were 0.020, 0.039, 0.123 and 0.610 $\mu\text{g/ml}$ for $G\alpha_{16}$ QL, 2A-QL, 3A-QL and 5A-QL, respectively. These results indicate a progressive decrease of the efficiencies for activating PLC β when one or more of the putative PLC β -interacting domains on $G\alpha_{16}$ were mutated to alanine. Mutations at the $\alpha 3$ helix (2A) might be particularly detrimental to PLC β activity, because the maximal stimulatory response of 2A-QL was always lower than those of $G\alpha_{16}$ QL and the other two alanine mutants. Since 5A-QL could attain a higher maximal PLC β response than 2A-QL, the incorporation of 3A mutations in the switch III region apparently relieved the functional impairment associated with the 2A mutations in the $\alpha 3$ helix of $G\alpha_{16}$.

Differential Regulations of Downstream Effectors

Diverse downstream effectors have been found to be regulated by $G\alpha_{16}$, including ERK, STAT3, NF- κ B and JNK [7,9,11,23], but their dependencies on PLC β activation have not been well defined. To further investigate the functions of the selected amino acids in regulating these downstream effectors, WT and QL forms of $G\alpha_{16}$ and the three mutants were examined for their regulations of these downstream effectors when expressed in HEK293 cells. We employed HEK293 cells in this part of the study because of the regulations of ERK, STAT3, and NF- κ B had been fully characterized in this cellular background [7,11] and all of the mutants could be efficiently expressed in these cells (data not shown). As illustrated in Figure 3A, no observable alteration in the total ERK1/2 levels was found in cells transfected with different cDNA constructs. The induction of ERK phosphorylation by the QL forms of

the alanine mutants was significantly higher than their corresponding WT controls, and they were all similar to $G\alpha_{16}$ QL. It should be noted that 2A-QL and 5A-QL consistently generated a slightly lower level of ERK phosphorylation than 3A-QL and $G\alpha_{16}$ QL.

As both PLC β and ERK can serve as upstream regulators of STAT3 [7], we examined the ability of the mutants to induce STAT3 Tyr⁷⁰⁵ phosphorylation. All three mutants were capable of stimulating STAT3 phosphorylation at Tyr⁷⁰⁵ albeit weaker than that induced by $G\alpha_{16}$ QL (Figure 3B). The phosphorylation triggered by 2A-QL, 3A-QL and 5A-QL were not significantly different between each other. Given that $G\alpha_{16}$ QL-induced STAT3 phosphorylation involves PLC β and PKC [7], these results were not surprising because the alanine mutants have impaired PLC β regulation (Figure 2).

Regulation of NF- κ B by $G\alpha_{16}$ has been demonstrated in recombinant systems as well as in human lymphoblastoma REH cells [11]. The phosphorylation of NF- κ B induced by constitutively active alanine mutants was examined. As compared to $G\alpha_{16}$, the expression levels of total NF- κ B were unaffected in cells transfected with the alanine mutants (Figure 3C). The levels of NF- κ B phosphorylation caused by the constitutively active alanine mutants were only slightly reduced (but not statistically different) when compared to the $G\alpha_{16}$ QL responses. There was also no obvious difference between 2A, 3A and 5A mutants.

Since the background phosphorylation of JNK in transfected HEK293 cells are relatively high (Chan AS and Wong YH, unpublished observation), *in vitro* kinase assay was adopted to determine the activity of JNK activity in COS-7 cells instead. COS-7 cells were co-transfected with $G\alpha_{16}$ mutants together with HA-tagged JNK. The expression level of HA-tagged JNK were similar in all transfected cells (Figure 3D). $G\alpha_{16}$ QL enhanced JNK-mediated [³²P]-labeling of GST-c-Jun significantly by 3.6-fold as compared to that of $G\alpha_{16}$. Surprisingly, none of the constitutively active alanine mutants was capable of inducing JNK activation (Figure 3D). It appeared that the sequence integrity on the switch III region and $\alpha 3$ helix is more critical for the regulation of JNK activation.

Differential Associations with PLC β and TPRI

As the alanine mutations of the two clusters caused reduction of PLC β activation, one of the possible explanations would be the impairment of cognate recognition of PLC β . Co-immunoprecipitation experiments were performed to study the possible alteration of the mutants and PLC $\beta 2$, a PLC β isoform commonly expressed and known to interact with different G_q family members (Figure 4A). Expression levels of different $G\alpha_{16}$ constructs were similar when co-




Figure 3

Role of $G\alpha_{16}$ mutants in STAT3, ERK1/2, NF- κ B and c-Jun phosphorylation. HEK293 cells were transfected with pcDNA3, wild type or QL mutants of $G\alpha_{16}$, 2A, 3A or 5A. The transfectants were deprived of serum overnight, and cell lysates were prepared for SDS-PAGE separation. Phosphorylated form or native ERK1/2 (A), STAT3 (B), and NF- κ B (C) were detected by Western blotting as indicated. (D) For measuring $G\alpha_{16}$ -triggered JNK activity, COS-7 cells were transiently transfected with each of the constructs mentioned together with a plasmid encoding HA-tagged JNK. Serum starvation was performed as mentioned and JNK assay was performed as described in Methods. Expression of tagged JNK was determined by anti-HA antibody. The activation of JNK was monitored by detecting the phosphorylation of GST-fused-cjun. The fold induction of the phosphorylation of various effectors were quantified and plotted on the right hand side for comparisons. * QL counterparts of the mutants stimulate phosphorylation of the detected proteins significantly over cells expressing wild type complements (Tukey-Kramer's test, $p < 0.05$). ‡ QL counterparts of 2A-, 3A- and 5A-stimulated phosphorylation were significantly lower than that of $G\alpha_{16}$ QL (Tukey-Kramer's test, $p < 0.05$).




Figure 4

The alanine mutants co-immunoprecipitate with TPR1 and PLCβ₂. (A) HEK293 cells were co-transfected with PLCβ₂ and the Gα constructs as indicated at the bottom of the blots. Total cell lysates (TCL) from each condition were subjected to immunoprecipitation using either anti-PLCβ₂ or anti-Gα₁₆ antiserum followed by protein G-agarose. Well-washed immunoprecipitates were subjected to SDS-PAGE and the proteins of interest were detected using specific antibodies as indicated. TCL were separately run on blots for detecting the corresponding protein expressions. Two separate sets of transfected cells yielded similar results. (B) Similar procedures were applied for detecting the interaction between Flag-tagged TPR1 and the Gα constructs as indicated. TCL were subjected to immunoprecipitation using either anti-Flag agarose gel or anti-Gα₁₆ antiserum followed by protein G-agarose. Two separate sets of transfected cells yielded similar results. Band intensities were quantified and figures on the lanes of QL mutants are the fold increase compared with the band on the lane of their corresponding wild-type counterparts.

expressed with recombinant PLC β 2, as revealed in the total cell lysates. The endogenously expressed PLC β 2 could be detected (lane 1 of the bottom strip in Figure 4A) but the level was too low for co-immunoprecipitation assays. Both G α_{16} and G α_{16} QL could be precipitated with PLC β 2 using either anti-G α_{16} or anti-PLC β 2 antibodies, while G α_{16} QL showed much stronger interaction, indicating that the activation of G α_{16} facilitated its interaction with PLC β 2. Similarly, the QL forms of either one of the three mutants showed better interactions with PLC β 2 when compared with their corresponding wild-type forms. However, the fold increases between the three alanine mutants and their corresponding QL forms were much lower than the G α_{16} pair (values on top of each lane as indicated in Figure 4A). Such reduction was apparently due to the slight enhancement of the interactions between the alanine mutants and PLC β 2 as compared with G α_{16} . No significant difference of the fold changes was observed between the three mutant pairs. Nonetheless, the results clearly indicated that the G α_{16} /PLC β interaction was impaired by both clusters of alanine mutations.

One of the distinctive features of G α_{16} -mediated signaling is the association with TPR1, which facilitates the accumulation of active Ras [13] and in turn activate the Raf/MEK/ERK cascade. To study the impact of the alanine mutations on the interaction between G α_{16} and TPR1, a Flag-tagged full-length TPR1 was co-expressed with each of the G α_{16} constructs as indicated in Figure 4. Co-immunoprecipitation of the proteins of interest was performed using anti-Flag tag or anti-G α_{16} antibodies. Both G α_{16} and G α_{16} QL could be co-immunoprecipitated with TPR1 (Figure 4); G α_{16} QL was bound to TPR1 more efficiently under our experimental condition. A truncated form of TPR1 lacking the C-terminal tail did not bind to G α_{16} (Rico K. Lo, Andrew M. Liu and Yung H. Wong, data not shown). The three alanine mutants could also be co-immunoprecipitated with TPR1. Similar to the G α_{16} -PLC β 2 interactions, the QL versions of the alanine mutants showed stronger interactions with TPR1 than their corresponding non-active counterparts. Fold enhancements between the QL form of the alanine mutants and the non-active forms were weaker than G α_{16} pair. The 5A-QL mutant showed the weakest enhancement of TPR1 interaction. Expression levels of 5A and 5A-QL might be less in the experiment as showed in the total cell lysates, but the fold enhancement was still the least among the others. More obvious results were obtained when the TPR1-G α_{16} complexes were immunoprecipitated with anti-G α_{16} antibody. The results suggested that the interactions between TPR1 and G α_{16} were also dependent on the identities of the two amino acid clusters. Alanine mutations at both clusters caused the greatest reduction of the interaction.

Perturbation of Receptor-Mediated Regulation of PLC β

G α_{16} is well-known for its receptor coupling promiscuity [4], but it also exhibits different degrees of coupling efficiencies to various GPCRs [22,24]. Since the receptor- and effector-interacting domains overlap partially on the surface of G α subunits, it is possible that the alanine mutations can also affect the receptor coupling of G α_{16} . To test this possibility, a panel of GPCRs was examined for their functional coupling to G α_{16} and the alanine mutants using the IP accumulation assay. These GPCRs included G $_i$ -coupled adenosine A $_1$ receptor (A $_1$ R), complement C5aR receptor, formyl peptide receptor (fMLPR), and G $_s$ -coupled adenosine A $_{2A}$ and A $_{2B}$ receptors (A $_{2A}$ R, A $_{2B}$ R) and dopamine D $_1$ receptor (D $_1$ R). All of these receptors are capable of utilizing G α_{16} to elicit intracellular calcium mobilization in FLIPR assays [24].

The selected G $_i$ - or G $_s$ -coupled receptors all induced IP accumulation significantly in the presence of G α_{16} , but with variable efficiencies (Figure 5). These receptors were also capable of interacting with the alanine mutants. In general, IP accumulation mediated by 2A and 3A were ~50% and 35%, respectively, of that induced by G α_{16} . In some cases, such as C5aR and A $_{2A}$ R, the differences between the responses of 2A and 3A mutants were not significant, but it may be due to the weaker overall responses when compared with other receptors. Contrastingly, the differences between the phenotypes of these two mutants in the coupling to A $_1$ R, A $_{2B}$ R and D $_1$ R were more obvious ($\geq 50\%$). Differences in the absolute responses of various receptors have been demonstrated previously in the measurement of intracellular Ca $^{2+}$ mobilization (see Table 1 of [24]). The differential coupling efficiencies of the tested GPCRs with G α_{16} allowed us to detect the differences on GPCR coupling between the alanine mutants. Agonist-induced activation of 5A could only marginally increase IP accumulation in cases of A $_1$ R, fMLPR and D $_1$ R, while the other receptors were totally unable to elicit a response via 5A (Figure 5). The defective phenotype indicated that the two clusters were essential for the effective receptor-mediated PLC β activation through G α_{16} . The switch III region (harboring the 3A mutations) appeared to be more influential to the productive receptor coupling events than the α 3 helix.

Upon closer examination of the basal levels of the cells expressing G α_{16} or the 3 mutants, it was noticed that the basal levels of IP accumulation in the cells expressing A $_1$ R, A $_{2A}$ R or A $_{2B}$ R with G α_{16} were higher than with the mutants (Figure 5). This trend of decreasing basal IP accumulation was similar to the receptor-activated responses, with G α_{16} >2A>3A>5A. The results here showed that the basal activities of all three adenosine receptors were suppressed by the expression of the alanine mutants, particularly for 5A. Such evidence further implied that the alanine




Figure 5

Coupling of alanine mutants of $G\alpha_{16}$ to different categories of G protein-coupled receptors. Individual GPCRs were coexpressed with $G\alpha_{16}$, 2A, 3A or 5A in COS-7 cells. Transfected cells were labeled with [3 H]myo-inositol and treated with PTX overnight. Cells were then treated with the appropriate receptor agonists (1 μ M CHA for A₁R, 100 nM C5a for C5aR, 100 nM fMLP for fMLPR, 10 μ M adenosine for both A_{2A}R and A_{2B}R, 10 μ M dopamine for D₁R) for an hour before extraction of accumulated labeled IP. *Agonist significantly stimulated IP production as compared to their corresponding untreated counterparts; Tukey-Kramer's test, $p < 0.05$. ‡ Agonist-induced responses in mutant-expressing cells were significantly lower than that obtained with $G\alpha_{16}$; Tukey-Kramer's test, $p < 0.05$.

mutants might associate with the receptors, but such receptor-G protein complexes were defective and the mutants actually sequestered the spontaneous activities of the receptors (also see Discussion).

5A Mutant was Defective in Receptor-Mediated Regulation of Type II Adenylyl Cyclase


Type 2 adenylyl cyclase (AC2) can be synergistically stimulated via the G $\beta\gamma$ complex released from G α in the presence of activated G α_s [25,26]. Measurement of AC2-mediated cAMP production in such circumstance is useful for the detection of receptor-mediated release of G $\beta\gamma$ subunits. Functional interaction of the alanine mutants with G $_i$ -coupled receptors, if any, would be expected to stimulate AC2 upon agonist triggering. HEK293 cells were co-transfected with empty vector pCDNA3 and cDNAs encoding G α_s QL, fMLPR and AC2. Activation of the fMLPR resulted in the stimulation of endogenous G $_i$ and release G $\beta\gamma$ which led to the activation of AC2 (Figure 6). PTX pretreatment attenuated the corresponding increase of cAMP level. In cells co-expressing G α_{16} , G α_s QL, fMLPR and AC2, activation of fMLPR led to an increase in cAMP formation in a PTX-insensitive manner (Figure 6), indicating the functional coupling of fMLPR with G α_{16} . Both 2A and 3A mutants exhibited comparable significant increases in cAMP production as G α_{16} . However, 5A was totally incapable of stimulating AC2, which suggested that the receptor-mediated activation of G $_{16}$ and the subsequent G $\beta\gamma$ release might be severely impaired by the mutations on both switch III and α 3 helix.

Discussion

There are at least two paths through which G α_{16} can transmit signals to downstream effectors, via PLC β or TPR1. Their relative contributions to well-established downstream effectors such as ERK, STAT3, NF- κ B and JNK have not been clearly addressed. This study attempted to decipher the differential roles of the two streams of signals by mutating putative "PLC β -interacting" residues on switch III and α 3 helix of G α_{16} . All three mutants (2A, 3A, and 5A) exhibited impaired ability to stimulate PLC β . Differential regulations of ERK, STAT3, NF- κ B and JNK by the mutants have been observed, suggesting that G α_{16} regulates these downstream signals through overlapping but discrete pathways. Interactions between the mutants and PLC β 2 or TPR1 were reduced by the alanine mutations, and mutations of both clusters caused the greatest reduction of interactions. Furthermore, the 5A mutant with both clusters mutated was severely impaired in receptor coupling, as was observed in receptor-mediated activation of PLC β and AC2, and the sequestration of constitutive receptor activity. Collectively, this study provided evidence that the integrity of the switch III region and α 3 helix of G α_{16} was critical for both PLC β -and TPR1-dependent signaling events.

G α_q and G α_{16} belong to the same subfamily of heterotrimeric G α subunits and both regulate PLC β in a similar fashion. Presumably, the molecular interactions between G α_q and G α_{16} with PLC β would share some common features. Based on the previous study on G α_q [1], the five homologous residues on G α_{16} were mutated into alanine, and the resultant mutants showed partial impairment on PLC β regulation (Figures 2). Employment of Q212L mutation bypassed the possible influence on receptor coupling as well as the effect of G $\beta\gamma$ complex release, which may also activate PLC β in its own. The incomplete impairment suggests that the G α -effector interactions may involve multiple contact sites, as documented in the studies on other G α subunits [15,17,18]. Co-immunoprecipitation experiments showed that the physical association between PLC β and G α_{16} was not completely abolished by the mutations of either clusters (Figure 4). The mere association between the two proteins was apparently not sufficient to exert the full activating effect. Mutants bearing the 2A mutations on the surface of α 3 helix of G α_{16} were less effective in PLC activation (Figure 2). These two residues were more likely to contribute to the direct interaction and activation of PLC β , and so have a greater effect. The switch III region where 3A mutations were located has been suggested to influence the interdomain interactions between the helical and GTPase domain of G α_s subunit, which in turn affect the GTP-induced activation [27]. Assuming the activation mechanisms of the homologous G α subunits were very similar, the weaker suppression on PLC β activation by 3A mutations might be an indirect effect. As all of the mutants were expressed to a comparable level (Figure 2), the differences in PLC β activation could not be attributed to insufficient expression of a particular mutant.

We have previously proposed that G α_{16} QL-induced STAT3 phosphorylation is mediated via c-Src/JAK and ERK pathways, with PLC β /PKC serving as upstream modulators [7,11]. Reduction of PLC β activation caused by the mutations may disturb the signaling cascades at multiple points, and the modulation of downstream effectors may be altered differentially depending on the strength of upstream signals to be integrated. The present study shows that the alanine mutants partially inhibited the phosphorylation of STAT3 (Figure 3B), and the inhibitory profiles are reminiscent of the PLC β activities (Figure 2B). However, the same mutants had very little effect on ERK phosphorylation (Figure 3A). These results suggest that G α_{16} QL-induced PLC β signal is important for STAT3 phosphorylation, but not ERK phosphorylation. With the discovery of TPR1, G α_{16} may stimulate the Raf/MEK/ERK axis through Ras. However, Ras-mediated activation of ERK is not the only ERK-regulating signal. Other streams of signals contributing to ERK activation include the PLC β -mediated Ca $^{2+}$ mobilization and PKC activation, as

**Figure 6**

Receptor-induced activation of type II adenylyl cyclase (AC2) mediated by Gα₁₆ and its mutants. HEK293 cells were cotransfected with cDNAs encoding the AC2 (3 μg/ml), Gα_sQL (0.015 μg/ml) and fMLPR (3 μg/ml) together with Gα₁₆, 2A, 3A or 5A. Cells were labeled with [³H]adenine and treated with or without PTX (100 ng/ml) overnight. cAMP accumulation was assayed in response to the treatment with 100 nM fMLP for 1 h. * cAMP accumulation was significantly increased as compared with their corresponding basal value; Tukey-Kramer's test, *p* < 0.05.

well as possibly c-Src and JAK [10,11]. Furthermore, various molecular scaffolds for ERK pathway intermediates, GRKs and other signaling components may also contribute to the specificity of the ERK regulation [28,29] induced by the activation of Gα₁₆, which have not been rigorously studied here. The dependence of STAT3 activity on ERK is somewhat controversial, because ERK has been shown to negatively regulate STAT3 [30]. JNK, on the other hand, appears to be necessary for Src-mediated activation of STAT3 [31]. In this regard, it is interesting to note that all three mutants exhibited defective regulations of JNK (Figure 3D). It remains to be determined if JNK/Src signals are critical for the regulation of STAT3 tyrosine phosphorylation.

Gα₁₆QL is known to activate JNK in various cell types [8,9]. The complete loss of the ability of all three alanine

mutants to stimulate JNK suggests that Gα₁₆-induced PLCβ activation is essential for this pathway. PLCβ-triggered intracellular Ca²⁺ release can lead to the activation of guanine exchange factors (e.g. Sos) for small G proteins like Rac, which then activates JNK. The heavy dependence of JNK on the Ca²⁺ signal has been previously demonstrated: treatment of Ca²⁺ chelator BAPTA-AM completely abolished the bradykinin-induced JNK activity in HepG2 cells [32]. Although the PLCβ activation was not completely attenuated by the alanine mutants, diminished flux of intracellular Ca²⁺ might be already enough to suppress the JNK activity. Several studies from our laboratory have shown that Gβγ complex is a mediator of JNK activation by various GPCRs [32-36], but such regulation is abolished upon suppression of the Ca²⁺-activated SOS/Rac pathway [32,33]. The preservation of ERK activation did not alleviate the suppressive effects of the mutants on

JNK activation, which further implied the distinctive regulatory mechanisms of ERK and JNK activation by $G\alpha_{16}$.

Phosphorylation of NF- κ B induced by $G\alpha_{16}$ was much less affected by the alanine mutations (Figure 3C). Regulation of NF- κ B activation is mainly through the IKK/I κ B pathway [12], and both Ca^{2+} and MAPKs play important regulatory roles on IKK/I κ B activity. Apparently, PLC β / Ca^{2+} signals did not play a dominating role in the NF- κ B regulation. The very similar profile of the activation of ERK and NF- κ B suggested that ERK might be the key regulator of $G\alpha_{16}$ -mediated NF- κ B activation. The essential role of ERK on the regulation of NF- κ B has been studied in other cellular contexts [37,38]. The unique interaction of $G\alpha_{16}$ with TPR1, which promotes the Ras-mediated ERK activation [13], might be one of the possible explanations. Various monomeric G proteins, including Ras and Rac1, can activate NF- κ B activity [39].

Mutations of both amino acid clusters resulted in almost complete loss of GPCR-induced PLC β activation. Consistently, the effect of 3A mutation on the switch III region appeared to be more detrimental than 2A on the α 3 helix, and that seemed to be contrary to the QL mutants. The apparent discrepancies may be due to the different functional impacts of the two mutated clusters towards receptor-mediated activation of $G\alpha_{16}$ versus the mutation-induced spontaneous activation of $G\alpha_{16}$ QL. It has been shown previously that the intramolecular interactions along the interface between the helical and GTPase domains of $G\alpha_s$, of which switch III is involved, is critical for the receptor-mediated activation of $G\alpha_s$ [27,40]. The impact of the mutations of switch III (i.e. 3A mutant) could be revealed when studying the receptor-dependent signaling events, like those in Figures 5 and 6. The results indicated that $G\alpha_{16}$ made use of similar molecular architecture as $G\alpha_s$, wherein the alterations of the interdomain interactions (3A mutant) exerted a greater effect on the receptor-mediated activation than the mutations in the middle of α 3 helix (2A mutant). However, in the presence of Q212L mutation (the constitutively active mutant), $G\alpha_{16}$ is simply 'locked' into a GTP-bound state with certain intramolecular conformational adaptations. Such behavior might mask the effect of 3A mutation on the activation process of $G\alpha_{16}$. Instead, the functional defects caused by the mutations on the exposed surface (2A mutant) became more prominent.

The co-expression of $G\alpha_{16}$ and GPCRs often leads to an elevated basal IP accumulation [4,24], presumably because the spontaneous activity of receptors becomes amplified through the formation of GPCR/ $G\alpha_{16}$ complexes. Agonist-independent constitutive activation of A_1 R [41,42] and A_{2A} R [43,44] have been previously demonstrated. Although not much evidence is available to illustrate the constitutive activity of A_{2B} R, inverse agonists

for A_{2B} R have been identified [45]. Expression of the three alanine mutants resulted in the reduction of the basal IP accumulation, with the effect of 5A being the most prominent, indicating that the G protein complex formed by the alanine mutants reversed the constitutive activity of the adenosine receptors. Such hypothesis is not over-speculative, as it has been shown that the affinity of G_s for β_2 -adrenergic receptor can be increased by replacing 5 $G\alpha_s$ residues in the α 3/ β 5 loop region with the homologous $G\alpha_{12}$ residues [17]. Our previous study also showed that $G\alpha_{16/z}$ chimeras exhibit robust constitutive activity when co-expressed with various GPCRs [24], indicating that $G\alpha_{16}$ could associate with GPCR and transduce spontaneous receptor activity.

AC2 activation assay was employed to study the release of $G\beta\gamma$ complex from coupled G protein after receptor activation (Figure 6). The assay readout is less sensitive to the functional effect of $G\alpha$ subunits, and hence mostly reflects the efficiency of receptor-mediated $G\beta\gamma$ release from the activated $G\alpha$ subunit. One of our previous studies [21] showed that mutations on $G\alpha_{16}$ causing defective $G\beta\gamma$ interaction significantly reduced its ability to stimulate AC2. In this study, mutations of both but not either one of the clusters destroyed the receptor coupling of $G\alpha_{16}$ to fMLP receptor. Defective functional release of $G\beta\gamma$ upon receptor activation would suggest that the G protein trimer could not transmit the proper conformational changes from activated receptor or it could not even detect the activated receptor. One possibility is that the pre-associated receptor-G protein complex prefers to stay in an inactivated and associated form, which is echoed by the observation of the reduction of the constitutive receptor activities of the adenosine receptors (Figure 6).

Conclusion

The two clusters of putative PLC β -interacting residues on $G\alpha_{16}$ were indeed crucial for both receptor coupling and regulation of various downstream effectors. The two residues on the α 3 helix were more exposed and important for PLC β activation, whereas the clusters of three residues on the switch III region affected the receptor-mediated activation of $G\alpha_{16}$ more. Mutations of either clusters suppressed the activation of STAT3 and, to a greater extent, JNK activation but not ERK and NF- κ B signaling. The physical associations between $G\alpha_{16}$ and PLC β or TPR1 were sensitive to the mutations, and both clusters were both determinative to the interactions. Further studies on the functional impacts of the mutual interactions between $G\alpha_{16}$, PLC β and TPR1 may provide more insights on the characteristics of G_{16} -mediated signaling events.

Methods

Materials

Human TPR1 cDNA was kindly provided by Richard D. Ye (Department of Pharmacology, College of Medicine, Uni-

versity of Illinois, Chicago, IL). Human embryonic kidney 293 (HEK293) and monkey kidney fibroblast COS-7 cells were purchased from American Type Culture Collection (ATCC CRL-1573 and 1651). Restriction endonucleases were from Roche Applied Sciences. DNA purification columns were from Qiagen. DNA modification enzymes, custom primers and cell culture and transfection reagents were from Invitrogen. [³H]adenine and chemiluminescence detection kit for Western blotting were from GE Healthcare. [γ -³²P]ATP and [³H]myo-inositol was from PerkinElmer Inc. Antibodies against various molecules are from: Cell Signaling Technology (Danvers, MA, USA) for ERK1/2 (Thr²⁰²/Tyr²⁰⁴), STAT3 (Tyr⁷⁰⁵) and NF- κ B (Ser⁵³⁶) and their total forms; Torrey Pines Biolabs (C-terminus; Houston, TX, USA) and Gramsch Laboratories (N-terminus, custom-made; Schwabhausen, Germany) for $G\alpha_{16}$; sc-206, Santa Cruz Biotechnology (Santa Cruz, CA, USA) for anti-PLC β 2; anti-FLAG antiserum and anti-FLAG affinity gel were from Sigma (St. Louis, MO, USA). Protein G-agarose and cross-linking reagent dithiobis [succinimidylpropionate] (DSP) were from Pierce Biotechnology (IL, USA). Pertussis toxin was purchased from List Biological Laboratories. All other chemicals were obtained from Sigma.

Sequence alignment and molecular modeling

Complete amino acid sequence alignment of $G\alpha$'s was generated by CLUSTAL X 1.83 [46]. Molecular models were created by SWISS-MODEL web server [47] using the coordinates of the crystal structures of $G\alpha_{11}$ [48] and $G\alpha_{11}$ [49] in their corresponding heterotrimers retrieved from Protein Data Bank maintained by Research Collaboratory for Structural Bioinformatics [50] as the modeling templates. The model was modified and visualized with UCSF Chimera [51].

Construction of $G\alpha_{16}$ Mutants

Polymerase chain reactions (PCR) were performed to construct the mutants. Amplified cDNA fragment was subcloned into pcDNA3 vector, which possessed T7 and SP6 promoter regions for primer attachment in PCR reaction. Human $G\alpha_{16}$ WT and $G\alpha_{16}$ QL [21] were used as a template to amplify mutants. A pair of overlapping sense and anti-sense primers was designed encoding the desired mutations to replace particular amino acids into alanines. Primers are listed (from 5' to 3') with the mismatch nucleotides underlined: 16-2A/S: CTC GCA TTG TTT GCG GCG ATC CTG GAA CTA CCC; 16-2A/AS: TTC CAG GAT CGC CGC AAA CAA TGC GAG GCT; 16-3A/S: GAG GAG AAC GCA GCT GCA AAC CGC ATG AAG GAG; 16-3A/AS: CAT GCG GTT TGC AGC TGC GTT CTC CTC CAG GCA. The corresponding cDNA fragments were amplified using the desired primers with either T7 or SP6 primer by thermal cycling at 94°C (60 sec)/53°C (60 sec)/72°C (90 sec) for 35 cycles in Thermocycler 40 from Stratagene. Subse-

quently, the two fragments were pooled together as templates and the full-length mutated cDNAs were generated with the flanking (T7 and SP6) primers. The same thermal cycle program was used for second PCR to amplify full-length products. Full-length PCR products were subcloned in *Xba* I site of pcDNA3. DNA sequences of mutants were verified by autosequencing using BigDye® Terminator v3.1 cycle sequencing kit and ABI PRISM® Genetic Analyser (Applied Biosystems) and restrictive enzyme digestion. 5A and 5A-QL were constructed by using 3A and 3A-QL as templates, respectively.

Cell Culture and Transfection

HEK293 and COS-7 cells were cultured in Earle's modified essential medium (MEM) and Dulbecco's modified Eagle's medium (DMEM), respectively, with 10% fetal bovine serum (FBS; vol/vol), 50 units/ml penicillin and 50 μ g/ml streptomycin. Cells were incubated at 37°C in humidified air with 5% CO₂. At the day before transfection, COS-7 cells were seeded on 12-well plates at a density of 1×10^5 cells/well. For western blot analysis, HEK293 cells were used instead, wherein 2×10^5 cells were seeded on 6-well plates. For adenylyl cyclase assay, 4×10^5 of HEK293 cells were transferred to 12-well plates. For c-Jun N-terminal kinase assay, 3×10^5 of COS-7 cells were seeded on 6-well plates. cDNA transfection was achieved using Lipofectamine™ and PLUS™ reagents following the manufacturer's protocol. 50–75% of the cell population will take up the cDNAs as indicated by co-expressing β -galactosidase as a reporter.

Inositol Phosphate (IP) Accumulation Assay

750 μ l of inositol-free DMEM containing 5% FBS and 2.5 μ Ci/ml myo-[³H]-inositol was added to each well of transfected COS-7 cells and incubated for 18–24 hr. The labeling media were subsequently replaced by 1 ml of assay medium (DMEM with 20 mM HEPES, pH 7.5 and 10 mM LiCl) for 10 min and then 1 ml of assay medium containing the appropriate agonist was added to the cells for another 1 h at 37°C. Reaction was stopped by adding 750 μ l of ice-cold 20 mM formic acid after aspiration and the plates were stored at 4°C for 30 min. [³H]-IP were separated from labeled inositol by ion-exchange chromatography as described previously [52].

cAMP Accumulation Assay

HEK293 cell were labeled with [³H]-adenine (1 μ Ci/ml) in MEM containing 1% FBS (vol/vol) one day after transfection. After 18–24 h the labeling media were replaced by 1 ml of 20 mM HEPES-buffered MEM containing 1 mM isobutylmethylxanthine (IBMX) and the appropriate drugs and incubated at 37°C for 30 min. The treatment was terminated with 1 ml ice-cold 5% trichloroacetic acid (wt/vol) with 1 mM ATP after aspiration and stored at 4°C for 30 min. [³H]cAMP was extracted from the pool of

labeled nucleotides by sequential ion-exchange chromatography as described [53].

Data Analysis

For cAMP and IP accumulation assay, absolute values for cAMP or IP accumulations varied between experiments, but variability within a given experiment was in general <10%. The cAMP levels were interpreted as the ratios of the counts per minute of [³H]cAMP fractions to those of the total labeled nucleotide fractions and expressed as cAMP/(cAMP + Total). Similarly, IP levels were expressed as IP/(IP + Total). Data shown in the figures were the mean ± S.D. of triplicates within one single experiment. At least three individual experiments yielded similar results. One-way ANOVA with Tukey-Kramer's test were performed using GraphPad Prism 3.03 to verify the significance between different treatment groups within the experiments.

Cross-linking and Coimmunoprecipitation

Transfected HEK293 cells were washed with PBS twice and then treated in the same buffer containing 0.5 mM DSP for 15 min at room temperature to cross-link the membrane proteins. Cells were washed as above and maintained in quenching solution (50 mM glycine in PBS, pH 7.4) for 5 min, and then lysed in RIPA buffer (25 mM HEPES, pH 7.5, 0.1% SDS, 1% Nonidet P-40, 0.5% sodium deoxycholate, 1 mM dithiothreitol, 200 μM Na₃VO₄, 0.7 μg/ml pepstatin, 4 μg/ml aprotinin, 100 μM PMSF, and 2 μg/ml leupeptin). For coimmunoprecipitation, cell lysates were incubated with anti-Gα₁₆ (4 μg/sample) or anti-PLCβ2 antiserum (0.4 μg/sample), followed by the incubation with 30 μl of protein G-agarose (50% slurry) at 4°C for 2 h, or 30 μl anti-FLAG affinity agarose gel (50% slurry) at 4°C overnight. Immunoprecipitates were washed by 400 μl RIPA buffer 4 times, and then resuspended in 50 μl RIPA buffer gel loading buffer and boiled for 5 min. Gα₁₆ and FLAG-TPR1 proteins in the immunoprecipitates were detected by specific primary antisera and horseradish peroxidase-conjugated secondary antisera using Western blotting analysis.

Western Blotting Analysis

COS-7 cells were grown on 100-mm dishes to 70–80% confluence. Transfection was performed as in 12-well plates with proper adjustments to the surface area of the plate and the amounts of the reagents used. After 48 h in normal growth conditions, cells were washed with Ca²⁺/Mg²⁺-free Dulbecco's PBS (DPBS) and harvested with 3 ml DPBS containing 10 mM EDTA. The following procedures were performed at 4°C. Cells were spun down briefly (200 g, 5 min), resuspended in hypotonic lysis buffer and lysed by one cycle of freeze-thawing followed by 10 passages through a 27-gauge needle. Nuclei were removed by brief spinning and membranes were collected by spinning the supernatants at 15,000 g for 15 min. Membrane pel-

lets were finally resuspended in lysis buffer. Detergent-compatible protein assay system (Bio-Rad) was employed to determine the protein contents in cell lysates or crude membrane preparations. Protein samples were separated by 10% SDS-PAGE. Separated proteins were transferred to nitrocellulose membrane by electroblotting. Antisera against specific proteins were applied and the detection was facilitated by ECL chemiluminescence reagents (GE Healthcare). Band intensities on the autoradiographs were analyzed and quantified using ImageJ 1.40 [54].

In vitro c-Jun N-terminal Kinase (JNK) Assay

Details of the assays were basically identical as described in our previous study [55]. Briefly, COS-7 cell were transfected with the appropriate constructs and serum-starved overnight before assay. Cells were lysed by chilled detergent-containing lysis buffer with agitation on ice. 50 μl of each supernatant was used for the detection of JNK-HA expression after removing the cell debris by centrifugation, and the remaining was incubated for 1 h at 4°C with anti-HA antibody (2 μg/sample), followed by incubation with 30 μl of protein A-agarose at 4°C for 1 h. Immunoprecipitates were washed twice with lysis buffer and twice with kinase assay buffer, resuspended in kinase assay buffer with 5 μg of GST-c-Jun, and the kinase reactions were initiated by the addition of 10 μl of ATP buffer (50 μM ATP containing 2 μCi of [γ-³²P]-ATP per sample). After 30 min incubation at 30°C with occasional shaking, the reactions were terminated by adding sample buffer, and the samples were resolved by 12% SDS-PAGE. The gel was fixed for 30 min and the radioactivity incorporated to GST-c-Jun was detected and quantified by PhosphorImager (Molecular Dynamics 445 SI).

Abbreviations

A₁R, A_{2A}R, A_{2B}R: types 1, 2A and 2B adenosine receptor; AC: adenylyl cyclase; CHA: N⁶-cyclohexyladenosine; D₁R: type 1 dopamine receptor; ERK: extracellular signal-regulated kinases 1 and 2; fMLP: N-formyl-methionyl-leucyl-phenylalanine; GPCR: G protein coupled receptor; GTPase: GTP hydrolase; IBMX: isobutylmethylxanthine; IKK: inhibitor of κB kinase; IP: inositol phosphates; JNK: c-jun N-terminal kinase; MAPK: mitogen-activating protein kinase; MEK: MAPK/ERK kinase; MEM: Eagle's minimal essential medium; MKK: MAPK kinase; NF-κB: nuclear factor κB; PKC: protein kinase C; PLCβ: phospholipase Cβ isoform; PTX: pertussis toxin; STAT3: type 3 signal transducers and activator of transcription protein; TPR1: tetratricopeptide repeat protein 1.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MY carried out most of the experiments, except the co-immunoprecipitation, which was performed by AMFL.

MKCH helped to design the mutant constructs, performed molecular model constructions, analyzed and interpreted the results, and drafted the manuscript. YHW conceived of the study and participated in its design and coordination.

Acknowledgements

We thank Dr. Richard Ye for the generous donation of human TPRI cDNA, Lisa Yung for technical assistance and Dr. Anthony Chan for helpful discussion and sharing of unpublished results. The study was supported in part by grants from the Research Grants Council of Hong Kong (HKUST 6120/04M and 3/03C), the University Grants Committee (AoE/B-15/01), and the Hong Kong Jockey Club. YHW was a recipient of the Croucher Senior Research Fellowship.

References

- Venkatakrishnan G, Exton JH: **Identification of determinants in the alpha-subunit of Gq required for phospholipase C activation.** *J Biol Chem* 1996, **271**:5066-5072.
- Coleman DE, Berghuis AM, Lee E, Linder ME, Gilman AG, Sprang SR: **Structures of active conformations of Gi alpha 1 and the mechanism of GTP hydrolysis.** *Science* 1994, **265**:1405-1412.
- Lambright DG, Noel JP, Hamm HE, Sigler PB: **Structural determinants for activation of the alpha-subunit of a heterotrimeric G protein.** *Nature* 1994, **369**:621-628.
- Offermanns S, Simon MI: **G alpha 15 and G alpha 16 couple a wide variety of receptors to phospholipase C.** *J Biol Chem* 1995, **270**:15175-15180.
- Willkie TM, Scherle PA, Strathmann MP, Slepak VZ, Simon MI: **Characterization of G-protein alpha subunits in the Gq class: expression in murine tissues and in stromal and hematopoietic cell lines.** *Proc Natl Acad Sci USA* 1991, **88**:10049-10053.
- Shyamala V, Khoja H: **Interleukin-8 receptors R1 and R2 activate mitogen-activated protein kinases and induce c-fos, independent of Ras and Raf-1 in Chinese hamster ovary cells.** *Biochemistry* 1998, **37**:15918-15924.
- Lo RK, Cheung H, Wong YH: **Constitutively active Galpha16 stimulates STAT3 via a c-Src/JAK- and ERK-dependent mechanism.** *J Biol Chem* 2003, **278**:52154-52165.
- Heasley LE, Storey B, Fanger GR, Butterfield L, Zamarripa J, Blumberg D, Maue RA: **GTPase-deficient G alpha 16 and G alpha q induce PC12 cell differentiation and persistent activation of cJun NH2-terminal kinases.** *Mol Cell Biol* 1996, **16**:648-656.
- Higashita R, Li L, Van Putten V, Yamamura Y, Zarinetchi F, Heasley L, Nemenoff RA: **Galpha16 mimics vasoconstrictor action to induce smooth muscle alpha-actin in vascular smooth muscle cells through a Jun-NH2-terminal kinase-dependent pathway.** *J Biol Chem* 1997, **272**:25845-25850.
- Lo RK, Wong YH: **Transcriptional activation of c-Fos by constitutively active Galpha(16)QL through a STAT1-dependent pathway.** *Cell Signal* 2006, **18**:2143-2153.
- Liu AM, Wong YH: **G16-mediated activation of nuclear factor kappaB by the adenosine A1 receptor involves c-Src, protein kinase C, and ERK signaling.** *J Biol Chem* 2004, **279**:53196-53204.
- Yang M, Sang H, Rahman A, Wu D, Malik AB, Ye RD: **G alpha 16 couples chemoattractant receptors to NF-kappa B activation.** *J Immunol* 2001, **166**:6885-6892.
- Marty C, Browning DD, Ye RD: **Identification of tetratricopeptide repeat 1 as an adaptor protein that interacts with heterotrimeric G proteins and the small GTPase Ras.** *Mol Cell Biol* 2003, **23**:3847-3858.
- Bae H, Anderson K, Flood LA, Skiba NP, Hamm HE, Graber SG: **Molecular determinants of selectivity in 5-hydroxytryptamine 1B receptor-G protein interactions.** *J Biol Chem* 1997, **272**:32071-32077.
- Berlot CH, Bourne HR: **Identification of effector-activating residues of Gs alpha.** *Cell* 1992, **68**:911-922.
- Cai K, Itoh Y, Khorana HG: **Mapping of contact sites in complex formation between transducin and light-activated rhodopsin by covalent crosslinking: use of a photoactivatable reagent.** *Proc Natl Acad Sci USA* 2001, **98**:4877-4882.
- Grishina G, Berlot CH: **A surface-exposed region of G(salpha) in which substitutions decrease receptor-mediated activation and increase receptor affinity.** *Mol Pharmacol* 2000, **57**:1081-1092.
- Ho MK, Wong YH: **The amino terminus of Galpha(z) is required for receptor recognition, whereas its alpha4/beta6 loop is essential for inhibition of adenylyl cyclase.** *Mol Pharmacol* 2000, **58**:993-1000.
- Lee CH, Katz A, Simon MI: **Multiple regions of G alpha 16 contribute to the specificity of activation by the C5a receptor.** *Mol Pharmacol* 1995, **47**:218-223.
- Skiba NP, Bae H, Hamm HE: **Mapping of effector binding sites of transducin alpha-subunit using G alpha t/G alpha ii chimeras.** *J Biol Chem* 1996, **271**:413-424.
- Ho MK, Yung LY, Wong YH: **Disruption of receptor-mediated activation of G protein by mutating a conserved arginine residue in the switch II region of the alpha subunit.** *J Neurochem* 1999, **73**:2101-2109.
- Ho MK, New DC, Wong YH: **Co-expressions of different opioid receptor types differentially modulate their signaling via G(16).** *Neurosignals* 2002, **11**:115-122.
- Wu EH, Lo RK, Wong YH: **Regulation of STAT3 activity by G16-coupled receptors.** *Biochem Biophys Res Commun* 2003, **303**:920-925.
- Liu AM, Ho MK, Wong CS, Chan JH, Pau AH, Wong YH: **Galpha(16/z) chimeras efficiently link a wide range of G protein-coupled receptors to calcium mobilization.** *J Biomol Screen* 2003, **8**:39-49.
- Federman AD, Conklin BR, Schrader KA, Reed RR, Bourne HR: **Hormonal stimulation of adenylyl cyclase through Gi-protein beta gamma subunits.** *Nature* 1992, **356**:159-161.
- Tang WJ, Gilman AG: **Type-specific regulation of adenylyl cyclase by G protein beta gamma subunits.** *Science* 1991, **254**:1500-1503.
- Grishina G, Berlot CH: **Mutations at the domain interface of Gsalpha impair receptor-mediated activation by altering receptor and guanine nucleotide binding.** *J Biol Chem* 1998, **273**:15053-15060.
- Shaul YD, Seger R: **The MEK/ERK cascade: from signaling specificity to diverse functions.** *Biochim Biophys Acta* 2007, **1773**:1213-1226.
- Werry TD, Sexton PM, Christopoulos A: **"Ins and outs" of seven-transmembrane receptor signalling to ERK.** *Trends Endocrinol Metab* 2005, **16**:26-33.
- Jain N, Zhang T, Fong SL, Lim CP, Cao X: **Repression of Stat3 activity by activation of mitogen-activated protein kinase (MAPK).** *Oncogene* 1998, **17**:3157-3167.
- Turkson J, Bowman T, Adnane J, Zhang Y, Djeu JY, Sekharam M, Frank DA, Holzman LB, Wu J, Sebti S, Jove R: **Requirement for Ras/Rac1-mediated p38 and c-Jun N-terminal kinase signaling in Stat3 transcriptional activity induced by the Src oncoprotein.** *Mol Cell Biol* 1999, **19**:7519-7528.
- Chan AS, Wong YH: **Gbetagamma signaling and Ca2+ mobilization co-operate synergistically in a Sos and Rac-dependent manner in the activation of JNK by Gq-coupled receptors.** *Cell Signal* 2004, **16**:823-836.
- Chan AS, Yeung WW, Wong YH: **Integration of G protein signals by extracellular signal-regulated protein kinases in SK-N-MC neuroepithelioma cells.** *J Neurochem* 2005, **94**:1457-1470.
- Kam AY, Chan AS, Wong YH: **Kappa-opioid receptor signals through Src and focal adhesion kinase to stimulate c-Jun N-terminal kinases in transfected COS-7 cells and human monocytic THP-1 cells.** *J Pharmacol Exp Ther* 2004, **310**:301-310.
- Kam AY, Chan AS, Wong YH: **Phosphatidylinositol-3 kinase is distinctively required for mu-, but not kappa-opioid receptor-induced activation of c-Jun N-terminal kinase.** *J Neurochem* 2004, **89**:391-402.
- Kam AY, Chan AS, Wong YH: **Rac and Cdc42-dependent regulation of c-Jun N-terminal kinases by the delta-opioid receptor.** *J Neurochem* 2003, **84**:503-513.
- Chen BC, Lin WW: **PKC- and ERK-dependent activation of I kappa B kinase by lipopolysaccharide in macrophages: enhancement by P2Y receptor-mediated CaMK activation.** *Br J Pharmacol* 2001, **134**:1055-1065.
- Ho RC, Hirshman MF, Li Y, Cai D, Farmer JR, Aschenbach WG, Witczak CA, Shoelson SE, Goodyear LJ: **Regulation of I kappa B kinase and NF-kappa B in contracting adult rat skeletal muscle.** *Am J Physiol Cell Physiol* 2005, **289**:C794-801.

39. Frost JA, Swantek JL, Stippec S, Yin MJ, Gaynor R, Cobb MH: **Stimulation of NFkappa B activity by multiple signaling pathways requires PAK1.** *J Biol Chem* 2000, **275**:19693-19699.
40. Marsh SR, Grishina G, Wilson PT, Berlot CH: **Receptor-mediated activation of Galpha: evidence for intramolecular signal transduction.** *Mol Pharmacol* 1998, **53**:981-990.
41. Bakker RA, Casarosa P, Timmerman H, Smit MJ, Leurs R: **Constitutively active Gq/iI-coupled receptors enable signaling by co-expressed G(i/o)-coupled receptors.** *J Biol Chem* 2004, **279**:5152-5161.
42. Shryock JC, Ozeck MJ, Belardinelli L: **Inverse agonists and neutral antagonists of recombinant human A1 adenosine receptors stably expressed in Chinese hamster ovary cells.** *Mol Pharmacol* 1998, **53**:886-893.
43. Klinger M, Kuhn M, Just H, Stefan E, Palmer T, Freissmuth M, Nanoff C: **Removal of the carboxy terminus of the A2A-adenosine receptor blunts constitutive activity: differential effect on cAMP accumulation and MAP kinase stimulation.** *Naunyn Schmiedeberg's Arch Pharmacol* 2002, **366**:287-298.
44. Maenhaut C, Van Sande J, Libert F, Abramowicz M, Parmentier M, Vanderhaegen JJ, Dumont JE, Vassart G, Schiffmann S: **RDC8 codes for an adenosine A2 receptor with physiological constitutive activity.** *Biochem Biophys Res Commun* 1990, **173**:1169-1178.
45. Varani K, Gessi S, Merighi S, Vincenzi F, Cattabriga E, Benini A, Klotz KN, Baraldi PG, Tabrizi MA, Lennan SM, et al.: **Pharmacological characterization of novel adenosine ligands in recombinant and native human A2B receptors.** *Biochem Pharmacol* 2005, **70**:1601-1612.
46. Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, Thompson JD: **Multiple sequence alignment with the Clustal series of programs.** *Nucleic Acids Res* 2003, **31**:3497-3500.
47. Schwede T, Kopp J, Guex N, Peitsch MC: **SWISS-MODEL: An automated protein homology-modeling server.** *Nucleic Acids Res* 2003, **31**:3381-3385.
48. Wall MA, Coleman DE, Lee E, Iniguez-Lluhi JA, Posner BA, Gilman AG, Sprang SR: **The structure of the G protein heterotrimer Gi alpha 1 beta 1 gamma 2.** *Cell* 1995, **83**:1047-1058.
49. Lambright DG, Sondek J, Bohm A, Skiba NP, Hamm HE, Sigler PB: **The 2.0 A crystal structure of a heterotrimeric G protein.** *Nature* 1996, **379**:311-319.
50. Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, Bourne PE: **The Protein Data Bank.** *Nucleic Acids Res* 2000, **28**:235-242.
51. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE: **UCSF Chimera – a visualization system for exploratory research and analysis.** *J Comput Chem* 2004, **25**:1605-1612.
52. Tsu RC, Chan JS, Wong YH: **Regulation of multiple effectors by the cloned delta-opioid receptor: stimulation of phospholipase C and type II adenylyl cyclase.** *J Neurochem* 1995, **64**:2700-2707.
53. Wong YH: **Gi assays in transfected cells.** *Methods Enzymol* 1994, **238**:81-94.
54. Girish V, Vijayalakshmi A: **Affordable image analysis using NIH Image/ImageJ.** *Indian J Cancer* 2004, **41**:47.
55. Chan AS, Wong YH: **Regulation of c-Jun N-terminal kinase by the ORL(1) receptor through multiple G proteins.** *J Pharmacol Exp Ther* 2000, **295**:1094-1100.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

