


FSHD myotubes



Gene expression during normal and FSHD myogenesis

Tsumagari *et al.*

RESEARCH ARTICLE

Open Access

Gene expression during normal and FSHD myogenesis

Koji Tsumagari¹, Shao-Chi Chang¹, Michelle Lacey^{2,3}, Carl Baribault^{2,3}, Sridar V Chittur⁴, Janet Sowden⁵, Rabi Tawil⁵, Gregory E Crawford⁶ and Melanie Ehrlich^{1,3*}

Abstract

Background: Facioscapulohumeral muscular dystrophy (FSHD) is a dominant disease linked to contraction of an array of tandem 3.3-kb repeats (D4Z4) at 4q35. Within each repeat unit is a gene, *DUX4*, that can encode a protein containing two homeodomains. A *DUX4* transcript derived from the last repeat unit in a contracted array is associated with pathogenesis but it is unclear how.

Methods: Using exon-based microarrays, the expression profiles of myogenic precursor cells were determined. Both undifferentiated myoblasts and myoblasts differentiated to myotubes derived from FSHD patients and controls were studied after immunocytochemical verification of the quality of the cultures. To further our understanding of FSHD and normal myogenesis, the expression profiles obtained were compared to those of 19 non-muscle cell types analyzed by identical methods.

Results: Many of the ~17,000 examined genes were differentially expressed (> 2-fold, $p < 0.01$) in control myoblasts or myotubes vs. non-muscle cells (2185 and 3006, respectively) or in FSHD vs. control myoblasts or myotubes (295 and 797, respectively). Surprisingly, despite the morphologically normal differentiation of FSHD myoblasts to myotubes, most of the disease-related dysregulation was seen as dampening of normal myogenesis-specific expression changes, including in genes for muscle structure, mitochondrial function, stress responses, and signal transduction. Other classes of genes, including those encoding extracellular matrix or pro-inflammatory proteins, were upregulated in FSHD myogenic cells independent of an inverse myogenesis association. Importantly, the disease-linked *DUX4* RNA isoform was detected by RT-PCR in FSHD myoblast and myotube preparations only at extremely low levels. Unique insights into myogenesis-specific gene expression were also obtained. For example, all four Argonaute genes involved in RNA-silencing were significantly upregulated during normal (but not FSHD) myogenesis relative to non-muscle cell types.

Conclusions: *DUX4*'s pathogenic effect in FSHD may occur transiently at or before the stage of myoblast formation to establish a cascade of gene dysregulation. This contrasts with the current emphasis on toxic effects of experimentally upregulated *DUX4* expression at the myoblast or myotube stages. Our model could explain why *DUX4*'s inappropriate expression was barely detectable in myoblasts and myotubes but nonetheless linked to FSHD.

Background

Differentiation of myoblasts to myotubes is one of the best cell culture models for vertebrate differentiation. However, there has been only limited expression profiling of well characterized myoblast cell strains and of myoblasts differentiated in vitro to myotubes [1-3]. In this study, we profiled expression of control myoblasts

and myotubes as well as analogous cells from patients with facioscapulohumeral muscular dystrophy (FSHD). Importantly, we were able to compare control and FSHD myoblasts and myotubes with 19 different non-muscle cell types subjected to identical expression profiling. The data are directly comparable because the same experimental and computational techniques were used for all the cell types. This allowed us to identify myogenesis-specific as well as disease-associated differences in expression. We are particularly interested in regenerative myogenesis [4], as opposed to embryonic

* Correspondence: ehrlich@tulane.edu

¹Human Genetics Program, Tulane Medical School, New Orleans, LA, USA
Full list of author information is available at the end of the article

myogenesis [5], because of its role in limiting atrophy due to muscle damage, aging, and disease.

FSHD is a dominant disease whose pathogenesis is still perplexing despite new insights into its genetic linkage [6-8]. It is progressively debilitating and painful and mainly affects skeletal muscle. FSHD is linked to contraction at 4q35 of a tandem array of 3.3-kb repeats, D4Z4, from about 11-100 to 1-10 copies [9]. It is usually diagnosed in the second decade, and the patient's lifespan is generally not affected. Initially, the pathology is limited to a small set of skeletal muscles, often asymmetrically. There is apparently no involvement of smooth muscle. No efficacious treatment is available.

Although other expression profiling studies of FSHD vs. normal- or disease-control muscle biopsies have been done [6,10-13], no clear consensus has emerged as to the genes that lead to the muscle pathology. Usually, only modest up- or downregulation of gene expression was observed. FSHD is likely to involve defects in muscle cell precursors [10]; therefore, studies of FSHD myoblasts and myotubes should also elucidate normal myogenesis. In analyses of muscle tissue, myogenesis-specific, disease-related changes in expression are obscured by the very low percentages of (activated) satellite cells. Upon expression profiling of FSHD and control myoblasts (but not myotubes) in 2003, Winokur et al. found ~20 genes were FSHD-dysregulated; among them were genes involved in the response to oxidative stress [14]. Accordingly, they demonstrated and Barro et al. confirmed [14,15] that FSHD myoblasts are significantly more sensitive to the lethal effects of drug-induced oxidative stress than normal-control and disease-control myoblasts. Nonetheless, Barro et al. demonstrated that this hypersensitivity did not affect growth rates or the ability of myoblasts to differentiate to myotubes. A recent expression profiling study of FSHD and control myoblasts and myotubes by Cheli et al. [16] provided no characterization of the purity of the myoblast or myotube samples and paradoxically reported no muscle-related terms among 177 functional terms for genes with differential expression in normal-control myoblast vs. normal-control myotube preparations, which is very different from what we have found, as described below.

A number of 4q35 genes have been considered as candidates for the initially dysregulated gene during FSHD pathogenesis, namely, *FRG1*, *DUX4*, *DUX4C*, *ANT1* (*SLC25A4*), *FRG2*, *TUBB4Q*, and *FAT1* [6,17-24]. Recently implicated in FSHD pathogenesis from genetic mapping is *DUX4*, a 1.6-kb gene that resides within each 3.3-kb repeat unit of D4Z4. *DUX4* encodes a protein containing two homeodomains [25,26]. The protein is strongly pro-apoptotic when highly overexpressed in experimental models [21,27-29]. *DUX4* transcripts are

normally difficult to detect probably because of heterochromatinization of normal long D4Z4 arrays inhibiting their transcription [30,31] and the lack of a polyadenylation signal within *DUX4* [32], which generally leads to *DUX4* mRNA being unstable. However, in patients, a polyadenylation signal is provided for the most distal *DUX4* gene in the D4Z4 array by a common SNP located immediately distal to D4Z4 and specific to 4q35, but not to the non-pathogenic 10q26 D4Z4 array [6,7]. Therefore, expression of the FSHD-linked *DUX4* RNA isoform, *DUX4-fl* (full length), generally requires both array contraction at 4q35 and this SNP. Exceptions to the requirement for array contraction for generation of detectable *DUX4-fl* transcript were seen in normal testis and in myoblasts and myotubes from patients with a variant of FSHD called FSHD2, which is associated with inappropriate expression of *DUX4-fl* RNA from 4q35 despite a lack of contraction of the D4Z4 array [8]. Both of these exceptions may involve D4Z4 chromatin loosening in normal-length arrays due to partial hypomethylation of D4Z4 DNA [31,33].

DUX4-fl RNA was found in all examined FSHD myotube preparations, several FSHD myoblast cell strains, and two FSHD fibroblast cell strains, but not in the analogous control cells [7,8]. However, there are still fundamental questions about the disease mechanisms. Firstly, in view of the convincing genetic data linking FSHD to the *DUX4-fl* RNA isoform [6-8], why was *DUX4-fl* RNA expression in FSHD myoblasts and myotubes extraordinarily infrequent, e.g., ~ 1 in 1000 myoblast nuclei positive [8]? Why did two of five FSHD myoblast preparations lack detectable *DUX4-fl* RNA by nested RT-PCR [8] despite the finding that FSHD myoblasts have a phenotype of hypersensitivity to oxidative stress [14,15]? Why is FSHD essentially only a muscle-specific disease even though FSHD fibroblasts and FSHD myotubes display similar (very low) levels of *DUX4-fl* RNA isoform [8]? And lastly, why do C2C12 myoblasts with induced expression of human *D4Z4-fl* RNA [27,28] undergo apoptosis or display diminished differentiation to myotubes, unlike FSHD myoblasts, which grow and differentiate normally? Although apoptosis has been noted in FSHD muscle samples, it was mostly in late-stage muscle biopsies [34].

In the current study, using exon-based expression microarrays, we found differences in expression of diverse categories of genes in control myoblasts and myotubes relative to various non-muscle cell types. In addition, FSHD muscle progenitors had a unique pattern of gene dysregulation compared with analogous control cells. A distinctive expression profile was observed in FSHD myoblasts as well as in FSHD myotubes even though *DUX4-fl* RNA was present at extremely low levels or was undetectable in these cells.

Methods

Cell culture

Myoblast cell strains from FSHD patients and controls and the other cell types are described in Additional File 1, Additional File 2, and Additional File 3. FSHD myoblast cell strains from moderately affected muscle were no more difficult to generate than control myoblast cell strains. Duly signed patient consent forms were obtained that had been approved by the Institutional Review Boards of Tulane Health Science Center, the University of Rochester School of Medicine, and the University of Mississippi Medical Center in Jackson. Myoblasts were propagated by collagenase and dispase dispersion of the tissue, establishment of the cell strain in F10 medium with 50% MRC-5 F10 conditioned medium, pre-plating to remove contaminating fibroblasts, growth in the presence of 10 ng/ml bFGF and 1 μ M dexamethasone, and differentiation by limiting serum (2% horse serum for 1 day and 15% horse serum for 4-6 days), as described in more detail in Additional File 4. Each batch of cells was checked by immunocytochemistry for desmin (Thermo, RB-9014-P), a marker for muscle cells. The extent of myotube formation was determined by immunostaining for desmin and myosin heavy chain (MF20 monoclonal antibody from Stephen Hauschka) and determining the percentage of nuclei in multinucleated cells.

Microarray expression analysis

Myoblast cultures at ~70% confluence and myotube preparations harvested 5 - 7 days after induction of differentiation were snap-frozen and stored in liquid nitrogen. Total RNA was isolated by standard methods (TRIzol extraction and RNeasy column, QIAGEN), that included DNaseI digestion. RNA was checked for quality (Nanodrop and Agilent Bioanalyzer), and 1 μ g was labeled by a standard procedure that included a riboreduction step (Whole Transcript Sense Target labeling protocol, Affymetrix). The fragmented biotin-labeled cDNA was hybridized for 16 h to Affymetrix Exon 1.0 ST arrays and scanned (Scanner 3000 7 G, AGCC software, Affymetrix). After confirming the quality of the resulting .cel files (Affymetrix Expression Console software), they were imported into GeneSpring GX10. The data were quantile-normalized using only core-level transcripts (RMA) and baseline transformed to the median of all samples. The probe sets were further filtered to exclude ones that lie in the bottom 20th percentile across all samples. The raw data were deposited in the GEO database under series GSE26145.

qRT-PCR

RNA was isolated from snap-frozen myoblasts and myotubes (RNeasy Mini kit; QIAGEN). After treatment of

the RNA (RNA Quality Indicator ≥ 9 ; Experion, Bio-Rad) with DNaseI (Turbo DNA free, Ambion), cDNA was prepared using 1 μ g of total RNA (Protoscript M-MuLV First Strand cDNA synthesis kit with Random Primer Mix; NEB). RT-PCR was performed with SYBR Green detection (RT² SYBR[®] Green qPCR master mixes, SABiosciences; iCycler MyiQ, Bio-Rad). Unless otherwise noted, amplifications were done in duplicate using 20 - 40 ng of cDNA and 0.2 μ M primer (Invitrogen) and the following parameters: 95°C, 30 s; annealing at an optimized temperature, 30 s; 72°C, 30 s for 45 cycles. Primers were designed <http://frodo.wi.mit.edu/primer3/> and checked for the predicted specificity <http://www.ncbi.nlm.nih.gov/tools/primer-blast/>. Calibration curves for primer-pairs with serial 10-fold dilutions of a mixture of FSHD and control myotube and myoblast cDNAs gave slopes of -3.3 ± 0.4 and correlation coefficients of ≥ 0.98 . Melting curves from PCR products were confirmed to give a single peak. The normalization standard for qRT-PCR was *M6PR*, which, according to the microarray data, had the following average expression ratios for FSHD vs. control myotubes and FSHD vs. control myoblasts: 0.97 ($p = 0.9$) and 1.27 ($p = 0.2$), respectively. *M6PR* was the most stably expressed gene from a set of seven standards (*M6PR*, *MSN*, *POLR1D*, *PPIA*, *B2M*, *EIF4A2*, and *MAT2A*) tested on FSHD vs. control myotube samples [35] <http://medgen.ugent.be/~jvdesomp/genorm>.

Statistical analysis

ANOVA models for the comparison of gene expression levels across sample groups were fit using the limma package [36] in Bioconductor <http://www.bioconductor.org>, and heat maps were generated using the R gplots package <http://CRAN.R-project.org/package=gplots>. The Benjamini-Hochberg correction was used to adjust for multiple testing, and p -values of < 0.01 were considered significant. Chi-square tests for association and regression analyses were performed using R version 2.10 <http://www.R-project.org>.

Results

Cell samples for analysis

FSHD myoblasts were generated from moderately affected muscle (See Additional File 1 for details about samples). The control myoblasts were from two normal individuals and one disease-control patient (sporadic inclusion body myositis). The myoblast preparations contained 90-98% desmin-positive cells before differentiation to myotubes. Therefore, there was minimal contamination with muscle-derived fibroblasts. The minor portion of fibroblasts may nonetheless have influenced the behaviour of the myoblasts by cell-signaling and

might have made a contribution to the array signal, although only a minor one, as indicated by the consistent differences between FSHD and control samples. The myotube preparations had 72-80% of the nuclei in multinucleated myotubes (more than two nuclei per cell) after differentiation in comparison to the myoblast preparations with < 1% multinucleated cells (Figure 1). The growth medium for propagating myoblasts contained 10 ng/ml of basic fibroblast growth factor (bFGF) plus 1 μ M dexamethasone, as recommended in studies of normal and FSHD myoblasts [37,38] and as used in recent studies of *DUX4* expression [7,8]. For differentiation to myotubes, cells were incubated in the absence of bFGF and dexamethasone and with replacement of the normal 20% fetal bovine serum by 2% horse serum for one day, and then 15% horse serum for 4-6 days. Both the control and FSHD myotube preparations had cells with a more normal appearance and a higher percentage of myotubes due to the increase in the horse serum concentration after the first day of serum limitation. Dexamethasone improved the growth, the maximum number of cell population doublings, and appearance of control and FSHD myoblasts, as did regular additions of basic fibroblast growth factor. FSHD and control myoblasts and myotubes looked similar by phase microscopy

(Figure 1), and the FSHD and control myoblasts grew and differentiated to myotubes equally well.

Myogenesis-specific differences in gene expression

We used an exon-based microarray (Affymetrix Exon 1.0 ST) to profile the expression of myoblast cultures harvested at ~70% confluence from FSHD and control myoblasts and from myotube preparations (three each). Upon differentiation of control myoblasts to myotubes, predominantly upregulation of gene expression was seen (Figure 2). Setting the fold change (FC) threshold for control myotubes vs. control myoblasts of > 2.0 at a significance level of $p < 0.01$, 511 genes were upregulated and 224 downregulated; all p -values for array data were adjusted for multiple comparisons using the Benjamini-Hochberg correction. As expected, the group of genes that was upregulated in control myotubes vs. myoblasts contained a very strong overrepresentation of muscle genes. Seventeen out of 20 of the most overrepresented GO terms for this set of genes were specifically related to muscle and the other three were actomyosin structure organization, cell adhesion, and biological adhesion (DAVID functional analysis tool). The analogous downregulated group of genes had only several strong associations with GO terms, and all involved the plasma membrane.





Figure 1 Examples of myogenesis-specific immunostaining of FSHD and control myoblasts and myotubes used for the analysis. Green, desmin; red, myosin heavy chain (MF20); blue, DAPI. The last panels in both rows were myotube preparations doubly stained with MF20 and DAPI. The occasional small bright green or red dots are a staining artifact. Fibroblasts do not stain with the desmin antibody nor with MF20.




Also available for determination of myogenesis-specific gene expression was the powerful resource of our expression profiling of 19 heterologous non-muscle cell types as part of the ENCODE project (See Additional File 2 for descriptions of the cell types; Crawford GE, unpub. data). The diverse non-muscle cell cultures, which included fibroblasts, melanocytes, hepatocytes, and astrocytes, were profiled using the same type of exon microarrays and identical methods as for the myoblasts and myotubes. An analysis of variance (ANOVA) model was fit to determine expression differences between myoblasts or myotubes and these non-muscle cell types. The majority of the gene expression differences ($FC > 2$, $p < 0.01$) between control myoblasts and non-muscle cells involved increases in expression (Figure 3A). The 20 GO terms most overrepresented among the myoblast-upregulated genes were associated with the plasma membrane, muscle, the actin cytoskeleton, cell adhesion, and enzyme-linked receptor protein signaling; also prominent in this set of genes were GO terms for extracellular structure organization, calcium ion binding, response to oxygen levels, and GTPase regulator activity. The GO terms associated with the group of myoblast-downregulated genes were more diverse but featured sterol metabolism, cell cycling, and cell-cell junction. As expected, among the genes upregulated in control myotubes vs. non-muscle cell types, muscle GO

terms predominated but extracellular structure organization, cell adhesion, actin cytoskeleton, transmembrane receptor, and GTPase regulator activity were also prominent. For genes downregulated in control myotubes vs. non-muscle cell types (Figure 3A), almost all the top 20 overrepresented GO terms were related to cell division.

Extensive dysregulation of gene expression in FSHD myoblasts and myotubes

Comparing FSHD and control myoblasts, 1.7% of the ~17,000 genes represented on the microarray displayed $FC > 2.0$ in expression at a significance level of $p < 0.01$. Remarkably, only 10% of these 295 FSHD myoblast-dysregulated genes were downregulated (Figure 2). At the myotube stage there were almost three times as many FSHD-dysregulated genes (797 genes; 4.7% of the genes on the microarray) and a much higher percentage was downregulated (37%). FSHD dysregulation at the myoblast stage was significantly associated with dysregulation at the myotube stage. For example, in FSHD-to-control comparisons, 266 and 502 genes were upregulated more than 2-fold in myoblasts and myotubes, respectively, and 130 genes were found in both sets of FSHD-upregulated genes ($p < 0.00001$, Chi-square test). Therefore, 49% of the genes upregulated with a $FC > 2$ and $p < 0.01$ in FSHD vs. control myoblast samples (3 each) were similarly upregulated in



FSHD vs. control myotube samples (3 each). Similarly 39% of the 29 genes downregulated more than 2-fold in FSHD vs. control myoblasts were also downregulated more than 2-fold in FSHD vs. control myotubes.

It was not expected that 100% of the genes would behave similarly because of the extensive differences in control myoblast-specific vs. control myotube-specific transcription described above.

Previously reported evidence suggested that 4q35.2 genes *FRG1*, 4q35.1 gene *ANT1* (*SLC25A4*), and the 5q31 gene *PITX1*, which were represented on our microarrays, might be involved in FSHD [6,17-22]. In accord with other qRT-PCR studies [23,24,39], we did not observe dysregulation of *FRG1* or *PITX1*. Although detection of transcripts from *FRG1* is complicated by its cross-homology to many transcribed genomic sequences, even in the most specific probe-sets for the gene located in the first exon, similar signal intensities were seen for FSHD and control myogenic cells. We did observe that *ANT1* was upregulated 2-fold in FSHD vs. control myoblasts ($p = 0.006$), although not in myotubes. Other 4q35.2 genes represented on the microarray were *FAT1* (the only 4q35.2 gene with muscle-lineage specific DNaseI hypersensitivity sites [24]), *F11*, *CYP4V2*, *MTNR1A*, *ZFP42*, *TRIML1*, and *TRIML2*. They showed no differences in RNA signal between FSHD and control myogenic cells. In addition, the muscular atrophy-associated gene *FBXO32* (atrogin-1) showed no significant change in FSHD vs. control myogenic cells. Another atrophy-associated gene *TRIM63* (*MURF1*) had decreased, not increased, RNA signal in FSHD myotubes vs. control myotubes (FC = -2.3, $p < 0.0001$). No significant FSHD-related changes were seen for the muscle-differentiation inhibitory *MSTN* (myostatin), *ID1*, *ID2*, or *ID3* genes.

By qRT-PCR, using mostly cDNAs not analyzed on the microarray (Table 1 Figure 4A and 4B, Additional File 3 for sources of cDNA, and Additional File 5 for primers), we validated twelve sets of microarray-

determined FSHD-associated differences in expression. Some of these genes were part of functionally related sets that showed similar FSHD-related changes in expression in the microarray data. For example, we quantified *MYOM1* expression by qRT-PCR (Table 1), and, according to the microarray data, the muscle-associated myomesin genes, *MYOM1*, *MYOM2*, and *MYOM3* were all downregulated in expression in FSHD vs. control myotubes (FC = - 8.9, -8.1, and -3.8, respectively, each $p < 0.0005$). Nonetheless, these genes still had significantly more expression in FSHD myotubes than in the non-muscle cell samples (FC = 3.2, 1.5, and 3.8, respectively, each $p < 0.0001$).


The gene chosen for normalization of the qRT-PCR data was *M6PR* on the basis of its essentially identical average expression levels in control and FSHD myoblasts and myotubes, as determined from the microarray expression data. The often-used *GAPDH*, *PPIA*, and *B2M* genes were not optimal standards because of difficulty in finding unique primer-pairs due to closely related genes or pseudogenes elsewhere in the genome (*GAPDH* and *PPIA*) or FSHD-linked differences in expression levels (*B2M* and several *GAPDH*-related glycolytic enzymes). Our finding that ~800 genes showed more than 2-fold differences in expression in FSHD vs. control myotubes ($p < 0.01$) and many more showed significant differences that were less than 2-fold indicates the importance of using expression profiling data to choose gene standards for qRT-PCR that will not give artifacts in determining relative expression levels of test genes.

Table 1 Validation by qRT-PCR of FSHD dysregulation of gene expression

Gene	Microarray profiling		qRT-PCR		No. of FSH samples	No. of Ctl samples
	FSH/Ctl FC ^b	p-value	FSH/Ctl FC ^b	p-value		
Myoblasts						
<i>PCOLCE</i> ^a	5.9	4E-05	3.5	0.04	5	6
<i>RUVBL2</i>	2.2	5E-04	1.7	0.01	7	8
<i>TGFB2</i>	-2.8	4E-04	-1.9	0.02	9	8
<i>EIF2C2</i>	-1.6	0.007	-1.6	0.03	8	8
Myotubes						
<i>BGN</i>	11.0	< 1E-06	16.8	0.04	8	9
<i>DCN</i>	6.4	6E-04	7.5	0.01	8	9
<i>PCOLCE</i>	10.2	< 1E-06	3.7	0.03	6	7
<i>MYOM1</i>	-8.9	2E-06	-2.9	0.05	4	4
<i>SFRS9</i>	3.5	< 1E-06	3.5	0.01	6	6
<i>CAPG</i>	3.0	0.002	3.2	0.02	8	9
<i>IGFBP6</i>	2.5	0.02	3.5	0.01	8	9
<i>FRAS1</i>	-3.02	0.001	-2.2	0.001	4	4

^aFull gene names and primer-pairs are given in Additional File 5 TableS4

^bFC, fold change in RNA signal was determined by exon microarray analysis or qRT-PCR



We also assayed relative levels of *DUX4-fl* RNA, the RNA isoform associated with FSHD, using very high amounts of cDNA as template, namely, 400 ng, as per the method of Lemmers et al. [7]. We could detect *DUX4-fl* RNA in some of the FSHD myotube and myoblast samples (Figure 4C and 4D). None of the control myoblasts and myotubes gave appreciable signal. The Ct values from positive FSHD myoblasts and myotubes were usually much higher, namely, 33-36, than for the other tested amplicons with the same cDNAs despite the use of 10-20 times more cDNA for the *DUX4* RT-PCR. We do not have microarray results for comparison to the *DUX4* RT-PCR data because *DUX4* is not represented on the microarray due to its extensive cross-hybridization in the genome [40].

Dampening of muscle-specific transcription changes in FSHD vs. control myogenic cells

Downregulated expression in FSHD vs. control myogenic cells was associated with upregulated expression in control myogenic cells vs. non-muscle cell types ($p < 0.01$; Figure 5). The analogous inverse relationship was seen for genes upregulated in FSHD vs. control myogenic cells. Often, the downregulation in FSHD myoblasts or myotubes relative to the analogous control cells was a dampening of myogenesis-linked expression increases, rather than the absence of these normal myogenesis-related expression changes (See Additional File 6 and Additional File 7 for fitted regression analysis). This dampening of myogenesis-specific expression is evidenced by the finding that a much higher percentage of genes showed myogenesis specificity




Figure 5 Inverse relationships between myoblast- and myotube-specific gene expression changes and FSHD-specific changes. **A.** Genes up- or downregulated in FSHD vs. control myoblasts often showed opposite changes in expression in control myoblasts vs. 19 non-muscle cell types. This heat map of log₂ expression ratios for genes upregulated > 2-fold ($p < 0.01$) in FSHD vs. control myoblasts illustrates that these genes were often downregulated in control myoblasts vs. diverse non-muscle cell cultures and that there was the converse relationship for genes downregulated in FSHD vs. control myoblasts and upregulated in control myoblasts vs. non-muscle cell cultures. Both associations were significant ($p < 0.01$). **B.** This heat map is analogous to that of Panel **A** except that it is for myotubes instead of myoblasts.

in FSHD myogenic cells vs. non-muscle cell types when the expression differences had to meet the criterion of $p < 0.01$ but no fold-change threshold of 2.0 was set as for the control myogenic cells (Figure 3B vs. 3A). Examples of genes with dampening of myogenesis-specific expression changes in FSHD cells are shown in Figure 6B. However, other myogenesis-associated genes, e.g., the critical myogenesis-specific transcription factors *MYOD1* and *MYOG*, displayed no change in RNA levels in FSHD vs. control myoblasts or myotubes (Figure 6A). Some genes were dysregulated in FSHD myoblasts or myotubes but did not

exhibit inverse expression changes in myogenic vs. non-muscle cells. For example, 5% of the genes upregulated in FSHD vs. control myotubes ($FC > 2$, $p < 0.01$) were also upregulated in control (and FSHD) myotubes vs. non-muscle cell types. In this group there was an overrepresentation of genes associated with inflammation or encoding extracellular proteins.

Transcriptional dysregulation in FSHD

Some transcription regulatory genes were significantly dysregulated in FSHD vs. control myogenic cells (Figure

Gene	Gene description	Cell type	Control myogenesis		FSHD myogenesis		FSHD-specificity	
			Mb or Mt/Non-muscle cells		Mb or Mt/Non-muscle cells		FSH vs. Ctl Mb or Mt	
			Fold change	P-value	Fold change	P-value	Fold change	P-value
A. Examples of muscle-associated genes without significant dysregulation of expression in FSHD myogenic cells								
<i>MYOD1</i>	myogenic differentiation 1	Mb	12	<1E-06	15	<1E-06	1.3	0.42
<i>MYOG</i>	myogenin (myogenic factor 4)	Mt	59	<1E-06	56	<1E-06	-1.1	0.94
<i>MYF5</i>	myogenic factor 5	Mb	255	<1E-06	233	<1E-06	-1.1	0.84
<i>MYF6</i>	myogenic factor 6 (herculin)	Mb	19	<1E-06	21	<1E-06	1.1	0.92
<i>CHRNA1</i>	cholinergic receptor, nicotinic, alpha 1 (muscle)	Mt	131	<1E-06	115	<1E-06	-1.1	0.60
<i>KBTBD10</i>	kelch repeat and BTB (POZ) domain containing 10	Mt	336	<1E-06	243	<1E-06	-1.4	0.27
<i>TTN</i>	titin	Mt	590	<1E-06	304	<1E-06	-1.9	0.31
B. Examples of muscle-associated genes with significant downregulation of expression in FSHD myogenic cells								
<i>CLCN4</i>	chloride channel 4	Mt	5.2	<1E-06	1.6	0.003	-3.1	5E-05
<i>MYPN</i>	myopalladin	Mt	40	<1E-06	9.0	<1E-06	-4.4	8E-05
<i>SGCG</i>	sarcoglycan, gamma	Mt	65	<1E-06	12	<1E-06	-5.4	0.002
<i>SRL</i>	sarcalumenin	Mt	83	<1E-06	15	<1E-06	-5.7	0.002
<i>MYL1</i>	myosin, light chain 1, alkali; skeletal, fast	Mt	112	<1E-06	19	<1E-06	-5.9	0.006
<i>NRAP</i>	nebulin-related anchoring protein	Mt	4.4	<1E-06	-1.4	0.08	-6.0	<1E-06
<i>MYH7</i>	myosin, heavy chain 7, cardiac muscle, beta	Mt	48	<1E-06	6.1	<1E-06	-7.9	5E-05
<i>XIRP2</i>	xin actin-binding repeat containing 2	Mt	35	<1E-06	4.0	3E-05	-8.8	4E-05
<i>MYOT</i>	myotilin	Mt	21	<1E-06	1.8	0.001	-11.7	<1E-06
<i>ATP1B4</i>	ATPase, Na ⁺ /K ⁺ transporting, beta 4 polypeptide	Mt	103	<1E-06	8.5	<1E-06	-12.2	<1E-06
<i>MYBPC1</i>	myosin binding protein C, slow type	Mt	100	<1E-06	4.7	<1E-06	-21.3	<1E-06

Figure 6 Some muscle-associated genes are downregulated in FSHD myogenic cells while others are not. Fold changes in RNA signal were determined by identical expression profiling of FSHD and control myoblast and myotube preparations and 19 diverse non-muscle cell types (See Additional File 1 and Additional File 2 for detailed descriptions of samples). All *p*-values for microarray data were adjusted for multiple comparisons; pink, significantly upregulated at *p* < 0.01; green, significantly downregulated at *p* < 0.01. Myotube (Mt) data are shown for the genes with much stronger expression at the myotube stage than at the myoblast stage. Myoblast data are given for *MYOD1*, *MYF5*, and *MYF6*.

7). *MEF2A*, *RB1*, *MKL2*, and *KLHL31*, which are associated with transcription control during myogenesis [41,42], were downregulated in FSHD vs. control myotubes. FSHD myotubes relative to those of controls displayed upregulation of *JUNB* and *CREB3L1*, which are transcription factors with protective roles for muscle or cellular stress [43,44]. In FSHD vs. control myoblasts, *DPY30*, *RUVBL2*, *DRAP1*, *PMF1*, *HMGN3*, and *LMO3* were among the genes involved in the control of transcription that were significantly dysregulated.

To obtain additional evidence for dysregulation of steady-state RNA levels in FSHD being governed, in part, by differential activity of transcription factors, we analyzed the distribution of predicted transcription factor binding sites (TFBS) among the promoter regions of a set of 826 genes. These were approximately equally divided between genes that were upregulated and those that were downregulated in FSHD vs. control myotubes. Four of the 126 analyzed TFBS motifs were significantly skewed toward either FSHD up- or downregulation (*p* < 0.01; Table 2). *MEF2A* and *E4BP4* DNA-binding motifs were enriched among FSHD-downregulated genes, which is consistent with the downregulation of *MEF2A*

RNA in FSHD vs. control myotubes (Figure 7) and its role in myogenesis and the role of *E4BP4/NFIL3* as a transcription repressor. The motifs for *TP53*, which is associated with inflammation, and *PPARG*, which is associated with stress response and lipid metabolism, were significantly correlated with genes that were upregulated in FSHD vs. control myotubes. There was no change in *TP53* RNA levels and only a ~1.5-fold increase (*p* = 0.02) in *PPARG* RNA levels; however, expression of the protein products, their modification, and interactions might have changed in FSHD cells.

Evidence for post-transcriptional dysregulation of RNA levels in FSHD

Genes encoding the components of the RNA-induced silencing (RNAi) machinery, including all four Argonaute genes, were significantly downregulated in FSHD vs. control myotubes and two of them were also downregulated in FSHD vs. control myoblasts (Figure 7). The downregulation of *EIF2C2* (Argonaute 2) in FSHD myoblasts was confirmed by qRT-PCR (Figure 4A). Surprisingly, these genes showed significantly higher expression in control myogenic cells vs. 19 non-muscle cell types

Gene	Gene description	Fold change					
		FSH/Ctl		Ctl/non-muscle		FSH/non-muscle	
		Myob	Myot	Myob	Myot	Myob	Myot
Transcription factors, chromatin modification and remodeling proteins							
<i>KLHL31</i>	Kelch-like 31(Drosophila)	-1.1	<i>-7.1</i>	1.9	<i>23.5</i>	1.7	<i>3.3</i>
<i>LMO3</i>	LIM domain only 3	<i>-9.6</i>	<i>-5.6</i>	<i>33.6</i>	4.1	3.5	<i>-1.4</i>
<i>MYF6</i>	myogenic factor 6 (herculin)	1.1	<i>-3.3</i>	<i>19.5</i>	<i>24.9</i>	<i>20.9</i>	<i>7.5</i>
<i>MEF2A</i>	Myocyte enhancer factor 2A	-1.1	<i>-2.3</i>	2.5	7.5	2.3	<i>3.3</i>
<i>RB1</i>	Retinoblastoma 1	-1.4	<i>-2.1</i>	2.6	<i>3.2</i>	1.9	<i>1.5</i>
<i>MKL2</i>	MKL/myocardin-like 2	-1.1	<i>-1.9</i>	1.4	<i>2.1</i>	1.2	1.1
<i>DPY30</i>	Dpy-30 homlog	<i>3.2</i>	<i>3.5</i>	<i>-2.1</i>	<i>-2.8</i>	1.5	1.3
<i>CREB3L1</i>	cAMP responsive element binding 3L1	1.7	2.7	1.4	1.6	2.4	<i>4.3</i>
<i>RUVBL2</i>	RuvB-like 2 (E. coli)	<i>2.2</i>	<i>2.2</i>	<i>-1.9</i>	<i>-3.1</i>	1.2	<i>-1.4</i>
<i>MEIS2</i>	Meis homobox 2	1.6	2.1	<i>-2.1</i>	<i>-2.3</i>	-1.3	<i>-1.1</i>
<i>DRAP1</i>	DR1-associated protein 1	<i>2.7</i>	2.1	<i>-1.8</i>	<i>-2.6</i>	1.5	<i>-1.3</i>
<i>JUNB</i>	Jun B proto-oncogene	1.2	<i>1.9</i>	<i>-1.5</i>	<i>-1.3</i>	<i>-1.2</i>	<i>1.4</i>
<i>HMGN3</i>	HMG nucleosomal binding domain 3	<i>2.6</i>	1.8	<i>-1.7</i>	1.5	1.5	1.3
<i>VPS72</i>	Vacuolar protein sorting 72 homlog	<i>2.2</i>	1.8	<i>-2.3</i>	<i>-1.8</i>	1.0	1.0
<i>PMF1</i>	Polyamine-modulated factor 1	1.9	<i>1.6</i>	<i>-1.5</i>	<i>-1.6</i>	<i>1.3</i>	1.0
RNAi machinery							
<i>EIF2C1</i>	Euk. transln. initiation factor 2C, 1	-1.1	<i>-1.5</i>	1.2	<i>1.4</i>	1.1	<i>-1.1</i>
<i>EIF2C2</i>	Euk. transln. initiation factor 2C, 2	<i>-1.6</i>	<i>-1.8</i>	1.9	<i>2.0</i>	1.2	1.1
<i>EIF2C3</i>	Euk. transln. initiation factor 2C, 3	-1.5	<i>-1.6</i>	<i>2.4</i>	<i>2.5</i>	<i>1.6</i>	<i>1.6</i>
<i>EIF2C4</i>	Euk. transln. initiation factor 2C, 4	-1.1	<i>-2.3</i>	1.4	<i>3.2</i>	1.3	1.4
<i>DICER</i>	Dicer 1, ribonuclease type III	<i>-2.1</i>	<i>-1.9</i>	<i>2.2</i>	<i>1.9</i>	1.0	1.0
<i>DDX6</i>	DEAD box polypeptide6	-1.5	<i>-1.7</i>	1.6	<i>1.7</i>	1.1	1.0
<i>TNRC6B</i>	Trinucleotide repeat containing 6B	-1.1	<i>-1.6</i>	<i>1.7</i>	<i>2.2</i>	<i>1.5</i>	<i>1.4</i>

Figure 7 Transcription control and RNAi machinery genes dysregulated in FSHD cells. Pink or green highlighting indicates significant up- or downregulation, respectively, for the indicated comparison at $p < 0.01$; italics, $p < 0.001$. The FSHD/Ctl difference in expression for *MEIS2* was the exception with $p = 0.02$. *KLHL31*, *RUVBL2*, and *EIF2C2* had been tested and validated for FSHD-associated dysregulation by qRT-PCR (Table 1).

Table 2 TFBS motifs associated preferentially with FSHD up- vs. downregulated genes in myotubes.

TFBS motif	No. of genes with the motif from 826 promoter regions ^a			<i>p</i> -value: skewing of up- or dnreg.	Prominent functional annotation ^b	
	FSH upreg.	FSH dnreg.	Upreg./dnreg.		FSH upreg.	FSH dnreg.
MEF2A	39	101	0.4	< 0.0001	PI3K signaling, 8 genes	muscle phenotype, 16 genes
E4BP4	16	42	0.4	0.008	cell adhesion, 5 genes, GO:0007155	none
PPARG	100	43	2.3	0.0001	extracellular matrix, 13 genes, GO:0031012; arachidonic acid, 15 genes	cardiac hypertrophy, 7 genes
p53	84	37	2.3	0.0009	inflammation, 29 genes; superoxide, 29 genes	Activation of cAMP-dependent PKA, 5 genes

^aPredicted transcription factor binding site motifs (TFBS) in the upstream (promoter) region for 403 and 423 genes that displayed the strongest up- or downregulation ($p < 0.01$), respectively, in FSHD vs. control myotubes. These sites were identified by GeneCards <http://genecards.org> and were analyzed by Chi-square tests for association with FSHD up- vs. downregulation in myotubes. Of the 126 TFBS scored, the four shown in this table had the strongest association with either FSHD up- or downregulated genes. For PPARG, PPARG1 and PPARG2 motifs were pooled.

^bFunctional annotation from GeneDecks <http://genecards.org>

(Figure 7). In FSHD myogenic cells compared with the non-muscle cell types, these genes were not downregulated; most of them were just not upregulated.

Exon-based microarrays can reveal the presence of cell type-specific differences in RNA isoforms from a given gene by analysis of exon-normalized probe data instead of the gene-normalized data described above. By gel electrophoresis, we examined oligo(dT)₂₃-primed RT-PCR products (0.2 - 0.5 kb) from five of the genes for which the exon-normalized array data suggested FSHD-specific RNA isoforms. None of these representative genes (*FAT1*, *SCUBE3*, *ILF3*, *TFPI2*, or *SFRS7*) was validated as giving the predicted FSHD-specific differences in RNA sizes. In addition, we looked for previously reported FSHD-associated RNA isoforms from *FXR1P*, *TNNT3*, and *MTMR1* [17,45] but did not see any evidence for them in our cell populations.

Functional terms associated with genes dysregulated in FSHD myoblasts and myotubes

A bioinformatics analysis was done to look for functional terms associated with genes significantly dysregulated in FSHD myoblasts or myotubes ($p < 0.01$). Functional terms overrepresented among FSHD-upregulated genes (e.g., mitochondrial terms, extracellular matrix, Rho) were mostly very different from those for FSHD-downregulated genes (e.g., myofibril, RNA-induced silencing complex; Table 3). Many of the genes that were upregulated in both FSHD myoblasts and myotubes were related to the response to cell stress, such as *GSTP1*, *HSP90AA1*, *HSP90AB1*, *HSPA1A*, *HSPC152*, and *DNAJC4* (See Additional File 8 for gene lists). However for cell stress genes, as for most of the overrepresented functional terms among FSHD-dysregulated genes, the up- or downregulation in FSHD vs. control myogenic cells was often associated with down- or upregulation, respectively, in control myogenic cells vs. non-muscle cells. This is illustrated in Additional File 9 for FSHD-dysregulated pro- and anti-apoptosis genes.

Discussion

Our study is the first to examine in parallel many different human cell types and compare them to well-characterized FSHD and control myoblasts and myotubes and thereby demonstrate extensive FSHD-linked dysregulation of gene expression. Importantly, we confirmed by RT-PCR that the disease-associated RNA isoform of *DUX4*, *DUX4-fl* RNA, is expressed at extraordinarily low levels in FSHD (but not detectable in control) myoblasts and myotubes despite the hundreds of genes dysregulated more than 2-fold in FSHD vs. control myoblasts and myotubes. Our findings of extremely infrequent expression of *DUX4-fl* RNA in FSHD


Table 3 Some pathways and functional terms overrepresented among FSHD-dysregulated genes.

Functional terms ^a	Mb or Mt	No. of up- or dnreg. genes	Ratio ^b
RNA-induced silencing complex	Mt	4 dn	0.80
	Mb	1 dn	0.20
Fatty acid elongation in mitochondria	Mb	7 up	0.44
	Mt	2 up	0.13
Extracellular matrix	Mt	35 up, 1 dn	0.30
	Mb	4 up, 2 dn	0.05
NRF2-mediated oxidative stress response	Mt	15 up, 7 dn	0.13
	Mb	7 up	0.04
HIF1 α Signaling	Mt	6 up, 7 dn	0.12
	Mb	5 up	0.05
Regulation of actin-based motility by Rho	Mt	9 up, 3 dn	0.10
	Mb	6 up	0.07
Mitochondrial matrix	Mt	18 up, 5 dn	0.11
	Mb	22 up, 1 dn	0.11
Oxidative phosphorylation	Mt	18 up	0.16
	Mb	11 up	0.09
Induction of apoptosis	Mt	13 up, 5 dn	0.09
	Mb	8 up, 2 dn	0.05
Anti-apoptosis	Mt	16 up, 3 dn	0.10
	Mb	4 up	0.02

^aOverrepresented functional terms for genes up- or downregulated ($p < 0.01$) in FSHD vs. control myoblasts (Mb) or myotubes (Mt) were identified by bioinformatics programs. Details and the names of the dysregulated genes are given in Additional File 8 Table S5.

^bThe number of genes for a given pathway or GO-term that were up- or downregulated in FSHD vs. control myogenic cells/the total number of genes for that pathway or GO-term that were included in the microarray.

myoblasts and myotubes and undetectable levels in some of these FSHD cell populations are consistent with previous reports. This transcript was detected in only ~1 out of 1000 FSHD myotube nuclei and was observed less frequently in FSHD myoblasts than in myotubes [8]. Similarly, its detection by one round of real-time PCR required much higher-than-normal amounts of FSHD myotube cDNA [7]. Because FSHD myoblasts and myotubes had a strong transcription dysregulation profile (this study) and FSHD myoblasts are hypersensitive to oxidative stress [14,15], if *DUX4* is the first pathologically dysregulated gene in FSHD, then it must be expressed much more extensively, but transiently, at an earlier stage in myogenesis. We propose that *DUX4-fl* RNA initiates a cascade of gene dysregulation at or before activation of FSHD satellite cells to form myoblasts (Figure 8). *DUX4-fl* transcripts in myotubes and myoblasts would then represent a rare re-activation of inappropriate *DUX4* expression that is not central to pathogenesis. This model contrasts with the current



emphasis on the rare expression of *DUX4-fl* RNA in myoblasts and myotubes.

Induction of *DUX4-fl* transcription in transduced C2C12 myoblasts caused apoptosis; inhibited differentiation to myotubes; gave dramatic changes in cell shape; and, even at sublethal concentrations, inhibited transcription of *MYOD1* and *MYOG* and increased that of *CDKN1A* (p21) [27]. In addition, *DUX4-fl* RNA injected into *Xenopus* or zebrafish embryos is highly cytotoxic [21,46]. In contrast, we found only minimal apoptosis in FSHD myoblasts and myotubes and no more than in controls (~5% of nuclei as detected by staining with ethidium bromide and acridine orange [47] and no internucleosomal fragmentation of DNA [48], data not shown). In our comparison of FSHD and control

myotubes, more anti-apoptotic than pro-apoptotic changes in gene expression were seen and the dysregulation of most of these genes can be explained by FSHD-associated changes in the normal myogenesis program (See Additional File 9). The good growth and efficient differentiation to myotubes of FSHD myoblasts observed by us and others [15] are consistent with our finding that the myogenic regulatory factors *MYOD1*, *MYOG*, *MYF5*, and *MYF6* were equally highly expressed at the RNA level in FSHD vs. control myoblasts and only *MYF6* was significantly downregulated in FSHD vs. control myotubes. This result argues against FSHD-related differences in posttranscriptional processing of the products of these four genes in myoblasts because their expression is autoregulatory [49-52]. In addition, we

observed no differences between FSHD and control myogenic cells in RNA levels for E-box protein heterodimer partners of these myogenic regulatory factors (data not shown). In support of our model of the proposed non-cytotoxic expression of *DUX4* very early in myogenesis (Figure 8), the early-myogenesis transcription factors *PAX3* and *PAX7* can partly counteract the deleterious effects of overexpression of *DUX4* in C2C12 cells [27]. Moreover, *DUX4-fl* transcripts are normally rather abundant in testis [8].

Previously used arrays for transcription profiling of well-characterized FSHD and control myogenic samples [10-12,14] did not have multiple probe-sets per exon for each transcript nor probes for exons other than the 3' exon and so are much less representative of the transcript populations. Given this major difference and the use of muscle tissue rather than myoblasts (a very minor component of muscle) in most of the previously published FSHD RNA profiling studies, it is not surprising that there was only minimal overlap between genes reported as dysregulated in FSHD in previous studies and genes that we observed to be dysregulated in FSHD vs. control myogenic precursors. An example of such infrequent overlap is the upregulation of the vascular smooth muscle-associated *CTGF*, *ENG*, and *TAGLN* genes in FSHD vs. control muscle [12] and, in this study, in FSHD vs. control myotubes (2- to 3-fold upregulation, $p < 0.001$ for all comparisons). In the previous expression profiling of well-characterized FSHD and control myoblasts, a "vacuolar/necrotic phenotype" was noted for "the majority" of FSHD myoblasts, which were "markedly swollen." That morphological phenotype might be due to FSHD myoblasts being more sensitive to stress than analogous controls. Because we used only moderately affected muscle to generate myoblast cell strains and FSHD is characteristically a slowly progressive disease in which disease muscle biopsies look relatively normal at the time of clinical onset [53], the normal appearance of FSHD myoblasts and myotubes under our optimized growth conditions is likely to be relevant to understanding pathogenesis. Moreover, the equally good generation, propagation, growth, and differentiation of FSHD and control myoblasts also argue against the possibility that we selected a non-representative sub-phenotype of FSHD myoblasts. Similarly, these observations fit the high degree of correlation of the overall expression profiles of all the FSHD and control myogenic samples with each other when compared to the 19 non-muscle cell populations despite the hundreds of significant more-than-two-fold differences in RNA signal for individual genes in comparisons of FSHD and control myoblasts or myotubes.

There was a recent report by Cheli et al. [16] about exon array-based expression profiling of FSHD and

control myoblasts and myotubes but it included no characterization of the percentage myoblasts in the studied cell populations nor the efficiency of differentiation to myotubes. The extent of contamination of untransformed myoblast cultures with non-muscle cells can vary dramatically between different myoblast cell strains and even at different passage numbers and, thereby, have a major impact on expression profiling. Cheli et al. reported < 4% overlap between several hundred genes with dysregulation in FSHD vs. control cells at the myoblast stage and those dysregulated at the myotube stage, unlike the present study in which we found 48% overlap between genes with FSHD dysregulation ($p < 0.01$, $FC > 2$) in myoblasts and myotubes. The difference in the results from the study of Cheli et al. and the present study might be due to the unknown percentage of cells that differentiated in their experiment vs. 72-80% in ours.

Our analysis of normal myogenesis from a comparison of expression profiles of control myoblasts and myotubes and 19 non-muscle cell types indicated the prominent role of upregulation of genes generally involved in the actin cytoskeleton, organization of the extracellular matrix, cell adhesion, and GTPase regulator activity, in addition to the expected muscle-specific genes. One unexpected functional class of genes that was more highly expressed in control myoblasts and myotubes than in non-muscle cells was the RNA silencing machinery genes. These same RNA silencing machinery genes were lacking upregulation during FSHD myogenesis, which may contribute to the observed excess of FSHD-upregulated vs. FSHD-downregulated genes in myogenic precursors.

Our expression profiling of FSHD vs. control myoblasts suggests an explanation for the FSHD myoblasts' hypersensitivity to external oxidative stress [14,15] as well as an imbalance in the redox system in muscle [54]. The observed FSHD-related upregulation of many transcripts from oxidative phosphorylation genes could result in an increase in endogenous reactive oxygen species and might eventually result in apoptosis in some severely affected muscles. Accordingly, increases in H_2O_2 were seen in FSHD vs. control muscle [54] and upregulation of some mitochondrial oxidative phosphorylation proteins and oxidative stress-response proteins was observed in affected and also in unaffected FSHD skeletal muscle vs. normal-control muscle [20]. Similarly, we found FSHD-associated upregulation of RNA for oxidative stress-response and oxidative phosphorylation proteins, including several of the same proteins (*SOD1* and *HSPB1*) whose FSHD-upregulation was seen in muscle [20]. The hypothesized inappropriate expression of *DUX4* very early during regenerative myogenesis would help explain why even unaffected muscle showed

these abnormalities in protein levels [20] and why myoblasts from unaffected FSHD muscle samples displayed an FSHD-associated hypersensitivity to oxidative stress [15]. Moreover, it would be consistent with our finding that myoblast cell strains from diverse, moderately affected FSHD muscle samples displayed FSHD-related changes in gene expression.

The most prominent feature of the transcription dysregulation in FSHD myoblasts and myotubes was the decrease in the up- and downregulation of RNA levels associated with normal myogenesis, which can account for most of the FSHD-related dysregulation. Expression profiling of other muscular dystrophies [55-58] has not revealed such a widespread dampening of normal myogenesis-associated transcription changes in various functional gene categories. Some classes of genes, including those encoding extracellular matrix or pro-inflammatory proteins, were strongly enriched in FSHD-upregulation in myogenic cells independent of any inverse myogenesis association. The proliferation of FSHD myoblasts and their differentiation to myotubes *in vitro* was unaffected by these changes in gene expression. Apparently, there is also not a large disease-related depletion of satellite cells in FSHD patients because of the above-mentioned finding that generating myoblast cell strains from moderately affected muscle biopsies of FSHD patients was no more difficult than from control muscle. Moreover, although Reed et al. [59] observed abnormal spatial relationships of the sarcolemma with the underlying contractile apparatus in affected FSHD muscle, the structure of the contractile apparatus itself appeared normal. The observed FSHD-associated gene dysregulation may have been heightened in the FSHD myoblasts and myotubes relative to their *in-vivo* counterparts due to the effects of cell culture and the use of the myoblast-stimulatory [37,38] dexamethasone in the culture medium for both FSHD and control myoblasts [60,61]. If extensive cell culture promoted increased gene dysregulation in FSHD myoblasts, this could be relevant to the disease *in vivo* because usually it is only slowly progressive. In addition, glucocorticoids are relevant *in vivo* because of the effects of endogenous glucocorticoids in traumatic or muscle wasting conditions [62,63] and the therapeutic use of glucocorticoids. Moreover, if only a small fraction of the extent of gene dysregulation that we saw exists *in vivo*, this could lead to atrophy by interfering with effective regenerative myogenesis. For example, all three of the skeletal muscle-associated myomesin genes showed downregulation in FSHD vs. control myotubes of about 4 to 9 fold, and the products of these genes bind to other muscle structural proteins in a dose-dependent manner as major components of the myofibrillar M-band.

It is not yet clear whether the dysregulated gene expression in FSHD myoblasts is due to disease-related differences in transcription regulatory or RNA-processing proteins, cell signaling (e.g., TGF β or RHO/mTOR pathways [64,65]), indirect effects on transcription from overexpression of extracellular proteins [66], indirect effects of mitochondrial dysfunction [67], subtle differences in timing of expression of some myogenesis-specific gene(s), and/or disease-specific epigenetic differences. At the myotube stage, the increase in the number of genes that were dysregulated in FSHD cells may be partly due to the FSHD-associated decreases in expression of the transcription regulatory *MYF6* and *MEF2A* genes and abnormal increases in expression of *MEIS2* after induction of differentiation to myotubes.

Conclusions

Given the extremely low rate of FSHD-associated inappropriate expression of *DUX4* at the myoblast, myotube, and muscle stages, many of the FSHD-dysregulated transcription-regulatory or cell signaling genes revealed by our expression profiling may be more effective targets for developing pharmacologically-based or gene therapy-based treatment of FSHD than *DUX4* itself. Our findings point to FSHD being a differentiation-associated disease, and so study of this enigmatic muscular dystrophy is likely to elucidate new aspects of normal myogenesis as well as pathogenesis. In addition, our comparison of transcription profiles of control myoblasts and myotubes and those of 19 other cell types that were examined identically showed how very extensive gene expression changes are upon formation of myoblasts and upon their differentiation to myotubes.

Additional material

Additional file 1: Table S1. Description of myoblast and myotube samples for microarray analysis

Additional file 2: Table S2. Descriptions of non-muscle cell types for expression profiling

Additional file 3: Table S3. Description of myoblast and myotube samples for qRT-PCR

Additional file 4: Additional Methods: Generation, propagation, and differentiation of myoblast cultures from human skeletal muscle.

Additional file 5: Table S4. qPCR primers and full names of genes tested.

Additional file 6: Figure S1. FSHD-downregulated genes: relationship between gene expression in FSHD myotubes vs. non-muscle cell types to that in control myotubes vs. non-muscle cell types.

Additional file 7: Figure S2. FSHD-upregulated genes: relationship between gene expression in FSHD myotubes vs. non-muscle cell types to that in control myotubes vs. non-muscle cell types.

Additional file 8: Table S5. Some pathways and functional terms overrepresented among FSHD-dysregulated genes.

Additional file 9: Table S6. Most of the pro- or anti-apoptotic genes that were dysregulated in FSHD vs. control myotubes were apparently

up- or downregulated in FSHD myotubes because of dampening of their normal myogenesis-associated changes in expression.

Acknowledgements

We are grateful to Drs. V. Vedanarayanan and Anthony Romeo for several of the muscle samples from which myoblast cell strains were generated, to Jerry Simbarashe Zifodya and Zhangcheng Zhang for help with bioinformatics analysis, to Lingyun Song for growth of the non-muscle cell types, and to Stephen Hauschka for generously donating the MF20 antibody and for advice on culturing conditions for optimal myotube formation. This work was supported by the National Institutes of Health [NS04885 and AA3768G2 to ME, HG003169 to GEC.], the FSHD Global Research Foundation and the FSH Society [ME], and the Fields Center for FSHD and Neuromuscular Research [RT].

Author details

¹Human Genetics Program, Tulane Medical School, New Orleans, LA, USA.

²Department of Mathematics, Tulane University, New Orleans, LA, USA.

³Tulane Cancer Center, New Orleans, LA, USA. ⁴Biomedical Sciences, University at Albany-SUNY, Albany, NY, USA. ⁵Department of Neurology, University of Rochester School of Medicine and Dentistry, Rochester, NY, USA. ⁶Institute for Genome Sciences & Policy, Duke University, Durham, NC, USA.

Authors' contributions

KT prepared and characterized the myoblasts and myotubes and prepared the RNA and cDNA for qRT-PCR. S-CC did the qRT-PCR analyses. SVC did the hybridization and generated the initial data sets from the microarrays. ML did the statistical analyses of the expression profiling data. CB and ME did the analyses of functional categories of dysregulated genes. RB obtained the muscle biopsies for most of the myoblast cell cultures and JS generated most of the myoblast cell strains from them. GEC and SVC did the expression profiling of the 19 non-muscle cell types. ME wrote the MS. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 19 July 2011 Accepted: 27 September 2011

Published: 27 September 2011

References

- Bortoli S, Renault V, Eveno E, Auffray C, Butler-Browne G, Pietu G: **Gene expression profiling of human satellite cells during muscular aging using cDNA arrays.** *Gene* 2003, **321**:145-154.
- Sterrenburg E, Turk R, t Hoen PA, van Deutekom JC, Boer JM, van Ommen GJ, den Dunnen JT: **Large-scale gene expression analysis of human skeletal myoblast differentiation.** *Neuromuscul Disord* 2004, **14**(8-9):507-518.
- Raymond F, Metairon S, Kussmann M, Colomer J, Nascimento A, Mormeneo E, Garcia-Martinez C, Gomez-Foix AM: **Comparative gene expression profiling between human cultured myotubes and skeletal muscle tissue.** *BMC Genomics* 2010, **11**:125.
- Wang J, Conboy I: **Embryonic vs. adult myogenesis: challenging the 'regeneration recapitulates development' paradigm.** *J Mol Cell Biol* 2010, **2**(1):1-4.
- Yokoyama S, Asahara H: **The myogenic transcriptional network.** *Cell Mol Life Sci* 2011, **68**(11):1843-1849.
- Dixit M, Anseau E, Tassin A, Winokur S, Shi R, Qian H, Sauvage S, Matteotti C, van Acker AM, Leo O, et al: **DUX4, a candidate gene of facioscapulohumeral muscular dystrophy, encodes a transcriptional activator of PITX1.** *Proc Natl Acad Sci USA* 2007, **104**(46):18157-18162.
- Lemmers RJ, van der Vliet PJ, Klooster R, Sacconi S, Camano P, Dauwerse JG, Snider L, Straasheijm KR, Jan van Ommen G, Padberg GW, et al: **A unifying genetic model for facioscapulohumeral muscular dystrophy.** *Science* 2010, **329**:1650-1653.
- Snider L, Geng LN, Lemmers RJ, Kyba M, Ware CB, Nelson AM, Tawil R, Filippova GN, van der Maarel SM, Tapscott SJ, et al: **Facioscapulohumeral dystrophy: incomplete suppression of a retrotransposed gene.** *PLoS Genet* 2010, **6**(10):e1001181.
- Tawil R: **Facioscapulohumeral muscular dystrophy.** *Neurotherapeutics* 2008, **5**(4):601-606.
- Winokur ST, Chen YW, Masny PS, Martin JH, Ehmsen JT, Tapscott SJ, van der Maarel SM, Hayashi Y, Flanigan KM: **Expression profiling of FSHD muscle supports a defect in specific stages of myogenic differentiation.** *Hum Mol Genet* 2003, **12**(22):2895-2907.
- Celegato B, Capitanio D, Pescatori M, Romualdi C, Pacchioni B, Cagnin S, Viganò A, Colantoni L, Begum S, Ricci E, et al: **Parallel protein and transcript profiles of FSHD patient muscles correlate to the D4Z4 arrangement and reveal a common impairment of slow to fast fibre differentiation and a general deregulation of MyoD-dependent genes.** *Proteomics* 2006, **6**(19):5303-5321.
- Osborne RJ, Welle S, Venance SL, Thornton CA, Tawil R: **Expression profile of FSHD supports a link between retinal vasculopathy and muscular dystrophy.** *Neurology* 2007, **68**(8):569-577.
- Eisenberg I, Eran A, Nishino I, Moggio M, Lamperti C, Amato AA, Lidov HG, Kang PB, North KN, Mitrani-Rosenbaum S, et al: **Distinctive patterns of microRNA expression in primary muscular disorders.** *Proc Natl Acad Sci USA* 2007, **104**(43):17016-17021.
- Winokur ST, Barrett K, Martin JH, Forrester JR, Simon M, Tawil R, Chung SA, Masny PS, Figlewicz DA: **Facioscapulohumeral muscular dystrophy (FSHD) myoblasts demonstrate increased susceptibility to oxidative stress.** *Neuromuscul Disord* 2003, **13**:322-333.
- Barro M, Carnac G, Flavie S, Mercier J, Vassetzky Y, Laoudj-Chenivresse D: **Myoblasts from affected and non affected FSHD muscles exhibit morphological differentiation defects.** *J Cell Mol Med* 2010, **275**:289.
- Cheli S, Francois S, Bodega B, Ferrari F, Tenedini E, Roncaglia E, Ferrari S, Ginelli E, Meneveri R: **Expression profiling of FSHD-1 and FSHD-2 cells during myogenic differentiation evidences common and distinctive gene dysregulation patterns.** *PLoS one* 2011, **6**(6):e20966.
- Gabellini D, D'Antona G, Moggio M, Prella A, Zecca C, Adami R, Angeletti B, Ciscato P, Pellegrino MA, Bottinelli R, et al: **Facioscapulohumeral muscular dystrophy in mice overexpressing FRG1.** *Nature* 2006, **439**(7079):973-977.
- Wright TJ, Wijmenga C, Clark LN, Frants RR, Williamson R, Hewitt JE: **Fine mapping of the FSHD gene region orientates the rearranged fragment detected by the probe p13E-11.** *Hum Mol Genet* 1993, **2**(10):1673-1678.
- Bosnakovski D, Lamb S, Simsek T, Xu Z, Belayew A, Perlingeiro R, Kyba M: **DUX4c, an FSHD candidate gene, interferes with myogenic regulators and abolishes myoblast differentiation.** *Exp Neurol* 2008, **214**(1):87-96.
- Laoudj-Chenivresse D, Carnac G, Bisbal C, Hugon G, Bouillot S, Desnuelle C, Vassetzky Y, Fernandez A: **Increased levels of adenine nucleotide translocator 1 protein and response to oxidative stress are early events in facioscapulohumeral muscular dystrophy muscle.** *J Mol Med* 2005, **83**(3):216-224.
- Wuebbles RD, Long SW, Hanel ML, Jones PL: **Testing the effects of FSHD candidate gene expression in vertebrate muscle development.** *Int J Clin Exp Pathol* 2010, **3**(4):386-400.
- van Geel M, van Deutekom JC, van Staalduijn A, Lemmers RJ, Dickson MC, Hofker MH, Padberg GW, Hewitt JE, de Jong PJ, Frants RR: **Identification of a novel beta-tubulin subfamily with one member (TUBB4Q) located near the telomere of chromosome region 4q35.** *Cytogenet Cell Genet* 2000, **88**(3-4):316-321.
- Klooster R, Straasheijm K, Shah B, Sowden J, Frants R, Thornton C, Tawil R, van der Maarel S: **Comprehensive expression analysis of FSHD candidate genes at the mRNA and protein level.** *Eur J Hum Genet* 2009, **17**(12):1615-1624.
- Xu X, Tsumagari K, Sowden J, Tawil R, Boyle AP, Song L, Furey TS, Crawford GE, Ehrlich M: **DNase hypersensitivity at gene-poor, FSH dystrophy-linked 4q35.2.** *Nucleic Acids Res* 2009, **37**(22):7381-7393.
- Hewitt JE, Lyle R, Clark LN, Valleley EM, Wright TJ, Wijmenga C, van Deutekom JC, Francis F, Sharpe PT, Hofker M, et al: **Analysis of the tandem repeat locus D4Z4 associated with facioscapulohumeral muscular dystrophy.** *Hum Mol Genet* 1994, **3**(8):1287-1295.
- Winokur ST, Bengtsson U, Feddersen J, Mathews KD, Weiffenbach B, Bailey H, Markovich RP, Murray JC, Wasmuth JC, Altherr MR, et al: **The DNA rearrangement associated with facioscapulohumeral muscular dystrophy involves a heterochromatin-associated repetitive element: implications for a role of chromatin structure in the pathogenesis of the disease.** *Chromosome Res* 1994, **2**(3):225-234.

27. Bosnakovski D, Xu Z, Gang EJ, Galindo CL, Liu M, Simsek T, Garner HR, Agha-Mohammadi S, Tassin A, Coppee F, et al: **An isogenetic myoblast expression screen identifies DUX4-mediated FSHD-associated molecular pathologies.** *EMBO J* 2008, **27**(20):2766-2779.
28. Kowaljow V, Marcowycz A, Anseau E, Conde CB, Sauvage S, Matteotti C, Arias C, Corona ED, Nunez NG, Leo O, et al: **The DUX4 gene at the FSHD1A locus encodes a pro-apoptotic protein.** *Neuromuscul Disord* 2007, **17**:611-623.
29. Wallace LM, Garwick SE, Mei W, Belayew A, Coppee F, Ladner KJ, Guttridge D, Yang J, Harper SQ: **DUX4, a candidate gene for facioscapulohumeral muscular dystrophy, causes p53-dependent myopathy in vivo.** *Ann Neurol* 2010.
30. Zeng W, de Greef JC, Chen YY, Chien R, Kong X, Gregson HC, Winokur ST, Pyle A, Robertson KD, Schmiesing JA, et al: **Specific loss of histone H3 lysine 9 trimethylation and HP1gamma/cohesin binding at D4Z4 repeats is associated with facioscapulohumeral dystrophy (FSHD).** *PLoS Genet* 2009, **5**(7):e1000559.
31. de Greef JC, Lemmers RJ, van Engelen BG, Sacconi S, Venance SL, Frants RR, Tawil R, van der Maarel SM: **Common epigenetic changes of D4Z4 in contraction-dependent and contraction-independent FSHD.** *Hum Mutat* 2009, **30**:1-11.
32. Gabriels J, Beckers MC, Ding H, De Vriese A, Plaisance S, van der Maarel SM, Padberg GW, Frants RR, Hewitt JE, Collen D, et al: **Nucleotide sequence of the partially deleted D4Z4 locus in a patient with FSHD identifies a putative gene within each 3.3 kb element.** *Gene* 1999, **236**(1):25-32.
33. Tsieng F, Sun B, Hopkins NE, Vedanarayanan V, Figlewicz D, Winokur S, Ehrlich M: **Hypermethylation of the FSHD syndrome-linked subtelomeric repeat in normal and FSHD cells but not in ICF syndrome cells.** *Molec Gen Metab* 2001, **74**:322-331.
34. Sandri M, El Meslemani AH, Sandri C, Schjerling P, Vissing K, Andersen JL, Rossini K, Carraro U, Angelini C: **Caspase 3 expression correlates with skeletal muscle apoptosis in Duchenne and facioscapulo human muscular dystrophy. A potential target for pharmacological treatment?** *J Neuropathol Exp Neurol* 2001, **60**(3):302-312.
35. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F: **Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes.** *Genome Biol* 2002, **3**(7).
36. Smyth GK: **Limma: linear models for microarray data.** In *Bioinformatics and Computational Biology Solutions using R and Bioconductor*. Edited by: Gentleman R, Carey VS, Dudoit S, Irizarry R, Huber W. New York: Springer; 2005:397-420.
37. Guerriero V, Florini JR: **Dexamethasone effects on myoblast proliferation and differentiation.** *Endocrinology* 1980, **106**(4):1198-1202.
38. Vilquin JT, Marolleau JP, Sacconi S, Garcin I, Lacassagne MN, Robert I, Ternaux B, Bouazza B, Larghero J, Desnuelle C: **Normal growth and regenerating ability of myoblasts from unaffected muscles of facioscapulohumeral muscular dystrophy patients.** *Gene Ther* 2005, **12**(22):1651-1662.
39. Jiang G, Yang F, van Overveld PG, Vedanarayanan V, van der Maarel S, Ehrlich M: **Testing the position-effect variegation hypothesis for facioscapulohumeral muscular dystrophy by analysis of histone modification and gene expression in subtelomeric 4q.** *Hum Mol Genet* 2003, **12**:2909-2921.
40. Lyle R, Wright TJ, Clark LN, Hewitt JE: **The FSHD-associated repeat, D4Z4, is a member of a dispersed family of homeobox-containing repeats, subsets of which are clustered on the short arms of the acrocentric chromosomes.** *Genomics* 1995, **28**(3):389-397.
41. Blais A, van Oevelen CJ, Margueron R, Acosta-Alvear D, Dynlacht BD: **Retinoblastoma tumor suppressor protein-dependent methylation of histone H3 lysine 27 is associated with irreversible cell cycle exit.** *J Cell Biol* 2007, **179**(7):1399-1412.
42. Abou-Elhamd A, Cooper O, Munsterberg A: **Klhl31 is associated with skeletal myogenesis and its expression is regulated by myogenic signals and Myf-5.** *Mech Dev* 2009, **126**(10):852-862.
43. Raffaello A, Milan G, Masiero E, Carnio S, Lee D, Lanfranchi G, Goldberg AL, Sandri M: **JunB transcription factor maintains skeletal muscle mass and promotes hypertrophy.** *J Cell Biol* 2010, **191**(1):101-113.
44. Vellanki RN, Zhang L, Guney MA, Rocheleau JV, Gannon M, Volchuk A: **OASIS/CREB3L1 induces expression of genes involved in extracellular matrix production but not classical endoplasmic reticulum stress response genes in pancreatic beta-cells.** *Endocrinology* 2010, **151**(9):4146-4157.
45. Davidovic L, Sacconi S, Bechara EG, Delplace S, Allegra M, Desnuelle C, Bardoni B: **Alteration of expression of muscle specific isoforms of the fragile X related protein 1 (FXR1P) in facioscapulohumeral muscular dystrophy patients.** *J Med Genet* 2008, **45**(10):679-685.
46. Snider L, Asawachaicharn A, Tyler AE, Geng LN, Petek LM, Maves L, Miller DG, Lemmers RJ, Winokur ST, Tawil R, et al: **RNA transcripts, miRNA-sized fragments and proteins produced from D4Z4 units: new candidates for the pathophysiology of facioscapulohumeral dystrophy.** *Hum Mol Genet* 2009, **18**(13):2414-2430.
47. Mishell BB, Shiiqi SM, Henry C, Chen EL, North J, Gallily R, Slomich M, Miller K, Marbrook J, Parks D, et al: **Apoptosis: quantitative analysis techniques.** *Selected Methods in Cellular Immunology* San Francisco WH Freeman; 1980, 21-22.
48. Abul-Milth M, Wu Y, Lau B, Lingwood CA, Barnett Foster D: **Induction of epithelial cell death including apoptosis by enteropathogenic Escherichia coli expressing bundle-forming pili.** *Infect Immun* 2001, **69**(12):7356-7364.
49. Berkes CA, Tapscott SJ: **MyoD and the transcriptional control of myogenesis.** *Semin Cell Dev Biol* 2005, **16**(4-5):585-595.
50. Sartorelli V, Caretti G: **Mechanisms underlying the transcriptional regulation of skeletal myogenesis.** *Curr Opin Genet Dev* 2005, **15**(5):528-535.
51. Wilson EM, Rotwein P: **Control of MyoD function during initiation of muscle differentiation by an autocrine signaling pathway activated by insulin-like growth factor-II.** *J Biol Chem* 2006, **281**(40):29962-29971.
52. Zingg JM, Pedraza-Alva G, Jost JP: **MyoD1 promoter autoregulation is mediated by two proximal E-boxes.** *Nucleic Acids Res* 1994, **22**(12):2234-2241.
53. Fitzsimons RB: **Retinal vascular disease and the pathogenesis of facioscapulohumeral muscular dystrophy. A signalling message from Wnt?** *Neuromuscul Disord* 2011, **21**(4):263-271.
54. Macaione V, Aguenouz M, Rodolico C, Mazzeo A, Patti A, Cannistraci E, Colantone L, Di Giorgio RM, De Luca G, Vita G: **RAGE-NF-kappaB pathway activation in response to oxidative stress in facioscapulohumeral muscular dystrophy.** *Acta Neurol Scand* 2007, **115**(2):115-121.
55. Chen YW, Zhao P, Borup R, Hoffman EP: **Expression profiling in the muscular dystrophies: identification of novel aspects of molecular pathophysiology.** *J Cell Biol* 2000, **151**(6):1321-1336.
56. Saenz A, Azpitarte M, Armananzas R, Leturcq F, Alzualde A, Inza I, Garcia-Bragado F, De la Herran G, Corcuera J, Cabello A, et al: **Gene expression profiling in limb-girdle muscular dystrophy 2A.** *PLoS one* 2008, **3**(11):e3750.
57. Sterrenburg E, van der Wees CG, White SJ, Turk R, de Menezes RX, van Ommen GJ, den Dunnen JT, t Hoen PA: **Gene expression profiling highlights defective myogenesis in DMD patients and a possible role for bone morphogenetic protein 4.** *Neurobiol Dis* 2006, **23**(1):228-236.
58. Du H, Cline MS, Osborne RJ, Tuttle DL, Clark TA, Donohue JP, Hall MP, Shiue L, Swanson MS, Thornton CA, et al: **Aberrant alternative splicing and extracellular matrix gene expression in mouse models of myotonic dystrophy.** *Nat Struct Mol Biol* 2011, **17**(2):187-193.
59. Reed P, Porter NC, Strong J, Pumplun DW, Corse AM, Luther PW, Flanigan KM, Bloch RJ: **Sarcolemmal reorganization in facioscapulohumeral muscular dystrophy.** *Ann Neurol* 2006, **59**(2):289-297.
60. Giorgino F, Pedrini MT, Matera L, Smith RJ: **Specific increase in p85alpha expression in response to dexamethasone is associated with inhibition of insulin-like growth factor-I stimulated phosphatidylinositol 3-kinase activity in cultured muscle cells.** *J Biol Chem* 1997, **272**(11):7455-7463.
61. Shah OJ, Kimball SR, Jefferson LS: **Among translational effectors, p70S6k is uniquely sensitive to inhibition by glucocorticoids.** *Biochem J* 2000, **347**(Pt 2):389-397.
62. Hasselgren PO: **Glucocorticoids and muscle catabolism.** *Curr Opin Clin Nutr Metab Care* 1999, **2**(3):201-205.
63. Combaret L, Taillandier D, Dardevet D, Bechet D, Ralliere C, Claustre A, Grizard J, Attaix D: **Glucocorticoids regulate mRNA levels for subunits of the 19 S regulatory complex of the 26 S proteasome in fast-twitch skeletal muscles.** *Biochem J* 2004, **378**(Pt 1):239-246.
64. Lee CH, Hong CH, Yu HS, Chen GS, Yang KC: **Transforming growth factor-beta enhances matrix metalloproteinase-2 expression and activity**

through AKT in fibroblasts derived from angiofibromas in patients with tuberous sclerosis complex. *Br J Dermatol* 2010, **163**(6):1238-1244.

65. Bryan BA, Li D, Wu X, Liu M: **The Rho family of small GTPases: crucial regulators of skeletal myogenesis.** *Cell Mol Life Sci* 2005, **62**(14):1547-1555.
66. Goetsch SC, Hawke TJ, Gallardo TD, Richardson JA, Garry DJ: **Transcriptional profiling and regulation of the extracellular matrix during muscle regeneration.** *Physiol Genomics* 2003, **14**(3):261-271.
67. Cunningham JT, Rodgers JT, Arlow DH, Vazquez F, Mootha VK, Puigserver P: **mTOR controls mitochondrial oxidative function through a YY1-PGC-1alpha transcriptional complex.** *Nature* 2007, **450**(7170):736-740.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1755-8794/4/67/prepub>

doi:10.1186/1755-8794-4-67

Cite this article as: Tsumagari et al.: Gene expression during normal and FSHD myogenesis. *BMC Medical Genomics* 2011 **4**:67.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

