



SHORT GENOME REPORT

Open Access

High-quality permanent draft genome sequence of the *Parapiptadenia rigida*-nodulating *Cupriavidus* sp. strain UYPR2.512

Sofie E De Meyer¹, Elena Fabiano², Rui Tian¹, Peter Van Berkum³, Rekha Seshadri⁴, TBK Reddy⁴, Victor Markowitz⁵, Natalia N Ivanova⁴, Amrita Pati⁴, Tanja Woyke⁴, John Howieson¹, Nikos C Kyrpides^{4,6} and Wayne Reeve^{1*}

Abstract

Cupriavidus sp. strain UYPR2.512 is an aerobic, motile, Gram-negative, non-spore-forming rod that was isolated from a root nodule of *Parapiptadenia rigida* grown in soils from a native forest of Uruguay. Here we describe the features of *Cupriavidus* sp. strain UYPR2.512, together with sequence and annotation. The 7,858,949 bp high-quality permanent draft genome is arranged in 365 scaffolds of 369 contigs, contains 7,411 protein-coding genes and 76 RNA-only encoding genes, and is part of the GEBA-RNB project proposal.

Keywords: Root-nodule bacteria, Nitrogen fixation, Rhizobia, Betaproteobacteria, GEBA-RNB

Introduction

Legumes establish symbiotic associations with a group of soil bacteria, rhizobia, able to fix atmospheric nitrogen (N₂). Rhizobia elicit the formation of a symbiotic organ called a nodule comprising differentiated plant and bacterial cells. Differentiated rhizobia within nodules are termed bacteroids, and acquire the ability to fix nitrogen. Rhizobia are phylogenetically diverse including genera from the *Alphaproteobacteria* (*Allorhizobium*, *Azorhizobium*, *Bradyrhizobium*, *Ensifer*, *Mesorhizobium*, *Rhizobium*, etc.) as well as from the *Betaproteobacteria* (*Burkholderia*, *Cupriavidus*) [1,2].

The biological nitrogen fixation process significantly contributes to the development of sustainable agriculture reducing the use of supplies dependent on fuel and alleviating environmental impacts produced by the addition of chemical fertilizer [3]. Moreover, forestation with leguminous trees associated with rhizobia, “nitrogen-fixing trees”, has been successfully used for recovering degraded soils [4].

Parapiptadenia rigida (Benth.) Brenan, is a “nitrogen-fixing tree” belonging to the Piptadenia group from the Mimosoideae subfamily [5]. It is a multipurpose tree,

very appreciated because of its timber and therefore used in high quality furniture and construction. It is also used for gums, tannins and essential oil extraction, has medicinal properties and is included in agroforestry and reforestation programs [4,6,7]. Taulé *et al.* [8] demonstrated that this species could be nodulated either by Alpha-rhizobia (*Rhizobium*) or by Beta-rhizobia (*Burkholderia* and *Cupriavidus*) with *Burkholderia* being the preferred natural symbiont of this legume. In the case of *Cupriavidus* sp. UYPR2.512, this strain was isolated from a nodule of a *P. rigida* plant grown in soils collected from Mandiyú native forest in Artigas, Uruguay. Isolated bacterial colonies of *Cupriavidus* sp. UYPR2.512 were able to nodulate and to promote the growth of *P. rigida*, as well as *Mimosa pudica* plants [8].

To our knowledge, the only published sequenced genome of a Beta-rhizobia belonging to the *Cupriavidus* genus so far is that of *C. taiwanensis* LMG 19424^T [9]. Interestingly, the closest relative of *Cupriavidus* sp. UYPR2.512 is *C. necator* ATCC 43291^T [8]. Here, we present the description of the *Cupriavidus* sp. UYPR2.512 high-quality permanent draft genome sequence and its annotation.

Organism information

Classification and features

Cupriavidus sp. strain UYPR2.512 is a motile, Gram-negative, non-spore-forming rod (Figure 1 Left, Center)

* Correspondence: W.Reeve@murdoch.edu.au

¹Centre for Rhizobium Studies, Murdoch University, Murdoch, Western Australia

Full list of author information is available at the end of the article




Figure 1 Images of *Cupriavidus* sp. strain UYPR2.512 using scanning (Left) and transmission (Center) electron microscopy and the appearance of colony morphology on solid media (Right).

in the order *Burkholderiales* of the class *Betaproteobacteria*. The rod-shaped form varies in size with dimensions of 0.5-0.7 μm in width and 0.9-1.2 μm in length (Figure 1 Left). It is fast growing, forming 0.5-0.8 mm diameter colonies after 24 h when grown on TY [10] at 28°C. Colonies on TY are white-opaque, slightly domed, moderately mucoid with smooth margins (Figure 1 Right).

Figure 2 shows the phylogenetic relationship of *Cupriavidus* sp. strain UYPR2.512 in a 16S rRNA gene sequence based tree. This strain is the most similar to *Cupriavidus necator* ATCC 43291^T, *Cupriavidus oxalaticus* DSM 1105^T and *Cupriavidus taiwanensis* LMG 19424^T based on the 16S rRNA gene alignment with sequence identities of 99.32%, 98.49% and 98.42%, respectively, as determined using the EzTaxon-e server [11]. *Cupriavidus necator* ATCC 43291^T has been isolated from soil and is a non-obligate predator causing lysis of various Gram-positive and Gram-negative bacteria in the soil [12]. *Cupriavidus taiwanensis* LMG 19424^T is a plant symbiont and was isolated from root nodules of *Mimosa pudica* collected from three fields at Ping-Tung Country in the southern part of Taiwan [1]. Minimum Information about the Genome Sequence (MIGS) is provided in Table 1 and Additional file 1: Table S1.

Symbiotaxonomy

Cupriavidus sp. strain UYPR2.512 was isolated from *Parapiptadenia rigida*, a Mimosoideae legume native to Uruguay [8]. This tree is native to South America, including south Brazil, Argentina, Paraguay, and Uruguay, and used by locals for timber and as a source of gums, tannins and essential oils [8]. *Cupriavidus* sp. strain UYPR2.512 is able to renodulate its original host and is highly efficient in fixing nitrogen with this host [8]. A selection of other host plants, including *Trifolium repens*, *Medicago sativa*, *Peltophorum dubium* and *Mimosa pudica* were tested for their ability to nodulate with UYPR2.512. Of these plants, strain UYPR2.512

was only able to nodulate and fix nitrogen effectively with *M. pudica* [8].

Genome sequencing information

Genome project history

This organism was selected for sequencing on the basis of its environmental and agricultural relevance to issues in global carbon cycling, alternative energy production, and biogeochemical importance, and is part of the Genomic Encyclopedia of Bacteria and Archaea, The Root Nodulating Bacteria chapter (GEBA-RNB) project at the U.S. Department of Energy, Joint Genome Institute [25]. The genome project is deposited in the Genomes OnLine Database [14] and the high-quality permanent draft genome sequence in IMG [26]. Sequencing, finishing and annotation were performed by the JGI using state of the art sequencing technology [27]. A summary of the project information is shown in Table 2.

Growth conditions and DNA isolation

Cupriavidus sp. strain UYPR2.512 was grown to mid logarithmic phase in TY rich media [10] on a gyratory shaker at 28°C. DNA was isolated from 60 mL of cells using a CTAB (Cetyl trimethyl ammonium bromide) bacterial genomic DNA isolation method [29].

Genome sequencing and assembly

The draft genome of *Cupriavidus* sp. UYPR2.512 was generated at the DOE Joint Genome Institute [27]. An Illumina Std shotgun library was constructed and sequenced using the Illumina HiSeq 2000 platform which generated 29,312,424 reads totaling 4,396.9 Mbp [30]. All general aspects of library construction and sequencing performed at the JGI can be found at the JGI web site [31]. All raw Illumina sequence data was passed through DUK, a filtering program developed at JGI, which removes known Illumina sequencing and library preparation artifacts (Mingkun L,




Figure 2 Phylogenetic tree highlighting the position of *Cupriavidus* sp. strain UYPR2.512 (shown in blue print) relative to other type and non-type strains in the *Cupriavidus* genus using a 1,034 bp internal region of the 16S rRNA gene. Several Alpha-rhizobia sequences were used as an outgroup. All sites were informative and there were no gap-containing sites. Phylogenetic analyses were performed using MEGA, version 5.05 [13]. The tree was built using the maximum likelihood method with the General Time Reversible model. Bootstrap analysis with 500 replicates was performed to assess the support of the clusters. Type strains are indicated with a superscript T. Strains with a genome sequencing project registered in GOLD [14] are shown in bold and have the GOLD ID mentioned after the strain number, otherwise the NCBI accession number has been provided. Finished genomes are designated with an asterisk.

Table 1 Classification and general features of *Cupriavidus* sp. strain UYPR2.512 in accordance with the MIGS recommendations [15] published by the Genome Standards Consortium [16]

MIGS ID	Property	Term	Evidence code
	Current classification	Domain <i>Bacteria</i>	TAS [17]
		Phylum <i>Proteobacteria</i>	TAS [18,19]
		Class <i>Betaproteobacteria</i>	TAS [20]
		Order <i>Burkholderiales</i>	TAS [21]
		Family <i>Burkholderiaceae</i>	TAS [22]
		Genus <i>Cupriavidus</i>	TAS [23]
		Species <i>Cupriavidus</i> sp.	IDA
	Gram stain	Negative	IDA [23]
	Cell shape	Rod	IDA
	Motility	Motile	IDA
	Sporulation	Non-sporulating	IDA [23]
	Temperature range	mesophile	IDA [23]
	Optimum temperature	28°C	IDA
	pH range; Optimum	Not reported	
	Carbon source	Not reported	
MIGS-6	Habitat	Soil, root nodule on host	IDA
MIGS-6.3	Salinity	Not reported	
MIGS-22	Oxygen requirement	aerobic	IDA
MIGS-15	Biotic relationship	Symbiotic	IDA
MIGS-14	Pathogenicity	Non-pathogenic	NAS
MIGS-4	Geographic location	Uruguay	IDA
MIGS-5	Nodule collection date	2006	IDA
MIGS-4.1	Latitude	-30.507	IDA
MIGS-4.2	Longitude	-57.71	IDA
MIGS-4.4	Altitude	58 m	IDA

Evidence codes – IDA: Inferred from Direct Assay; TAS: Traceable Author. Statement (i.e., a direct report exists in the literature); NAS: Non-traceable Author. Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [24].

Copeland A, Han J. unpublished). Artifact filtered sequence data was then screened and trimmed according to the k-mers present in the dataset. High-depth k-mers, presumably derived from MDA amplification bias, cause problems in the assembly, especially if the k-mer depth varies in orders of magnitude for different

Table 2 Genome sequencing project information for *Cupriavidus* sp. strain UYPR2.512

MIGS ID	Property	Term
MIGS-31	Finishing quality	Permanent-draft
MIGS-28	Libraries used	Illumina Std PE
MIGS-29	Sequencing platforms	Illumina HiSeq 2000
MIGS-31.2	Fold coverage	106.8 X Illumina
MIGS-30	Assemblers	Velvet 1.1.04, ALLPATHS-LG V. r41043
MIGS-32	Gene calling methods	Prodigal 1.4
	Locus Tag	A3A5
	Genbank ID	ARBE00000000
	Genbank Date of Release	September 16, 2013
	GOLD ID	Gp0009663 [28]
	BIOPROJECT	PRJNA165301
MIGS-13	Source Material Identifier	UYPR2.512
	Project relevance	Symbiotic N ₂ fixation, agriculture

regions of the genome. Reads with high k-mer coverage (>30x average k-mer depth) were normalized to an average depth of 30x. Reads with an average kmer depth of less than 2x were removed. Following steps were then performed for assembly: (1) normalized Illumina reads were assembled using Velvet version 1.1.04 [32] (2) 1–3 Kbp simulated paired end reads were created from Velvet contigs using wgsim [33] (3)


Table 3 Genome statistics for *Cupriavidus* sp. strain UYPR2.512

Attribute	Value	% of total
Genome size (bp)	7,858,949	100
DNA coding (bp)	6,709,332	85.37
DNA G + C (bp)	5,128,158	65.25
DNA scaffolds	365	
Total genes	7,487	100
Protein coding genes	7,411	98.98
RNA genes	76	1.02
Pseudo genes	0	0
Genes in internal clusters	419	5.6
Genes with function prediction	5,663	75.64
Genes assigned to COGs	4,807	64.20
Genes with Pfam domains	5,959	79.59
Genes with signal peptides	696	9.30
Genes with transmembrane helices	1,545	20.64
CRISPR repeats	1	

normalized Illumina reads were assembled with simulated read pairs using Allpaths-LG (version r41043)[34]. Parameters for assembly steps were: 1) Velvet (velvet: 63 –shortPaired and velvetg: –very clean yes –exportFiltered yes –min contig lgth 500 –scaffolding no –cov cutoff 10) 2) wgsim (–e 0 –1 100 –2 100 –r 0 –R 0 –X 0) 3) Allpaths-LG (PrepareAllpathsInputs: PHRED 64 = 1 PLOIDY = 1 FRAG COVERAGE = 125 JUMP COVERAGE = 25 LONG JUMP COV = 50, RunAllpathsLG: THREADS = 8 RUN = std_shredpairs TARGETS = standard VAPI_WARN_ONLY = True OVERWRITE = True). The final draft assembly contained 369 contigs in 365 scaffolds. The total size of the genome is 7.9 Mbp and the final assembly is based on 839.6 Mbp of Illumina data, which provides an average of 106.8x coverage.

Genome annotation

Genes were identified using Prodigal [35], as part of the DOE-JGI genome annotation pipeline [36,37] followed by a round of manual curation using GenePRIMP [38] for finished genomes and Draft genomes in fewer than 10 scaffolds. The predicted CDSs were translated and used to search the National Center for Biotechnology Information (NCBI) non-redundant database, UniProt, TIGRFam, Pfam, KEGG, COG, and InterPro databases. The tRNAscanSE tool [39] was used to find tRNA genes, whereas ribosomal RNA genes were found by searches against models of the ribosomal RNA genes built from SILVA [40]. Other non-coding RNAs such as the RNA components of the protein secretion complex and the RNase P were identified by searching the genome for the corresponding Rfam profiles using



INFERNAL [41]. Additional gene prediction analysis and manual functional annotation was performed within the Integrated Microbial Genomes-Expert Review (IMG-ER) system [42] developed by the Joint Genome Institute, Walnut Creek, CA, USA.

Genome properties

The genome is 7,858,949 nucleotides with 65.25% GC content (Table 3) and comprised of 365 scaffolds and 369 contigs (Figure 3). From a total of 7,487 genes, 7,411 were protein encoding and 76 RNA only encoding genes. The majority of genes (75.64%) were assigned a putative function whilst the remaining genes were annotated as hypothetical. The distribution of genes into COG functional categories is presented in Table 4.

Conclusion

Cupriavidus sp. UYPR2.512 belongs to a group of Beta-rhizobia isolated from *Parapiptadenia rigida*, a native

tree from Uruguay belonging to the Mimosoideae legume group [8]. This tree is also native to the south of Brazil, Argentina and Paraguay [8]. Greenhouse experiments from previous studies have shown that *Cupriavidus* sp. UYPR2.512 is also able to nodulate and fix nitrogen with *Mimosa pudica*, an invasive species in many regions around the world [8]. Phylogenetic analysis revealed that UYPR2.512 is the most closely related to *Cupriavidus necator* ATCC 43291^T, *Cupriavidus oxalaticus* DSM 1105^T and *Cupriavidus taiwanensis* LMG 19424^T. In contrast to the other two strains, *Cupriavidus taiwanensis* LMG 19424^T is a microsymbiont that is able to nodulate and fix nitrogen in association with *Mimosa* species [43]. In total five *Cupriavidus* strains (AMP6, LMG 19424^T, STM6018, STM6070 and UYPR2.512), which can form a symbiotic association have now been sequenced. A comparison of these strains reveals that UYPR2.512 has the largest genome (7.9 Mbp), with the highest KOG count (1398), the lowest G + C (65.25%)

Table 4 Number of protein coding genes of *Cupriavidus* sp. strain UYPR2.512 associated with the general COG functional categories

Code	Value	% of total (5,426)	COG Category
J	183	3.37	Translation, ribosomal structure and biogenesis
A	1	0.02	RNA processing and modification
K	526	9.69	Transcription
L	192	3.54	Replication, recombination and repair
B	3	0.06	Chromatin structure and dynamics
D	35	0.65	Cell cycle control, Cell division, chromosome partitioning
Y	0	0.00	Nuclear structure
V	56	1.03	Defense mechanisms
T	210	3.87	Signal transduction mechanisms
M	277	5.11	Cell wall/membrane/envelope biogenesis
N	99	1.82	Cell motility
Z	0	0.00	Cytoskeleton
W	0	0.00	Extracellular structures
U	140	2.58	Intracellular trafficking, secretion, and vesicular transport
O	181	3.34	Posttranslational modification, protein turnover, chaperones
C	434	8.00	Energy production and conversion
G	268	4.94	Carbohydrate transport and metabolism
E	487	8.98	Amino acid transport and metabolism
F	89	1.64	Nucleotide transport and metabolism
H	194	3.58	Coenzyme transport and metabolism
I	337	6.21	Lipid transport and metabolism
P	272	5.01	Inorganic ion transport and metabolism
Q	236	4.35	Secondary metabolite biosynthesis, transport and catabolism
R	661	12.18	General function prediction only
S	545	10.04	Function unknown
-	2680	35.80	Not in COGS

and signal peptide (9.3%) percentages in this group. All of these genomes share the nitrogenase-RXN MetaCyc pathway catalyzed by a multiprotein nitrogenase complex. Out of five *Cupriavidus* strains (AMP6, LMG 19424^T, STM6018, STM6070 and UYPR2.512), which contain the N-fixation pathway, only *Cupriavidus* sp. UYPR2.512 has been shown to nodulate and fix effectively with *Parapiptadenia rigida*. The genome attributes of *Cupriavidus* sp. UYPR2.512 will therefore be important for ongoing molecular analysis of the plant microbe interactions required for the establishment of leguminous tree symbioses with this host.

Additional file

Additional file 1: Associated MIGS Record.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

EF supplied the strain and background information for this project, PVB supplied DNA to JGI, TR performed all imaging, SDM and WR drafted the paper, JH provided financial support and all other authors were involved in sequencing the genome and editing the final manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Berkeley National Laboratory under contract No. DE-AC02-05CH11231, Lawrence Livermore National Laboratory under Contract No. DE-AC52-07NA27344, and Los Alamos National Laboratory under contract No. DE-AC02-06NA25396.

Author details

¹Centre for Rhizobium Studies, Murdoch University, Murdoch, Western Australia. ²Instituto de Investigaciones Biológicas Clemente Estable, Montevideo, Uruguay. ³Soybean Genomics and improvement laboratory Bldg 006, BARC-West USDA ARS 10300 Baltimore Blvd, Beltsville, MD 20705, USA. ⁴DOE Joint Genome Institute, Walnut Creek, CA, USA. ⁵Biological Data Management and Technology Center, Lawrence Berkeley National Laboratory, Berkeley, CA, USA. ⁶Department of Biological Sciences, King Abdulaziz University, Jeddah, Saudi Arabia.

Received: 26 November 2014 Accepted: 5 December 2014

Published: 11 April 2015

References

1. Chen WM, Laevens S, Lee TM, Coenye T, De Vos P, Mergeay M, et al. *Ralstonia taiwanensis* sp. nov., isolated from root nodules of *Mimosa* species and sputum of a cystic fibrosis patient. *Int J Syst Evol Microbiol*. 2001;51:1729–35.
2. Moulin L, Munive A, Dreyfus B, Boivin-Masson C. Nodulation of legumes by members of the beta-subclass of Proteobacteria. *Nature*. 2001;411:948–50.
3. Crews TE, Peoples MB. Legume versus fertilizer sources of nitrogen: ecological tradeoffs and human needs. *Agric Ecosyst Environ*. 2004;102:279–97.
4. Farias JA, Hoppe JM, Vivian JAC. Comportamento de mudas de *Parapiptadenia rigida* (Benth) Brenan, submetidas a diferentes índices de luminosidade e em função de diferentes dimensões de recipientes. *Caderno de Pesquisa Série Biologia*. 2005;17:69–80.
5. Jobson RW, Luckow M. Phylogenetic study of the genus *Piptadenia* (Mimosoideae : Leguminosae) using plastid *trnL-F* and *trnK/matK* sequence data. *Syst Bot*. 2007;32:569–75.
6. Schmidt CA, Murillo R, Bruhn T, Bringmann G, Goettert M, Heinzmann B, et al. Catechin derivatives from *Parapiptadenia rigida* with in vitro wound-healing properties. *J Nat Prod*. 2010;73:2035–41.
7. de Souza GC, Haas AP, von Poser GL, Schapoval EE, Elisabetsky E. Ethnopharmacological studies of antimicrobial remedies in the south of Brazil. *J Ethnopharmacol*. 2004;90:135–43.
8. Taule C, Zabaleta M, Mareque C, Platero R, Sanjurjo L, Sicardi M, et al. New betaproteobacterial rhizobium strains able to efficiently nodulate *Parapiptadenia rigida* (Benth.) Brenan. *Appl Environ Microbiol*. 2012;78:1692–700.
9. Amadou C, Pascal G, Mangenot S, Glew M, Bontemps C, Capela D, et al. Genome sequence of the beta-rhizobium *Cupriavidus taiwanensis* and comparative genomics of rhizobia. *Genome Res*. 2008;18:1472–83.
10. Beringer JE. R factor transfer in *Rhizobium leguminosarum*. *J Gen Microbiol*. 1974;84:188–98.
11. Kim O-S, Cho Y-J, Lee K, Yoon S-H, Kim M, Na H, et al. Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *Int J Syst Evol Microbiol*. 2012;62:716–21.
12. Makkar NS, Casida LE. *Cupriavidus necator* gen. nov., sp. nov. - a nonobligate bacterial predator of bacteria in soil. *Int J Syst Evol Microbiol*. 1987;37:323–6.
13. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular Evolutionary Genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol*. 2011;28:2731–9.
14. Pagani I, Liolios K, Jansson J, Chen IM, Smirnova T, Nosrat B, et al. The Genomes OnLine Database (GOLD) v. 4: status of genomic and metagenomic projects and their associated metadata. *Nucleic Acids Res*. 2012;40:D571–579.
15. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, et al. Towards a richer description of our complete collection of genomes and metagenomes "Minimum Information about a Genome Sequence" (MIGS) specification. *Nat Biotechnol*. 2008;26:541–7.
16. Field D, Amaral-Zettler L, Cochrane G, Cole JR, Dawyndt P, Garrity GM, et al. The Genomic Standards Consortium. *PLoS Biol*. 2011;9:e1001088.
17. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc Nat Acad Sci USA*. 1990;87:4576–9.
18. Chen WX, Wang ET, Kuykendall LD. *The Proteobacteria*. New York: Springer - Verlag; 2005.
19. Validation of publication of new names and new combinations previously effectively published outside the IJSEM. *Int J Syst Evol Microbiol*. 2005;55:2235–8.
20. Garrity GM, Bell JA, Lilburn TE. Class II. Betaproteobacteria. In: Garrity GM, Brenner DJ, Krieg NR, Staley JT, editors. *Bergey's Manual of Systematic Bacteriology*. Volume 2. 2nd ed. New York: Springer - Verlag; 2005.
21. Garrity GM, Bell JA, Lilburn TE. Order 1. *Burkholderiales*. In: Garrity GM, Brenner DJ, Krieg NR, Staley JT, editors. *Bergey's Manual of Systematic Bacteriology*. Volume 2. 2nd ed. New York: Springer - Verlag; 2005.
22. Garrity GM, Bell JA, Lilburn TE. Family I. *Burkholderiaceae*. In: Garrity GM, Brenner DJ, Krieg NR, Staley JT, editors. *Bergey's Manual of Systematic Bacteriology*. Volume 2. 2nd ed. New York: Springer - Verlag; 2005.
23. Balkwill DL. Genus I. *Cupriavidus*. In: Garrity GM, Brenner DJ, Krieg NR, Staley JT, editors. *Bergey's Manual of Systematic Bacteriology*. Volume 2. 2nd ed. New York: Springer - Verlag; 2005.
24. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet*. 2000;25:25–9.
25. Reeve W, Ardley J, Tian R, Eshragi L, Yoon J, Ngamwisetkun P, et al. A genomic encyclopedia of the root nodule bacteria: Assessing genetic diversity through a systematic biogeographic survey. *Stand Genomic Sci*. 2015;10:14.
26. Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Pillay M, et al. IMG 4 version of the integrated microbial genomes comparative analysis system. *Nucleic Acids Res*. 2014;42:D560–7.
27. Mavromatis K, Land ML, Brettin TS, Quest DJ, Copeland A, Clum A, et al. The fast changing landscape of sequencing technologies and their impact on microbial genome assemblies and annotation. *PLoS ONE*. 2012;7:e48837.
28. GOLD ID *Cupriavidus* sp. UYPR2.512. [https://gold.jgi-psf.org/projects?id=9663]
29. CTAB DNA extraction protocol. [http://jgi.doe.gov/collaborate-with-jgi/pmo-overview/protocols-sample-preparation-information/]

30. Bennett S. Solexa Ltd. *Pharmacogenomics*. 2004;5:433–8.
31. JGI Website. [<http://www.jgi.doe.gov>]
32. Zerbino D, Birney E. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res*. 2008;18:821–9.
33. wgsim. [<https://github.com/lh3/wgsim>]
34. Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, et al. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci*. 2011;108:1513–8.
35. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics*. 2010;11:119.
36. Mavromatis K, Ivanova NN, Chen IM, Szeto E, Markowitz VM, Kyrpides NC. The DOE-JGI Standard Operating Procedure for the annotations of microbial genomes. *Standards Genomic Sci*. 2009;1:63–7.
37. Chen IM, Markowitz VM, Chu K, Anderson I, Mavromatis K, Kyrpides NC, et al. Improving microbial genome annotations in an integrated database context. *PLoS ONE*. 2013;8:e54859.
38. Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, et al. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods*. 2010;7:455–7.
39. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res*. 1997;25:955–64.
40. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, et al. SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res*. 2007;35:7188–96.
41. INFERNAL. Inference of RNA alignments. [<http://infernal.janelia.org>]
42. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpides NC. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics*. 2009;25:2271–8.
43. Chen WM, Moulin L, Bontemps C, Vandamme P, Bena G, Boivin-Masson C. Legume symbiotic nitrogen fixation by beta-proteobacteria is widespread in nature. *J Bacteriol*. 2003;185:7266–72.

doi:10.1186/1944-3277-10-13

Cite this article as: De Meyer *et al.*: High-quality permanent draft genome sequence of the *Parapiptadenia rigida*-nodulating *Cupriavidus* sp. strain UYPR2.512. *Standards in Genomic Sciences* 2015 **10**:13.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

