

POSTER PRESENTATION

Combination of a novel oncolytic immunotherapeutic agent, CAVATAK (coxsackievirus A21) and immune-checkpoint blockade significantly reduces tumor growth and improves survival in an immune competent mouse melanoma model

Darren Shafren^{1*}, Min Quah², Yvonne Wong², Robert HI Andtbacka³, Howard L Kaufman⁴, Gough G Au²

From Society for Immunotherapy of Cancer 29th Annual Meeting National Harbor, MD, USA. 6-9 November 2014

Coxsackievirus A21 (CAVATAK™) is a bio-selected oncolytic immunotherapy virus. Following intratumoral (i.t) injection, CAVATAK selectively infects ICAM-1expressing tumor cells, resulting in tumor cell lysis and a systemic immune-mediated anti-tumor response. A Phase II trial of i.t delivered CAVATAK (NCT01227551) in advanced melanoma patients has highlighted antitumor activity in both injected and distant non-injected lesions. Such responses have occurred at times when no circulating infectious CAVATAK was detected in patient serum and in an environment of high levels of anti-CAVATAK neutralizing antibodies. In further support of the generation of CAVATAK-mediated immune anti-tumor activity is the identification of a possible novel serum cytokine signature of elevated levels of IL-8 and IFN- γ in treated patients associated with tumor inflammation and systemic tumor response. Blockade of programmed death-1 (PD-1) in patients with metastatic melanoma has resulted in substantial tumor responses via a mechanism involving reversal of tumor-induced T cell suppression. We hypothesized that a combination of CAVATAK and PD-1 blockade may enhance anti-tumor responses, potentially leading to improved clinical activity. Preclinical studies in C57BL mice were conducted to assess the anti-tumor activity of CAVATAK and anti-mouse PD-1 (mPD-1) mAb in a B16-ICAM-1 melanoma immune competent mouse model. B16-ICAM-1 cells are murine melanoma B16 cells stably transfected to express human ICAM-1 allowing CAVATAK binding and cell infection. CAVA-TAK was administered i.t, while anti mPD-1 mAb was delivered intraperitoneally. Following treatment of the primary cutaneous B16-ICAM-1 tumor with 8 cycles of CAVATAK injections and 4 cycles of anti-PD-1mAb, mice were challenged with additional subcutaneous administration of B16 cells. Significant single agent antitumor activities against the primary B16-ICAM-1 tumor were observed in mice treated with either CAVATAK or anti-PD-1 mAb relative to saline controls. Combination of CAVATAK and anti-PD-1 mAb mediated significantly greater anti-tumor activity and offered greater survival benefit when compared to use of either agent alone. Of particular interest was the finding that a combination of CAVATAK and anti-PD-1 mAb was able to noticeably delay the onset of palpable tumor development following B16 cell challenge when compared to all other single agent treatment regimes. The significant anti-tumor activity mediated by the combination of CAVATAK and the checkpoint inhibitor antibody (anti-PD-1) observed in the presented murine melanoma model supports clinical evaluation of such an immunotherapeutic combination treatment regimen in patients with advanced melanoma.

¹Viralytics, Sydney, NSW, Australia

Full list of author information is available at the end of the article



© 2014 Shafren et al.; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

Authors' details

¹Viralytics, Sydney, NSW, Australia. ²School of BioMedical Sciences and Pharmacy, The University of Newcastle, New Lambton Heights, Australia. ³Huntsman Cancer Institute, Salt Lake City, UT, USA. ⁴Rutgers Cancer Institute of New Jersey, New Brunswick, NJ, USA.

Published: 6 November 2014

doi:10.1186/2051-1426-2-S3-P125

Cite this article as: Shafren *et al.*: Combination of a novel oncolytic immunotherapeutic agent, CAVATAK (coxsackievirus A21) and immune-checkpoint blockade significantly reduces tumor growth and improves survival in an immune competent mouse melanoma model. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):P125.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

BioMed Central

Submit your manuscript at www.biomedcentral.com/submit