

Review

# Global approaches to understanding ubiquitination

Peter Kaiser\* and Lan Huang<sup>†‡</sup>

Addresses: \*Department of Biological Chemistry, School of Medicine, <sup>†</sup>Department of Physiology and Biophysics, and <sup>‡</sup>Department of Developmental and Cell Biology, University of California, Irvine, CA 92697-1700, USA.

Correspondence: Peter Kaiser. E-mail: pkaiser@uci.edu

Published: 29 September 2005

*Genome Biology* 2005, **6**:233 (doi:10.1186/gb-2005-6-10-233)

The electronic version of this article is the complete one and can be found online at <http://genomebiology.com/2005/6/10/233>

© 2005 BioMed Central Ltd

## Abstract

Ubiquitination - the linkage of one or more molecules of the protein ubiquitin to another protein - regulates a wide range of biological processes in all eukaryotes. We review the proteome-wide strategies that are being used to study aspects of ubiquitin biology, including substrates, components of the proteasome and ubiquitin ligases, and deubiquitination.


Ubiquitin, a small protein of 76 amino acids, is highly conserved in all eukaryotes. In a multi-step process, ubiquitin is covalently linked to lysine residues of substrate proteins. If a single molecule of ubiquitin is linked to a protein, this is referred to as mono-ubiquitination, a process that is of particular importance for protein trafficking but has also been shown to regulate retrovirus budding and to modulate protein function directly [1]. A lysine residue of a ubiquitin molecule attached to a substrate can itself serve as an acceptor for an additional ubiquitin molecule, and this process can be repeated so that poly-ubiquitinated proteins form. Poly-ubiquitin chains serve as recognition signals for the 26S proteasome, the major regulator of protein abundance in cells, and poly-ubiquitination thus often initiates proteolysis of the substrate. But poly-ubiquitination can also regulate protein function directly without affecting stability, in ways similar to mono-ubiquitination and other post-translational modifications. The mechanisms underlying proteolysis-independent regulation by poly-ubiquitination are only poorly understood but might function by changing conformation or adding or obscuring a binding site (Figure 1; for reviews see [1-3]).

The transfer of ubiquitin is a multi-step process that involves at least three classes of enzymes: ubiquitin-activating enzymes, generally called E1 enzymes; ubiquitin-conjugating enzymes or E2s; and ubiquitin ligases, E3s (Figure 1). E3 ubiquitin ligases are of particular importance because they

confer substrate specificity to the system by interacting directly with substrate proteins and thereby directing the transfer of ubiquitin. The human genome encodes an estimated 500-600 ubiquitin ligases, a number comparable to the 518 predicted kinases [4,5]. If you consider that each ubiquitin ligase is active on several substrates, you can get some impression of the complexity and importance of the ubiquitin system.

Ubiquitination is a highly dynamic process and is balanced by deconjugation of ubiquitin by deubiquitinating enzymes (DUBs). The more than 70 DUBs that are estimated to be encoded in the human genome are responsible for the reversible nature of ubiquitin modifications and have important roles in recycling ubiquitin from proteasome substrates, in stabilizing proteins by counteracting their poly-ubiquitination, and in opposing the proteolysis-independent regulatory roles of ubiquitin modifications (for reviews see [6,7]). DUBs together with E1, E2 and E3 enzymes and the proteasome make up the ubiquitin-proteasome system.

The large number of proteins that constitute the ubiquitin-proteasome system and the enormous number of ubiquitination substrates mean that global approaches are required if we are to understand fully the role of ubiquitination in cell biology, development, and disease. Large-scale studies of the entire system are still in their early stages, but they have

**Figure 1**

The ubiquitin proteasome system. **(a)** Ubiquitin is activated by a ubiquitin-activating enzyme (E1) and transferred onto substrate proteins by ubiquitin-conjugating enzymes (E2) and ubiquitin ligases (E3), resulting in **(b)** either attachment of a single ubiquitin molecule (mono-ubiquitination), attachment of multiple ubiquitin units to several substrate lysine residues on the same protein (multi-ubiquitination) or synthesis of ubiquitin chains (poly-ubiquitination). **(c)** Many poly-ubiquitinated proteins are subsequently degraded by the 26S proteasome, which consists of the catalytic 20S complex and the regulatory 19S particles. Degradation substrates are either delivered to the proteasome by soluble ubiquitin receptors or recognized by the intrinsic ubiquitin-binding activity of the 19S particle. At the 19S proteasome the ubiquitin chain is disassembled, and the substrate is unfolded before it can enter the cavity of the 20S subunit where proteolysis takes place. Finally, proteolytic fragments exit the proteasome in a poorly understood way. **(d)** Ubiquitination can also directly regulate protein function in a proteolysis-independent manner, via mono-, multi- or poly-ubiquitinated proteins.

already made important contributions to the field. Here, we review the approaches taken in some of these studies and their findings.

### Proteomic approaches to characterizing the ubiquitin-proteasome system

Multi-protein complexes and protein-protein interactions have important roles in the ubiquitin-proteasome system. Both the 26S proteasome (see Figure 1c) and E3 ubiquitin ligases have been studied extensively using protein-complex purification coupled with mass-spectrometric protein identification [8-11]. Studies of the subunit composition of proteasomes from various organisms have revealed that the 26S proteasome complex consists of the 20S complex (made up of seven  $\alpha$  and seven  $\beta$  subunits) and the 19S complex (made up of six ATPase and twelve non-ATPase subunits) [12]. The 20S complex is well characterized as forming the catalytic core; the 19S regulatory complex is believed to be responsible for substrate recognition and unfolding (Figure 1c), but the specific functions of most of the 19S subunit components are still not well understood. There is accumulating evidence for a non-proteolytic role for the proteasome in processes such as transcription, chromatin packaging, and DNA repair

[13,14]. Additional subunits found in the majority of proteasome complexes have been identified following the development of new protein-purification and protein-identification techniques [8,15]. Recently, hybrid 26S proteasome complexes have been characterized, in which one copy of the 19S is present at one end of the 20S core and the other end is capped by Bml10, a newly characterized HEAT repeat protein in yeast. Bml10 and its mammalian ortholog PA28 function as 20S activators [16,17]. These complexes seem to have labile structures that cannot be preserved during the purification steps. Although the hybrid complexes reconstituted *in vitro* have higher peptidase activity than the classic versions, their roles in the degradation of ubiquitinated substrates *in vivo* are unclear [16,17]. Given the heterogeneous population and functional diversity of the various proteasome complexes, it remains a challenging task to purify and identify the subpopulations of proteasome complexes and to correlate the differences in their composition with their distinct functions *in vivo*.

A diverse groups of proteasome-interacting proteins, including ubiquitin ligases, DUBs, heat-shock proteins and many other proteins have been identified by affinity purification and mass spectrometry as well as from genome-wide two-hybrid

screens for protein-protein interactions [8,15,18-21]. Many of the identified interactions seem to be labile under conditions of active ATP hydrolysis by the 19S regulatory complex because addition of ATP coincides with the release of the interacting proteins [8]. This modulation has been suggested to be a part of the mechanism of protein degradation [22]. New methodologies are needed if we are to identify and characterize proteasome-interacting proteins fully, to understand how the different interacting proteins influence the catalytic cycle, and to clarify how they link the ubiquitin-proteasome system to other biological processes.

Mass-spectrometric approaches have also contributed much to our current understanding of the complex composition of E3 ubiquitin ligases. Two types of ubiquitin ligase that play important roles in cell-cycle regulation have been extensively investigated: SCFs and the anaphase-promoting complex/cyclosome (APC/C). SCF ubiquitin ligases are named after three of their four subunits - Skp1, Cdc53 (also known as Cul1) and one member of the F-box protein family - and they also include the Ring-H2 protein Hrt1 (also known as Roc1 or Rbx1) [23]. The substrate specificity of SCFs depends on the different F-box proteins that are tethered to the Cdc53-Hrt1 ubiquitin-ligase module by Skp1. Using sequential rounds of epitope tagging, affinity purification and mass spectrometry (a procedure called SEAM), Skp1 was found to form a variety of complexes, including some that are most likely to have functions other than ubiquitination [9,24,25].

In comparison, proteomic approaches have shown that the APC/C, which regulates mitosis, has a more complex structure than SCFs with at least 13 components [26]. Despite some success in identifying the subunits and also the modifications of APC/C, the molecular functions of the individual subunits are largely unknown. Exceptions are the RING-finger subunit Apc11 and the cullin-like subunit Apc2, which are believed to have a direct role in ubiquitin transfer [27].

Perhaps because of the large number of ubiquitin ligases present in the genome, ubiquitin-ligase proteomics is still in its infancy. Affinity purification coupled with mass spectrometry has promised great advances in the study of the composition of protein complexes and the identification of their interacting partners. But further advancements in proteomic research are expected to provide more information on the protein complexes involved in the ubiquitin-proteasome system, including their post-translational modifications, the stoichiometry of their subunits and how they are assembled.

### Identification of ubiquitination substrates *in vitro*

The large number of putative E3 ubiquitin ligases makes systematic characterization of their substrates a formidable task, but it is one that will be important if we are to gain a global view of the dynamics of the ubiquitin system. E3-substrate

interactions are generally only transient, and substrates are usually either degraded by the proteasome and/or released from the E3 ligase after the transfer of ubiquitin. This makes detection of E3-ligase-substrate interactions difficult. Two-hybrid assays have successfully identified some substrates of ubiquitin ligases [28], but identification of proteins that can interact with E3 ligases does not necessarily pinpoint substrates. A more effective strategy is to identify E3 substrates by their ubiquitination or degradation by the 26S proteasome.

One of the first effective large-scale attempts used *Xenopus* oocyte extracts to identify substrates of the APC/C, a ubiquitin ligase that regulates mitosis [29-32]. The approach exploited the unique regulation of APC/C activity: it is inactive during interphase but active during mitosis. When added to mitotic extracts, APC/C substrates are ubiquitinated and rapidly degraded by the proteasome, but the same substrates are unchanged in interphase lysates. In a large-scale approach *Xenopus* cDNA clones that had been *in vitro*-translated and labeled were divided into small pools and incubated with interphase and mitotic oocyte extracts. Proteins that disappeared specifically from mitotic extracts were isolated, and this led to the identification of several important APC/C substrates, including cyclin B, the DNA-replication inhibitor geminin, and the anaphase inhibitor securin [29-32].


A different *in vitro* approach was applied to identifying the potential substrates of the ubiquitin ligase that is formed by a heterodimer of BRCA1 and BARD1 [33,34]; the BRCA1/BARD1 heterodimer functions as a tumor suppressor that is important for protection from breast and ovarian cancer, and its ubiquitin-ligase activity has been linked to its protective function [35]. Sato and colleagues [33] immunoprecipitated BRCA1/BARD1 complexes, which were added to a ubiquitination reaction *in vitro*, that used ubiquitin tagged with the FLAG epitope. The rationale behind the approach was that substrates of ubiquitination by BRCA1 should be bound to the immunoprecipitated BRCA1 and subsequently ubiquitinated with FLAG-ubiquitin *in vitro*. Proteins conjugated to FLAG-ubiquitin were purified and identified by mass spectrometry [33]. A more directed strategy restricted the hunt for BRCA1-BARD1 substrates to components of the centrosome [34], because BRCA1 has been implicated in the regulation of centrosome duplication [36]. Starita and colleagues [34] incubated mammalian centrosome-containing cell fractions with recombinant BRCA1-BARD1 ligase complexes and biotinylated ubiquitin. Ubiquitinated proteins were detected through the biotin tag on ubiquitin and subsequently identified by mass spectrometry [34]. Both of these strategies [33,34] identified promising candidate BRCA1-BARD1 substrates - nucleoplasmin/B23 [33] and  $\gamma$ -tubulin [34] - that might be connected to the tumor-suppressor function of BRCA1. A high-throughput strategy has also been used to identify substrates of the yeast ubiquitin ligase Rsp5 *in vitro*. A luminescent assay involving biotinylated ubiquitin was used to screen several hundred purified yeast

proteins for Rsp5-dependent ubiquitination *in vitro*. Previously known, as well as new, candidate substrates of Rsp5 were identified [37].

A more general *in vitro* approach [38] used total HeLa cell lysates for large-scale identification of ubiquitinated proteins. Cell lysates were incubated with ubiquitin tagged with six histidines (6xHis-ubiquitin) and an ATP-regenerating system to sustain ubiquitination *in vitro*. The 6xHis-ubiquitin was covalently attached to proteins by the E1, E2, and E3 enzymes present in the cell lysates, and this allowed purification of ubiquitinated proteins on the basis of the affinity of 6xHis-ubiquitin to Ni<sup>2+</sup> ions (Ni-chelate chromatography) [38]. Over 100 ubiquitin-linked proteins were identified by mass spectrometry, of which a relatively high proportion was already implicated in the ubiquitin proteasome pathway, such as E2 and E3 enzymes and proteasome subunits. Because relatively mild purification conditions were chosen, both ubiquitinated proteins and proteins associated with them were identified. This is illustrated by the identification of 16 out of 18 subunits of the 19S proteasome; covalent modification of 19S proteasome subunits with ubiquitin has so far not been reported, but an intrinsic affinity of the 19S proteasome for poly-ubiquitin chains is well known [39] and is most likely to be responsible for the identification of these proteins in the study [38]. *Bona fide* ubiquitinated proteins can be distinguished from associated, copurifying proteins by fractionation strategies that use highly denaturing conditions and break non-covalent interactions. Such stringent purification conditions have been widely used to demonstrate covalent attachment of ubiquitin to specific proteins [40], as well as in proteome-wide approaches to identifying ubiquitinated proteins, as discussed below.

### Ubiquitination substrates *in vivo*

Identification of all ubiquitinated proteins in a cell under a given growth condition or developmental state is an ambitious aim, but it no longer seems impossible given the tremendous pace at which mass-spectrometry-based proteomics is developing (reviewed in [41]). Ubiquitin profiling was pioneered by Peng and colleagues [42] and usually involves expression of 6xHis-tagged ubiquitin in cells (Figure 2). The cellular ubiquitin system conjugates 6xHis-ubiquitin to target proteins and allows their purification by Ni-chelate chromatography. Because Ni-chelate purification is compatible with fully denaturing conditions, proteins that are associated with ubiquitinated proteins but are not ubiquitination substrates themselves can efficiently be removed. The purified ubiquitinated proteins are fragmented by trypsin (or similar proteases) to generate peptides, which can be used for mass-spectrometric identification of the proteins present in the purified fraction. More than 1,000 candidate ubiquitination substrates were identified using this method in the relatively simple eukaryote *Saccharomyces cerevisiae* [42], whose genome



**Figure 2**

Global strategies that use mass spectrometry (MS) to study ubiquitination. **(a)** Diagram of the lysine residues in ubiquitin; the carboxy-terminal Arg-Gly-Gly (RGG) motif is also indicated. **(b)** In ubiquitin profiling, 6xHis-tagged ubiquitin expressed in cells is conjugated to substrate proteins, and this facilitates purification of ubiquitinated proteins under denaturing conditions by Ni-chelate chromatography, in which histidine-tagged proteins bind specifically to immobilized Ni<sup>2+</sup> ions. Purified ubiquitinated proteins are digested with trypsin and the resulting peptides are analyzed by mass spectrometry to identify the proteins present in the sample. **(c)** Precise ubiquitination sites can be determined by mass spectrometry because of a characteristic mass shift caused by diglycine that is retained on ubiquitinated lysine residues within peptides after trypsin digestion. **(d)** A similar strategy allows differentiation between the various types of ubiquitin chain linkage that can lead to diverse ubiquitin-chain topologies. Depending on the lysine residue in ubiquitin that was used for the ubiquitin-ubiquitin linkage, different linkage-specific signature peptides with characteristic masses are produced by trypsin digestion. These signature peptides can be detected and distinguished by mass spectrometry.

encodes roughly 5,800 proteins. Surprisingly, most of the well-studied (and less abundant) ubiquitinated proteins were absent from the list, suggesting that many more yeast

proteins than the identified 1,000 candidates are ubiquitination substrates.

At first glance, it seems that a surprisingly large fraction of the proteome is ubiquitinated. But misfolded proteins, which can be generated by translation inaccuracy, folding problems or oxidative damage, are ubiquitinated and degraded as part of the protein quality-control pathway [43]. One can therefore expect that at least a small fraction of any protein will be ubiquitinated, and that sufficiently sensitive analytical methods might find that all proteins can be ubiquitination substrates. It is important to bear in mind that current ubiquitin-profiling experiments can indicate only whether any of a given protein is ubiquitinated but cannot give any estimate of what fraction of the protein is ubiquitinated. This imposes some limitations on how the results of large-scale studies can be interpreted.

To find more specific substrates of the ubiquitin-proteasome system, recent proteomic approaches have focused on specific parts of the system. Ubiquitin profiling has been used successfully to study the endoplasmic reticulum associated degradation pathway (ERAD) [44]. Membrane-enriched fractions from yeast cells expressing 6xHis-ubiquitin were used as a starting material for purification of ubiquitinated proteins and their subsequent identification by mass spectrometry. More than 80 candidate ERAD substrates were identified [44].

Mayor and colleagues [45] enriched for proteasome substrates on a poly-ubiquitin-binding protein resin and followed this with denaturing Ni-chelate chromatography in order to purify ubiquitinated proteins from yeast cells expressing 6xHis-ubiquitin. Remarkably, by profiling a yeast strain with a mutation in the proteasomal ubiquitin receptor Rpn10, they could identify 54 candidate ubiquitination substrates that require Rpn10 for degradation [45]. Among them were the transcription factor Gcn4 and the cell cycle regulator Sic1, two known proteasome substrates whose abundance is low. This study [45] demonstrates how subtractive ubiquitin profiling can help to define substrates of particular pathways of the ubiquitin-proteasome system. It is not hard to imagine that a similar strategy, in which cells defective in a particular E3 ligase are compared with wild-type cells, could be used for large-scale identification of the specific substrates of individual ubiquitin ligases. Furthermore, the introduction to proteomic analyses of various mass-spectrometric strategies that use stable isotope labeling promises to transform ubiquitin-profiling experiments by enabling detection of quantitative changes in ubiquitin profiles [46-48].

### Ubiquitination sites and ubiquitin-chain topology

The pioneering ubiquitin-profiling experiments of Peng and colleagues [42] demonstrated the feasibility of large-scale identification of ubiquitin-attachment sites in substrate proteins. This is possible because, after trypsin digestion, the

two carboxy-terminal residues of ubiquitin remain attached to the lysine residue of the substrate protein (Figure 2c). These two additional glycine residues lead to a characteristic 114 Da increase in the mass of the ubiquitinated substrate peptide, which is diagnostic for the ubiquitinated residue and can be monitored by mass spectrometry [42]. Over 100 precise ubiquitin attachment sites have been identified by analyzing peptide-mass data from global ubiquitin-profiling experiments [42,44]. Bioinformatic analyses of these data sets showed that ubiquitination sites are almost exclusively exposed on the protein surface, and located preferentially in a sequence environment that is predicted to form a loop structure [49]. No conserved ubiquitination motif could be defined, however.

A related strategy allowed detection of different ubiquitin chain topologies *in vivo* [42,50]. Formation of a poly-ubiquitin chain requires isopeptide linkages between the terminal carboxyl group of a free ubiquitin molecule and one of seven lysine residues present in a substrate-attached ubiquitin (Figure 2a,d). The most important chain topology is formed through the lysine in position 48 of ubiquitin [51]. Chains linked through Lys48 are the principal recognition signals for the proteasome and generally induce substrate degradation [52]. Chains linked through Lys63 do not induce substrate degradation but have direct effects on protein activity [53,54]; the biological role of other ubiquitin chain topologies is unclear. From the analytical perspective, the ubiquitin chain linkage can be regarded as a specific example of a ubiquitination site in a substrate: the substrate in this case is ubiquitin itself. Chain linkage can therefore be determined by the characteristic 114 Da mass shift, as described above (Figure 2c,d) [41].


Rather surprisingly, analysis of mass data from large-scale ubiquitin-profiling experiments [42] has revealed that all seven lysine residues in ubiquitin are used to form ubiquitin chains *in vivo*. The abundance of the different chain-linkage types was ranked and suggested that linkage through Lys48 is the most abundant topology, followed by Lys63 and Lys11 chains and the less frequent linkages through Lys33, Lys27, Lys6, and Lys29 (the latter is detected only in combination with Lys33 linkage) [42]. These results emphasize the complexity of ubiquitin biology. At the same time, interpretation of these experiments [42] is somewhat limited because they can describe only the linkage between two ubiquitins and cannot determine to which protein the chain was attached or whether the chain was attached to a substrate at all (Figure 2d). Similarly, it is unclear whether ubiquitin chains are homogenous or can contain mixed linkage types. The only rigorously studied example so far of a substrate-attached ubiquitin chain *in vivo* demonstrated the presence of a homogenous ubiquitin chain [50]. Many more studies are necessary, however, before we can decide whether mixed chains exist *in vivo* and whether they encode biologically important information.

### Chemistry-based and global *in vivo* approaches to deubiquitination

DUBs are an important component of the ubiquitin-proteasome system. They are proteases and can therefore be targeted by activity-dependent probes, which form covalent bonds with their active sites and have been successfully applied to other classes of proteases [55,56]. An elegant strategy using an activity-based approach to target DUBs has led to the identification of numerous deubiquitinating activities in cell lysates and the discovery of the new class of DUBs that contain an OTU domain (a domain characteristic of the ovarian tumor superfamily of proteins) [57]. Briefly, ubiquitin fused to a hemagglutinin (HA) epitope tag at its amino terminus and to one of various cysteine-reactive probes (which react with cysteine proteases) at the carboxyl terminus is incubated with total cell lysates. The active site of each DUB forms a covalent bond with the cysteine-reactive group on the HA-ubiquitin probe (Figure 3a) and can therefore be immunopurified using the HA tag and subsequently identified by mass spectrometry (Figure 3b) [58]. Activity-based ubiquitin probes have also been used to generate profiles of DUB activity in different cell lines and tissues and to identify proteins that interact with DUBs (Figure 3b) [58,59].

The problem of identifying DUBs that react with a specific ubiquitinated protein has been elegantly addressed using a collection of RNA-interference (RNAi) vectors that knock down the expression of more than 50 DUBs in mammalian cells [60]. Because the steady-state level of ubiquitin conjugates reflects the balance between ubiquitination and deubiquitination, knockdown of the activity of a specific DUB increases the fraction of the ubiquitinated form of its substrates. This strategy helped to identify the DUB USP1 as the deubiquitinating activity that acts on the mono-ubiquitinated version of FANCD2 (a protein defective in the Fanconi anemia complementation group D2) [60]. A similar collection of small interfering RNAs (siRNAs) has enabled the identification of the familial cylindromatosis tumor suppressor gene (*CYLD*) as a DUB involved in regulation of the NF $\kappa$ B transcriptional control pathway [61].

More than 25 years have passed since the initial discovery of the ubiquitin system. Ubiquitin has since extended its role from a protein-degradation signal to a regulatory protein modification that affects all areas of biology. The importance of the ubiquitin-proteasome system in biology was acknowledged with the 2004 Chemistry Nobel Prize to Aaron Ciechanover, Avram Herschko, and Irwin Rose, who first established its main features. The complexity and significance of the ubiquitin-proteasome system has started to attract global approaches that are beginning to make important contributions to our understanding of the system. Chemistry-based approaches to deubiquitination have demonstrated the effectiveness of these strategies, and analogous activity-based probes for studying the ubiquitin transfer will be of similar



**Figure 3**

Activity-based profiling of deubiquitinating enzymes and interacting proteins. **(a)** Ubiquitin fused to an amino-terminal epitope tag (for example hemagglutinin, HA) and a carboxy-terminal reactive group forms a covalent conjugate with deubiquitinating enzymes (DUBs; for details of the generation of these ubiquitin probes, see [57]). **(b)** The DUB-ubiquitin conjugates can be immunopurified using the HA epitope. Immunopurification under native conditions allows identification of DUBs and their interacting proteins by mass spectrometry (MS). The immunopurified fractions can be further separated by gel electrophoresis, and DUB-ubiquitin conjugates can be detected by anti-HA immunoblotting. Proteins corresponding to HA-reactive bands can be eluted from silver-stained gels (not shown) and the DUBs can be identified by mass spectrometry.

importance. Proteomics using mass spectrometry has had a tremendous impact on the field, as it has helped to describe the nature and regulation of multi-protein complexes that themselves regulate the ubiquitin-proteasome system. Large-scale ubiquitin-profiling experiments have highlighted the involvement of the system in a wide range of processes and demonstrated the complexity of ubiquitin-chain topology. Mass-spectrometric approaches promise to be particularly powerful in the future because one of the previous limitations - the inherent non-quantitative nature of these experiments - has been overcome by stable-isotope-based quantification strategies [48-50].

Some of the global approaches described here for the study of the ubiquitin system have also been applied to the study of other ubiquitin-like proteins, such as SUMO, ISG15, and Nedd8 [62,63]. The strategies that have been proven to be effective for studying ubiquitin biology will be just as important for rapidly advancing our understanding of the role of the growing family of ubiquitin-like modifiers.

## Acknowledgements

We thank K. Flick for critically reading the manuscript. Work in the laboratory of P.K. is supported by grants from NIH (GM-66164) and the California Breast Cancer Research Program (11NB-0177). L.H. acknowledges support from NIH (GM074830) and the DOD (PC-041126).

## References

- Hicke L: **Protein regulation by monoubiquitin.** *Nat Rev Mol Cell Biol* 2001, **2**:195-201.
- Hershko A, Ciechanover A: **The ubiquitin system.** *Annu Rev Biochem* 1998, **67**:425-479.
- Pickart CM: **Back to the future with ubiquitin.** *Cell* 2004, **116**:181-190.
- Wilkinson KD, Ventii KH, Friedrich KL, Mullally JE: **The ubiquitin signal: assembly, recognition and termination.** *EMBO Rep* 2005, **6**:815-820.
- Johnson SA, Hunter T: **Kinomics: methods for deciphering the kinome.** *Nat Methods* 2005, **2**:17-25.
- Wilkinson KD: **Regulation of ubiquitin-dependent processes by deubiquitinating enzymes.** *FASEB J* 1997, **11**:1245-1256.
- Wilkinson KD: **Ubiquitination and deubiquitination: targeting of proteins for degradation by the proteasome.** *Semin Cell Dev Biol* 2000, **11**:141-148.
- Verma R, Chen S, Feldman R, Schieltz D, Yates J, Dohmen J, Deshaies RJ: **Proteasomal proteomics: identification of nucleotide-sensitive proteasome-interacting proteins by mass spectrometric analysis of affinity-purified proteasomes.** *Mol Biol Cell* 2000, **11**:3425-3439.
- Seol JH, Shevchenko A, Deshaies RJ: **Skp1 forms multiple protein complexes, including RAVE, a regulator of V-ATPase assembly.** *Nat Cell Biol* 2001, **3**:384-391.
- Yoon HJ, Feoktistova A, Wolfe BA, Jennings JL, Link AJ, Gould KL: **Proteomics analysis identifies new components of the fission and budding yeast anaphase-promoting complexes.** *Curr Biol* 2002, **12**:2048-2054.
- Zachariae W, Shevchenko A, Andrews PD, Ciosk R, Galova M, Stark MJ, Mann M, Nasmyth K: **Mass spectrometric analysis of the anaphase-promoting complex from yeast: identification of a subunit related to cullins.** *Science* 1998, **279**:1216-1219.
- Pickart CM, Cohen RE: **Proteasomes and their kin: proteases in the machine age.** *Nat Rev Mol Cell Biol* 2004, **5**:177-187.
- Russell SJ, Reed SH, Huang W, Friedberg EC, Johnston SA: **The 19S regulatory complex of the proteasome functions independently of proteolysis in nucleotide excision repair.** *Mol Cell* 1999, **3**:687-695.
- Muratani M, Tansey WP: **How the ubiquitin-proteasome system controls transcription.** *Nat Rev Mol Cell Biol* 2003, **4**:192-201.
- Leggett DS, Hanna J, Borodovsky A, Crosas B, Schmidt M, Baker RT, Walz T, Ploegh H, Finley D: **Multiple associated proteins regulate proteasome structure and function.** *Mol Cell* 2002, **10**:495-507.
- Cascio P, Call M, Petre BM, Walz T, Goldberg AL: **Properties of the hybrid form of the 26S proteasome containing both 19S and PA28 complexes.** *EMBO J* 2002, **21**:2636-2645.
- Schmidt M, Haas W, Crosas B, Santamaria PG, Gygi SP, Walz T, Finley D: **The HEAT repeat protein Blm10 regulates the yeast proteasome by capping the core particle.** *Nat Struct Mol Biol* 2005, **12**:294-303.
- Ho Y, Gruhler A, Heilbut A, Bader GD, Moore L, Adams SL, Millar A, Taylor P, Bennett K, Boutillier K, et al.: **Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry.** *Nature* 2002, **415**:180-183.
- Gavin AC, Bosche M, Krause R, Grandi P, Marzioch M, Bauer A, Schultz J, Rick JM, Michon AM, Cruciat CM, et al.: **Functional organization of the yeast proteome by systematic analysis of protein complexes.** *Nature* 2002, **415**:141-147.
- Davy A, Bello P, Thierry-Mieg N, Vaglio P, Hitti J, Doucette-Stamm L, Thierry-Mieg D, Reboul J, Boulton S, Walhout AJ, et al.: **A protein-protein interaction map of the *Caenorhabditis elegans* 26S proteasome.** *EMBO Rep* 2001, **2**:821-828.
- Cagney G, Uetz P, Fields S: **Two-hybrid analysis of the *Saccharomyces cerevisiae* 26S proteasome.** *Physiol Genomics* 2001, **7**:27-34.
- Babbitt SE, Kiss A, Deffenbaugh AE, Chang YH, Bailly E, Erdjument-Bromage H, Tempst P, Buranda T, Sklar LA, Baumlner J, et al.: **ATP hydrolysis-dependent disassembly of the 26S proteasome is part of the catalytic cycle.** *Cell* 2005, **121**:553-565.
- Petroski MD, Deshaies RJ: **Function and regulation of cullin-RING ubiquitin ligases.** *Nat Rev Mol Cell Biol* 2005, **6**:9-20.
- Kaplan KB, Hyman AA, Sorger PK: **Regulating the yeast kinetochore by ubiquitin-dependent degradation and Skp1p-mediated phosphorylation.** *Cell* 1997, **91**:491-500.
- Galan JM, Wiederkehr A, Seol JH, Haguenaer-Tsapis R, Deshaies RJ, Riezman H, Peter M: **Skp1p and the F-box protein Rcy1p form a non-SCF complex involved in recycling of the SNARE Snc1p in yeast.** *Mol Cell Biol* 2001, **21**:3105-3117.
- Harper JW, Burton JL, Solomon MJ: **The anaphase-promoting complex: it's not just for mitosis any more.** *Genes Dev* 2002, **16**:2179-2206.
- Zachariae W, Nasmyth K: **Whose end is destruction: cell division and the anaphase-promoting complex.** *Genes Dev* 1999, **13**:2039-2058.
- Uetz P, Giot L, Cagney G, Mansfield TA, Judson RS, Knight JR, Lockshon D, Narayan V, Srinivasan M, Pochart P, et al.: **A comprehensive analysis of protein-protein interactions in *Saccharomyces cerevisiae*.** *Nature* 2000, **403**:623-627.
- Lustig KD, Stukenberg PT, McGarry TJ, King RW, Cryns VL, Mead PE, Zou LI, Yuan J, Kirschner MW: **Small pool expression screening: identification of genes involved in cell cycle control, apoptosis, and early development.** *Methods Enzymol* 1997, **283**:83-99.
- Zou H, McGarry TJ, Bernal T, Kirschner MW: **Identification of a vertebrate sister-chromatid separation inhibitor involved in transformation and tumorigenesis.** *Science* 1999, **285**:418-422.
- McGarry TJ, Kirschner MW: **Geminin, an inhibitor of DNA replication, is degraded during mitosis.** *Cell* 1998, **93**:1043-1053.
- King RW, Lustig KD, Stukenberg PT, McGarry TJ, Kirschner MW: **Expression cloning in the test tube.** *Science* 1997, **277**:973-974.
- Sato K, Hayami R, Wu W, Nishikawa T, Nishikawa H, Okuda Y, Ogata H, Fukuda M, Ohta T: **Nucleophosmin/B23 is a candidate substrate for the BRCA1-BARD1 ubiquitin ligase.** *J Biol Chem* 2004, **279**:30919-30922.
- Starita LM, Machida Y, Sankaran S, Elias JE, Griffin K, Schlegel BP, Gygi SP, Parvin JD: **BRCA1-dependent ubiquitination of gamma-tubulin regulates centrosome number.** *Mol Cell Biol* 2004, **24**:8457-8466.
- Narod SA, Foulkes WD: **BRCA1 and BRCA2: 1994 and beyond.** *Nat Rev Cancer* 2004, **4**:665-676.
- Deng CX: **Roles of BRCA1 in centrosome duplication.** *Oncogene* 2002, **21**:6222-6227.
- Kus BM, Gajadhar A, Stanger K, Cho R, Sun W, Rouleau N, Lee T, Chan D, Wolting C, Edwards AM, et al.: **A high throughput screen to identify substrates for the ubiquitin ligase Rsp5.** *J Biol Chem* 2005, **280**:29470-29478.
- Gururaja T, Li W, Noble WS, Payan DG, Anderson DC: **Multiple functional categories of proteins identified in an *in vitro* cellular ubiquitin affinity extract using shotgun peptide sequencing.** *J Proteome Res* 2003, **2**:394-404.
- Elsasser S, Finley D: **Delivery of ubiquitinated substrates to protein-unfolding machines.** *Nat Cell Biol* 2005, **7**:742-749.
- Kaiser P, Flick K, Wittenberg C, Reed SI: **Regulation of transcription by ubiquitination without proteolysis: Cdc34/SCF(Met30)-mediated inactivation of the transcription factor Met4.** *Cell* 2000, **102**:303-314.
- Kirkpatrick DS, Denison C, Gygi SP: **Weighing in on ubiquitin: the expanding role of mass-spectrometry-based proteomics.** *Nat Cell Biol* 2005, **7**:750-757.
- Peng J, Schwartz D, Elias JE, Thoreen CC, Cheng D, Marsischky G, Roelofs J, Finley D, Gygi SP: **A proteomics approach to understanding protein ubiquitination.** *Nat Biotechnol* 2003, **21**:921-926.

43. McClellan AJ, Tam S, Kaganovich D, Frydman J: **Protein quality control: chaperones culling corrupt conformations.** *Nat Cell Biol* 2005, **7**:736-741.
44. Hitchcock AL, Auld K, Gygi SP, Silver PA: **A subset of membrane-associated proteins is ubiquitinated in response to mutations in the endoplasmic reticulum degradation machinery.** *Proc Natl Acad Sci USA* 2003, **100**:12735-12740.
45. Mayor T, Lipford JR, Graumann J, Smith GT, Deshaies RJ: **Analysis of polyubiquitin conjugates reveals that the Rpn10 substrate receptor contributes to the turnover of multiple proteasome targets.** *Mol Cell Proteomics* 2005, **4**:741-751.
46. Ong SE, Foster LJ, Mann M: **Mass spectrometric-based approaches in quantitative proteomics.** *Methods* 2003, **29**:124-130.
47. Ong SE, Blagoev B, Kratchmarova I, Kristensen DB, Steen H, Pandey A, Mann M: **Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics.** *Mol Cell Proteomics* 2002, **1**:376-386.
48. Gygi SP, Rist B, Gerber SA, Turecek F, Gelb MH, Aebersold R: **Quantitative analysis of complex protein mixtures using isotope-coded affinity tags.** *Nat Biotechnol* 1999, **17**:994-999.
49. Catic A, Collins C, Church GM, Ploegh HL: **Preferred in vivo ubiquitination sites.** *Bioinformatics* 2004, **20**:3302-3307.
50. Flick K, Ouni I, Wohlschlegel JA, Capati C, McDonald WH, Yates JR, Kaiser P: **Proteolysis-independent regulation of the transcription factor Met4 by a single Lys 48-linked ubiquitin chain.** *Nat Cell Biol* 2004, **6**:634-641.
51. Finley D, Sadis S, Monia BP, Boucher P, Ecker DJ, Crooke ST, Chau V: **Inhibition of proteolysis and cell cycle progression in a multiubiquitination-deficient yeast mutant.** *Mol Cell Biol* 1994, **14**:5501-5509.
52. Pickart CM: **Targeting of substrates to the 26S proteasome.** *FASEB J* 1997, **11**:1055-1066.
53. Spence J, Sadis S, Haas AL, Finley D: **A ubiquitin mutant with specific defects in DNA repair and multiubiquitination.** *Mol Cell Biol* 1995, **15**:1265-1273.
54. Wang C, Deng L, Hong M, Akkaraju GR, Inoue J, Chen ZJ: **TAK1 is a ubiquitin-dependent kinase of MKK and IKK.** *Nature* 2001, **412**:346-351.
55. Liu Y, Patricelli MP, Cravatt BF: **Activity-based protein profiling: the serine hydrolases.** *Proc Natl Acad Sci USA* 1999, **96**:14694-14699.
56. Greenbaum D, Medzihradsky KF, Burlingame A, Bogoy M: **Epoxide electrophiles as activity-dependent cysteine protease profiling and discovery tools.** *Chem Biol* 2000, **7**:569-581.
57. Hemelaar J, Galardy PJ, Borodovsky A, Kessler BM, Ploegh HL, Ovaa H: **Chemistry-based functional proteomics: mechanism-based activity-profiling tools for ubiquitin and ubiquitin-like specific proteases.** *J Proteome Res* 2004, **3**:268-276.
58. Borodovsky A, Ovaa H, Kolli N, Gan-Erdene T, Wilkinson KD, Ploegh HL, Kessler BM: **Chemistry-based functional proteomics reveals novel members of the deubiquitinating enzyme family.** *Chem Biol* 2002, **9**:1149-1159.
59. Ovaa H, Kessler BM, Rolen U, Galardy PJ, Ploegh HL, Masucci MG: **Activity-based ubiquitin-specific protease (USP) profiling of virus-infected and malignant human cells.** *Proc Natl Acad Sci USA* 2004, **101**:2253-2258.
60. Nijman SM, Huang TT, Dirac AM, Brummelkamp TR, Kerkhoven RM, D'Andrea AD, Bernards R: **The deubiquitinating enzyme USPI regulates the Fanconi anemia pathway.** *Mol Cell* 2005, **17**:331-339.
61. Brummelkamp TR, Nijman SM, Dirac AM, Bernards R: **Loss of the cylindromatosis tumour suppressor inhibits apoptosis by activating NF-kappaB.** *Nature* 2003, **424**:797-801.
62. Denison C, Kirkpatrick DS, Gygi SP: **Proteomic insights into ubiquitin and ubiquitin-like proteins.** *Curr Opin Chem Biol* 2005, **9**:69-75.
63. Welchman RL, Gordon C, Mayer RJ: **Ubiquitin and ubiquitin-like proteins as multifunctional signals.** *Nat Rev Mol Cell Biol* 2005, **6**:599-609.