

MINI-REVIEW

Open Access



Central nervous system effects of 5-HT₇ receptors: a potential target for neurodegenerative diseases

Alejandro Quintero-Villegas^{1,2} and Sergio Iván Valdés-Ferrer^{1,2,3*} 

Abstract

5-HT₇ receptors (5-HT₇R) are the most recently identified among the family of serotonin receptors. Their role in health and disease, particularly as mediators of, and druggable targets for, neurodegenerative diseases, is incompletely understood. Unlike other serotonin receptors, for which abundant preclinical and clinical data evaluating their effect on neurodegenerative conditions exist, the available information on the role of the 5-HT₇R receptor is limited. In this review, we describe the signaling pathways and cellular mechanisms implicated in the activation of the 5-HT₇R; also, we analyze different mechanisms of neurodegeneration and the potential therapeutic implications of pharmacological interventions for 5-HT₇R signaling.

Keywords: 5-HT₇, Alzheimer disease, Parkinson disease, Dementia, Neurodegeneration, Neuroprotection

Introduction

5-hydroxytryptamine (serotonin) 7 receptors (5-HT₇R) are members of the family of 5-HT receptors identified in 1993, but their functional and pathological implications are incompletely understood (Bard et al. 1993; Lovenberg et al. 1993; Ruat et al. 1993). Like other serotonin receptors, their activation is mediated by G-protein-coupled receptor (GPCR) signaling pathways. 5-HT₇R is broadly expressed in the central nervous system (CNS), gastrointestinal tract, and other organs, where they potentially regulate different physiological functions including the sleep–wake cycle, body temperature, nociception, and gastrointestinal motility, to name a few (Sanger 2008; Ciranna and Catania 2014; Chang-Chien et al. 2015). The 5-HT₇R gene is located in the chromosome 10 (q21-124), which contains 3 introns in the coding region, (Gellynck et al. 2013) moreover, 5-HT₇R is expressed in

four different isoforms, being 5-HT₇R_(A), 5-HT₇R_(B) and 5-HT₇R_(D) the ones isolated in humans, and 5-HT₇R_(A), 5-HT₇R_(B) and 5-HT₇R_(C) in rats, with no apparent functional distinction between each isoform (Heidmann et al. 1997). Of interest, 5-HT₇R presents high homology between species (90%) but little homology with other 5-HT receptors (as low as 50%) (Hannon and Hoyer 2008).

Experimental data suggest that 5-HT₇R may be an amenable therapeutic target in neurodegenerative disorders. However, no clinical studies have evaluated the role of 5-HT₇R in neurodegenerative processes, although targeting other serotonin receptors (with drugs such as selective serotonin reuptake inhibitors) has shown little clinical benefit in neurodegenerative conditions (Hüll et al. 2006; Lalut et al. 2017). Here, we aimed to review the potential role of 5-HT₇R in neurodegeneration and their potential therapeutic implications, based on different *in vivo* and *in vitro* pre-clinical studies (Table 1).

*Correspondence: sergio.valdesf@incmnsz.mx

¹ Department of Neurology, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Table 1 Common agonists and antagonists of 5-HT₇R used in preclinical studies

Name	Action mechanism	Administration route (dose)	References
AS-19	Selective full agonist	s.c (5 mg/kg), i.t. (5 µL at 100 µM), i.p. (10 mg/kg)	McDaid et al. (2020), Fields et al. (2015), Albayrak et al. (2013)
LP-12	Selective full agonist	i.t. (10 µL at 0.02–0.2 nM), cultures (300 nM)	Godínez-Chaparro et al. (2012), Samara-jeewa et al. (2014)
LP-44	Selective full agonist	i.p. (1,5 and 10 mg/kg)	Demirkaya et al. (2016)
LP-211	Selective full agonist	i.p. (1,5 and 10 mg/kg), i.p. (0.25 mg/kg), i.p. (0.003–0.3 mg/kg), i.c.v. (0.2 µL at 2–6 mM)	Demirkaya et al. (2016), Liu et al. (2021), Norouzi-Javidan et al. (2016), (Monti et al. 2014)
Methiothepin maleate	Non-specific 5-HT _{1/6/7} R agonist	Culture (10 µM)	Soga et al. (2007)
8-OH-DPAT	Non-specific 5-HT _{1A/7} R agonist	i.p. (0.2–0.4 mg/kg and 1.0 mg/kg)	Cassaday and Thur (2019), Odland et al. (2019)
SB-269970	Competitive selective antagonist, quasi-full inverse agonist	i.p. (10 mg/kg)	Perez-García and Meneses (2005), Liu et al. (2021)
SB-258741	Competitive selective antagonist, partial inverse agonist	s.c. (2.3 mg/kg and 3.5 mg/kg)	Pouzet (2002)
SB-258719	Competitive selective antagonist	i.p. (5 mg/kg)	Brenchat et al. (2011)
HBK-15	Competitive non-selective 5-HT _{1A/3/7} R antagonist	i.p. (1.25 mg/kg) i.v. (1.25 mg/kg)	Pytka et al. (2018)
Lurasidone	Competitive non-selective 5-HT _{2A/7} R antagonist	Microdialysis (3 mg/kg/d)	Okada et al. (2021)

Methods

We performed a comprehensive search of the PubMed database English language literature to identify all original research and review articles regarding 5-HT₇R localization, signaling pathways, effectors, its role in health in the central nervous system, and the pathology of selected neurodegenerative diseases. For that purpose, we included the following Medical Subject Headings (MeSH) and main keywords for searches: 5-HT₇, LP-211, LP-44, LP-21, AS-19, SB-269970, SB-656104-A, 5-HT₇R mechanism of action, 5-HT₇R signaling pathway, 5-HT₇R effect, 5-HT₇R distribution, 5-HT₇R neuroprotection, excitotoxicity, oxidative stress, apoptosis, necrosis, unfolded protein response, endoplasmic reticle stress, amyloid β, tau tangles, tau oligomers, α-synuclein, inflammation, Alzheimer's, Parkinson's, Huntington's, Frontotemporal dementia, dementia, and neurodegeneration. We also reviewed the articles cited in the reference lists of the articles identified during the search. The authors independently reviewed the selected articles. The search included articles available from 1993 (when the receptor was originally cloned) to March 2022.

Distribution in the CNS

5-HT₇R is broadly expressed in different cell types in the CNS, including neurons, astrocytes, and microglia (Shimizu et al. 1998; Mahé et al. 2005; Guseva et al. 2014). CNS expression of 5-HT₇R is not homogeneous; the highest expression occurs in the hippocampus,

amygdala, and hypothalamus (Thomas and Hagan 2004), and comparatively lower density occurs in the dorsal raphe, caudate nucleus, putamen, and substantia nigra (Martín-Cora and Pazos 2004). In the amygdala, 5-HT₇R are present primordially in GABAergic interneurons; in other structures, their expression in specific neurons is not clear, hence of critical importance for therapeutic purposes (Kusek et al. 2021). Of biological and clinical relevance, the regional expression of 5-HT₇R is evolutionarily conserved in mammals (Martín-Cora and Pazos 2004). 5-HT₇R are also expressed in the digestive tract, aorta, and other tissues, exerting immunomodulatory effects as well as other organ-specific effects (Quintero-Villegas and Valdés-Ferrer 2019).

Cellular mechanisms and signaling pathways

Like other GPCR, 5-HT₇R are found in the cell membrane, where they form homodimers and homooligomers, with no known relevant differences between their biological functions (Smith et al. 2011; Guseva et al. 2014). In addition, 5-HT₇R can form heterodimers and heterooligomers with 5-HT_{1A} receptors (5-HT_{1A}R), which in turn lead to diminished activity and increased internalization of 5-HT_{1A} receptors without a discernible effect on 5-HT₇R signaling or activity (Renner et al. 2005; Prasad et al. 2019). This is of biological relevance, as activation of 5-HT_{1A}R results in activation of G_{αi} and reducing levels of cyclic adenosine monophosphate (cAMP) and mitogen-activated protein kinase (MAPK), also known as an extracellular signal-regulated kinase (ERK), hence

antagonizing the effects of the 5-HT₇R-Gas signaling cascade (Zhou et al. 2019).

The activation of 5-HT₇R leads to the initiation of two possible signaling pathways: the *canonical* one, described when the receptor was originally cloned (Lovenberg et al. 1983) acts via Gas (Fig. 1). The activation of this pathway, like in other GPCRs results in the phosphorylation of different adenylyl cyclases (AC) (Baker et al. 1998). This leads to cAMP production, activation of protein kinase A, and, finally, phosphorylation of different proteins, like ERK 1 and ERK 2, Akt, and tropomyosin-related kinase B (Trkb) (Errico et al. 2001; Johnson-Farley et al. 2005; Samarajeewa et al. 2014).

The *Non-canonical* signaling pathway of 5-HT₇R acts via Gα12 (Guseva et al. 2014). This leads to the activation of Rho, Rac, and cell division control protein 42 (Cdc42) all part of the Rho family of small GTPases, which in neurons promote dendrite sprouting, formation of filopodia, and synaptogenesis (Hart, et al. xxx; Kobe et al. 2012a; Speranza et al. 2013; Speranza et al. 2015; Marin and Dityatev 2017). Of relevance, Trkb expression (a brain-derived neurotrophic factor [BDNF] receptor) appears to be enhanced by both Gas and Gα12 (Fig. 1) (Samarajeewa et al. 2014). These signaling pathways may be

of therapeutic relevance for neurodegenerative diseases, although few studies have so far evaluated these effects (Hashemi-Firouzi et al. 2017; Costa et al. 2018; Quintero-Villegas et al. 2019).

Biopathology of neurodegeneration

The biological mechanisms leading to neurodegeneration include many neuronal and glial molecular pathways that result in neuronal damage. Below, we briefly summarize the most common mechanisms involved in neurodegeneration and how they might associate with the known 5-HT₇R effects (Fig. 2; Table 2).

Amyloid β-mediated neurodegeneration

Amyloid β (Aβ) is a key mediator of neurodegeneration in Alzheimer’s disease (AD), inducing damage through multiple pathways (Querfurth and Laferla 2018), some of which overlap with other mechanisms discussed below. Besides AD, Aβ may also play a role in diseases such as frontotemporal dementia (FTD), cerebral amyloid angiopathy, and cerebral amyloidosis (Miller-Thomas et al. 2016). Aβ is a critical source of reactive oxygen species (ROS) and reactive nitrogen species in AD, causing neuronal lipoperoxidation in neurons and thus,





Fig. 1 5-HT₇ and 5-HT_{1A} receptor signaling pathways and oligo/heterodimer formation. 5-HT₇ receptor monomers (in yellow) can form homodimers or homologomers, with the same signaling pathways and cellular effects. 5-HT₇ can also form heterodimers or heterologomers with 5-HT_{1A} (in teal), resulting in the inhibition of the 5-HT_{1A} signaling pathway, with no net effect downstream of 5-HT₇. When activated, 5-HT₇ activates Gas (canonical pathway) with a subsequent signaling cascade that results in the activation of ERK (also known as MAPK) and Akt; in contrast, the activation of Ga12 activates mTOR and different Rho family small GTPases. As illustrated, the phosphorylation of Trkb is mediated by both G proteins. AC adenylyl cyclase, cAMP cyclic adenosine monophosphate, Cdc42 cell division control protein 42 homolog, ERK extracellular signal-regulated kinases, MAPK mitogen-activated protein kinases, mTOR mammalian target of rapamycin, Trkb Tropomyosin receptor kinase B



neurodegeneration (Bernal-Mondragón et al. 2013). Also, Aβ helps the formation of voltage-independent, cation channels in the lipid membranes, which could lead to excitotoxicity-mediated neurodegeneration (Arispe et al. 1993). Chronically Aβ-stimulated microglia releases multiple pro-inflammatory cytokines, such as interleukin (IL)-1, IL-6, and tumor necrosis factor (TNF)-α, which

induces pathological changes in the CNS (Heppner et al. 2015). Aβ-stimulated microglia increases neuronal damage and further accumulation of Aβ, an effect mediated by receptors for advanced glycation end products (Fang et al. 2010). Although not directly associated with neuronal death, Aβ impairs synaptic function and synaptogenesis and dysregulates neurotransmitter levels in the synaptic cleft, contributing to

Table 2 Mechanisms of neuronal damage, and possible beneficial effects of 5-HT₇ receptors agonism

Neurodegeneration mechanism	5-HT ₇ possible role	References
Excitotoxicity	Activation of MAPK/ERK and PI3/Akt/GSK3b protects against glutamate-induced damage Decreased expression of NR2B and NR1 subunits of NMDA glutamate receptors Increased expression of superoxide dismutase and glutathione	Jiang et al. (2000); Pi et al. (2004); Li et al. (2005) Vasefi et al. (2013a) Yuksel et al. (2019)
Oxidative stress	No study evaluated effect in the CNS In a sepsis-induced lung injury, 5-HT ₇ receptor agonism decreased ROS burden 5-HT ₇ antagonism decreased oxidative burden in bleomycin-induced pulmonary fibrosis 5-HT ₇ activation enhances microsome stability towards oxidative metabolism (Lacivita et al., 2016a) ERK and Akt protect PC12 cells from oxidative damage	Cadirci et al. (2013) Tawfik and Makary (2017) Lacivita et al. (2016a) Ong et al. (2016)
Apoptosis	5-HT ₇ receptor agonism reduces apoptosis in the streptozotocin-induced AD model	Hashemi-Firouzi et al. (2017)
Long term depression/ potentiation impairment	5-HT ₇ KO mice display LTP impairment 5-HT ₇ agonism reduces mGluR-dependent LTD	Roberts et al. (2004) Costa et al. (2012)
Synaptic impairment	5-HT ₇ agonism increases dendritic density and synaptogenesis in the cortical and striatal forebrain 5-HT ₇ agonism induces dendritic sprouting and neurite enlargement	Speranza et al. (2013, 2015, 2017) Kvachnina et al. (2005); Canese et al. (2015)
Neurotrophin depletion	5-HT ₇ agonism increases PDGF-β 5-HT ₇ agonism increases the expression and affinity of trk-B	Vasefi et al. (2013b) Samarajeewa et al. (2014)

the symptoms in AD (Querfurth and Laferla 2018; Cai 2019; Ding et al. 2019).

5-HT₇R agonism with LP-211 (a highly selective agonist) reverts neuronal damage and cognitive impairment induced by Aβ (Quintero-Villegas et al. 2019). Aβ induces neurotoxicity through several mechanisms including apoptosis, excitotoxicity, and oxidative stress (Querfurth and Laferla 2018; Bernal-Mondragón et al. 2013). In a streptozotocin-mediated neurodegeneration murine model for AD, the intracerebroventricular (ICV) treatment of AS-19 (a 5-HT₇R selective agonist) reduced long-term potentiation (LTP) impairment and apoptosis in hippocampal (Hashemi-Firouzi et al. 2017).

The exact mechanism of 5-HT₇R-mediated neuroprotection in Aβ-induced neurodegeneration is currently under investigation, an effect that is likely to be mediated through multiple mechanisms.

Excitotoxicity

This is an important cause of neuronal damage in neurodegenerative diseases, including AD, stroke, Huntington's disease (HD), and Parkinson's disease (PD) (Martire et al. 2013; Lai et al. 2014; Iovino et al. 2020). Persistent excitatory -mainly glutamatergic- stimuli lead to altered calcium homeostasis, resulting in oxidative stress, mitochondrial dysfunction, disturbances in protein turnover, inflammation, and caspase-mediated apoptosis (Binvignat and Olloquequi 2020). In vitro studies suggest that the MAPK/ERK and phosphatidylinositol-3/

Akt/Glycogen synthase kinase 3b pathways are closely associated with protection against glutamate-induced damage (Jiang et al. 2000; Pi et al. 2004; Li et al. 2005); modulation of these kinases via 5-HT₇R Gαs could have therapeutic implications. 5-HT₇R activation leads to a decrease in the NR2B and NR1 subunits of the N-methyl-D-aspartate (NMDA) glutamate receptors, thus protecting against glutamate-mediated excitotoxicity (Vasefi et al. 2013a). Treatment with LP-44, a 5-HT₇R-specific agonist, protects human neuroblastoma SH-SY5Y cells against glutamate-mediated damage in an in-vitro model, also increasing superoxide dismutase and glutathione while decreasing TNF-α, and caspase-3 and -9 (Yuksel et al. 2019). Moreover, 5-HT₇R modulate glutamate-NMDA activity in a time-dependent manner; while the acute activation of 5-HT₇R by LP-12, a selective agonist of 5-HT₇R, enhances NMDA activity, the chronic activation inhibits its activity (Vasefi et al. 2013b).

Oxidative stress

This can lead to membrane damage and neuronal death. (Bernal-Mondragón et al. 2013) Although, oxidative phosphorylation in mitochondria produces ROS, reactive nitrogen species, carbon-centered radicals, and sulfur-centered radicals (Pero et al. 1990), and these by-products are considered necessary for neuronal function and development (Salim 2017); increase of their levels, beyond a physiological threshold, are considered deleterious. Multiple in vitro studies have demonstrated that

high levels of ROS reduce long-term potentiation, synaptic signaling, and brain plasticity (Salim 2017; O'Dell et al. 1991; Stevens and Wang 1993). Moreover, oxidative stress damages macromolecules, mainly lipid-rich structures such as membranes, via lipoperoxidation (Salim 2017).

Although no study so far has evaluated the potential effect of 5-HT₇R in oxidative stress damage in the CNS, a study evaluating sepsis-induced lung injury demonstrated that 5-HT₇R agonism decreased ROS burden (Cadirci et al. 2013). In contrast, the antagonism of the 5-HT₇R by SB-269970 decreased oxidative burden in bleomycin-induced pulmonary fibrosis (Tawfik and Makary 2017). Also, 5-HT₇R activation by LP-44 enhances microsome stability towards oxidative metabolism (Lacivita et al. 2016a). As mentioned before, 5-HT₇R regulates the activation of ERK and Akt, and these kinases have biological effects on oxidative stress injury protection in PC12 cells (Ong et al. 2016). Thus, 5-HT₇R may have therapeutic implications in ROS-induced neurodegeneration, something that needs to be experimentally assessed.

Apoptosis

Neuronal cell death is a major pathological characteristic of every neurodegenerative disease, whether via apoptosis or necrosis. In AD, both phosphorylated tau protein and A β aggregates induce apoptosis in *in vitro* studies, with contradictory effects in tissular studies. Other proteins, including α -Synuclein (in PD or Lewy body dementia), or mutant huntingtin protein (in HD) also induce neuronal cell death via multiple mechanisms, including apoptosis (Chi et al. 2018).

It is also important to note, that apoptosis is closely related to excitotoxicity and oxidative stress; thus, these pathological effects are somewhat overlapped (Yuksel et al. 2019; Loh et al. 2006; Nicholls et al. 2007). The 5-HT₇R agonist AS-19 reduces apoptosis in the streptozotocin-induced AD model (Hashemi-Firouzi et al. 2017). While no other study has yet evaluated this effect in the CNS, methiothepin maleate (a non-specific 5-HT_{1/6/7}R agonist) prevents monocyte activation via ERK 1/2 and Nuclear factor- κ B (Soga et al. 2007).

Long-term potentiation and long-term depression (LTD) impairment

Impairment in LTP and LTD have been extensively described in many types of dementia, such as AD (Skaper et al. 2017), PD (Marinelli et al. 2017), and HD (Filippo et al. 2007), to name a few, with a strong correlation with the cognitive symptoms in each disease. LTP and LTD are crucial for memory formation, and impairment in these are associated with amnesic and psychiatric symptoms (Loprinzi 2020).

Like other CNS effects, the role of 5-HT₇R in LTP is controversial. Because chronically stimulated neurons by 5-HT₇R agonists show a reduction in the expression of NMDA glutamate receptors, 5-HT₇R has been associated with a reduction in LTP (Kobe et al. 2012b), however, 5-HT₇R knock-out mice also display an impairment in LTP, suggesting that 5-HT₇R receptors at a baseline state are necessary for LTP (Roberts et al. 2004). Regarding LTD, 5-HT₇R agonism by 8-OH-DPAT (a 5-HT_{1A/7}R agonist) reduced mGluR mediated LTD (Costa et al. 2012). Finally, LP-211 induces LTD on the parallel fiber-Purkinje cell synapse via Protein kinase C-MAPK pathway while impairing LTP, and pharmacological antagonism with SB-269970 decreases LTD (Lippiello et al. 2016).

Synaptogenesis and brain plasticity reduction

In many neurodegenerative diseases, reductions in synaptogenesis, brain plasticity, and dendritic sprouting are hallmarks of severity and progression (Querfurth and LaFerla 2010). So far, no treatment strategies have been shown to reverse that.

In vitro, activation of 5-HT₇R in cortical and striatal forebrain neurons using LP-211 increases dendritic spine density and synaptogenesis, an effect that is abrogated by the genetical or pharmacological blockade of 5-HT₇R (Speranza et al. 2013; Speranza et al. 2015; Speranza et al. 2017). 5-HT₇R stimulation induces dendritic sprouting and neurite enlargement (Kvachnina et al. 2005; Rojas et al. 2014; Canese et al. 2015), an effect probably mediated by effectors of both, Gas and Ga12 signaling pathways (Volpicelli et al. 2014).

Neurotrophin depletion

Neurotrophins like BDNF and platelet-derived growth factor (PDGF)- β are necessary for the development and correct function of the CNS. Depletion or altered signaling occurs in neurodegenerative diseases (Kashyap et al. 2018) AD, PD, HD, and FTD are associated with a reduction in the expression of BDNF and other neurotrophins, where the modulation of BDNF could have a potential therapeutic role (Palasz et al. 2020; Schulte-Herbrüggen et al. 2008; Alberch et al. 2002; Huey et al. 2020). Interestingly, 5-HT₇R modulates both neurotrophins and their receptors. 5-HT₇R activation by LP-12 leads to an increase in PDGF- β , with increased protection against glutamate-mediated excitotoxicity (Vasefi et al. 2013a). In addition, 5-HT₇R agonism by LP-12 increases the expression and affinity of the tropomyosin-related kinase B receptor, one of the receptors for BDNF (Samarajeewa et al. 2014).

Immune-mediated damage

The specific role of 5-HT₇R as neuro-immune mediators is still debated. 5-HT₇R is expressed broadly by different immune cells, including monocytes, lymphocytes, and dendritic cells, but its anti-inflammatory potential has been shown in some but not all studies (Quintero-Villegas and Valdés-Ferrer 2019).

Potential disease-specific role of 5-HT₇R in neuropsychiatric illness

Neuronal hyperexcitability and seizures

Epilepsy-induced neuronal damage shares pathophysiological neurodegenerative features with dementias and other CNS diseases, such as increased inflammation and excitotoxicity (Nikiforuk 2015). Epilepsy is prevalent in sufferers of CNS diseases, including AD, PD, or HD, and, when present, indicates a higher burden of neurodegeneration (Cano et al. 2021).

5-HT₇R manipulation has shown controversial results in pre-clinical studies of epilepsy. Non-specific pharmacological blockade of 5-HT₇R reduces the prevalence of audiogenic seizures in a DBA/2J rat model (Bourson et al. 1997).

Additionally, SB-258719, an antagonist of 5-HT₇R showed a benefit in reducing spontaneous epileptic activity in a WAG/Rij rat model of absence seizures (Graf et al. 2004). Finally, in a pilocarpine-induced rat model of temporal lobe epilepsy, AS-19 increased epileptic activity, whereas SB-269970, an antagonist of 5-HT₇R reduced seizure activity. Interestingly, the expression of 5-HT₇R was higher in the epilepsy group, compared to the control group (Yang et al. 2012). However, in a pilocarpine-induced model of epilepsy, the density of 5-HT₇R decreased in the hippocampus, especially in the dentate gyrus (Núñez-Ochoa et al. 2021). Hence, the role of 5-HT₇R in epilepsy is still unclear but avidly explored. The controversial finding may represent that 5-HT₇R plays different roles in different models of epilepsy. Evaluation of 5-HT₇R tissue expression in specimens obtained from patients undergoing surgical excision of epileptic foci may help to clarify this controversy.

Mood disorders

The relationship between mood disorders and cognitive disorders is a topic of continuous research. The prevalence of depression and anxiety is higher among patients with dementia, and patients with depression have a higher prevalence of dementia (Lyketsos et al. 2002). Of therapeutic relevance, even mild levels of depression can impact substantially the functionality and quality of life of patients with dementia. Thus, treating these symptoms may be crucial in the management of dementia (Gutzmann and Qazi 2015).

In a rat model of stress using forced swim and tail suspension, pharmacological and genetic blockade of 5-HT₇R reduced depressive symptoms and improved REM sleep (Hedlund et al. 2005). 5-HT₇R KO mice show improved mobility in the Porsolt swim test; however, the pharmacological blockade by SB-258719 only had the same results when rats were tested in the dark (Guscott et al. 2005). Similar results were observed with SB-269970 in depression and anxiety, with an effect similar to the one observed with imipramine (Wesołowska et al. 2006). In experimental depression, pharmacological blockade of 5-HT₇R seems to have a rapid effect in reducing depressive symptoms (Mnie-Filali et al. 2011). Altogether, while more data is needed before moving to the clinical trial setting, these experimental models suggest that 5-HT₇R may be a druggable target for depression.

The calcium-binding protein S100B interacts with 5-HT₇R and negatively regulates inducible cAMP accumulation; its overexpression in transgenic mice is associated with depressive-like symptoms, which are reversed by the administration of SB269970 (Stroth and Svenningsson 2015).

The non-specific blockade of 5-HT₇R with HBK-15 (5-HT₇R, 5-HT_{1A}R, and 5-HT₃R antagonist) (Pytko et al. 2018), aryl sulfonamide derivatives of dihydro benzofuran oxyethyl piperidines (a2 and 5-HT₇R antagonist) (Canale et al. 2021), and lurasidone (5-HT_{2A}R and 5-HT₇R antagonist) (Woo et al. 2013), to name a few, have similar effects regarding depression and anxiety.

The contradictory memory effect of 5-HT₇R

Evidence about the effect of 5-HT₇R on learning and memory is still contradictory. Systemic administration of LP-211 and AS-19 revert scopolamine-induced amnesia and enhanced auto-shaping training learning respectively in rats; whereas pharmacologic blockade with SB-269970 has the opposite effect (Perez-García and Meneses 2005). Accordingly, LP-211 administrated intraperitoneally improves long-term memory, with conditioned responses up to 80%, compared to 20–30% in the control groups, with no effect on short-term memory, and reverts scopolamine-induced memory impairment, something associated with increased cAMP levels (Meneses et al. 2015). Moreover, 5-HT₇R agonism appears to revert the memory impairment mediated by 5-HT_{1A}R, based on the fact that the co-administration of SB-269970 and 8-OH-DPAT (a 5-HT₇R/5-HT_{1A}R agonist) caused greater performance impairment in contextual learning than the administration of 8-OH-DPAT alone (Eriksson et al. 2008).

On the other hand, SB-269970 improved reference memory in a radial arm maze task (Gasbarri et al. 2008). In agreement, the administration of SB-656104-A (a

5-HT₇R-specific antagonist) reverses dizocilpine-induced memory impairment induced, with significant differences in thigmotaxic swimming in the Morris water maze test (Horisawa et al. 2011). We have observed that intracerebroventricular injection of LP-211 reverts memory impairment induced by A β without a discernible effect on healthy rats (Quintero-Villegas et al. 2019). The answer to this paradox is inconclusive with the current evidence (Meneses 2014; Stiedl et al. 2015; Zareifopoulos and Papatheodoropoulos 2016), suggesting a memory type-specific role of 5-HT₇R.

Lack of clinical trial-derived data

No clinical trials with 5-HT₇R-specific modulation for neurodegenerative studies have been performed. However, the potential effect of non-specific serotonin activation in neurodegenerative diseases is a highly studied topic and the outflow of human data may start shortly. Interestingly, lurasidone, a 5-HT₇R antagonist, with an effect on other serotonin receptors and D2 dopamine receptor (Meltzer et al. 2020), has been FDA approved for the treatment of schizophrenia (Okubo et al. 2021).

Conclusion

At this point, no 5-HT₇R-specific drugs have been evaluated for neurodegenerative diseases, and few studies have experimentally evaluated their potential therapeutic effects. Hence, elucidating the effect of 5-HT₇R in health and (predominantly neurodegenerative) diseases may have vast translational and therapeutic implications. 5-HT₇R agonists may be neuroprotective by acting at multiple levels, including the reduction of excitotoxicity and oxidative stress, synaptic remodeling, regulation of neurotrophic factors, or immunomodulation, to name a few. The need for more studies, both experimental and clinical, before reaching conclusions about a therapeutic role for this serotonin receptor cannot be overemphasized.

Abbreviations

5-HT: 5-Hydroxytryptamine/serotonin; 5-HT_{1A}R: 5-HT_{1A} receptor; 5-HT₇R: 5-HT₇ receptor; AD: Alzheimer's disease; A β : Amyloid β ; BDNF: Brain-derived neurotrophic factor; cAMP: Cyclic adenosine monophosphate; Cdc42: Cell division control protein 42; CNS: Central nervous system; ERK: Extracellular signal-regulated kinase; FTD: Frontotemporal dementia; GPCR: G-protein-coupled receptor; HD: Huntington's disease; ICV: Intracerebroventricular; IL: Interleukin; LTD: Long-term depression; LTP: Long-term potentiation; MAPK: Mitogen-activated protein kinase; PD: Parkinson's disease; PDGF: Platelet-derived growth factor; ROS: Reactive oxygen species; TNF: Tumor necrosis factor; Trkb: Tropomyosin-related kinase B.

Acknowledgements

The figures on this manuscript were created with biorender.com.

Author contributions

Both SIVF and AQV contributed in an equal manner to the making of this article. Both the authors read and approved the final manuscript.

Author's information

AQV: Medical resident and Assistant Investigator, Laboratory of Neurobiology of Systemic Illness, Department of Neurology Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City. SIVF: Head and Principal Investigator, Laboratory of Neurobiology of Systemic Illness, Department of Neurology Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, and Institute of Bioelectronic Medicine, Feinstein Institutes for Medical Research, Manhasset, NY.

Funding

This manuscript was supported by Grants 289788 and A1-S-18342 from Consejo Nacional de Ciencia y Tecnología (CONACyT), Mexico (to SIVF).

Availability of data and materials

All data were extracted from the PubMed database using the MeSH established in the "Methods" section.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

All the authors gave their approval for submission.

Competing interests

The authors declare no conflicts of interest while making this article.

Author details

¹Department of Neurology, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico. ²Department of Infectious Diseases, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico. ³Institute of Bioelectronic Medicine, Feinstein Institutes for Medical Research, Manhasset, NY, USA.

Received: 29 March 2022 Accepted: 9 June 2022

Published online: 20 June 2022

References

- Albayrak A, et al. Inflammation and peripheral 5-HT₇ receptors: the role of 5-HT₇ receptors in carrageenan induced inflammation in rats. *Eur J Pharmacol.* 2013;715:270–9.
- Alberch J, Pérez-Navarro E, Canals JM. Neuroprotection by neurotrophins and GDNF family members in the excitotoxic model of Huntington's disease. *Brain Res Bull.* 2002;57:817–22.
- Arispe N, Pollard HB, Rojas E. Giant multilevel cation channels formed by Alzheimer disease amyloid β -protein [A β P-(1–40)] in bilayer membranes. *Proc Natl Acad Sci USA.* 1993;90:10573–7.
- Baker LP, et al. Stimulation of type 1 and type 8 Ca²⁺/calmodulin-sensitive adenylyl cyclases by the G(s)-coupled 5-hydroxytryptamine subtype 5-HT(7A) receptor. *J Biol Chem.* 1998;273:17469–76.
- Bard JA et al. Cloning of a novel human serotonin receptor (5-HT₇) positively linked to adenylyl cyclase. 1993;268: 23422–23426.
- Bernal-Mondragón C, Rivas-Arancibia S, Kendrick KM, Guevara-Guzmán R. Estradiol prevents olfactory dysfunction induced by A- β 25–35 injection in hippocampus. *BMC Neurosci.* 2013;14.
- Binvignat O, Olloquequi J. Excitotoxicity as a target against neurodegenerative processes. *Curr Pharm Des.* 2020;26:1251–62.
- Bourson A, et al. Correlation between 5-HT₇ receptor affinity and protection against sound-induced seizures in DBA/2J mice. *Naunyn-Schmiedeberg Arch Pharmacol.* 1997;356:820–6.
- Brenchat A, Ejarque M, Zamanillo D, Vela JM, Romero L. Potentiation of morphine analgesia by adjuvant activation of 5-HT₇ receptors. *J Pharmacol Sci.* 2011;116:388–91.
- Cadirci E, et al. Peripheral 5-HT₇ receptors as a new target for prevention of lung injury and mortality in septic rats. *Immunobiology.* 2013;218:1271–83.

- Cai M et al. Spinosin attenuates Alzheimer's disease-associated synaptic dysfunction via regulation of plasmin activity. 2019;6: 1–6.
- Canale V, et al. Design, sustainable synthesis and biological evaluation of a novel dual $\alpha 2A/5$ -HT7 receptor antagonist with antidepressant-like properties. *Molecules*. 2021;26:828.
- Canese R, et al. Persistent modification of forebrain networks and metabolism in rats following adolescent exposure to a 5-HT7 receptor agonist. *Psychopharmacology*. 2015;232:75–89.
- Cano A, et al. Epilepsy in neurodegenerative diseases: related drugs and molecular pathways. *Pharmaceuticals (basel)*. 2021;14:1057.
- Cassaday HJ, Thur KE. Intraperitoneal 8-OH-DPAT reduces competition from contextual but not discrete conditioning cues. *Pharmacol Biochem Behav*. 2019;187: 172797.
- Chang-Chien CC, Hsin LW, Su MJ. Activation of serotonin 5-HT7 receptor induces coronary flow increase in isolated rat heart. *Eur J Pharmacol*. 2015;748:68–75.
- Chi H, Chang H-Y, Sang T-K. Neuronal cell death mechanisms in major neurodegenerative diseases. *Int J Mol Sci*. 2018;19:3082.
- Ciranna L, Catania MV. 5-HT7 receptors as modulators of neuronal excitability, synaptic transmission and plasticity: physiological role and possible implications in autism spectrum disorders. *Front Cell Neurosci*. 2014;8:1–17.
- Costa L, et al. Activation of 5-HT7 serotonin receptors reverses metabotropic glutamate receptor-mediated synaptic plasticity in wild-type and *fmr1* knockout mice, a model of fragile X syndrome. *Biol Psychiat*. 2012;72:924–33.
- Costa L, et al. Activation of serotonin 5-HT7 receptors modulates hippocampal synaptic plasticity by stimulation of adenylate cyclases and rescues learning and behavior in a mouse model of fragile X syndrome. *Front Mol Neurosci*. 2018;11:1–14.
- Demirkaya K, et al. Selective 5-HT7 receptor agonists LP 44 and LP 211 elicit an analgesic effect on formalin-induced orofacial pain in mice. *J Appl Oral Sci*. 2016;24:218–22.
- Di Filippo M, Tozzi A, Picconi B, Ghiglieri V, Calabresi P. Plastic abnormalities in experimental Huntington's disease. *Curr Opin Pharmacol*. 2007;7:106–11.
- Ding Y, et al. Amyloid beta oligomers target to extracellular and intracellular neuronal synaptic proteins in Alzheimer's disease. *Front Neurol*. 2019;10:1–16.
- Eriksson TM, Golkar A, Ekström JC, Svenningsson P, Ögren SO. 5-HT7 receptor stimulation by 8-OH-DPAT counteracts the impairing effect of 5-HT1A receptor stimulation on contextual learning in mice. *Eur J Pharmacol*. 2008;596:107–10.
- Errico M, Crozier RA, Plummer MR, Cowen DS. 5-HT7 receptors activate the mitogen activated protein kinase extracellular signal related kinase in cultured rat hippocampal neurons. *Neuroscience*. 2001;102:361–7.
- Fang, et al. RAGE-dependent signaling in microglia contributes to neuroinflammation, A β accumulation, and impaired learning/memory in a mouse model of Alzheimer's disease. *FASEB J*. 2010;24:1043–55.
- Fields DP, Springborn SR, Mitchell GS. Spinal 5-HT7 receptors induce phrenic motor facilitation via EPAC-mTORC1 signaling. *J Neurophysiol*. 2015;114:2015–22.
- Gasbarri A, Cifariello A, Pompili A, Meneses A. Effect of 5-HT7 antagonist SB-269970 in the modulation of working and reference memory in the rat. 2008; 195: 164–170.
- Gellynck E, et al. The serotonin 5-HT7 receptors: two decades of research. *Exp Brain Res*. 2013;230:555–68.
- Godínez-Chaparro B, López-Santillán FJ, Orduña P, Granados-Soto V. Secondary mechanical allodynia and hyperalgesia depend on descending facilitation mediated by spinal 5-HT₄, 5-HT₆ and 5-HT₇ receptors. *Neuroscience*. 2012;222:379–91.
- Graf M, Jakus R, Kantor S, Levay G, Bagdy G. Selective 5-HT1A and 5-HT7 antagonists decrease epileptic activity in the WAG/Rij rat model of absence epilepsy. *Neurosci Lett*. 2004;359:45–8.
- Guscott M, et al. Genetic knockout and pharmacological blockade studies of the 5-HT7 receptor suggest therapeutic potential in depression. *Neuropharmacology*. 2005;48:492–502.
- Guseva D, Wirth A, Ponimaskin E. Cellular mechanisms of the 5-HT₇ receptor-mediated signaling. *Front Behav Neurosci*. 2014;8:1–8.
- Gutzmann H, Qazi A. Depression associated with dementia. *Z Gerontol Geriatr*. 2015;48:305–11.
- Hannon J, Hoyer D. Molecular biology of 5-HT receptors. *Behav Brain Res*. 2008;195:198–213.
- Hart MJ et al. Exchange activity of p115 RhoGEF by Ga. 1998;280: 0–2.
- Hashemi-Firouzi N, Komaki A, Soleimani Asl S, Shahidi S. The effects of the 5-HT7 receptor on hippocampal long-term potentiation and apoptosis in a rat model of Alzheimer's disease. *Brain Res Bull*. 2017;135:85–91.
- Hedlund PB, Huitron-Resendiz S, Henriksen SJ, Sutcliffe JG. 5-HT7 receptor inhibition and inactivation induce antidepressantlike behavior and sleep pattern. *Biol Psychiatry*. 2005;58:831–7.
- Heidmann DE, Metcalf MA, Kohen R, Hamblin MW. Four 5-hydroxytryptamine7 (5-HT7) receptor isoforms in human and rat produced by alternative splicing: species differences due to altered intron–exon organization. *J Neurochem*. 1997;68:1372–81.
- Heppner FL, Ransohoff RM, Becher B. Immune attack: the role of inflammation in Alzheimer disease. *Nat Rev Neurosci*. 2015;16:358–72.
- Horisawa T, et al. The effects of selective antagonists of serotonin 5-HT7 and 5-HT1A receptors on MK-801-induced impairment of learning and memory in the passive avoidance and Morris water maze tests in rats: mechanistic implications for the beneficial effects of the novel. *Behav Brain Res*. 2011;220:83–90.
- Huey ED, et al. Effect of functional BDNF and COMT polymorphisms on symptoms and regional brain volume in frontotemporal dementia and corticobasal syndrome. *J Neuropsychiatry Clin Neurosci*. 2020;32:362–9.
- Hüll M, Berger M, Heneka M. Disease-modifying therapies in Alzheimer's disease: how far have we come? *Drugs*. 2006;66:2075–93.
- Iovino L, Tremblay ME, Civiero L. Glutamate-induced excitotoxicity in Parkinson's disease: the role of glial cells. *J Pharmacol Sci*. 2020;144:151–64.
- Jiang Q, Gu Z, Zhang G, Jing G. Diphosphorylation and involvement of extracellular signal-regulated kinases (ERK1/2) in glutamate-induced apoptotic-like death in cultured rat cortical neurons. *Brain Res*. 2000;857:71–7.
- Johnson-Farley NN, Kertesz SB, Dubyak GR, Cowen DS. Enhanced activation of Akt and extracellular-regulated kinase pathways by simultaneous occupancy of Gq-coupled 5-HT2A receptors and Gs-coupled 5-HT7A receptors in PC12 cells. *J Neurochem*. 2005;92:72–82.
- Kashyap MP, Roberts C, Waseem M, Tyagi P. Drug targets in neurotrophin signaling in the central and peripheral nervous system. *Mol Neurobiol*. 2018;55:6939–55.
- Kobe F, et al. 5-HT7R/G12 signaling regulates neuronal morphology and function in an age-dependent manner. *J Neurosci*. 2012a;32:2915–30.
- Kobe F, et al. 5-HT 7R/G 12 signaling regulates neuronal morphology and function in an age-dependent manner. *J Neurosci*. 2012b;32:2915–30.
- Kusek M, et al. 5-HT7 receptors enhance inhibitory synaptic input to principal neurons in the mouse basal amygdala. *Neuropharmacology*. 2021;198: 108779.
- Kvachnina E, et al. 5-HT7 receptor is coupled to Ga subunits of heterotrimeric G12-protein to regulate gene transcription and neuronal morphology. *J Neurosci*. 2005;25:7821–30.
- Lacivita E, et al. Structural modifications of the serotonin 5-HT7 receptor agonist *N*-(4-cyanophenylmethyl)-4-(2-biphenyl)-1-piperazinehexanamide (LP-211) to improve in vitro microsomal stability: a case study. *Eur J Med Chem*. 2016a;120:363.
- Lai TW, Zhang S, Wang YT. Excitotoxicity and stroke: identifying novel targets for neuroprotection. *Prog Neurobiol*. 2014;115:157–88.
- Lalut J, Karila D, Dallemagne P, Rochais C. Modulating 5-HT4 and 5-HT6 receptors in Alzheimer's disease treatment. *Future Med Chem*. 2017;9:781–95.
- Li W, et al. Novel dimeric acetylcholinesterase inhibitor bis(7)-tacrine, but not donepezil, prevents glutamate-induced neuronal apoptosis by blocking *N*-methyl-D-aspartate receptors. *J Biol Chem*. 2005;280:18179–88.
- Lippiello P, et al. The 5-HT7 receptor triggers cerebellar long-term synaptic depression via PKC-MAPK. *Neuropharmacology*. 2016;101:426–38.
- Liu B, et al. Effects of 5-HT7 receptors on circadian rhythm of mice anesthetized with isoflurane. *Chronobiol Int*. 2021;38:38–45.
- Loh KP, Huang SH, Silva RD, Tan BKH, Zhu YZ. Oxidative stress: apoptosis in neuronal injury. *Curr Alzheimer Res*. 2006;3:327–37.
- Loprinzi PD. Effects of exercise on long-term potentiation in neuropsychiatric disorders. *Adv Exp Med Biol*. 2020;1228:439–51.
- Lovenberg TW et al. A novel adenylyl cyclase-activating serotonin receptor (5-HT7) implicated in the regulation of mammalian circadian rhythms. 1993;11: 449–458.

- Lyketsos CG, et al. Prevalence of neuropsychiatric symptoms in dementia and mild cognitive impairment: results from the cardiovascular health study. *JAMA*. 2002;288:1475–83.
- Mahé C, et al. Serotonin 5-HT7 receptors coupled to induction of interleukin-6 in human microglial MC-3 cells. *Neuropharmacology*. 2005;49:40–7.
- Marin P, Dityatev A. 5-HT7 receptor shapes spinogenesis in cortical and striatal neurons: an editorial highlight for 'Serotonin 5-HT7 receptor increases the density of dendritic spines and facilitates synaptogenesis in forebrain neurons'. *J Neurochem*. 2017;141:644–6.
- Marinelli L, Quartarone A, Hallett M, Frazzitta G, Ghilardi MF. The many facets of motor learning and their relevance for Parkinson's disease. *Clin Neurophysiol off J Int Fed Clin Neurophysiol*. 2017;128:1127–41.
- Martín-Cora FJ, Pazos A. Autoradiographic distribution of 5-HT7 receptors in the human brain using [3H]mesulergine: comparison to other mammalian species. *Br J Pharmacol*. 2004;141:92–104.
- Martire A, et al. BDNF prevents NMDA-induced toxicity in models of Huntington's disease: the effects are genotype specific and adenosine A2A receptor is involved. *J Neurochem*. 2013;125:225–35.
- McDaid J, Mustaly-Kalimi S, Stutzmann GE. Ca²⁺ dyshomeostasis disrupts neuronal and synaptic function in Alzheimer's disease. *Cells*. 2020;9(12):1–25.
- Meltzer HY, Share DB, Jayathilake K, Salomon RM, Lee MA. Lurasidone improves psychopathology and cognition in treatment-resistant schizophrenia. *J Clin Psychopharmacol*. 2020;40:240–9.
- Meneses A. Memory formation and memory alterations: 5-HT6 and 5-HT7 receptors, novel alternative. *Rev Neurosci*. 2014;25:325–56.
- Meneses A, et al. 5-HT7 receptor activation: procognitive and anti-amnesic effects. *Psychopharmacology*. 2015;232:595–603.
- Miller-Thomas MM, et al. Multimodality review of amyloid-related diseases of the central nervous system. *Radiographics*. 2016;36:1147–63.
- Mnie-Filali O, et al. Pharmacological blockade of 5-HT7 receptors as a putative fast acting antidepressant strategy. *Neuropsychopharmacol off Publ Am Coll Neuropsychopharmacol*. 2011;36:1275–88.
- Monti JM, Leopoldo M, Jantos H. Systemic administration and local microinjection into the central nervous system of the 5-HT7 receptor agonist LP-211 modify the sleep-wake cycle in the rat. *Behav Brain Res*. 2014;259:321–9.
- Nicholls DG, Johnson-Cadwell L, Vesce S, Jekabsons M, Yadava N. Bioenergetics of mitochondria in cultured neurons and their role in glutamate excitotoxicity. *J Neurosci Res*. 2007;85:3206–12.
- Nikiforuk A. Targeting the serotonin 5-HT7 receptor in the search for treatments for CNS disorders: Rationale and progress to date. *CNS Drugs*. 2015;29:265–75.
- Norouzi-Javidan A, et al. Effect of 5-HT7 receptor agonist, LP-211, on micturition following spinal cord injury in male rats. *Am J Transl Res*. 2016;8:2525–33.
- Núñez-Ochoa MA, Chiprés-Tinajero GA, Medina-Ceja L. Evaluation of the hippocampal immunoreactivity of the serotonin 5-HT1A, 5-HT2 and 5-HT7 receptors in a pilocarpine temporal lobe epilepsy rat model with fast ripples. *NeuroReport*. 2021;32:306–11.
- O'Dell TJ, Hawkins RD, Kandel ER, Arancio O. Tests of the roles of two diffusible substances in long-term potentiation: evidence for nitric oxide as a possible early retrograde messenger. *Proc Natl Acad Sci USA*. 1991;88:11285–9.
- Odland AU, Jessen L, Fitzpatrick CM, Andreassen JT. 8-OH-DPAT induces compulsive-like deficit in spontaneous alternation behavior: reversal by MDMA but not citalopram. *ACS Chem Neurosci*. 2019;10:3094–100.
- Okada M, Matsumoto R, Yamamoto Y, Fukuyama K. Effects of subchronic administrations of vortioxetine, lurasidone, and escitalopram on thalamocortical glutamatergic transmission associated with serotonin 5-HT7 receptor. *Int J Mol Sci*. 2021;22:1351.
- Okubo R, Hasegawa T, Fukuyama K, Shiroyama T, Okada M. Current limitations and candidate potential of 5-HT7 receptor antagonism in psychiatric pharmacotherapy. *Front Psych*. 2021;12:623684.
- Ong Q, et al. The timing of Raf/ERK and AKT activation in protecting PC12 cells against oxidative stress. *PLoS ONE*. 2016;11:e0153487.
- Palasz E, et al. BDNF as a promising therapeutic agent in Parkinson's disease. *Int J Mol Sci*. 2020;21:1170.
- Perez-García GS, Meneses A. Effects of the potential 5-HT7 receptor agonist AS 19 in an autoshaping learning task. *Behav Brain Res*. 2005;163:136–40.
- Pero RW, Roush GC, Markowitz MM, Miller DG. Oxidative stress, DNA repair, and cancer susceptibility. *Cancer Detect Prev*. 1990;14:555–61.
- Pi R, et al. Minocycline prevents glutamate-induced apoptosis of cerebellar granule neurons by differential regulation of p38 and Akt pathways. *J Neurochem*. 2004;91:1219–30.
- Pouzet B. SB-258741: a 5-HT7 receptor antagonist of potential clinical interest. *CNS Drug Rev*. 2002;8:90–100.
- Prasad S, Ponimaskin E, Zeug A. Serotonin receptor oligomerization regulates cAMP-based signaling. *J Cell Sci*. 2019.
- Pytka K, et al. Single administration of HKB-15-a triple 5-HT(1A), 5-HT(7), and 5-HT(3) receptor antagonist-reverses depressive-like behaviors in mouse model of depression induced by corticosterone. *Mol Neurobiol*. 2018;55:3931–45.
- Querfurth HW, LaFerla FM. Alzheimer's disease. *N Engl J Med*. 2010;362:329–44.
- Querfurth HW, LaFerla FM. Alzheimer's disease. *N Engl J Med*. 2018;362:329–44.
- Quintero-Villegas A, Valdés-Ferrer SI. Role of 5-HT7 receptors in the immune system in health and disease. *Mol Med*. 2019;26:4–11.
- Quintero-Villegas A, Manzo-Alvarez HS, Valenzuela Almada MO, Bernal-Mondragón C, Guevara-Guzmán R. Procognitive and neuroprotective effect of 5-HT7 agonists in an animal model by intracerebroventricular amyloid-β injection. *SciFed J Alzheimer's Dementia*. 2019;2019(1):1–10.
- Renner U, et al. Heterodimerization of serotonin receptors 5-HT1A and 5-HT7 differentially regulates receptor signalling and trafficking. *J Cell Sci*. 2005;2:2486–99.
- Roberts AJ, et al. Mice lacking 5-HT7 receptors show specific impairments in contextual learning. *Eur J Neurosci*. 2004;19:1913–22.
- Rojas PS, Neira D, Muñoz M, Lavandero S, Fiedler JL. Serotonin (5-HT) regulates neurite outgrowth through 5-HT1A and 5-HT7 receptors in cultured hippocampal neurons. *J Neurosci Res*. 2014;92:1000–9.
- Ruat M, Traiffort E, Leurs ROB, Tardivel-Iacombe J, Diaz J. Molecular cloning, characterization, and localization of a high-affinity serotonin receptor (5-HT7) activating cAMP formation. 1993;90:8547–8551.
- Salim S. Oxidative stress and the central nervous system. *J Pharmacol Exp Ther*. 2017;360:201–5.
- Samarajeewa A, et al. 5-HT7 receptor activation promotes an increase in TrkB receptor expression and phosphorylation. *Front Behav Neurosci*. 2014;8:1–9.
- Sanger GJ. 5-Hydroxytryptamine and the gastrointestinal tract: Where next? *Trends Pharmacol Sci*. 2008;29:465–71.
- Schulte-Herbrüggen O, Jockers-Scherübl MC, Hellweg R. Neurotrophins: from pathophysiology to treatment in Alzheimer's disease. *Curr Alzheimer Res*. 2008;5:38–44.
- Shimizu M, Nishida A, Zensho H, Miyata M, Yamawaki S. Agonist-induced desensitization of adenylyl cyclase activity mediated by 5-hydroxytryptamine7 receptors in rat frontocortical astrocytes. *Brain Res*. 1998;784:57–62.
- Skaper SD, Facci L, Zusso M, Giusti P. Synaptic plasticity, dementia and Alzheimer disease. *CNS Neurol Disord Drug Targets*. 2017;16:220–33.
- Smith C, Toohey N, Knight JA, Klein MT, Teitler M. Risperidone-induced inactivation and clozapine-induced reactivation of rat cortical astrocyte 5-hydroxytryptamine 7 receptors: evidence for in situ G protein-coupled receptor homodimer protomer cross-talk. 2011; 79:318–325.
- Soga F, Katoh N, Inoue T, Kishimoto S. Serotonin activates human monocytes and prevents apoptosis. *J Investig Dermatol*. 2007;127:1947–55.
- Speranza L, et al. The serotonin receptor 7 promotes neurite outgrowth via ERK and Cdk5 signaling pathways. *Neuropharmacology*. 2013;67:155–67.
- Speranza L, et al. Activation of 5-HT7 receptor stimulates neurite elongation through mTOR, Cdc42 and actin filaments dynamics. *Front Behav Neurosci*. 2015;9:1–14.
- Speranza L, et al. Serotonin 5-HT7 receptor increases the density of dendritic spines and facilitates synaptogenesis in forebrain neurons. *J Neurochem*. 2017;141:647–61.
- Stevens CF, Wang Y. Reversal of long-term potentiation by inhibitors of haem oxygenase. *Nature*. 1993;364:147–9.
- Stiedl O, Pappa E, Konradsson-Geuken Å, Ögren SO. The role of the serotonin receptor subtypes 5-HT1A and 5-HT7 and its interaction in emotional learning and memory. *Front Pharmacol*. 2015;6:162.
- Stroth N, Svenningsson P. S100B interacts with the serotonin 5-HT7 receptor to regulate a depressive-like behavior. *Eur Neuropsychopharmacol J Eur Coll Neuropsychopharmacol*. 2015;25:2372–80.

- Tawfik MK, Makary S. 5-HT₇ receptor antagonism (SB-269970) attenuates bleomycin-induced pulmonary fibrosis in rats via downregulating oxidative burden and inflammatory cascades and ameliorating collagen deposition: Comparison to terguride. *Eur J Pharmacol.* 2017;814:114–23.
- Thomas D, Hagan J. 5-HT₇ receptors. *Curr Drug Target-CNS Neurol Disord.* 2004;3:81–90.
- Vasefi MS, Kruk JS, Heikkila JJ, Beazely MA. 5-Hydroxytryptamine type 7 receptor neuroprotection against NMDA-induced excitotoxicity is PDGFR β receptor dependent. *J Neurochem.* 2013a;125:26–36.
- Vasefi MS, et al. Acute 5-HT₇ receptor activation increases NMDA-evoked currents and differentially alters NMDA receptor subunit phosphorylation and trafficking in hippocampal neurons. *Mol Brain.* 2013b;6:24.
- Volpicelli F, Speranza L, di Porzio U, Crispino M, Perrone-Capano C. The serotonin receptor 7 and the structural plasticity of brain circuits. *Front Behav Neurosci.* 2014;8:318.
- Wesołowska A, Nikiforuk A, Stachowicz K. Potential anxiolytic and antidepressant effects of the selective 5-HT₇ receptor antagonist SB 269970 after intrahippocampal administration to rats. *Eur J Pharmacol.* 2006;553:185–90.
- Woo YS, Wang HR, Bahk W-M. Lurasidone as a potential therapy for bipolar disorder. *Neuropsychiatr Dis Treat.* 2013;9:1521–9.
- Yang Z, Liu X, Yin Y, Sun S, Deng X. Involvement of 5-HT₇ receptors in the pathogenesis of temporal lobe epilepsy. *Eur J Pharmacol.* 2012;685:52–8.
- Yuksel TN, et al. Protective effect of 5-HT₇ receptor activation against glutamate-induced neurotoxicity in human neuroblastoma SH-SY5Y cells via antioxidative and antiapoptotic pathways. *Neurotoxicol Teratol.* 2019;72:22–8.
- Zareifopoulos N, Papatheodoropoulos C. Effects of 5-HT-7 receptor ligands on memory and cognition. *Neurobiol Learn Mem.* 2016;136:204–9.
- Zhou X, Zhang R, Zhang S, Wu J, Sun X. Activation of 5-HT_{1A} receptors promotes retinal ganglion cell function by inhibiting the cAMP-PKA pathway to modulate presynaptic GABA release in Chronic Glaucoma. *J Neurosci.* 2019;39:1484–504.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

