

CORRECTION

Open Access



Correction to: MeioCapture: an efficient method for staging and isolation of meiocytes in the prophase I sub-stages of meiosis in wheat

Arun S. K. Shunmugam¹, Venkatesh Bollina¹, Stefanie Dukowic-Schulze², Pankaj K. Bhowmik¹, Chris Ambrose³, James D. Higgins⁴, Curtis Pozniak⁵, Andrew G. Sharpe^{1,6}, Kevin Rozwadowski⁷ and Sateesh Kagale^{1*}

Correction to: BMC Plant Biol (2018) 18:293
<https://doi.org/10.1186/s12870-018-1514-z>

Following publication of the original article [1], a reader spotted an incorrect citation of the reference 14 [2] in the 'Background'. The male meiocyte isolation work described in this article [2] was carried out in rice and not in Brassica as originally stated in the 'Background' [1]. Thus, the following amendment to the Background section should be noted:

Original sentence:

"Several methods have been previously described or proposed for the isolation of male meiocytes [6], such as micromanipulation (Plumbago [10, 11], Nicotiana [12], Brassica [13, 14], Arabidopsis [15, 16, 17] and sunflower [18]), capillary collection of meiocytes (CCM) (Arabidopsis [19, 20] and maize [21, 22]), laser capture microdissection (LCM) in rice [23, 24, 25, 26], Percoll gradient separation (Arabidopsis [27, 28], rice [29] and Brassica [30]) and isolation of nuclei tagged in specific cell types (INTACT) in Arabidopsis [31]."

Corrected sentence:

"Several methods have been previously described or proposed for the isolation of male meiocytes [6], such as micromanipulation (Plumbago [10, 11], Nicotiana [12], Brassica [13], rice [14], Arabidopsis [15, 16, 17] and sunflower [18]), capillary collection of meiocytes (CCM) (Arabidopsis [19, 20] and maize [21, 22]), laser capture microdissection (LCM) in rice [23, 24, 25, 26], Percoll gradient separation (Arabidopsis [27, 28], rice [29] and Brassica [30]) and isolation of nuclei tagged in specific cell types (INTACT) in Arabidopsis [31]."

Author details

¹National Research Council Canada, Saskatoon, SK, Canada. ²Department of Horticultural Science, University of Minnesota, St. Paul, MN, USA. ³Department of Biology, University of Saskatchewan, Saskatoon, SK, Canada. ⁴Department of Genetics and Genome Biology, University of Leicester, Leicester, UK. ⁵Department of Plant Sciences, University of Saskatchewan, Saskatoon, Canada. ⁶Global Institute for Food Security, University of Saskatchewan, Saskatoon, Canada. ⁷Agriculture and Agri-Food Canada, Saskatoon, SK, Canada.

Published online: 02 May 2019

References

1. Shunmugam ASK, Bollina V, Dukowic-Schulze S, et al. MeioCapture: an efficient method for staging and isolation of meiocytes in the prophase I sub-stages of meiosis in wheat. *BMC Plant Biol.* 2018;18(1):293.
2. Collado-Romero M, Alós E, Prieto P. Unravelling the proteomic profile of rice meiocytes during early meiosis. *Front Plant Sci.* 2014;5.

* Correspondence: Sateesh.Kagale@nrc-cnrc.gc.ca

¹National Research Council Canada, Saskatoon, SK, Canada

Full list of author information is available at the end of the article

