

CASE REPORT

Open Access



Bilateral vocal cord paralysis after endoscopic placement of fully covered self-expandable metal stent for palliative treatment of malignant proximal esophageal obstruction: two case reports

Y. Chiche¹, G. Beltramo^{1,2,3}, T. Degand⁴, A. Drouillard⁴, C. Foignot¹, N. Baudouin¹, P. Bonniaud^{1,2,3} and M. Georges^{1,3,5*} 

Abstract

Background: Oesophageal stents have several well-known respiratory complications, including aspiration pneumonia, fistula and airway compression. However, bilateral vocal cord paralysis has rarely been described.

Methods: We describe two patients who presented with refractory dysphagia due to malignant proximal oesophageal strictures. Both received palliative treatment consisting of fully covered self-expandable metal stents that were placed across the strictures.

Results: Both patients developed inspiratory stridor and acute hypoxemic respiratory failure shortly after the stent was placed. Flexible bronchoscopy revealed vocal cord paralysis in paramedian position, potentially due to extrinsic compression of the posterior branch of the recurrent laryngeal nerve following the progressive opening of the esophageal prosthesis. One patient recovered after the stent was removed.

Conclusions: Bilateral vocal cord paralysis is a rare but potentially fatal complication of proximal esophagus stenting.

Keywords: Vocal cord paralysis, Esophageal proximal stent, Complication

Background

The respiratory complications of esophageal stents, including aspiration pneumonia, fistula and airway compression, are well described in literature [1]. However, bilateral vocal cord paralysis has rarely been described.

Case presentation

First case report

An 88-year-old patient with a history of esophageal squamous cell cancer presented with refractory dysphagia. The malignant esophageal stricture identified 18 cm from the incisors was dilated, and a fully covered Niti-S esophageal prosthesis of 80x18mm (Taewoong, Busan, South Korea) was subsequently placed across the stricture.

The day after the procedure, the patient developed inspiratory stridor and acute hypoxemic respiratory failure. X-rays of the neck showed that the stent

* Correspondence: marjolaine.georges@chu-dijon.fr

¹Department of Respiratory Diseases and Intensive Care, Reference Center for Adult Rare Pulmonary Diseases, University Hospital of Dijon – Bourgogne, Dijon, France

³University of Burgundy Franche-Comté, Dijon, France

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

was correctly placed. Orotracheal intubation was required on day 4. A flexible bronchoscopy showed that the vocal cords were paralyzed in paramedian position, with no evidence of locoregional edema or laryngeal injury, particularly cricoarytenoid edema or external tracheal compression. The stent was removed on day 6, and the patient was extubated on day 7. A second flexible bronchoscopy showed complete restoration of right vocal cord mobility and partial improvement on the left side. The respiratory outcome was favorable and oxygen therapy was stopped at day 10.

Second case report

A 58-year-old patient with a history of right upper lobectomy for stage IA primary lung adenocarcinoma presented with dyspnea and aphagia. A CT scan revealed a proximal esophageal mass extending to the lower trachea with no evidence of lung cancer recurrence. A flexible bronchoscopy showed tracheal stenosis due to an apparently malignant endotracheal mass combined with external airway compression. The vocal cords were normal. A silicone tracheal prosthesis of 15 × 80 mm (Trachéobronxane, Novatech, La Ciotat, France) was placed across the stricture during rigid bronchoscopy. At the same time, gastroscopy revealed severe extrinsic esophageal compression 23 cm after the incisors, and so an esophageal prosthesis was placed (Taewoong, Busan, South Korea). Tachypnea and dyspnea appeared immediately after extubation. Flexible bronchoscopy showed that the vocal cords were in paramedian position without tracheal prosthesis

migration or laryngeal edema. The patient died 6 days following the placement of the prosthesis due to acute respiratory distress.

Discussion

The two mechanisms illustrated in Fig. 1 (adapted from Ceruse [2]) could explain the occurrence of bilateral vocal cord paralysis after the placement of proximal esophageal stents [3].

Firstly, direct compression from the esophageal prosthesis or local inflammation could induce neuropraxia of the posterior branch of the recurrent laryngeal nerve, which innervates the posterior cricoarytenoid muscles and transverse arytenoid muscles.

In addition, post-operative local inflammation may cause spasms in the interarytenoid muscles.

We presume that the progressive appearance of respiratory symptoms in our cases was due to the gradual opening of the esophageal prosthesis which increasingly compressed the posterior larynx in the days following the procedure. In previously published cases, the time to respiratory failure can vary from several hours to several days [4, 5].

Endoscopists should consider bilateral vocal cord paralysis as a rare but potentially fatal complication of proximal esophagus stenting. The diagnosis of bilateral vocal cord paralysis is a therapeutic emergency considering that patients can potentially recover once the stent is removed. Tracheostomy should also be considered as a treatment option.




Fig. 1 Descriptive anatomy of the larynx

Acknowledgments

The authors thank Suzan Ranking for reviewing the manuscript.

Authors' contributions

MG and NB had the original idea. GB, TD, AD, CF and NB were in charge of the patients reported in this manuscript. YC collected data. YC and MG wrote the manuscript. PB provided critical feedback and helped shape the manuscript and design the figure. All authors discussed the cases and contributed to the final version of the submitted manuscript. All authors have read and approved the manuscript in its final state.

Funding

No funding.

Availability of data and materials

All data analysed during this study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

Not applicable.

Consent for publication

At the moment we wrote this manuscript, the two patients reported had died. Relatives were informed for our work and gave their written consent for publication of case reports.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Respiratory Diseases and Intensive Care, Reference Center for Adult Rare Pulmonary Diseases, University Hospital of Dijon – Bourgogne, Dijon, France. ²INSERM U1231, University of Burgundy Franche-Comté, Dijon, France. ³University of Burgundy Franche-Comté, Dijon, France. ⁴Department of Hepato-Gastro-Enterology, University Hospital Dijon – Bourgogne, Dijon, France. ⁵Centre des Sciences du Goût et de l'Alimentation, UMR 6265 CNRS 1234 INRA, University of Burgundy Franche-Comté, Dijon, France.

Received: 1 March 2020 Accepted: 7 May 2020

Published online: 14 May 2020

References

1. Łochowski MP, Brzeziński D, Rębowski M, et al. Complications after treating esophageal strictures with prostheses and stents - 20 years' experience. *Wideochir Inne Tech Maloinwazyjne*. 2016;11:4.
2. Cérouse P, Ltaief-Boutrigou A, Buiret G, et al. Anatomie descriptive, endoscopique et radiologique du larynx. *EMC Oto rhino laryngologie*. 2012;7:2.
3. Friedman M, Toriumi DM. Esophageal stethoscope. Another possible cause of vocal cord paralysis. *Arch Otolaryngol Head Neck Surg*. 1989;115:1.
4. Moreels TG, De Schepper HU, Macken EJ, et al. Vocal cord paralysis due to self-expandable metal stent in the proximal esophagus. *Endoscopy*. 2014; 46(Suppl 1 UCTN):E155–6.
5. Gellad ZF, Hampton D, Tebbit CL, et al. Bilateral vocal cord paralysis following stent placement for proximal esophageal stricture. *Endoscopy*. 2008;40(Suppl 2):E150.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

