

RESEARCH

Open Access



# Prevalence of anemia and its associated factors among under-five age children in Shanan gibe hospital, Southwest Ethiopia

Destaw Kebede<sup>1\*</sup>, Fantahun Getaneh<sup>1</sup>, Kirubel Endalamaw<sup>1,2</sup>, Tariku Belay<sup>2</sup> and Abebe Fenta<sup>3</sup>

## Abstract

**Background:** Anemia is a major health problem in the worldwide. Because of health and socioeconomic problems, the prevalence of anemia is higher in developing countries. However, there was a limited finding in our study area. Therefore, the aim of this study was to determine the prevalence of anemia and its associated factors among under-five age children in Shanan Gibe Hospital (SGH), Southwest Ethiopia.

**Methods:** Institution based cross sectional study was conducted at SGH, Ethiopia using consecutive convenient sampling technique during 1 January to 30 April, 2021. Data was collected by interviewing and capillary blood was taken from the fingertip for hemoglobin determination by using HaemoCue digital photometer. Additionally, stool sample was processed using wet mount and formal-ether concentration technique. Then after, the data were entered to Epidata version 3.1 and analysed with Statistical Package for the Social Sciences (SPSS) version 20. Factors associated with anaemia were assessed by bivariable and multivariable logistic regression model by considering  $P < 0.05$  as statistical significance.

**Results:** A total of 368 under five children were recruited to the study and the current prevalence of anemia was 48.9%. Of this anemia, 25.0% mild, 15.8% moderate and 8.2% were severely anemic. More ever, being rural resident (AOR = 6.11; 95% CI = 1.49–8.99,  $P = 0.002$ ), family low income (AOR = 6.27, 95% CI = 1.35–11.43,  $P = 0.004$ ), family size greater than five (AOR = 3.12; 95% CI = 1.47–7.11,  $P = 0.002$ ) and intestinal parasite infections such as *Enteameoba histolytica* (AOR = 3.37; 95%CI = 2.16–11.31,  $P = 0.005$ ), Hookworm (AOR = 6.09; 95%CI = 2.37–11.56,  $P = 0.001$ ), and *Trichuris trichuria* (AOR = 2.79; 95%CI = 1.45–9.13,  $P = 0.002$ ) ( $P < 0.05$ ) were factors significantly associated with anemia among under five children.

**Conclusion:** The current prevalence of anemia among under five age children is relatively high. On the other hand, the rural residence, large family size, low family income, infection with *Enteameoba histolytica*, hookworm and *Trichuris trichuria* were the identified factors associated with anemia among under five children. Therefore, there should be massive and routine deworming program in addition to imperative targeting anemia prevention, and nutritional supplementation to reduce the burden of anemia.

**Keywords:** Anemia, Prevalence, Under five children, Associate factors, Southwest Ethiopia

## Background

Anemia is a low number of red blood cells or a low hemoglobin or hematocrit in which the hemoglobin content of the blood is lower than normal range as a result of deficiency of one or more essential nutrients [1] or

\*Correspondence: amaueldestaw@gmail.com

<sup>1</sup> Department of Diagnostic Laboratory at Shegaw Motta General Hospital, East Gojjam, P.O. Box 50, Motta Town, Ethiopia  
Full list of author information is available at the end of the article



due to heavy blood loss, parasitic infections and congenital hemolytic diseases. Types of anemia includes iron deficiency anemia, sickle cell anemia, Vitamine deficiency anemia, Aplastic anemia, hemolytic anemia and anemia of inflammation [2] According to World Health Organization a child less than five years old is anemic if the blood hemoglobin is less than 110g per liter (Hgb < 11.0 g/dl) [3].

Globally, approximately 1.62 billion people are affected by anemia [4] and approximately 36.4 to 61.9% of under five children in sub-Saharan Africa are also affected [5]. Similarly, it is considered to be a serious public health problem in Sub Saharan Africa in which, approximately 83.5 million children was affected and its prevalence of anemia was 67% [6]. The risk factors for anemia vary in different settings; they include intestinal worms, malaria, HIV infection, nutritional deficiencies, hematological malignancies and chronic diseases like sickle cell disease [6, 7].

Globally, anemia is a public health problem affecting people in both developed and developing countries with bad consequences of human health as well as social and economic development. It is also associated with increased morbidity and mortality [8]. Anemia affects all age group especially, under-five age children from low income families have a higher risk for developing anemia due to iron deficiency that occurs as a result of high demand for iron during the period of rapid growth [9].

In developing countries having low-and middle-income, the prevalence of anemia among 6–59 month age children was >20% based on latest demographic and health survey (DHS) report rounds between 2005–2018, and it is classified as severe public health problem [10]. The problem is alarming in Sub-Saharan African Countries such as Kenya 48.9% [11]; Mali 55.8% [12] and Tanzania 79.6% [13]. Lack of awareness among the mothers about the problem coupled with their low educational status [14], poor nutritional practices and unhealthy food habits [15], low iron bioavailability of the diet [16], decreased physical activities [17], malaria and parasitic infestations are additional factors associated with lower hemoglobin (Hgb) level in children [18]. Factors including family size, low socio-economic status, illiteracy and ignorance are associated with anemia among under five children. Infection with Hook worm and intestinal helminthes causes gastro-intestinal blood loss resulting in depletion of iron stores and consequently also impaired erythropoietin [7]. This leads to mal-absorption and inhibition of appetite, there by worsening micronutrient deficiency and children anemia [19]. The consequence of anemia in under five age children include: decrease mental performance, low tolerance to infection, death from anemic heart failure [17, 19].

Although anemia remains a widespread public health problem in developed countries, it contributes significant proportion of children death in most developing countries including Ethiopia. As such, various factors like parasite have impact on cognitive development and physical growth, studies on the magnitude of anemia among under-five age children have paramount importance. Meanwhile, there is a limited study on the prevalence and associated factor of anemia among under-five age children in our study area. Therefore, this study aimed to determine the prevalence of anemia and its associated factors among under-five children in Shanan Gibe Hospital, Southwest Ethiopia.

## Methods

### Study area, design, and period

Institution based cross sectional study was conducted in Shanan Gibe Hospital, Jimma town, Ethiopia from 1 January to 30 April, 2021.

### Population

All under-five age children who attended in SGH were the source population. However, under five age children who visited SGH during the study period were study population and the under five children who provided stool sample and blood were study participant in the current study.

### Sample size and sampling technique

The sample size was calculated by using single population proportion formula by taking 41.1% prevalence of anemia among under five children in in Gugufu, South Wollo, Northeast Ethiopia [20] using the assumption of 95% confidence level ( $z = Z_{\alpha/2} = 95\% = 1.96$ ), margin of error ( $d = 5\% = 0.05$ ), Then, sample size was determined as follow:

$$N = \frac{(Z_{\alpha/2})^2 p (1 - p)}{d^2} = 368$$

Therefore, the minimum of 368 study participants were selected by consecutive convenient sampling technique.

### Inclusion and exclusion criteria

All under five children attending to SGH and those provided stool sample were included to study. However, those under five children who did take drugs for deworming before one month of data collection were excluded to this study.

### Data collection and processing

English version questionnaire was translated in to Amharic version and finally it was translated back to

English version to check its consistency. The data were collected by a trained nurse and a principal investigator. Thus, Socio demographic and possible associated risk factors of anemia were collected by structured pre-tested Amharic version questionnaire using face to face interview with parents/ guardians based on an interviewer-administered semi structured questionnaire. At each data collection spot, sufficient explanation about the aim of the research was given to the parents or study participants before conducting the interview.

#### Sample collection and transportation

Whole blood and stool samples were collected aseptically. The whole blood was used for hemoglobin measurement immediately at site data collection in hospital, while the stool samples were transported from clinical services outlets to department of diagnostic laboratory in Shana Gibe Hospital (SGH).

#### Hemoglobin measurement

The Hgb concentration of each participant was measured by taking a finger-prick blood sample using a portable hemoglobin meter instrument (calibrated HaemoCue digital photometer). Hemoglobin value was displayed within 15–20s and interpreted as; Hgb < 11 g/dl were anemic and Hgb > 11 g/dl was defined as non-anemic for under five age children. The severity of anemia was categorized according to WHO cut-off value scheme as Hgb between 10.0 g/dl– 10.9 g/dl for mild anemia, Hgb between 7.0–9.9 g/dl for moderate anemia and Hgb less than 7 g/dl for severe anemia on under-five age children [21].

#### Parasitic examination

After the interviewer administered questionnaire, the participants were sent to laboratory and asked them to bring about 10-g (teaspoonful amount) stool specimen was collected with wide mouth screw capped containers labeled with their identification code number for each study participant following the standard operating procedures (SOPs). The stool samples were examined for intestinal parasites using wet mount preparation on 10x objective microscopy within 10–15 min after collection. The remaining samples were stored in a cool box (regulated the temperature to < 20°C) and on the next day, formal-ether concentration technique was employed to diagnosis few parasitic density [22]. Finally, report and were record each parasite in the prepared laboratory format.

#### Data analysis

Data was entered by EpiData version 3.1 and data analysis was performed using SPSS version 20. The prevalence

of anemia was determined by descriptive statistics. Multivariable logistic regression was done by entering the variables with  $p < 0.2$  in bivariable logistic regression to identify the factors associated with anemia. A  $P$  value < 0.05 considered as statistically significant association in the multivariable logistic regression.

#### Quality control

The data were collected by a trained nurse and principal investigators. Pre-test was done on 18 under five children at Jimma Medical Center before data collection and the actual study was performed to check acceptability of the questionnaire whether it contains the necessary information or not and if unnecessary, to make possible corrections..

Completeness of questionnaires was checked soon after collecting the data and calibrated HaemoCue digital photometer was used for Hgb concentration determination. The reagents were checked for their expired date before any test was performed. The patient's blood and stool sample was collected, prepared and tested according to SOP to get reliable result. In each step, pre analytical, analytical and post analytical phases were maintained for quality assurance. At the end, the results were checked and registered on the laboratory record format before delivery to the patients.

#### Ethical consideration

The study approved by College of Medicine and Health Science, Jimma University's Research Institutional Review Board with reference number RU 1986/07 and a permission letter was obtained from Shanan Gibe Hospital. The purpose and importance of the study was explained to the participants. Written informed consent from parent/guardian was obtained in accordance with the Declaration of Helsinki. Additionally, absence of link between the study and their service was explained and participation was entirely voluntary based. Furthermore, the confidentiality of study participant was kept and identification of study participant by name was avoided. Finally, all participants who were diagnosed positive for intestinal parasites and anemia were linked to Medical doctors in Hospital for further management.

#### Operational definitions

**Under five children:** the children whose ages are less than five years.

**Hemoglobin** – A red substance in the blood that carries oxygen and contains iron.

**Non anemic** - Hemoglobin concentrations greater than 11.0 g/dl in the blood of the individual.

**Anemic** - Children with Hgb level < 11.0 g/dL.

**Mild anemia** – Hemoglobin concentrations in between 10–10.9 g/dl in the blood of the individual.

**Moderate anemia** – Hemoglobin concentrations in between 7.0 g /dl-9.9 g/dl in the blood of the individuals.

**Severe anemia**– is hemoglobin concentrations which is less than 7.0 g /dl in the individual’s blood.

**Monthly income:** it was categorized based Ethiopian civil service proclamation as individual earning with low income (<3000 ETB), medium income (3000–7500 ETB) and high income (>7500 ETB).

**Results**

**Socio-demographic characteristics of study participants**

A total of 368 under-five age children were participated in the study. Of these, about 195 (52.9.0%) were females and 147 (39.9%) were within the age group of 2–3 years. According to educational status of their mothers, the highest (47.0%) of the children’s mothers were unable to read and write. However, the family monthly income with 247(67.1%) had low income and 1–5 family size accounted 228 (62.0%). The other socio demographic characteristics were summarized below (Table 1).

**Prevalence of anemia**

The prevalence of anemia in this study was 180 (48.9%) with [95% CI: 39.24–53.19]. The degree of Anemia was classified as mild, moderate, & severe anemia according to WHO cutoff values for grading anemia (43). Thus, from the total of 180 (48.9%) anemic children, 92 (24.9%)

and 58 (15.8%) and 30 (8.2%) were mild, moderate and severe anemia, respectively (Fig. 1).

**Proportion of intestine parasites**

The proportions of intestinal parasite infection was 115 (31.25%) with (95%CI=30.28–72.42). From these, the proportion of each intestinal parasites were recorded as *Ascaris lumbricoids* (*A. lumbricoide*) 42 (36.5%), *Gardia lamblia* (*G. lamblia*) 37 (32.2%), *Enteameba histolytica* (*E. histolytica*) 13(11.3%), Hookworm 3 (2.60%) and *Trichuris trichuria* (*T. trichuria*) 20 (17.40%) as shown below (Fig. 2).



**Factor associated with anemia**

All independent variables showing *P*-value <0.2 in the bivariable analysis including age, residence, mother monthly income, family size, *A. lumbricoide*, *G. lamblia*, *E. histolytica*, Hookworm and *T.trichuria* were entered in to multivariable logistic regression analysis to determine factors associated with anemia among under-five age children. Accordingly, only being rural residence (AOR=6.11; 95% CI=1.49–8.99, *P*=0.002), family low income (AOR=6.27, 95% CI=1.35–11.43, *P*=0.04) and family size greater than five (AOR=3.12; 95% CI =1.47–7.11, *P*=0.002) *P*<0.05 were demonstrated as significant association with anaemia among under five children. Thus, the adjusted odds of children with rural residence having anemia was 6.11 times higher compared to those from urban residents.. Similarly, the adjusted odds of children with low income mothers having anemia

**Table 1** Socio demographic distribution of under-five age children visited in Shanan Gibe Hospital, Southwestern Ethiopia, 1 January to 30 April, 2021

Characteristics	Category	Frequency	Percent (%)
Sex	Male	173	47.0
	Female	195	53.0
Age	0–1	87	23.6
	2–3	147	40.0
	4–5	134	36.4
Educational status (Mothers)	Unable to read and write	173	47.0
	Primary (1–8)	9	26.4
	Secondary (9–12)	28	7.6
	>Grade 12	70	19.0
Residence (mother)	Urban	212	57.6.0
	Rural	156	42.4
Monthly income (mother)	Low income (<3000 ETB)	247	67.1
	Medium income (3000–7500 ETB)	76	20.7
	High income (>7500 ETB)	45	12.2
Family size	1–5	228	62.0
	>5	140	38.0

Note: ETB Ethiopian Birr



was 6.27 times higher compared to those children from mothers having high monthly income. Additionally,, the adjusted odds of children with in from greater than five family size in home having anemia was 3.12 times higher compared to those children form family size less than five in home (Table 2).

On the other hand, some intestinal parasites were significantly associated with anemia. Hence, *E. histolytica* (AOR = 3.37; 95%CI = 2.16–11.31, *P* = 0.005), Hookworm (AOR = 6.09; 95%CI = 2.37–11.56, *P* = 0.001), and *T. trichuria* (AOR = 2.79; 95%CI = 1.45–9.13, *P* = 0.002) were the indentified intestinal parasites associated with anemia among under five children by considering *P* value less than 0.05 as statistical significantly association (Table 2).

**Discussion**

Anemia has been continued to be a health problem on under five children [23] which adversely affects mental, physical and social development of the children [20]. This burden also affected middle and low income countries including Ethiopia in particular our study area. Hence, the prevalence of anemia was 48.9% at 95% CI (39.24–53.19) in the current study. As the result, other previous studies were compared with our study by categorizing in to low, comparable and high prevalence on the bases of CI of this study.

Accordingly, the prevalence of anemia in the current study was higher than previous studies reported in Huaihua (29.73%) [24], Senegal (30.7%) [25], Uganda



**Table 2** Factors associated with anemia among under five age children in Shanan Gibe Hospital, Southwestern Ethiopia, 1 January to 30 April, 2021

Variables	Category	Anemic (n = 180) N (%)	Non-anemic (n = 188) N (%)	COR(95%CI) P value	AOR(95%CI)	P value
Sex	Male	82 (47.4)	91 (52.6)	1	–	–
	Female	98 (50.3)	97 (49.7)	3.21 (0.67–8.12)0.88		
Age	0–1	37 (42.5)	50 (57.5)	7.12 (1.93–11.32)0.001*	5.31 (0.9–10.13)	0.17
	2–3	77 (52.4)	70 (47.6)	3.27 (2.5–8.45)0.08*	4.62 (0.85–7.44)	0.44
	4–5	66 (49.3)	68 (50.7)	1	1	
Educational statuses (mother)	Unable read & write	93 (53.8)	80 (46.2)	4.12 (0.69–12.71)0.91	–	–
	primary (1–8)	41 (42.3)	56 (57.7)	5.15 (0.72–6.88)0.53		
	Secondary (9–12)	10 (35.7)	18 (64.3)	2.32 (0.93–5.78)0.31		
	>Grade 12	36 (51.4)	34 (48.6)	1		
Residence	Urban	114 (53.8)	98 (46.2)	3.57 (1.09–7.55)0.001*	6.11 (1.49–8.99)	0.002**
	Rural	66 (42.3)	90 (57.7)	1	1	
Monthly income (mother)	Low income	128 (51.8)	119 (48.2)	7.34 (1.77–11.9)0.06*	6.27 (1.35–11.43)	0.004**
	Medium income	28 (36.8)	48 (63.2)	4.51 (1.21–8.02)0.014*	2.17 (1.83–8.05)	0.41
	High income	24 (53.3)	21 (46.7)	1	1	
Family size	1–5	116 (50.9)	112 (49.1)	1	1	
	>5	64 (45.7)	76 (54.3)	4.55 (1.33–6.71)0.001*	3.12 (1.47–7.11)	0.002**
Utilize latrine	Yes	60 (51.3)	57 (48.7)	0.78 (0.42–5.38)0.58	–	–
	No	120 (47.8)	131 (52.2)	1		
Hand wash before meal	Yes	69 (60.0)	46 (40.0)	2.93 (0.39–7.09)0.73	–	–
	No	111 (43.9)	142 (56.1)	1		
Finger trimming	Yes	72 (54.1)	61 (45.9)	3.23 (0.99–8.02)0.44	–	–
	No	108 (46.0)	127 (54.0)	1		
<i>A. lumbricoide</i>	Yes	42 (91.3)	4 (8.7)	4.51 (2.13–7.54)0.006*	3.76 (0.82–5.98)	0.21
	No	138 (42.9)	184 (57.1)	1	1	
<i>G. lamblia</i>	Yes	37 (92.5)	3 (7.5)	1.35 (1.21–7.25)0.01*	1.45 (0.9–8.12)	0.57
	No	143 (43.6)	185 (56.4)	1	1	
<i>E. histolytica</i>	Yes	13 (86.7)	2 (13.3)	2.57 (2.1–9.13)0.001*	3.37 (2.16–11.31)	0.005**
	No	167 (47.3)	186 (52.7)	1	1	
Hookworm	Yes	3 (75.0)	1 (25.0)	4.75 (3.21–10.92)0.001*	6.09 (2.37–11.56)	0.001**
	No	177 (48.6)	187 (51.4)	1	1	
<i>T. trichuria</i>	Yes	20 (74.1)	7 (25.9)	3.53 (2.67–8.23)0.001*	2.79 (1.45–9.13)	0.002**
	No	160 (46.9)	181 (53.1)	1	1	

Key: AOR Adjusted Odd ratio, COR Crude Odd Ratio, N Frequency, % Percent, CI Confidence interval, 1 reference, \* = variables selected for multivariable logistic regression, \*\* = significantly associated variable

26.2% [15], Tanzania at Rombo district 37.9% [26]. The variation might be due to the variability of risk factors across different geographic regions, plus lower socio-economic, large family size and maternal education status of under-five age children those factor which contribute increase the prevalence of anemia.

The current anemia prevalence among under five children was analogous with studies indicated in Palestine 33.5% [27], South Kivu 39.6% [18], Ghana (41%) [28], Cape Verde 51.8% [29] and in Ethiopia (44.83%)

[30], Wollo (41.1%) [20], Filtu Town, Somali region (41.7%) [31], Duggina Fanigo District of Wolaita Zone 51.4% [32] and Amhara region (41.43%) [33]. In contrast, a high prevalence e of anemia had been reported in sub-Saharan Africa (64.1%) [34], East Africa (75%) [35], Brazil (56.6%) [36], Malawi 56.9% [4], Togo (70.9%) [37], Tanzania, Bugando Medical Centre (77.2%) [38], Arusha District in Tanzania (84.6%) [39], Kenyan Coast (76.3%) [40], Mali (58%), Tanzania (57%), and Mozambique (54%) [27], Ghana 78.4% [41]. This variation might be due to the mothers having the problem with

their low monthly income or parasitic infestations of children which are contributing factors associated with lower hemoglobin (Hgb) level in children.

According to factors in the current study, being rural residence (AOR=6.11;  $P=0.002$ ) was associated with anemia in the current study. Beside this, analogs association of rural residence finding was reported in Amhara region [33, 42]. As such, studies indicated that the distribution of anemia is more prevalent among children from rural resident compared to urban ones [43]. This is because of low socioeconomic status, low serving of iron-rich foods, lack of adequate nutrition information or dietary intake and due to a high number of illiterates in rural areas as compared to urban [44].

Family low income (AOR=6.27;  $P=0.004$ ) was the identified associated factor with anemia among under five children in current study which was previously reported in Ethiopia like Wollo [20] and Filtu Town, Somali region [31]. This factor could be associated with anemia due to high family income with higher wealth quintile are more likely to provide balanced macro and micronutrients (minerals and vitamins) to their children and children from a lower economic status are vulnerable to various nutritional disorders including anemia [45]. On the other hand, family size greater than five (AOR=3.12;  $P=0.002$ )  $P<0.05$  was another identified factors associated with anemia among under five children which was founded in sub-Saharan Africa [34]. So, having large family size was reported factor associated with as anemia due to an increase in the number of children might lead to a risk of communicable disease transmission, and competition for food, consequently, nutritional deficiencies [46] Thus, children with rural residence, low income family and family size greater than five in home were 6.11, 6.27 and 3.12 times more likely to develop anaemia among under five children by adjusting other confounding factors, respectively.

On the other hand, intestinal parasites such as *E. histolytica* (AOR =3.37;  $P=0.005$ ), Hookworm (AOR=6.09;  $P=0.001$ ), and *T. trichuria* (AOR=2.79;  $P=0.002$ ) contributed to anemia in children. Similar findings of *E. histolytica* was report in Health Center, North Ethiopia [47], while Hookworm findings also reported in Kenyan coast [39], Ethiopia particularly, Yirgacheffee [48] and Harbu Town [49]. This association might be due to significantly low level of serum iron in *E. histolytica* infected children due to *E. histolytica* needs great levels of iron to stay alive and reproduction [50]. *T. trichiura* and hookworm can cause massive amounts of blood loss and decreased hemoglobin levels [51]. So, anemia is positively associated with intestinal parasite which might be due to lack of hygiene related practices, Red Blood Cell (RBC) destruction and feeding, and loss of appetite caused by worms

[52]. Similarly, the blood loss that can occur in *T. trichiura* infection is likely to contribute to anemia [51, 53].

As the result, children infected with *E. histolytica*, Hookworm and *T. trichuria* were 3.37, 6.09 and 2.79 times more likely to develop anaemia by adjusting other confounding factors, respectively. This study is limited due to the fact that present study didn't differentiate the types of anemia as it was due to iron, vitamin B12, folic deficiency or the RBC morphological effect and hemoparasites were not assessed for factors associated with anemia among under five children.

## Conclusion

The prevalence of anemia among under five age children was high. The study also revealed that the most important associated factors for anemia are rural resident, family size greater than five in home, low family income, infection with *E. histolytica*, Hookworm and *T. trichuria* are the factors associated with anemia in this study. Therefore, there should be a continuous health education for their parents on factors of anemia in addition to massive and routine deworming for intestinal parasites to reduce the burden of anemia among children in hospitals.

## Abbreviations

DHS: Demographic and health survey; ETB: Ethiopian Birr; Hgb: Hemoglobin; IP: Intestinal Parasite; SGH: Shanan Gibe Hospital; SOP: Standard Operational Procedure; SPSS: Statistical Program for Social Science; STH: Soil Transmitted Helminthes; WHO: World Health Organization.

## Acknowledgements

We would like to forward our gratitude to Institute of Health Science, School of Medical Laboratory Science, Jimma University for providing us the ethical clearance. Then we acknowledged Shanan Gibe Hospital administrative for their role in providing permission letter for us to conduct this study. Finally, we would like to extend our deepest gratitude all the children, parents and Shanan Gibe Hospital staff for their role in identification of patients and support during sample collection in this study.

## Authors' contributions

DK had a contribution in conceptualization, investigation and methodology design, and acquisition of data, analysis and interpretation of data as well as took part in writing original drafting of the article and preparing tables and figures. KE had a substantial contribution to the conception, design of the work; the acquisition, analysis and interpretation of data. FG had a contribution on approving the submitted version. TB also participated in writing and revising manuscript critically for important intellectual content. Similarly, AF participated in writing and revising manuscript critically for important intellectual content. However, all authors approved the final manuscript version for publication.

## Authors' information

DK is a Medical Microbiologist in the Department of Diagnostic Laboratory Science at Shegaw Motta General Hospital, East Gojjam Zone, Motta town, Ethiopia. KE is a Medical Laboratory Technologist in the Department of Diagnostic Laboratory Science at Shegaw Motta General Hospital, East Gojjam Zone, Motta town, Ethiopia. FG is Medical Parasitologist in the Department of Diagnostic Laboratory Science at Shegaw Motta General Hospital, East Gojjam Zone, Motta town, Ethiopia. TB is Assistant Professor at Institute of Health, Faculty of Health Science, School of Medical Laboratory Science, Jimma University, Jimma Town, Ethiopia. AF is (Medical Parasitologist, MPH in Epidemiology)

and Lecturer at Department of Medical Laboratory science, Health Science College, Debre Markos University, Debre Markos Town, Ethiopia.

#### Funding

No fund was obtained for execution of the study.

#### Availability of data and materials

Results of this study are generated from the data collected and analyzed based on stated materials and methods. All data concerning to this study could be available upon the request of Destaw Kebede (Correspondence author; mobile +251911594675, email: [amauledestaw@gmail.com](mailto:amauledestaw@gmail.com)).

#### Declarations

##### Ethics approval and consent to participate

The ethical approval was obtained from Institute of Health Science, Jimma University's Institutional Review Board (Reference number RU 1986/07) and a permission letter was also obtained from Shanab Gibe Hospital. Additionally, written informed consents from guardians or parents of each study participants were obtained in accordance with the Declaration of Helsinki. Furthermore, the participation of study participants were entirely volunteer based and there confidentiality was kept by coding rather than naming for identification. Finally, all anemic study participants or/and positive results for intestinal parasites were linked to medical doctors in hospital for further management.

##### Consent for publication

Not applicable.

##### Competing interests

The authors declare that they have no competing interests for this work.

##### Author details

<sup>1</sup>Department of Diagnostic Laboratory at Shegaw Motta General Hospital, East Gojjam, P.O. Box 50, Motta Town, Ethiopia. <sup>2</sup>Institute of Health Science, School of Medical Laboratory Science, Jimma University, Jimma Town, Ethiopia.

<sup>3</sup>Department of Medical Laboratory Science, College of medicine and Health Science, Debre Markos University, Debre Markos Town, Ethiopia.

Received: 4 July 2021 Accepted: 12 November 2021

Published online: 03 December 2021

#### References

- DeMaeyer EA, Adiels-Tegman M. The prevalence of anaemia in the world. *World Health Stat Quart.* 1985;38(3):302–16.
- Vieth JT, Lane DR. Anemia. *Emerg Med Clin.* 2014;32(3):613–28.
- Zuffo CR, Osório MM, Taconeli CA, Schmidt ST, da Silva BH, Almeida CC. Prevalence and risk factors of anemia in children. *J de Pediatria (Versão em Português).* 2016;92(4):353–60.
- Gaston RT, Ramroop S, Habyarimana F. Joint modelling of malaria and anaemia in children less than five years of age in Malawi. *Heliyon.* 2021;7(5):e06899.
- Roberts DJ, Matthews G, Snow RW, Zewotir T, Sartorius B. Investigating the spatial variation and risk factors of childhood anaemia in four sub-Saharan African countries. *BMC Public Health.* 2020;20(1):126.
- Kassebaum NJ, Jasrasaria R, Naghavi M, Wulf SK, Johns N, Lozano R, et al. A systematic analysis of global anemia burden from 1990 to 2010. *Blood.* 2014;123(5):615–24.
- Bremner KC. Pathogenetic factors in experimental bovine oesophagostomiasis: II. Plasma iron, iron-binding capacity, and reticulocyte responses in bleed and infected calves. *Exp Parasitol.* 1969;24(2):184–93.
- Benoist B, McLean E, Cogswell M, Egli I, Wojdyla D. Worldwide prevalence of anemia 1993–2005. In: *World Health Organization global database on anemia.* Geneva: World Health Organization; 2008. p. 7–13.
- Yusakanshori M, Purbasari S. Determinant factors affecting the nutritional status of children in regional health Center of Gresik. *Indian J Publ Health Res Dev.* 2019;10(11):1738–43.
- Hasan MM, Magalhaes RJ, Ahmed S, Pervin S, Tariqujjaman M, Fatima Y, et al. Geographical variation and temporal trend in anemia among children 6–59 months of age in low-and middle-income countries during 2000–2018: forecasting the 2030 SDG target. *Public Health Nutr.* 2021;9:1–20.
- Neumann CG, Bwibo NO, Murphy SP, Sigman M, Whaley S, Allen LH, et al. DemmentAnimal source foods improve dietary quality, micronutrient status, growth and cognitive function in Kenyan school children: background, study design and baseline findings. *MWJ Nutr.* 2003;133(11 Suppl 2):3941S–9S.
- Hall A, Roschnik N, Ouattara F, Touré I, Maiga F, Sacko M, et al. A randomized trial in Mali of the effectiveness of weekly iron supplements given by teachers on the hemoglobin concentrations of schoolchildren. *Public Health Nutr.* 2002;5(3):413–8.
- Tatala SR, Kihamia CM, Kyungu LH, Svanberg U. TanzaniaRisk factors for anemia in schoolchildren in Tanga region. *Tanzania J Health Res.* 2008;10(4):189–202.
- Alaofè H, Zee J, Dossa R, O'Brien HT. Education and improved iron intakes for treatment of mild iron-deficiency anemia in adolescent girls in southern Benin. *Food Nutr Bull.* 2009;30(1):24–36.
- Kikafunda JK, Lukwago FB, Turyashemererwa F. Anemia and associated factors among under-fives and their mothers in Bushenyi district, Western Uganda. *Public Health Nutr.* 2009;12(12):2302–8.
- Hashizume M, Shimoda T, Sasaki S, Kunii O, Caypil W, Dauletbaev D, et al. Anaemia in relation to low bioavailability of dietary iron among school-aged children in the Aral Sea region, Kazakhstan. *Int J Food Sci Nutr.* 2004;55(1):37–43.
- Djokic D, Drakulovic MB, Radojicic Z, Radovic LC, Rakic L, Kocic S, et al. Risk factors associated with anemia among Serbian school-age children 7–14 years old: results of the first national health survey. *Hipokratia.* 2010;14(4):252.
- Bahati YL, Delanghe J, Balaluka GB, Kishabongo AS, Philippé J. Asymptomatic submicroscopic plasmodium infection is highly prevalent and is associated with anemia in children younger than 5 years in south kivu/democratic republic of Congo. *Am J Trop Med Hygiene.* 2020;102(5):1048.
- Ginzburg YZ, Glassberg J. Inflammation, hemolysis, and erythropoiesis lead to competitive regulation of hepcidin and possibly systemic iron status in sickle cell disease. *EBio Medicine.* 2018;34:8–9.
- Gebreweld A, Ali N, Ali R, Fisha T. Prevalence of anemia and its associated factors among children under five years of age attending at Gugufu health center, south Wollo, Northeast Ethiopia. *PLoS One.* 2019;14(7):e0218961.
- Cheesbrough M. *District laboratory practice in tropical countries, vol. 2.* 2nd ed. Cambridge: Cambridge University Press; 2005.
- Cheesbrough M. *District laboratory practice in tropical countries, vol. 1.* 2nd ed. Cambridge: Cambridge University Press; 2009.
- Pita GM, Jiménez S, Basabe B, García RG, Macías C, Selva L, et al. Anemia in children under five years old in eastern Cuba, 2005–2011. *Medic Rev.* 2014;16:16–23.
- Huang Z, Jiang FX, Li J, Jiang D, Xiao TG, Zeng JH. Prevalence and risk factors of anemia among children aged 6–23 months in Huaihua, Hunan Province. *BMC Public Health.* 2018;18(1):1–1.
- Tine RC, Ndiaye M, Hansson HH, Ndour CT, Faye B, Alifrangis M, et al. The association between malaria parasitaemia, erythrocyte polymorphisms, malnutrition and anaemia in children less than 10 years in Senegal: a case control study. *BMC Res Notes.* 2012;5(1):1–0.
- Mboya IB, Mamseri R, Leyaro BJ, George J, Msuya SE, Mgongo M. Prevalence and factors associated with anemia among children under five years of age in Rombo district, Kilimanjaro region, northern Tanzania. *F1000Research.* 2020;9(1102):1102.
- Sirdah MM, Yaghi A, Yaghi AR. Iron deficiency anemia among kindergarten children living in the marginalized areas of Gaza strip, Palestine. *Rev Bras Hematol Hemoter.* 2014;36:132–8.
- Ofori L, Manortey S, Vetsi O, Nartey C, Ugorji HO. Factors contributing to Anaemia in children under five years in the Ga east municipality, Ghana. *Int J Res Rep Hematol.* 2020;17:26–40.
- Semedo RM, Santos MM, Baião MR, Luiz RR, da Veiga GV. Prevalence of anaemia and associated factors among children below five years of age in Cape Verde, West Africa. *J Health Popul Nutr.* 2014;32(4):646.



30. Belachew A, Tewabe T. Under-five anemia and its associated factors with dietary diversity, food security, stunted, and deworming in Ethiopia: systematic review and meta-analysis. *Syst Rev*. 2020;9(1):31.
31. Gutema B, Adissu W, Asress Y, Gedefaw L. Anemia and associated factors among school-age children in Filu town, Somali region, Southeast Ethiopia. *BMC Hematol*. 2014;14(1):1–6.
32. Tiku YS, Mekonnen TC, Workie SB, Amare E. Does Anaemia have major public health importance in children aged 6–59 months in the Duggina Fanigo District of Wolaita zone, southern Ethiopia? *Ann Nutr Metab*. 2018;72(1):3–11.
33. Takele WW, Baraki AG, Wolde HF, Desyibelew HD, Derseh BT, Dadi AF, et al. Anemia and contributing factors in severely malnourished infants and children aged between 0 and 59 months admitted to the treatment centers of the Amhara region, Ethiopia: a multicenter chart review study. *Anemia*. 2021;2021.
34. Tesema GA, Worku MG, Tessema ZT, Teshale AB, Alem AZ, Yeshaw Y, et al. Prevalence and determinants of severity levels of anemia among children aged 6–59 months in sub-Saharan Africa: a multilevel ordinal logistic regression analysis. *PLoS One*. 2021;16(4):e0249978.
35. Chatterjee A, Bosch RJ, Kupka R, Hunter DJ, Msamanga GI. Predictors and consequences of anaemia among antiretroviral-naïve HIV-infected and HIV-uninfected children in Tanzania. *Public Health Nutr*. 2010;13:289–96.
36. Santos RF, Gonzalez ES, Albuquerque EC, Arruda IK, Diniz AD, Figueroa JN, et al. Prevalence of anemia in under five-year-old children in a children's hospital in Recife, Brazil. *Rev Bras Hematol Hemoter*. 2011;33:100–4.
37. Nambiema A, Robert A, Yaya I. Prevalence and risk factors of anemia in children aged from 6 to 59 months in Togo: analysis from Togo demographic and health survey data, 2013–2014. *BMC Public Health*. 2019;19(1):1–9.
38. Simbauranga RH, Kamugisha E, Hokororo A, Kidenya BR, Makani J. Prevalence and factors associated with severe anaemia amongst under-five children hospitalized at Bugando medical Centre, Mwanza, Tanzania. *BMC Hematol*. 2015;15(1):1–9.
39. Kejo D, Petrucka PM, Martin H, Kimanya ME, Moshia TC. Prevalence and predictors of anemia among children under 5 years of age in Arusha District, Tanzania. *Pediatric Health Med Therapeut*. 2018;9:9.
40. Brooker S, Peshu N, Warn PA, Mosobo M, Guyatt HL, Marsh K, et al. The epidemiology of hookworm infection and its contribution to anaemia among pre-school children on the Kenyan coast. *Trans R Soc Trop Med Hyg*. 1999;93(3):240–6.
41. Ewusie JE, Ahiadeke C, Beyene J, Hamid JS. Prevalence of anemia among under-5 children in the Ghanaian population: estimates from the Ghana demographic and health survey. *BMC Public Health*. 2014;14(1):1–9.
42. Akalu TY, Baraki AG, Wolde HF, Desyibelew HD, Derseh BT, Dadi AF, et al. Anemia and determinants among severely malnourished children admitted to Amhara regional referral hospitals, Northwest Ethiopia. *Open J Nutr Food Sci*. 2020;2(1):1007.
43. Aimone AM, Perumal N, Cole DC. A systematic review of the application and utility of geographical information systems for exploring disease-disease relationships in paediatric global health research: the case of anaemia and malaria. *Int J Health Geogr*. 2013;12(1):1–3.
44. Tesfaye TS, Tessema F, Jarso H. Prevalence of anemia and associated factors among "apparently healthy" urban and rural residents in Ethiopia: a comparative cross-sectional study. *J Blood Med*. 2020;11:89.
45. Habyarimana F, Zewotir T, Ramroop S. Structured additive quantile regression for assessing the determinants of childhood anemia in Rwanda. *Int J Environ Res Public Health*. 2017 Jun;14(6):652.
46. Khan JR, Awan N, Misu F. Determinants of anemia among 6–59 months aged children in Bangladesh: evidence from nationally representative data. *BMC Pediatr*. 2016;16(1):1–2.
47. Alemu M, Kinfe B, Tadesse D, Mulu W, Hailu T, Yizengaw E. Intestinal parasitosis and anaemia among patients in a health center, North Ethiopia. *BMC Res Notes*. 2017;10(1):1–7.
48. Molla E, Mamo H. Soil-transmitted helminth infections, anemia and undernutrition among schoolchildren in Yirgacheffee, South Ethiopia. *BMC Res Notes*. 2018;11(1):1–7.
49. Hailemeskel E, Erko B, Degarege A. Community-level epidemiology of intestinal helminth infections and anemia in Harbu town, northeastern Ethiopia. *Parasitol Res*. 2020;119(10):3451–7.
50. Zeki IN, Al-Warid HS. The prevalence of anemia among children infected with *Entamoeba histolytica* in Baghdad. *Iraqi J Sci*. 2019;29:2590–9.
51. Aryadnyani NP, Inderiati D, Ulfah F. Hemoglobin levels on *Trichuris trichiura* infection in children. *Med Lab Technol J*. 2020;6(1):79–83.
52. Stephenson L, Latham AE, Kinoti S, Pertet A. Weight gain of Kenyan school children infected with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* improved following once-or twice-yearly treatment with albendazole. *J Nutr*. 1993;123:656–65.
53. Stephenson LS, Holland CV, Cooper ES. The public health significance of *Trichuris trichiura*. *Parasitology*. 2000 Oct;121(S1):S73–95.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

