

RESEARCH

Open Access



Height is a predictor of hamstring tendon length and ACL graft characteristics in adolescents

Martijn Dietvorst^{1*}, M. C. Marieke van der Steen^{1,2}, Marijn van den Besselaar¹ and Rob PA Janssen^{1,3,4}

Abstract

Background Knowing the potential hamstring tendon length is relevant for planning ligament reconstructions in children and adolescents, as it is not uncommon to encounter small hamstring tendons intraoperatively. The aim of this study is to predict semitendinosus and gracilis tendon length based on anthropometric values in children and adolescents. The secondary aim is to analyse hamstring tendon autograft characteristics in a closed socket anterior cruciate ligament reconstructions and to evaluate the relationship with anthropometric variables. The hypothesis of this study was that height is predictor of hamstring tendon length and thereby graft characteristics.

Methods This observational study included two cohorts of adolescents undergoing ligament reconstructions between 2007–2014 and 2017–2020. Age, sex, height and weight were recorded preoperatively. Semitendinosus and gracilis tendon length and graft characteristics were measured intraoperatively. Regression analysis was performed on tendon length and anthropometric values. Subgroup analyses of the closed socket ACL reconstruction were performed and the relation between anthropometric values and graft characteristics were analysed.

Results The population consisted of 171 adolescents from 13 to 17 years of age, with a median age of 16 years [IQR 16–17]. The median semitendinosus tendon length was 29 cm [IQR 26–30] and gracilis tendon length was 27 cm [IQR 25–29]. Height was a significant predictor of semitendinosus and gracilis tendon length. Subgroup analysis of the closed socket ACL reconstructions showed that in 75% of the procedure, the semitendinosus tendon alone was sufficient to create a graft with a minimum diameter of 8.0 mm.

Conclusions Height is a significant predictor of semitendinosus and gracilis tendon length in adolescents between 13 and 17 years of age and outcomes are similar to data in adults. In 75% of the closed socket ACL reconstructions, the semitendinosus tendon alone is sufficient to create an adequate graft with a minimum diameter of 8 mm. Additional use of the gracilis tendon is more often necessary in females and shorter patients.

Level of evidence 3

Keywords Hamstring tendon, ACL reconstruction, graft, anthropometrics, adolescents

*Correspondence:

Martijn Dietvorst

martijn.dietvorst@mmc.nl

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Anterior cruciate ligament (ACL) injury in children and adolescents is a severe injury of the knee. The ACL reconstruction technique for children and adolescents depends on skeletal maturity and is also surgeon dependent [1, 2]. However, general principles, such as the use of a well-positioned soft-tissue autograft with an adequate size, diameter and fixation for ACL reconstruction in adults also apply for the younger patient population [1, 3]. Desired graft length depends on the type of ligament reconstruction, operative technique, for example full tunnels versus closed-sockets techniques, and fixation methods [4]. However, the graft diameter is important regardless of the technique, as a diameter of less than 8 mm is related to higher revision rates within the age category of 20 years and younger [5–11].

Quadrupled hamstring autograft is most commonly used as soft tissue graft for ACL reconstruction in children and adolescents [3]. Both semitendinosus (ST) as semitendinosus-gracilis (STG) grafts are used for ACL reconstructions. In some children and adolescents, the harvested hamstring graft might be too small to produce a graft with adequate specifications [8]. Preoperative knowledge of potential tendon dimensions could assist in graft planning for knee ligament surgery, as complex knee ligament reconstructions require specific tendon length and diameter [12]. Various studies on the anthropometric predictability of tendon dimensions have been conducted in adults, reporting that height is a predictor of hamstring tendon length [12–14]. The question remains whether these outcomes are applicable in adolescents, as adolescence is a period of growth, development and maturation. To this date, one study evaluated the predictability of hamstring tendon dimensions in children and adolescents based on magnetic resonance imaging (MRI) measurements [15]. Other studies evaluated the relationship of anthropometric values or MRI measurements with the diameter of a hamstring tendon autograft for a specific ACL reconstruction technique, but did not analyse the length of the hamstring tendons itself [16–19]. Predictability of hamstring tendon lengths in a paediatric and adolescent population has not been studied to this date, despite that tendon length is also an important aspect to be able to obtain the desired graft dimensions during ACL reconstruction or planning for multiligament reconstructions.

The primary aim of this study is to analyse the preoperative predictability of the ST and G tendon lengths based on anthropometric data in adolescents. We hypothesize that height will be a relevant anthropometric predictor. As a secondary aim this study analyses anthropometric characteristics in relation with graft characteristics, such as ST or STG graft, length and diameter in a closed

socket ACL reconstruction technique. We hypothesize that graft characteristics in a closed socket ACL reconstruction, such as additional use of G tendon for graft (ST versus STG graft), can also be predicted by height due to the predictability of tendon length by height.

Methods

In this retrospective study of prospectively obtained data, consecutive adolescent patients with ACL rupture, scheduled for ACL reconstruction between 2007–2014 and 2017–2020, were eligible for inclusion. In both periods, preoperative height, weight, BMI, sex and age were recorded and hamstring tendon lengths and graft characteristics were described intra-operatively. In the period from 2014 to 2017, the hamstring tendon lengths were not measured and patients from that period were therefore not included in this study. The patients included between 2007 and 2014 were also part of the data analysed in the study by Janssen et al. [12] and in the current study a subgroup analysis on the younger patients (<18 years) of that cohort was performed [12]. From 2007 to 2014, a full tunnel ACL reconstruction was used. From 2017, a closed socket technique (All-Inside, Arthrex®, Naples, USA) was used for ACL reconstructions. Anthropometric values, hamstring tendon lengths and graft characteristics were recorded again from 2017. All patients (<18 years) undergoing primary reconstruction with hamstring tendon autografts were eligible for inclusion. Exclusion criteria were ACL reconstruction with other types of allo- or autografts, previous harvest of the ipsilateral hamstring tendon, congenital limb deficiency that would affect total body weight and neuromuscular disorders. Preoperatively sex, age, height and weight of the patient were recorded as anthropometric variables.

Two orthopaedic surgeons (RJ and MvdB) performed all procedures using the same technique. From 2007 to 2014, a STG autograft with WasherLoc™ (Zimmer Biomet®, Warsaw, Indiana, USA) was used as the primary graft for a full-tunnel ACL reconstruction. The methodological description of that period has been published previously [12, 20].

From 2017, a closed socket technique (All-Inside, Arthrex®, Naples, USA) was used to reconstruct the ACL in young ACL reconstruction patients with the ST as autograft type of choice. The ST tendon was harvested and prepared in a standardized fashion according to the previous study [12, 20]. The available length of each tendon was measured with a ruler and recorded in cm, rounded off to the nearest 0.5 cm. A provisional ST graft was then created and the length and diameter of the graft were measured. The diameter of the hamstring graft was measured by soft tissue graft caliper (Arthrex®, Naples,

Florida, USA) with 0.5-mm increments between holes and the length of the graft is measured with a ruler. In case of insufficient graft diameter of the ST graft, the G tendon was also harvested. The primary goal was to create a hamstring autograft with a minimum length of 6 to 6.5 cm and diameter of ≥ 8 mm, preferably as a 4-strand ST graft (4-ST). In order to create a 4-ST graft of 6 cm, a minimum ST tendon length of 24 cm was necessary. Depending on length of the tendons and/or diameter of the graft, strand variations of the graft are possible with or without the use of the G tendon. The possible variations were a 3-strand ST, 4-strand ST, 5-strand ST(G), 6-strand STG, 7-strand STG or 8-strand STG graft.

Statistical analyses

Descriptive analyses were used to describe the groups based on the ACL reconstruction technique. Baseline parameters were compared between groups by medians of the Mann Whitney U test after tests for normality for the continuous variables and Chi square test for sex. Multivariate linear regression analyses were performed to predict the length of harvested ST and G tendons length. As variables of interest the anthropometric parameters sex, age, height, weight and BMI were considered. In order to create a standard 4-ST graft a minimum of 24 cm ST tendon length is necessary, therefore the division between significant anthropometric values and ST tendon length of 24 cm was analysed. Additional analyses were performed on the 2017–2020 closed socket technique subgroup. A logistic regression analysis was performed to analyse the necessity for additional use of the G tendon in the 2017–2020 group. Data analysis was performed in SPSS Statistics version 22.0 (IBM, Armonk, New York, USA). Significance was set at ≤ 0.05 in all analyses.

Ethical approval

The Institutional Review Board (IRB) of The Medical Ethical Committee Máxima Medisch Centrum determined

that this study was not subjected to the guidelines of the Medical Research Involving Human Subjects Acts (WMO) (N20.038).

Results

A total of 171 patients were included for analysis, of which 99 patients were included in the period from 2007–2014 and 72 patients from 2017–2020. See Table 1 for baseline characteristics and Fig. 1 for the age distribution with median heights. The height, weight and BMI of 2 patients were unknown and of 1 patient the ST length was unknown.

IQR interquartile range

Prediction of tendon lengths

The multivariate linear regression analyses in the total group on the ST and G tendon lengths are shown in Table 2. Height is a predictor for both the ST and G tendon length. For each centimetre increase in height, the predictive length of the ST and G tendon increase 0.18 cm and 0.14 cm respectively.

In order to create a standard 4-ST graft, the required ST tendon length was 24 cm. In Table 3 the division between ST tendon lengths of 24 cm or more and height categories are shown, as height was a significant predictor of ST tendon length.

Closed socket subgroup analysis

In the 2017–2020 group, 75% of the closed socket ACL reconstructions were performed with a ST tendon autograft only. In 18 (25%) cases, an additional G tendon autograft was necessary to achieve adequate graft dimension. Patients with a STG graft were significantly more often female, were shorter and lighter and had shorter ST tendons as is shown in Table 4.

Four of the 72 patients (6%) had a graft diameter of less than 8 mm (all had 7.5 mm), which were all ST grafts and these graft configurations were accepted intra-operatively. Four strand ST graft was used most

Table 1 Characteristics of the study population

	Total	2007–2014	2017–2020	<i>p</i> -value
	<i>N</i> = 171 Median [IQR]	<i>N</i> = 99 Median [IQR]	<i>N</i> = 72 Median [IQR]	
Age (years)	16.0 [16.0–17.0]	16.0 [16.0–17.0]	16.0 [15.0–17.0]	0.114
Sex, female %	47%	47%	49%	0.931
Height (cm)	174 [168–182]	174 [168–183]	172 [168–182]	0.354
Weight (kg)	65.0 [60.0–73.5]	65.0 [61.0–74.0]	65.0 [58.0–72.0]	0.388
BMI (kg/m ²)	21.5 [19.8–23.8]	21.5 [20.1–23.7]	21.4 [19.5–24.0]	0.664
ST length (cm)	29.0 [26.0–30.1]	29.0 [27.0–31.0]	28.5 [26.0–30.0]	0.238
G length (cm)	27.0 [25.0–29.0]	28.0 [25.0–29.5] <i>N</i> = 99	24.0 [23.0–27.0] <i>N</i> = 18	0.003




Fig. 1 Age distribution of females and males. Numbers above columns represent median height in cm [IQR] within the age-group

Table 2 Multivariate linear regression analysis of anthropometric values and ST or G length

Multivariate						
Model	Semitendinosus tendon length			Gracilis tendon length		
	Regression coefficients (95% CI)	P-value	R ²	Regression coefficients (95% CI)	P-value	R ²
Constant	0.702 (-11.848; 13.252)	0.912	0.337	5.223 (-12.456; 22.921)	0.559	0.212
Age (years)	-0.135 (-0.551; 0.281)	0.522		-0.164 (-0.807; 0.408)	0.616	
Sex (female)	-0.876 (-1.978; 0.225)	0.118		-0.619 (-2.132; 0.894)	0.420	
Height (cm)	0.180 (0.115; 0.245)	<0.001		0.140 (0.053; 0.226)	0.002	
Weight (kg)	-0.016 (-0.059; 0.028)	0.476		0.004 (-0.054; 0.062)	0.898	

CI Confidence interval

Table 3 Relation of height categories and ST tendon length

Height (cm)	ST tendon length	
	< 24 cm N (%)	≥ 24 cm N (%)
< 160	4 (44)	5 (56)
160–170	5 (13)	33 (87)
170–180	3 (4)	65 (96)
≥ 180	0 (0)	53 (100)

frequently. In case of using an STG graft, a six strand variation was used most frequently.

The results of the univariate logistic regression analyses of height and additional G tendon use as hamstring autograft is shown in Table 5. The STG group consisted of 18 patients and therefore no stable multivariate model could be created. Due to the significance of height in previous analyses, height was chosen as a factor of interest. According to the univariate logistic regression analysis, patient’s height is a statistically relevant predictor for additional need for the G tendon as autograft.

Patients with an STG graft were significantly more often females. Differences in height and weight could therefore be the result of differences in sex between groups and not to differences in BMI, as BMI was similar in both groups. Males had a significantly greater median height than females (180 cm [IQR 175–185] versus 168 cm [165–170], *p*-value<0.001). Males also were heavier than females (68 kg [60–79] versus 63 kg [55–67], *p*=0.009). Univariate logistic regression analysis of height and additional G tendon use within females (*n*=35, of which *n*=13 having a STG graft) showed no statistically significant value of height as is shown in Table 6.

Discussion

Height is a significant predictor of ST and G tendon length in the age category of 13 to 17 years. This study is the first study to analyse the relationship between anthropometrics, hamstring tendon lengths and graft characteristics in a closed-socket ACL reconstruction. Knowing the potential ST and G tendon length is relevant for planning the hamstring tendon autograft for ligament reconstructions in adolescents [8, 12]. Complex knee ligament

Table 4 Characteristics of patient with a ST and STG graft from the 2017–2020 closed socket group

	Patients with ST graft	Patients with STG graft	P-value
	N = 54 Median [IQR]	N = 18 Median [IQR]	
Age (years)	16.0 [15.0–17.0]	16.0 [15.8–17.0]	0.402
Sex, female %	41%	72%	0.021
Height (cm)	175 [169–182]	170 [165–175]	0.032
Weight (kg)	65.0 [60.0–74.0]	58.5 [53.5–67.8]	0.021
BMI (kg/m ²)	21.8 [19.6–24.1]	20.4 [19.1–23.1]	0.352
ST length (cm)	29.0 [26.0–30.0]	25.8 [24.0–28.1]	0.001
G length (cm)	NA	24.0 [23.0–27.0]	NA
Graft diameter (mm)	8.5 [8.0–9.0]	8.5 [8.0–9.0]	0.568
Strand type % (n)			
3-strand	2% (1)	0% (0)	
4-strand	85% (46)	0% (0)	
5-strand	13% (7)	0% (0)	
6-strand	0% (0)	73% (13)	
8-strand	0% (0)	22% (4)	
10-strand	0% (0)	6% (1)	

IQR Interquartile range, NA Not applicable

Table 5 Univariate logistic regression analysis of height and additional G tendon use (0=no additional G tendon use; 1 = additional G tendon use)

Univariate			
Model	Regression coefficient (95% CI)	P-value	Nagelkerke R ²
Constant	11.514	0.056	0.095
Height (cm)	-0.073 (0.868–0.996)	0.038	

CI Confidence interval

Table 6 Univariate logistic regression analysis of height and additional G tendon use (0=no additional G tendon use; 1 = additional G tendon use) within the female subgroup

Univariate			
Model	Regression coefficient (95% CI)	P-value	Nagelkerke R ²
Constant	12.202	0.264	0.056
Height (cm)	-0.076 (0.815–1.053)	0.244	

CI Confidence interval

reconstructions require specific tendon lengths to create the desired graft dimensions and there are concerns that in some adolescents, the harvested hamstring graft can be too small to produce a graft with an adequate diameter [8, 12]. To this date, no study previously analysed the prediction of hamstring tendon lengths focusing on an adolescent population. The results of this current study are in accordance with the first hypothesis and the study

by Janssen et al. [12], which showed that for each increase in 1 cm in body height, the ST and G length increase respectively with 0.20 cm and 0.16 cm in a population with a mean age of 28.7 years [12]. The importance of this study is that hamstring tendon length can be predicted by preoperative body height and hamstring tendons were in all cases large enough to create grafts with a diameter > 8 mm.

Considering that adolescents experience a growth spurt it might be surprising that age was not a significant predictor for tendon lengths in the multivariate regression analyses. However, considering the large variation in onset and duration of this growth spurt [21], maturation of adolescents seems best captured by height instead of age. Furthermore, the great majority of the population were 16 and 17 year-olds and may therefore have reached final height [22]. The increase in height is not only caused by growth of the lower extremities, but also by spinal growth. It is not known whether the increase in height of the lower extremities would result in a relatively equal increase in hamstring tendon lengths. There is limited evidence of the development of human tendons in vivo [23]. Current available evidence shows that throughout childhood and adolescence, the tendons seem to adapt in size and structure as the musculotendinous structures develop [24–27]. The influence of growth on hamstring tendon lengths has not been evaluated previously. However, the influence of growth on Achilles and patellar tendon lengths have been studied and showed that the lengths of both tendons in 14 year-olds boys are similar to adults [28, 29].

Current ACL reconstruction techniques allow the use of multiple-stranded hamstring autografts and depending on the reconstruction technique, different socket/tunnel lengths require different graft lengths [12]. For example, in order to create a 4-strand ST with a minimum length of 60 mm, a minimum ST tendon length of 24 cm (4×60 mm) is necessary. In 44% of the patients with a height of <160 cm the ST tendon was shorter than the required 24 cm. It is necessary in those cases to harvest an additional G tendon, as a 3-ST graft in most cases did not reach the required diameter. A recent systematic review showed that the hamstring tendon graft diameter should be ≥ 7 mm, but a threshold towards larger graft diameters should be considered for patients younger than 20 years [30]. High graft failure rates are problematic in this young population [31]. A graft diameter of less than 8 mm is related to higher revision rates within the age category of 20 years and younger [5–11]. For each increase of 0.5 mm in diameter within the 7.0 to 9.0 mm range, the likelihood of a revision was 0.82 lower [11]. It is therefore important to reduce the risks of graft failure by creating a graft with an adequate length and diameter [9]. Preoperative prediction of tendon length might therefore help in preparing the graft for ligament reconstructions.

In the current cohort of the closed socket technique, 25% of the reconstructed ACL autografts required an additional G tendon to create a graft of sufficient diameter. All STG grafts had a diameter of more than 8 mm and did not need augmentation of contralateral hamstring tendon autograft or allograft. This finding is somehow similar to the outcomes of the study by Stergios et al. [32], who found that the ST tendon alone was insufficient to create a 4-strand graft with a minimum diameter of 7 mm in one in five adult patients [32]. In the current study, logistic regression analysis showed that additional use of the G tendon can be predicted by height, which was in accordance with the second hypothesis. However, as there were significantly more females in the STG group who were significantly smaller than males, subgroup analysis of females showed no significant effect of height. The effect of height on additional use of the G tendon might be explained by the findings that more females needed an additional G tendon and females were smaller than males. This is in line with previous literature showing that females more often had an inadequate ST tendon length to create a ST 4-strand graft and alternative graft options should be considered [18, 32].

This study has several limitations. The first limitation is that the diameter of the tendons is not measured, although most likely both tendon length and diameter contribute to graft size. Recent studies showed that anthropometric data and CSA measurements of

hamstring tendon on MRI are correlated to the diameter of hamstring grafts [17–19, 33]. Another limitation of this study is that, similar to the study by Calvo et al. [16], measurements were based on chronological age and not on physiological age [16]. The number of children with remaining growth of the lower extremity is therefore not known. In these cohorts, Tanner stages were not obtained. As the majority of the cohort is 16 and 17 years of age and youngest patients are 13 years of age, it is expected that most patients (especially females) have reached Tanner stage 4 or 5, which means that no or very limited residual skeletal growth is expected. This is in accordance with the median heights in our cohorts, which were not statistically different between different age categories. The influence of physiological age and residual growth on tendon characteristics was not investigated, which would be an interesting topic for future research. Future research should also aim at skeletally immature children specifically of which bone age and remaining growth of the lower extremity is known. Finally, this study population consisted of Caucasian adolescents. Chiang et al. [13] concluded in their study that Caucasian patients had significantly longer hamstring tendons compared to Chinese Han population [12, 13]. The outcomes of this study might therefore not be extrapolated to adolescents of other ethnicities.

Conclusions

Height is a significant predictor of ST and G tendon length in adolescents between 13 and 17 years of age and outcomes are similar to data in adults. In 75% of the closed socket ACL reconstructions, the ST tendon alone is sufficient to create an adequate graft with a minimum diameter of 8 mm. Additional use of the G tendon is more often necessary in females and shorter patients.

Abbreviations

ACL	Anterior Cruciate Ligament
BMI	Body Mass Index
CM	Centimetre
CSA	Cross-Sectional Area
G	Gracilis
IQR	Interquartile Range
IRB	The Institutional Review Board
KG	Kilograms
M	Meter
MRI	Magnetic Resonance Imaging
N	Number
ST	Semitendinosus
STG	Semitendinosus Gracilis combined

Acknowledgements

The authors want to thank Miranda, Carla, Annie, Anke and all other scrub nurses for measuring and collecting the intraoperative data of the tendons and grafts.

Authors' contributions

MD was corresponding author, collected and analysed data; MvdS collected data and co-wrote the manuscript, MvdB collected data and co-wrote the manuscript, RJ was the initiator of the project, collected data and co-wrote the manuscript. All authors reviewed the manuscript.

Funding

There is no external funding for this research project.

Availability of data and materials

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Declarations**Ethics approval and consent to participate**

The Institutional Review Board (IRB) of The Medical Ethical Committee Máxima Medisch Centrum determined that this study was not subjected to the guidelines of the Medical Research Involving Human Subjects Acts (WMO) (N20.038). All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all subjects or, if subjects are under 16, from a parent and/or legal guardian.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Orthopaedic Surgery and Trauma, Máxima MC, Eindhoven, the Netherlands. ²Department of Orthopaedic Surgery and Trauma, Catharina Hospital Eindhoven, Eindhoven, the Netherlands. ³Dept. of Paramedical Sciences, Chair Value-Based Health Care, Fontys University of Applied Sciences, Eindhoven, the Netherlands. ⁴Department of Biomedical Engineering, Orthopaedic Biomechanics, Eindhoven University of Technology, Eindhoven, the Netherlands.

Received: 10 September 2022 Accepted: 7 July 2023

Published online: 11 July 2023

References

- Ardern CL, Ekås GR, Grindem H, et al. 2018 International Olympic Committee consensus statement on prevention, diagnosis and management of paediatric anterior cruciate ligament (ACL) injuries. *Br J Sports Med.* 2018;52(7):422–38.
- Dietvorst M, Reijman M, van Zutven R, et al. Current State of Care for Pediatric ACL Ruptures in the Netherlands: A Survey. *J Knee Surg.* 2019. <https://doi.org/10.1055/s-0039-1697626>.
- Seil R, Cucchi D, Ninulescu C, et al. Anatomic anterior cruciate ligament reconstruction for adolescent patients with open physis. *Ann Joint.* 2019;4:31.
- Connaughton AJ, Geeslin AG, Uggen CW. All-inside ACL reconstruction: How does it compare to standard ACL reconstruction techniques? *J Orthop.* 2017;14(2):241–6.
- Conte EJ, Hyatt AE, Gatt CJ Jr, et al. Hamstring autograft size can be predicted and is a potential risk factor for anterior cruciate ligament reconstruction failure. *Arthroscopy.* 2014;30(7):882–90.
- Magnussen RA, Lawrence JTR, West RL, et al. Graft size and patient age are predictors of early revision after anterior cruciate ligament reconstruction with hamstring autograft. *Arthroscopy.* 2012;28(4):526–31.
- Mariscalco MW, Flanigan DC, Mitchell J, et al. The influence of hamstring autograft size on patient-reported outcomes and risk of revision after anterior cruciate ligament reconstruction: a Multicenter Orthopaedic Outcomes Network (MOON) Cohort Study. *Arthroscopy.* 2013;29(12):1948–53.
- Pennock AT, Ho B, Parvanta K, et al. Does allograft augmentation of small-diameter hamstring autograft ACL grafts reduce the incidence of graft retear? *Am J Sports Med.* 2017;45(2):334–8.
- Perkins CA, Busch MT, Christino M, et al. Allograft augmentation of hamstring anterior cruciate ligament autografts is associated with increased graft failure in children and adolescents. *Am J Sports Med.* 2019;47(7):1576–82.
- Snaebjörnsson T, Hamrin-Senorski E, Svantesson E, et al. Graft diameter and graft type as predictors of anterior cruciate ligament revision: a cohort study including 18,425 patients from the Swedish and Norwegian national knee ligament registries. *J Bone Joint Surg Am.* 2019;101(20):1812–20.
- Spragg L, Chen J, Mirzayan R, et al. The effect of autologous hamstring graft diameter on the likelihood for revision of anterior cruciate ligament reconstruction. *Am J Sports Med.* 2016;44(6):1475–81.
- Janssen RPA, van der Velden MJ, van den Besselaar M, et al. Prediction of length and diameter of hamstring tendon autografts for knee ligament surgery in Caucasians. *Knee Surg Sports Traumatol Arthrosc.* 2017;25(4):199–204.
- Chiang ER, Ma HL, Wang ST, et al. Hamstring graft sizes differ between Chinese and Caucasians. *Knee Surg Sports Traumatol Arthrosc.* 2012;20(5):916–21.
- Treme G, Diduch DR, Billante MJ, et al. Hamstring graft size prediction: a prospective clinical evaluation. *Am J Sports Med.* 2008;36(11):2204–9.
- Tomaszewski R, Smyczek D, Woś-Cieśla I, et al. Developmental changes in ACLs and semitendinosus tendons dimensions according to age in children. *J Orthop Surg Res.* 2020;15(1):1–7.
- Calvo RR, Espinosa MI, Figueroa DP, et al. Prediction of semitendinosus-gracilis graft diameter in children and adolescents using anthropometric measures. *Knee Surg Sports Traumatol Arthrosc.* 2016;24(3):702–6.
- Bickel BA, Fowler TT, Mowbray JG, et al. Preoperative magnetic resonance imaging cross-sectional area for the measurement of hamstring autograft diameter for reconstruction of the adolescent anterior cruciate ligament. *Arthroscopy.* 2008;24(12):1336–41.
- Boisvert CB, Aubin ME, DeAngelis N. Relationship between anthropometric measurements and hamstring autograft diameter in anterior cruciate ligament reconstruction. *Am J Orthop (Belle Mead NJ).* 2011;40(6):293–5.
- Partan MJ, Stapleton EJ, Atlas AM, DiMauro JP. Predicting Autologous Hamstring Graft Diameter in the Pediatric Population Using Preoperative Magnetic Resonance Imaging and Demographic Data. *Am J Sports Med.* 2021;49(6):1482–91.
- Janssen RPA, van der Velden MJ, Pasmans HL, et al. Regeneration of hamstring tendons after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2013;21(4):898–905.
- Soliman A, De Sanctis V, Elalaily R, Bedair S. Advances in pubertal growth and factors influencing it: Can we increase pubertal growth? *Indian J Endocrinol Metab.* 2014;18(Suppl 1):S53–62.
- Schönbeck Y, Talma H, van Dommelen P, et al. The world's tallest nation has stopped growing taller: the height of Dutch children from 1955 to 2009. *Pediatr Res.* 2013;73(3):371–7.
- Mersmann F, Böhm S, Arampatzis A. Imbalances in the development of muscle and tendon as risk factor for tendinopathies in youth athletes: a review of current evidence and concepts of prevention. *Front Physiol.* 2017;8:987.
- Radnor JM, Oliver JL, Waugh CM, et al. The influence of growth and maturation on stretch-shortening cycle function in youth. *Sports Med.* 2018;48(1):57–71.
- Kubo K, Kanehisa H, Kawakami Y, et al. Elastic properties of muscle-tendon complex in long-distance runners. *Eur J Appl Physiol.* 2000;81(3):181–7.
- O'Brien TD, Reeves ND, Baltzopoulos V, et al. Mechanical properties of the patellar tendon in adults and children. *J Biomech.* 2010;43(6):1190–5.
- Waugh CM, Blazevich AJ, Fath F, et al. Age-related changes in mechanical properties of the Achilles tendon. *J Anat.* 2012;220(2):144–55.
- Kubo K, Teshima T, Hirose N, et al. Growth changes in morphological and mechanical properties of human patellar tendon in vivo. *J Appl Biomech.* 2014;30(3):415–22.

29. Kubo K, Teshima T, Hirose N, et al. A cross-sectional study of the plantar flexor muscle and tendon during growth. *Int J Sports Med.* 2014;35(10):828–34.
30. Alomar AZ, Nasser ASB, Kumar A, Kumar M, Das S, Mittal S. Hamstring graft diameter above 7 mm has a lower risk of failure following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2021. <https://doi.org/10.1007/s00167-021-06503-0>.
31. Faunø P, Rahr-Wagner L, Lind M. Risk for revision after anterior cruciate ligament reconstruction is higher among adolescents: results from the Danish registry of knee ligament reconstruction. *Orthop J Sports Med.* 2014;2(10):2325967114552405.
32. Stergios PG, Georgios KA, Konstantinos N, et al. Adequacy of semitendinosus tendon alone for anterior cruciate ligament reconstruction graft and prediction of hamstring graft size by evaluating simple anthropometric parameters. *Anat Res Int.* 2012; 424158.
33. An VV, Scholes C, Mhaskar VA, et al. Regression modelling combining MRI measurements and patient anthropometry for patient screening and prediction of graft diameter in hamstring autograft arthroscopic ACL reconstruction. *Asia Pac J Sports Med Arthrosc Rehabil Technol.* 2017;8:24–31.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

