

RESEARCH ARTICLE

Open Access



The transfer of maternal antibodies and dynamics of maternal and natural infection-induced antibodies against coxsackievirus A16 in Chinese children 0–13 years of age: a longitudinal cohort study

Jiaxin Zhou^{1†}, Yonghong Zhou^{1†}, Kaiwei Luo^{2†}, Qiaohong Liao^{1†}, Wen Zheng¹, Hui Gong¹, Huilin Shi¹, Shanlu Zhao², Kai Wang¹, Qi Qiu¹, Bingbing Dai³, Lingshuang Ren¹, Lili Wang¹, Lidong Gao², Meng Xu¹, Nuolan Liu¹, Wanying Lu¹, Nan Zheng¹, Xinhua Chen¹, Zhiyuan Chen¹, Juan Yang¹, Simon Cauchemez⁴ and Hongjie Yu^{1*}

Abstract

Background: A major hand-foot-and-mouth disease (HFMD) pathogen, coxsackievirus A16 (CVA16), has predominated in several of the last 10 years and caused the largest number of HFMD outbreaks between 2011 and 2018 in China. We evaluated the efficacy of maternal anti-CVA16 antibody transfer via the placenta and explored the dynamics of maternal and natural infection-induced neutralizing antibodies in children.

Methods: Two population-based longitudinal cohorts in southern China were studied during 2013–2018. Participants were enrolled in autumn 2013, including 2475 children aged 1–9 years old and 1066 mother-neonate pairs, and followed for 3 years. Blood/cord samples were collected for CVA16-neutralizing antibody detection. The maternal antibody transfer efficacy, age-specific seroprevalence, geometric mean titre (GMT) and immune response kinetics were estimated.

Results: The average maternal antibody transfer ratio was 0.88 (95% CI 0.80–0.96). Transferred maternal antibody levels declined rapidly (half-life: 2.0 months, 95% CI 1.9–2.2 months). The GMT decayed below the positive threshold (8) by 1.5 months of age. Due to natural infections, it increased above 8 after 1.4 years and reached 32 by 5 years of age, thereafter dropping slightly. Although the average duration of maternal antibody-mediated protection was < 3 months, the duration extended to 6 months on average for mothers with titres ≥ 64 .

[†]Jiaxin Zhou, Yonghong Zhou, Kaiwei Luo and Qiaohong Liao contributed equally to this work.

*Correspondence: yhj@fudan.edu.cn

¹School of Public Health, Fudan University, Key Laboratory of Public Health Safety, Ministry of Education, Shanghai, China

Full list of author information is available at the end of the article



Conclusions: Anti-CVA16 maternal antibodies are efficiently transferred to neonates, but their levels decline quickly. Children aged 0–5 years are the main susceptible population and should be protected by CVA16 vaccination, with the optimal vaccination time between 1.5 months and 1 year of age.

Keywords: Coxsackievirus A16 (CVA16), Transplacental transfer, Maternal antibody, Natural infection, Antibody kinetics

Background

Coxsackie A16 (CVA16) belongs to the picornavirus family, which can cause a wide spectrum of diseases, such as encephalitis, paralysis or myelitis, meningitis, hand-foot-and-mouth disease (HFMD), upper and lower respiratory tract diseases, pleurodynia, herpangina, and myopericarditis [1]. HFMD is a common infectious disease, with CVA16 as the major pathogen, mainly observed among children under 5 years old in the Asia-Pacific region, especially in China. CVA16 predominated in China in 2012, 2014, 2016 and 2018, accounting for 58.6% of HFMD outbreaks between 2011 and 2018. An increasing number of HFMD cases caused by CVA16 have been observed since 2017 [2]. Although most patients show minor symptoms, CVA16 can cause severe and fatal HFMD, and increasing numbers of reports of such outcomes have been noted in the US, France, Japan, mainland China, and Taiwan of China [3]. In Shenyang, northeastern China, 20.7% (19/92) of HFMD cases with neurological symptoms were caused by CVA16 infection, of which 2 patients had brainstem encephalitis and 1 had acute flaccid paralysis [4].

There is currently no specific treatment for HFMD, and the CVA16 HFMD vaccine is still in the preclinical phase, with promising results in animal models [3]. The only HFMD vaccine licenced in mainland China is the enterovirus A71 (EV-A71) monovalent HFMD vaccine, but this vaccine fails to confer cross-protection against HFMD caused by CVA16 or other enteroviruses [5–7]. All of these elements highlight the urgent need for the development of vaccines targeting CVA16.

Maternal antibodies have been recognized as an effective source to protect newborns against measles virus and influenza [8, 9]. Given the absence of a CVA16 vaccine, anti-CVA16 maternal antibodies, as the only source of passive immunity, may play an important role against CVA16 during the early months of life. After the waning of maternal antibodies, young children may become susceptible to infection. Serological studies are necessary to quantify maternal antibody transfer and anti-CVA16 antibody dynamics, including maternally transmitted antibodies as well as natural infection-induced antibodies. Such understanding may help select the target population for future CVA16 vaccination campaigns. Few studies have reported a positive correlation of


anti-CVA16 antibody levels in mother-neonate pairs [10–13]. Only one study reported the average transfer ratio of anti-CVA16 maternal antibodies [11], and the half-life of maternal antibodies was not reported. Limited cohort studies describing maternal antibody dynamics showed that seroprevalence in neonates decreased from 72% at birth to 0–26.4% at 6–7 months of age. However, these results may be affected by the small sample size [12] and limited follow-up [14, 15]. In addition, previous studies were cross-sectional studies, which cannot reveal the dynamics of natural infection-induced antibodies over time [16–20].

Here, we quantify the transfer of maternal antibodies and the dynamics of antibodies from birth to 13 years old using two population-based cohorts: one was a paired mother-neonate cohort (abbr. neonate cohort), and the other was a 1- to 9-year-old paediatric cohort (abbr. paediatric cohort). More specifically, we quantified concentrations of CVA16-specific antibodies in paired maternal and cord serum samples and assessed the transplacental transfer efficiency of maternal antibodies from a large cohort of mothers and neonates. We then analysed the antibody kinetics from birth to age 13 years (including the decline in maternal antibody levels and subsequent increases due to natural infection) using the paediatric cohort and neonate cohort together.

Methods

Study design and participants

Two population-based longitudinal cohorts were established in three townships (Tianzhuang, Jiangnan, and Qingtang) in Anhua County, Hunan Province, a rural area in southern China, including a paired mother-neonate cohort and a 1- to 9-year-old paediatric cohort. The neonate cohort design has been described previously [21]. Briefly, neonates who were born after Sep 20, 2013, and whose mothers resided in the area for at least 3 months were eligible for the mother-neonate paired cohort [21]. A total of 1066 pairs of mothers-neonates out of 3499 eligible mothers and 3532 eligible neonates were enrolled from Sep 20, 2013, to Oct 14, 2015, and we randomly selected half of these participants for this study (Fig. 1A). For the paediatric cohort, children who were 1–9 years old of age at enrolment from Sep to Nov 2013 and who resided in the study sites for more than 3



months were recruited. The 1-year age group was defined as those children aged 12–23 months; other age groups were defined in a similar way. Stratified random sampling of three townships was conducted from the eligible list to select children in each age group. Ultimately, 4188 children aged 1–9 years were enrolled out of 5996 eligible children, and 2475 children were selected to join this study (Fig. 1B). The details of the selection method are described in Additional file 1: Table S1. In addition, enhanced virological monitoring of HFMD was carried out locally in Anhua County from 2013 to 2016 [22].

Procedures

Taking the village as the unit, the parents led the children to the township health centre within the study period to participate in the research. If the participants were

unable or had no time to go to the township health centre, the well-trained project personnel and village doctors investigated the children by household survey as an auxiliary method. For the neonate cohort, a venous blood sample (2 ml) from mothers was collected around the time of delivery, and a cord blood sample (2 ml) was collected for each neonate at baseline. Mothers were invited to answer a questionnaire about basic demographic information and delivery information (such as gestational weeks, delivery method, and neonatal birthweight). After that, the neonates were followed up at 2, 4, 6, 12, 24, and 36 months. Details about the procedures of the neonate cohort can be found elsewhere [21].

For the paediatric cohort, blood samples were collected at baseline during Sep–Nov 2013. Each child was invited to attend the annual follow-up visit between

August and November during 2014–2016 for a total of 3 annual follow-ups. A subset of children was invited to attend another semiannual follow-up visit between February and March during 2014–2016 (total of 6 follow-ups). For both the neonatal and paediatric cohorts, blood samples (2 ml) and questionnaires about hygiene habits and vaccination history were collected at each follow-up by well-trained nurses and epidemiological investigators, respectively.

CVA16 activity was monitored through enhanced virological surveillance of HFMD in Anhua [22] and national notifiable disease surveillance data in Hunan [23].

Laboratory procedures

We performed a neutralization assay to measure serum neutralizing antibodies as previously described [24]. In brief, serum dilutions (ranging from 1:8 to 1:1024) were mixed with an equal volume of 100 TCID₅₀ of the virus strain, with a 2-fold serial dilution. All diluted samples were tested in duplicate, and the titration was double-checked. Antibody titres were defined as the reciprocal of the highest dilution capable of inhibiting 50% of the cytopathic effect and calculated by use of the Karber method [25] (Additional file 1: page 9, line 43–56).

Statistical analysis

The comparison of baseline demographic information between the two groups was conducted with the chi-square test for discrete variables and the Wilcoxon test for continuous variables. An antibody titre ≥ 8 was chosen as the positivity threshold; we considered 16 to indicate positivity in a sensitivity analysis. The antibody titre was log₂-transformed before analysis. Seroprevalence and the geometric mean titre (GMT) were calculated to estimate the maternal and neonatal immunity levels. The average transfer ratio of maternal antibody was defined as the geometric mean of the neonate-to-mother ratio, conditional on positivity of the mother [21]. All mother-neonate pairs (i.e., including those with seronegative mothers) were assessed in a sensitivity analysis.

Generalized linear regression was conducted to identify factors associated with neonatal titre, neonatal seropositivity and the transfer ratio. A generalized linear mixed model with a B-spline was used to estimate the kinetics of the immune response over age. The antibody dynamics specified by three maternal antibody levels (negative group: <8 , low level: $8-<64$, and high level: $64-512$) were described to explore the effect of maternal antibody level on the children's antibody dynamics, with the range of maternal antibody being 6 to 512. The Kaplan–Meier method was conducted to estimate the median time to losing protective maternal antibodies, and linear regression analysis was used to estimate

the relationship between the duration of protection and maternal antibody titres. Survival analysis considering interval censoring and interval regression were conducted as a sensitivity analysis. All analyses were performed in R (version 4.0.2) and SAS (version 9.4).

Results

A total of 3013 children, including 538 neonates and 2475 children aged 1–9 years old, were enrolled in our study between Sep 2013 and Sep 2018. The follow-up rate of the neonate cohort was above 60% (Fig. 1A). Over 80% of children in the paediatric cohort were followed up annually. The group with semiannual follow-up had at least an 85% follow-up rate from Sep 2013 to Sep 2016 (Fig. 1B). The local circulation of CVA16 during 2013–2018 in Anhua County, Hunan Province, is depicted in the Appendix (Additional file 1: page 6–8, line 6–42).

The general characteristics of the enrolled neonates and children at baseline are provided in Table 1. The median age of the participants was 3 years old (IQR: 1–6), and 1532 (51%) children were boys. The follow-up time ranged from 1 to 41 months (median: 22 months; IQR: 11–29 months). The median age of mothers at baseline was 24 years old (IQR: 26–29 years old). Ninety percent of the mothers' sera were collected at delivery (Additional file 1: Table S4 (a)). Comparisons between participants and local nonparticipants are presented in the appendix (Additional file 1: pages 12–16, line 118–125).

As shown in Fig. 2A, the baseline anti-CVA16 antibody titre of neonates at birth was highly correlated with the maternal anti-CVA16 antibody titre ($\rho=0.80$, 95% CI 0.77–0.83). The GMT and seroprevalence were similar between mothers and neonates at birth (mother vs. neonate: 9.82 (95% CI 9.14–10.55) vs. 9.61 (95% CI 8.97–10.30), $p=0.102$; 44.80% (95% CI 40.53–49.10) vs. 41.82% (95% CI 37.61–46.12), $p=0.349$, respectively).

The probability of neonatal positivity increased as the maternal antibody titre rose until the maternal antibody titre was >28 for CVA16, with the neonatal probability being close to 100% (Fig. 2B). The overall maternal anti-CVA16 antibody titre transfer ratio was 0.88 (95% CI 0.80–0.96). It decreased with maternal antibody levels (Fig. 2C), with an average value larger than 1 when maternal antibody titres were ≤ 13 . Although the transfer efficacy decayed, neonates can successively obtain positive maternal antibodies via placental transmission as long as the maternal antibody titre reaches the positive level (Fig. 2C, Additional file 1: Fig. S4).

Multivariate analysis (Additional file 1: Table S8) indicated that maternal anti-CVA16 antibody titres were highly correlated with neonates' titres at birth and the status of antibody titres (positive or not). For every 2-fold increase in maternal titres, neonatal titres increased by

Table 1 Participant demographic characteristics at baseline

Characteristics	Neonates (n=538)	1–9-year-old children (n=2475)	Total (n=3013)
Age at baseline (years)			
Median (IQR)	-	4 (2-6)	3 (1-6)
0	538 (100)	-	538 (18)
1	-	355 (14)	355 (12)
2	-	368 (15)	368 (12)
3	-	332 (13)	332 (11)
4	-	333 (13)	333 (11)
5	-	330 (13)	330 (11)
6	-	283 (11)	283 (9)
7	-	205 (8)	205 (7)
8	-	162 (7)	162 (5)
9	-	107 (4)	107 (4)
Male	286 (53)	1246 (50)	1532 (51)
Has twin siblings ^a	12 (2%)	-	12
Gestational age at birth (weeks)			
Median (IQR)	40 (39-41)	-	-
<37	10 (2)	105 (4)	115 (4)
37–42	508 (94)	2344 (95)	2852 (95)
>42	20 (4)	25 (1)	45 (1)
Delivery mode			
Transvaginal delivery	331 (62)	1577 (64)	1908 (63)
Caesarean	207 (38)	897 (36)	1104 (37)
Birthweight (g)			
Median (IQR)	3300 (3000–3600)	3250 (3000–3500)	3250 (3000–3500)
<2500	11 (2)	88 (4)	99 (3)
2500–<4000	491 (91)	2139 (86)	2630 (87)
≥4000	36 (7)	247 (10)	283 (9)
Annual family income (RMB, Yuan)			
<20,000	82 (15)	611 (25)	693 (23)
20,000–<50,000	297 (55)	1382 (56)	1679 (56)
≥50,000	158 (29)	482 (19)	640 (21)



Data are n (%). ^a4 pairs of twins, and 4 neonates had a twin sibling, but their twin sibling did not participate in our study

69.1% (95% CI 63.4–75.2). The probability of acquiring positive antibody levels for neonates at birth increased by 17.7 times (95% CI 15.5–20.0) for every 2-fold increase in maternal titres. Having a twin decreased the probability by 15.7% (95% CI 1.4–28.0) compared to being a single neonate. The transfer ratio decreased by 14.8% (95% CI 11.8–17.6%) for every 2-fold increase in maternal antibody titre. The transfer ratio was reduced by 16.4% (95% CI 2.8–28.1%) when the mother had liver system disease.

Sensitive analyses with a positive threshold at 16 are depicted in the appendix (Additional file 1: Fig. S7).

Maternal transmitted antibody levels in neonates diminished rapidly. This drop was followed by an increase in titres due to natural infection, as was apparent in both the GMT and seroprevalence stratified by


age groups (Fig. 3A, B). Forty-two per cent of neonates had positive antibody levels, with a GMT of 10 (95% CI 9–10). However, the GMT decayed below the positive threshold after 1.5 months (range: 1.0–2.0 months) and further decreased to a minimum at 6.8 months. The seroprevalence experienced a sharp drop, falling to 20% after 1.3 months, with the lowest point at 5.8 months. Then, due to natural infections, the seroprevalence increased rapidly from 6 months to 5 years old (60%), then slowly increased to 80% at 13 years old. The GMT increased above the positive threshold after 1.4 years (range: 1.2–1.5 years) and then reached 32 by the age of 5 years. It slightly dropped thereafter. For the sensitivity analysis with a higher threshold of 16, the kinetics of seroprevalence and the GMT were consistent



with those in the baseline analysis. The GMT increased above 16 by 2.7 (range: 2.4–3.1) years of age (Additional file 1: Fig. S7). The kinetics of antibodies from high-level, low-level and negative maternal antibody groups were significantly different within 6 months. The higher the level of maternal antibody, the longer the duration of neonatal titre decay to under the detective threshold (high-level vs. low-level: 5.3 months vs. 1.2 months) and to the bottom (high-level vs. low-level: 11.0 months vs. 8.1 months). The antibody dynamics in the lower level of maternal antibody groups, especially the negative group, rose earlier due to natural infection, and the corresponding times for the three groups to rise to 8 were 1.5, 2.0 and 2.1 years, respectively. However, the

difference was not obvious when age more than 1.5 years old (Additional file 1: Fig. S9).

The maternally transmitted antibody levels decayed faster than natural infection-induced antibody titres ($p < 0.001$), with corresponding antibody half-lives of 2.0 months (95% CI 1.9–2.2 months) and 13.9 months (95% CI 13.3–14.5 months), respectively (Additional file 1: Fig. S10). Half of the neonates who were seropositive at birth were seronegative by 2.4 months of age (95% CI 2.3–2.5 months) (Fig. 4A). The average time to loss of seropositivity was 2.8 months (95% CI 2.5–3.1 months). For every 2-fold increase in maternal antibody, the duration of protection was prolonged by approximately 38 days (95% CI 28–48 days). When the maternal antibody titre was



≥ 64 and >512 , the duration of protection for neonates was extended to 6 and 12 months, respectively (Fig. 4B). The sensitivity analyses are shown in Additional file 1: Table S13.

Discussion

Our study revealed that the average anti-CVA16 antibody titres and seroprevalence were similar in maternal and cord blood, with a mean transfer ratio of maternal antibodies of 0.88 (95% CI 0.80–0.96). Although the transfer ratio decreased as the maternal antibody titre increased, the positive maternal antibodies could still be successfully transmitted to neonates. However, due to the low maternal seroprevalence, only 42% of neonates acquired positive concentrations of anti-CVA16 antibodies from mothers, which declined rapidly, with a half-life of 2.0 months (95% CI 1.9–2.2 months). Among them, the average duration of protection from maternal antibodies was 2.8 months (95% CI 2.5–3.1 months), and the duration was extended to 6 months on average for mothers with titres of ≥ 64 . After the disappearance of maternal antibodies, antibody titres increased because of rising natural infections.

Transplacental antibody transfer from mother to foetus provides protection from infection in the first weeks of life, such as measles, rubella, diphtheria, and tetanus, with various transfer efficacy. A strong association between maternal anti-CVA16 antibodies and neonates with a correlation coefficient of 0.8 was observed in our study, which was close to 0.56–0.75 in previous studies [10–12]. The transfer of anti-CVA16 maternal antibody is efficient, with an overall transfer ratio of 0.88, which is similar to Fu's observation of 0.9 [11] and is lower than that for EV-A71 [21], measles, and poliomyelitis III antibodies [11, 21]. The trend of the transfer ratio decreasing as the maternal antibody titre increases has also been observed for measles [26] and EV-A71 [21], which may be due to the saturation of a finite number of Fc receptors [26, 27]. Maternal antibody level is the major factor associated with the transfer ratio, following the mother's health. Liver system disease in the mother was associated with a lower transfer ratio. Other factors associated with mothers, such as maternal age, gestational diabetes, gestational hypertension, parity and gravidity, as well as factors associated with neonates, such as gestational age and birth weight, were not found to be related to transfer in our study.

The seroprevalence and GMT in our study were minimal at the ages of 5.8 months and 6.8 months, respectively, in line with previous longitudinal observations [12, 14, 15, 28]. The neonatal seroprevalences at birth and 2–4 months of age were lower than those in the studies by Jiangsu, potentially because a higher prevalence

(90%) was observed in mothers in those studies [15, 29]. The time when the GMT reached 8 and when it reached its minimum value were similar to those in Guangzhou studies (at 3 and 6 months old) [12, 28]. No previous longitudinal study investigated the kinetics of the anti-CVA16 antibody level, and only cross-sectional studies reported age-specific seroprevalence and GMT. Our age-specific seroprevalence results between 6 months and 13 years were comparable with those of previous reports in China [30], Singapore [17], Korea [16] and Thailand [18].

Using a neonatal cohort, we showed in another study that the dynamics of anti-EV-A71 antibodies had a similar pattern [21]. With higher maternal seroprevalence, neonatal EV-A71 seroprevalence was significantly higher than that of CVA16 for neonates aged 0–4 months old, so it took slightly longer for the GMT to decay below the positive threshold. After the decay of maternal antibody, the seroprevalence was similar between CVA16 and EV-A71 within 4 months to 3-year-old children (unpublished study), although the EV-A71 GMT was higher than the CVA16 GMT. A similar pattern was also observed in previous studies [31–33], showing that EV-A71 might elicit stronger neutralizing antibody responses than CVA16. The capsid proteins of the two viruses have approximately 20% sequence differences, as well as antigenic variation caused by the different surface structures of the two viruses, which may explain this phenomenon [34].

After the rapid decay of maternal antibodies, the seroprevalence increased with age as a result of natural infection; however, 40–95% of children from 6 months to 5 years old were still seronegative. In addition, to curb the transmission of CVA16 infection, a threshold of 59–96% for herd immunity is needed [2, 35–37]. Meanwhile, 0- to 5-year-old children, accounting for 99% and 96.5% of severe and mild cases in Hunan Province, respectively, constitute the majority of HFMD cases associated with CVA16 (Additional file 1: Fig. S11). Among them, the 1-year-old group had the largest number of mild and severe cases, which was in line with a previous report [23]. The above evidence showed that children aged 0–5 years old represent the main susceptible population, which should be targeted for vaccination.

Overall, both avoiding potential maternal antibody interference and preventing the largest incidence, especially severe cases, in the 1-year-old group, we recommend that the optimal vaccination time be between 1.5 months and 1 year old. Complete vaccination during this period could prolong the positive antibody titre until 5 years old, although this hypothesis is based on the half-life of anti-CVA16 antibody induced by infection. Further research is required on antibody dynamics induced by vaccines. Thus, we call for accelerating the development

of a human CVA16 vaccine and recommend clinical trials to further assess doses, safety, and effectiveness in susceptible young children aged 1.5 months to 5 years old.

Vaccinating pregnant mothers to provide maternal antibodies to infants who are too young to be vaccinated could be an alternative intervention to protect these infants, which has proven efficacious for influenza, pertussis and COVID-19 [8, 38, 39]. As we estimated, anti-CVA16 maternal antibodies can be effectively transmitted to newborns, and high maternal antibody levels can prolong the duration of protection for infants, with protection lasting up to 6 months when the maternal antibody level is ≥ 64 . Meanwhile, the results of antibody dynamics specified by maternal antibody levels also indicated that a higher maternal antibody group was associated with later antibody decay to undetectable levels and later natural infection. It is possible to increase the mother's antibody titre to extend protection to 1 year of age. Given that, vaccination of mothers during pregnancy can be a solution to protect infants from disease and death in early life, especially those under 1 year old. This conclusion is based on observations of antibodies mounted upon natural infection rather than vaccination, and we therefore still require further research to elucidate the efficacy or effectiveness of maternal CVA16 vaccination, antibody response and duration of immunity produced by maternal CVA16 vaccination.

A study in Vietnam found that adults can have HFMD and contribute to household transmission [40]. In our study, the seroprevalence of mothers (44.8%) was lower than that of mothers (87–93%) and adults (65–83%) in some coastal provinces of China [11, 15], Germany [19] and Vietnam [40]. Given the absence of sex-specific and age-specific differences in CVA16 prevalence [19, 41, 42], males and adults of other ages are also susceptible in theory. Thus, vaccination for adults in close contact with infants could help reduce household transmission and provide indirect protection for vulnerable infants.

This study is a population-based longitudinal study with a large sample size, prolonged follow-up design and high follow-up rates, in contrast with those of previous studies [10, 11, 14–17]. The quality control of the neutralization test was stringent. The conclusion using children with complete follow-up is consistent with the main results (Additional file 1: Fig. S8). However, our study still had some limitations. First, we were unable to correlate antibody levels with clinical symptoms because only a few cases were reported during the follow-up period. Moreover, there are currently no studies confirming the association between antibody titre and symptom reduction or avoidance of infection. Thus, the positive threshold in this study is a threshold for detecting antibodies in neutralization assays. It is expected that a higher titre

is associated with higher protection; thus, a sensitivity analysis with 16 as the positive threshold was conducted, which showed a conclusion consistent with that of the baseline analysis. Second, we were unable to compare the demographic information between the study participants and the local eligible nonenrollees in the 1–9 years old cohort because they refused to provide personal information. Finally, the sample size of mothers with higher antibodies (≥ 256) in our study was relatively small, causing the relative decreasing trend of directly calculated prevalence when the maternal antibody titre was 512, while the predicted probability was stable due to logit transformation in logistic regression.

Conclusions

In conclusion, positive titres of anti-CVA16 antibodies were efficiently transferred to neonates from mothers through the placenta. However, titres declined rapidly to below the positive threshold by 1.5 months of age. Increasing the maternal antibody titre can effectively prolong the duration of protection. The majority of children aged 0–5 years old are susceptible, especially 1-year-old children with a high risk of developing severe disease. It is important to assess the feasibility of vaccinating children between 1.5 months and 1 year old, pregnant women or adults in close contact with infants to protect 0–5-year-old children.

Abbreviations

CVA16: Coxsackievirus A16; EV-A71: Enterovirus A71; GMT: Geometric mean titre; Neonate cohort: A paired mother-neonate cohort; Paediatric cohort: A 1- to 9-year-old paediatric cohort.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12916-022-02604-w>.

Additional file 1: Fig. S1. Time series and predicting epidemic magnitude of CVA16 activity in Anhua county during 2013–2016 (A) and that in Hunan province during 2009–2018 (B). **Fig. S2.** Participants' included time in baseline and follow-up times (A) and the epidemic seasons of CVA16 activity during 2009–2018 in Anhua, Hunan province (B). **Fig. S3.** Distribution of maternal antibody titre transfer ratio against CVA16 for all pairs of neonate-mother (A) and positive mothers with antibody titre ≥ 8 (B). **Fig. S4.** The aggregated maternal titre specified transfer ratio. **Fig. S5.** Maternal antibody titre transfers efficacy against CVA16. **Fig. S6.** Transfer ratio trend with all pairs of mother-neonate. **Fig. S7.** Dynamic of seroprevalence, neutralization antibody titre by age for neonates using 16 as positive threshold. **Fig. S8.** Dynamic of neutralization antibody titre by age for neonates using full follow-up subset. **Fig. S9.** Dynamics of neutralization antibody titer by age for neonates stratified by maternal antibody levels. **Fig. S10.** Waning rate of maternal transmitted CVA16 antibody and natural infection induced antibody. **Fig. S11.** Age distribution and cumulative distribution of HFMD associated to CVA16 by clinical severity in Hunan province during 2009–2018 and Anhua county during 2013–2016. **Table S1.** Distribution of participants aged 1–9 years and sampling ratio. **Table S2.** Seasonality comparison between Anhua county and Hunan province. **Table S3.** Epidemic duration for CVA16 circulation

in Anhua. **Table S4.** (a) Baseline characteristics of mothers and neonates in neonate cohort between participants and non-participants, selected participants and non-selected participants for CVA16 assay. (b) Baseline characteristics of neonates in neonate cohort between full follow-up participants and incomplete follow-up participants. **Table S5.** Comparison of baseline characteristics of 1-9 years old children between participants and non-selected objects through sampling, full follow-up and incomplete follow-up. **Table S6.** Seroprevalence and GMT of mothers and neonates at birth. **Table S7.** Sample size (n, %) for CVA16 antibody transfer ratio analysis. **Table S8.** Multivariate analysis with factors associated with neonates' seropositivity, antibody titre at birth and transfer ratio. **Table S9.** Univariate analysis with factors associated with neonates' seropositivity at birth. **Table S10.** Univariate analysis with factors associated with the transfer ratio of maternal antibody. **Table S11.** Univariate analysis with factors associated with neonates' antibody titre at birth. **Table S12.** Sensitive analysis for multivariate analysis with factors associated with maternal antibody transfer ratio. **Table S13.** Sensitive analysis of median age of neonate with positive antibody. **Appendix.**

Acknowledgements

We acknowledge that this work was a continuous research project conducted by H.Y., who was a former employee of the Chinese Center for Disease Control and Prevention and moved to Fudan University since May 2017. We thank staff members of the Anhua County-, Yiyang Prefecture-, and Hunan Provincial-level departments of health for providing assistance with administration and data collection; staff members at the Anhua County-level (Liang Zhao, Zhiyong Chen, Min He, Zhaohui Jiang, and Ping Liu), Yiyang Prefecture-level (Lizhi Xie, and Qing Li), and Hunan Provincial-level Centers for Disease Control and Prevention (Zhihong Deng, Fuqiang Liu, Zhifei Zhan, Liang Cai, Ge Zeng, Hao Yang, Yu Chen, Qianlai Sun, and Hengjiao Zhang), study hospitals (Anhua People's Hospital, Anhua Maternal and Child Health Hospital, Anhua Chinese Medicine Hospital, Anhua Second People's Hospital, Jiangnan Township Health Center, Qingtang Township Health Center and Tianzhuang Township Health Center) (Le Yang, Hong Liang, and Yi Zhang), China CDC (Fengfeng Liu, Li Luo, Yu Li, Junling Sun, and Chen Yuan), Hubei CDC (Qi Chen) for providing assistance with field investigation, administration, and data collection. We thank Yong Zhang and Shuangli Zhu from the China CDC for providing technical support for this work.

Disclaimer

The views expressed are those of the authors and do not necessarily represent the institutions with which the authors are affiliated.

Authors' contributions

H.Y. was responsible for its conception and design. K.L., Q.L., S.Z., B.D., L.G., L.R., X.C., J.Z., and J.Y. collected the data. Y.Z., W.Z., H.G., H.S., K.W., Q.Q., L.R., L.W., M.X., N.L., W.L., N.Z., X.C., and Z.C. performed the lab test. J.Z. and Y.Z. performed the analysis and prepared the figures and tables. J.Z. and Y.Z. wrote the first draft. H.Y., S.C., and J.Y. made critical revision of the manuscript for important intellectual content. All authors read and approved the final manuscript. H.Y. had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding

This work was supported by the Key Program of the National Natural Science Foundation of China (82130093), Li Ka Shing Oxford Global Health Programme (No. LG33), AXA Research Fund, the Investissement d'Avenir program, Laboratoire d'Excellence Integrative Biology of Emerging Infectious Diseases program (Grant ANR-10-LABX-62-IBEID), Models of Infectious Disease Agent Study of the National Institute of General Medical Sciences, INCEPTION project (PIA/ANR-16-CONV-0005), Chinese Preventive Medicine Association (No. 20101801), Hunan Provincial Natural Science Foundation (2019JJ80115), and the Scientific Research Project of Hunan Provincial Health Commission (B2019039). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Availability of data and materials

Data described in the manuscript are available on request due to privacy/ethical restrictions. The analytical code will be made available only to the

investigators who have participated in/contributed to the study. The study executive will consider specific requests for data analyses by non-contributing individuals upon reasonable request to the corresponding author.

Declarations

Ethics approval and consent to participate

The study was approved by the Institutional Review Board of the WHO Western Pacific Regional Office (2013.10.CHN.2.ESR), the Chinese Centre for Disease Control and Prevention (201224), and Fudan University (2019-05-0756). We obtained written informed consent from each child's guardian and all enrolled mothers themselves.

Consent for publication

Not applicable.

Competing interests

H.Y. has received research funding from Sanofi Pasteur, GlaxoSmithKline, Yichang HEC Changjiang and Shanghai Roche Pharmaceutical Company. None of the research funding is related to this work. All other authors report no competing interests.

Author details

¹School of Public Health, Fudan University, Key Laboratory of Public Health Safety, Ministry of Education, Shanghai, China. ²Hunan Provincial Center for Disease Control and Prevention, Changsha, China. ³Anhua County Center for Disease Control and Prevention, Yiyang, China. ⁴Mathematical Modelling of Infectious Diseases Unit, Institut Pasteur, Université de Paris, UMR2000, CNRS, 75015 Paris, France.

Received: 26 May 2022 Accepted: 11 October 2022

Published online: 09 November 2022

References

- Tapparel C, Siegrist F, Petty TJ, Kaiser L. Picornavirus and enterovirus diversity with associated human diseases. *Infect Genet Evol.* 2013;14:282–93.
- Zhang Z, Liu Y, Liu F, Ren M, Nie T, Cui J, et al. Basic Reproduction Number of Enterovirus 71 and Coxsackievirus A16 and A6: Evidence From Outbreaks of Hand, Foot, and Mouth Disease in China Between 2011 and 2018. *Clin Infect Dis.* 2021;73(9):e2552–9.
- Mao Q, Wang Y, Yao X, Bian L, Wu X, Xu M, et al. Coxsackievirus A16: epidemiology, diagnosis, and vaccine. *Hum Vaccin Immunother.* 2014;10(2):360–7.
- Xu W, Liu CF, Yan L, Li JJ, Wang LJ, Qi Y, et al. Distribution of enteroviruses in hospitalized children with hand, foot and mouth disease and relationship between pathogens and nervous system complications. *Virology.* 2012;9:8.
- Li R, Liu L, Mo Z, Wang X, Xia J, Liang Z, et al. An inactivated enterovirus 71 vaccine in healthy children. *N Engl J Med.* 2014;370(9):829–37.
- Zhu F, Xu W, Xia J, Liang Z, Liu Y, Zhang X, et al. Efficacy, Safety, and Immunogenicity of an Enterovirus 71 Vaccine in China. *N Engl J Med.* 2014;370(9):818–28.
- Zhu F-C, Meng F-Y, Li J-X, Li X-L, Mao Q-Y, Tao H, et al. Efficacy, safety, and immunology of an inactivated alum-adjunct enterovirus 71 vaccine in children in China: a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet.* 2013;381(9882):2024–32.
- World Health Organization. Vaccines against influenza WHO position paper — November 2012 = Note de synthèse de l'OMS concernant les vaccins antigrippaux — novembre 2012. *Weekly Epidemiol Rec = Relevé épidémiologique hebdomadaire.* 2012;87(47):461–76 <https://apps.who.int/iris/handle/10665/241993>.
- Cáceres VM, Strebel PM, Sutter RW. Factors Determining Prevalence of Maternal Antibody to Measles Virus throughout Infancy: A Review. *Clin Infect Dis.* 2000;31(1):110–9.
- Hong J, Li L, Liu YM, Ge HM, Wang XS, Hu JL, et al. Seroepidemiology of human enterovirus71 and coxsackievirusA16 in Jiangsu province, China. *Virology.* 2012;9:248.
- Fu C, Lu L, Wu H, Shaman J, Cao Y, Fang F, et al. Placental antibody transfer efficiency and maternal levels: specific for measles, coxsackievirus

- A16, enterovirus 71, poliomyelitis HIII and HIV-1 antibodies. *Sci Rep*. 2016;6:38874.
12. Cui M, Ya-Jing LI, Yang QY, Long LU, Qing HE, Dong ZQ, et al. Study on transition levels of maternal neutralizing antibodies against enterovirus 71 and coxsackievirus A16 in infants in Guangzhou. *Chin J Dis Control Prev*. 2016;20(3):236–9.
 13. Wei XIA, Yan LI, Yin DP, et al. Investigation on Neutralizing Antibody Level of Enterovirus Type 71 and Coxsackievirus Group Type 16 among Mothers and Newborns in Six Counties. *Chin J Vacc Immun*. 2013;19(02):164–167+174.
 14. Mao Q-y, Liao XY, Yu X, Li N, Zhu F-c, Zeng Y, et al. Dynamic change of mother-source neutralizing antibodies against enterovirus 71 and coxsackievirus A16 in infants. *Chin Med J (Engl)*. 2010;123(13):1679–84.
 15. Zhu FC, Liang ZL, Meng FY, Zeng Y, Mao QY, Chu K, et al. Retrospective study of the incidence of HFMD and seroepidemiology of antibodies against EV71 and CoxA16 in prenatal women and their infants. *PLoS One*. 2012;7(5):e37206.
 16. Kim S, Bae KS, Kim JH, Kang JH, Choi UY. Seroprevalence of Neutralizing Antibodies Against Candidate Serotypes of Enterovirus Vaccines Among Korean Children. *Viral Immunol*. 2021;34(2):62–7.
 17. Wei AL, Joanne T, Chee PM, Pu HJ, Jeffery C, Lyn J, et al. Seroepidemiology of Coxsackievirus A6, Coxsackievirus A16, and Enterovirus 71 Infections among Children and Adolescents in Singapore, 2008–2010. *PLoS One*. 2015;10(5):e0127999.
 18. Hatairat L, Jarunee P, Anek M, Chompunuch K, Piron, Journal NJV. Seroprevalence of antibodies to enterovirus 71 and coxsackievirus A16 among people of various age groups in a northeast province of Thailand. *Viol J*. 2018;15(1):158.
 19. Rabenau HF, Richter M, Doerr HW. Hand, foot and mouth disease: seroprevalence of Coxsackie A16 and Enterovirus 71 in Germany. *Med Microbiol Immunol*. 2010;199(1):45–51.
 20. Ni H, Yi B, Yin J, Fang T, He T, Du Y, et al. Epidemiological and etiological characteristics of hand, foot, and mouth disease in Ningbo, China, 2008–2011. *J Clin Virol*. 2012;54(4):342–8.
 21. Wei X, Yang J, Gao L, Wang L, Liao Q, Qiu Q, et al. The transfer and decay of maternal antibodies against enterovirus A71, and dynamics of antibodies due to later natural infections in Chinese infants: a longitudinal, paired mother-neonate cohort study. *Lancet Infect Dis*. 2021;21(3):418–26.
 22. Gao L, Zou G, Liao Q, Zhou Y, Liu F, Dai B, et al. Spectrum of Enterovirus Serotypes Causing Uncomplicated Hand, Foot, and Mouth Disease and Enteroviral Diagnostic Yield of Different Clinical Samples. *Clin Infect Dis*. 2018;67(11):1729–35.
 23. Xing W, Liao Q, Viboud C, Zhang J, Sun J, Wu JT, et al. Hand, foot, and mouth disease in China, 2008–12: an epidemiological study. *Lancet Infect Dis*. 2014;14(4):308–18.
 24. Qiu Q, Zhou J, Cheng Y, Zhou Y, Liang L, Cui P, et al. Kinetics of the neutralising antibody response in patients with hand, foot, and mouth disease caused by EV-A71: a longitudinal cohort study in Zhengzhou during 2017–2019. *EBioMedicine*. 2021;68:103398.
 25. Dang S, Gao N, Li Y, Li M, Wang X, Jia X, et al. Dominant CD4-dependent RNA-dependent RNA polymerase-specific T-cell responses in children acutely infected with human enterovirus 71 and healthy adult controls. *Immunology*. 2014;142(1):89–100.
 26. Hartter HK, Oyedele OI, Dietz K, Kreis S, Hoffman JP, Muller CP. Placental transfer and decay of maternally acquired antimeasles antibodies in Nigerian children. *Pediatr Infect Dis J*. 2000;19(7):635–41.
 27. Roopenian DC, Akilesh S. FcRn: the neonatal Fc receptor comes of age. *Nat Rev Immunol*. 2007;7(9):715–25.
 28. Fu C, Shen J, Lu L, Li Y, Cao Y, Wang M, et al. Pre-vaccination evolution of antibodies among infants 0, 3 and 6 months of age: A longitudinal analysis of measles, enterovirus 71 and coxsackievirus 16. *Vaccine*. 2017;35(31):3817–22.
 29. Mao QY, Liao XY, Xiang YU, Nan LI, Zhu FC, Zeng Y, et al. Dynamic change of mother-source neutralizing antibodies against enterovirus 71 and coxsackievirus A16 in infants. *Chin Med J (Engl)*. 2010;123(13):1679–84.
 30. D S. Serological survey of hand, foot and mouth disease among healthy people in Hebei District, Tianjin City, 2016 ~ 2019 Year. *Med Inform*. 2020;33(8):147–9.
 31. Gao F, Mao QY, Chen P, Bian LL, Yao X, Li JX, et al. Seroepidemiology of Coxsackievirus A6, Coxsackievirus A16, and Enterovirus 71 infections in infants and children: A prospective cohort study in Jiangsu, China. *J Infect*. 2016;73(5):509–12.
 32. Yu N, Guo M, He SJ, Pan YX, Chen XX, Ding XX, et al. Evaluation of human enterovirus 71 and coxsackievirus A16 specific immunoglobulin M antibodies for diagnosis of hand-foot-and-mouth disease. *Viol J*. 2012;9:12.
 33. Nguyet LA, Thanh TT, Nhan LNT, Hong NTT, Nhu LNT, Van HMT, et al. Neutralizing Antibodies against Enteroviruses in Patients with Hand, Foot and Mouth Disease. *Emerg Infect Dis*. 2020;26(2):298–306.
 34. Ren J, Wang X, Zhu L, Hu Z, Gao Q, Yang P, et al. Structures of Coxsackievirus A16 Capsids with Native Antigenicity: Implications for Particle Expansion, Receptor Binding, and Immunogenicity. *J Virol*. 2015;89(20):10500–11.
 35. Lim C, Jiang L, Ma S, James L, Ang LWJE. Infection: Basic reproduction number of coxsackievirus type A6 and A16 and enterovirus 71: estimates from outbreaks of hand, foot and mouth disease in Singapore, a tropical city-state. *Epidemiol Infect*. 2016;144(05):1028–34.
 36. Ma E, Fung C, Yip S, Wong C, Chuang SK. Tsang TJTPIDJ: Estimation of the basic reproduction number of enterovirus 71 and coxsackievirus A16 in hand, foot, and mouth disease outbreaks. *Pediatr Infect Dis J*. 2011;30(8):675–9.
 37. Takahashi S, Liao Q, Van Boeckel TP, Xing W, Sun J, Hsiao VY, et al. Hand, Foot, and Mouth Disease in China: Modeling Epidemic Dynamics of Enterovirus Serotypes and Implications for Vaccination. *PLoS Med*. 2016;13(2):e1001958.
 38. Mazzilli S, Tavoschi L, Lopcalo PL. Tdap vaccination during pregnancy to protect newborns from pertussis infection. *Annali Igiene : Med Prev Comunita*. 2018;30(4):346–63.
 39. Trostle ME, Agüero-Rosenfeld ME, Roman AS, Lighter JL. High antibody levels in cord blood from pregnant women vaccinated against COVID-19. *Am J Obstet Gynecol MFM*. 2021;3(6):100481.
 40. Hoang CQ, Nguyen HD, Ho NX, Vu THT, Pham TTM, Nguyen KT, et al. Incidence of Infection of Enterovirus 71 and Coxsackieviruses A6 and A16 among Household Contacts of Index Cases in Dong Thap Province, Southern Vietnam. *BioMed Res Int*. 2020;2020:9850351.
 41. Wang JX, Zhu SL, Jing W, Lin Y, Pei YW, Sun DP, et al. Ding SJPO: Sero-prevalence of Enterovirus A71 and Coxsackievirus A16 in Healthy People in Shandong Province, China. *PLoS One*. 2016;11(9):e0162373.
 42. Wang X, Xing M, Zhang C, Yang Y, Chi Y, Tang X, et al. Neutralizing antibody responses to enterovirus and adenovirus in healthy adults in China. *Emerg Microbes Infect*. 2014;3(5):e30.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

